

# Review of Australasian spider flies (Diptera, Acroceridae) with a revision of *Panops* Lamarck

Shaun L. Winterton<sup>1,†</sup>

<sup>1</sup> California State Collection of Arthropods, California Department of Food & Agriculture, Sacramento, California, USA

† [urn:lsid:zoobank.org:author:37F5AC48-EC3A-47ED-902B-2BD1467CCA72](http://urn:lsid:zoobank.org:author:37F5AC48-EC3A-47ED-902B-2BD1467CCA72)

Corresponding author: Shaun L. Winterton ([wintertonshaun@gmail.com](mailto:wintertonshaun@gmail.com))

---

Academic editor: T. Dikow | Received 9 August 2011 | Accepted 17 February 2012 | Published 1 March 2012

---

[urn:lsid:zoobank.org:pub:F365ECC1-1271-4740-B3FD-867D1C6A9532](http://urn:lsid:zoobank.org:pub:F365ECC1-1271-4740-B3FD-867D1C6A9532)

---

**Citation:** Winterton SL (2012) Review of Australasian spider flies (Diptera, Acroceridae) with a revision of *Panops* Lamarck. ZooKeys 172: 7–75. doi: 10.3897/zookeys.172.1889

---

## Abstract

The Australasian spider flies (Diptera: Acroceridae) are reviewed, with all eight currently recognized genera diagnosed and figured. The panopine genus *Panops* Lamarck, 1804 from Australia and Indonesia is revised with four new species described, increasing the total number of species in the genus to nine: *P. aurum* **sp. n.**, *P. danieli* **sp. n.**, *P. jade* **sp. n.** and *P. schlingeri* **sp. n.** Five species of *Panops* are redescribed: *Panops austra*e Neboiss, 1971, *P. baudini* Lamarck, 1804, *P. boharti* (Schlinger, 1959), **comb. n.**, *P. conspicuus* (Brunetti, 1926) and *P. grossi* (Neboiss, 1971), **comb. n.** The monotypic genera *Neopanops* Schlinger, 1959 and *Panocalda* Neboiss, 1971 are synonymized with *Panops*. Keys to genera of Australasian Acroceridae and species of *Panops*, *Helle* Osten Sacken, 1896 and Australasian *Pterodontia* Gray, 1832 are included.

## Keywords

cybertaxonomy, spider parasitoid

## Introduction

Spider flies (also known as small-headed flies) (Diptera: Acroceridae) are a distinctive group of lower brachyceran flies characterized by unusual adult body shape and highly specialized larval biology as parasitoids of spiders. Adults are recognized as important pollinators of angiosperms (Fig. 1), frequently as strong fliers with greatly elongate mouthparts for feeding in long corolla flowers, although some species have reduced or



**Figure 1.** *Panops baudini* Lamarck feeding on *Daviesia croniniana* F.Muell. (Fabaceae), photographed during September in Boorabbin National Park, Western Australia. Photograph by Dan Schoknecht (Western Australian Museum).

even vestigial mouthparts (Schlinger 1981, 1987). Acroceridae comprise approximately 520 species in 53 genera (Pape and Thompson 2010; Gillung and Winterton 2011) occupying most biogeographic regions. The family is presently classified into three extant subfamilies: Acrocerinae, Panopinae and Philopotinae (Schlinger 1981), although recent phylogenetic analyses using DNA sequence data suggest that Acrocerinae are polyphyletic and membership of that subfamily should be re-examined (Winterton et al. 2007). Larvae of Acroceridae are internal parasitoids of juvenile spiders, living internally within the opithsoma of the spider where they attach to the book-lungs of the host via their posterior spiracles. Upon completing development the mature, third instar larva emerges from the dead host before pupating (Schlinger 1987). There are exceptions though, with a Chilean species recorded as ectoparasitic on spiders (i.e. *Sphaerops appendiculata* Philippi, 1865 (Acrocerinae)) (Schlinger 1987), whilst Kerr and Winterton (2008) recently questioned the exclusivity of parasitism of spiders, describing a putative acrocerid planidium on an anystinid mite in Baltic Amber.

The Australasian acrocerid fauna comprises all three subfamilies, although represented by relatively few genera. Two acrocerine genera (*Ogcodes* Latreille, 1797 and *Pterodontia* Gray, 1832) are found throughout the region, and are considered cosmopolitan throughout all major biogeographic regions. Philopotinae are represented by an endemic genus in New Zealand (*Helle* Osten Sacken, 1896) (Paramonov 1955) and a recently described genus endemic to New Caledonia (*Schlengeriella* Gillung & Winterton, 2011).

Panopinae are well represented in the Australasian region. Six genera are described previously from New Zealand (*Apsona* Westwood, 1876), Indonesia (*Neopanops* Schlinger, 1959) and Australia (*Panocalda* Neboiss, 1971, *Panops* Lamarck, 1804, *Mesophysa* Macquart, 1838 and *Leucopsina* Westwood, 1876) (Paramonov 1955, 1957; Schlinger 1959; Neboiss 1971). *Pterodontia* has been considered by some authors to be placed in Panopinae based on the presence of tibial spines (Schlinger 1981, 1987, 2009), but most authors place it in Acrocerinae based on wing venation and antennal characteristics (e.g. Neboiss 1971) and molecular data (Winterton et al. 2007). *Panops* is the most species rich genus in the region and is revised herein. Three species were described previously and treated in the most recent revision of the genus by Neboiss (1971): *Panops austra*e Neboiss, 1971, *P. baudini* Lamarck, 1804, and *P. conspicuus* (Brunetti, 1926). An additional four species are described herein (*P. aurum* sp. n., *P. danieli* sp. n., *P. jade* sp. n. and *P. schlinger*i sp. n.) whilst another two species are moved from other genera (*P. boharti* (Schlinger, 1959), comb. n. and *P. grossi* (Neboiss, 1971), comb. n.). Discovery of these new species of *Panops* has expanded the concept of the genus, with various species exhibiting combinations of characteristics previously used to differentiate *Panops* from *Panocalda* and *Neopanops*—specifically length of the mouthparts and presence and distribution of eye pilosity. Consequently, *Neopanops* and *Panocalda* are newly synonymized with *Panops*. All Australasian acrocerid genera are diagnosed and figured. Four genera of Panopinae are now recognized from the Australasian region, *Apsona* (1 sp.), *Mesophysa* (4 spp.), *Panops* (9 spp.) and *Leucopsina* (2 spp.). Keys to genera of Australasian Acroceridae and species of *Panops*, *Helle* and Australasian *Pterodontia* are included.

## Material and methods

Terminology follows McAlpine (1981) and Schlinger (1981). In most acrocerids, two crossveins span the area between the radial and medial sectors. The proximal crossvein is r-m, while the distal crossvein bisecting cell  $r_{4+5}$  (between wing veins  $M_1$  and  $R_{4+5}$ , or rarely  $R_5$ ) is referred to here as 2r-m following Hardy (1946) and Gillung and Winterton (2011). Annotations of collection label data are included where appropriate in brackets. The following collection codens are cited in the text: Australian Museum (AMS), Australian National Insect Collection (ANIC), California Academy of Sciences (CAS), Canadian National Collection of Insects (CNC); Greg Daniels private collection [to be ultimately deposited in the Australian Museum] (GDCB/AMS), Museum

National d'Histoire Naturelle (MNHN), National Museum of Victoria (NMV), Oxford University Museum of Natural History (OUMNH), Queensland Museum (QM), Swedish Museum of Natural History (NHRS), South Australian Museum (SAM), The Natural History Museum (BMNH), Western Australian Museum (WAM). Descriptions were constructed using Lucid Builder 3.5, using a matrix database of character states, which were then exported using the natural language function into XML and a text document. Specimen images were taken at different focal points using a digital camera and subsequently combined into a serial montage image using Helicon Focus software. High-resolution digital images were deposited into Morphbank with embedded URL links within the document between descriptions and Morphbank images. All new nomenclatural acts and literature are registered in Zoobank (Pyle and Michel 2008).

## Taxonomy

### Key to genera of Australasian Acroceridae

- 1 Postpronotal lobes greatly enlarged, contiguous along midline to form collar for head ..... **Philopotinae, 2**
- Postpronotal lobes not greatly enlarged, widely separate along midline ..... **3**
- 2 Wing with cells d, br, bm, and cu-p present, venation relatively complete (Fig. 3A) ..... ***Helle Osten Sacken, 1896 (New Zealand)***
- Wing with only cell br present, venation reduced (Fig. 3B) .....  
..... ***Schlingeriella* Gillung & Winterton, 2011 (New Caledonia)**
- 3 Antenna usually styliform or rod-like with multiple terminal setae; wing venation reduced: at most three radial veins present, cells d and basal  $r_{4+5}$  merged or absent (Figs 3C, D); tibiae without spines (except *Pterodontia*) ....  
..... **Acrocerinae, 4**
- Antenna with elongate flagellum, cylindrical or flattened, without terminal styliform seta; wing venation complete: four radial veins present, cells d and basal  $r_{4+5}$  separate (Figs 2A–D); at least some tibiae with an apical spine on outer margin (absent in *Apsona*) ..... **Panopinae, 5**
- 4 Eye apilose, without setae; venation reduced with many veins absent or poorly defined, almost all cells weakly formed or absent; tibial spines absent (Figs 3C, 63–64) ..... ***Ogcodes* Latreille, 1797 (Cosmopolitan)**
- Eye pilose; all wing veins well defined to wing margin, discal cell and basal portion of  $r_{4+5}$  merged into single closed cell; tibial spines present (Figs 3D, 65–66) ..... ***Pterodontia* Gray, 1832 (Cosmopolitan)**
- 5 Eye strongly pilose; antennal flagellum slender and tapered to apex; tibial spines absent (Figs 2A, 4–6) ..... ***Apsona* Westwood, 1876 (New Zealand)**
- Eye apilose or weakly pilose; antennal flagellum thickened to apex; tibial spines present (Australia) ..... **6**

- 6 Eye apilose, or sparsely or partially pilose; wing hyaline; crossvein 2r-m joining to stem  $R_{4+5}$  (Fig. 2D) ..... ***Panops* Lamarck, 1804**
- Eye always apilose; wing at least partially infusate, particularly along anterior margin; crossvein 2r-m joining to vein  $R_5$  (Fig. 2B–C) ..... **7**
- 8 Dorsal profile of abdomen with swollen, rounded tergites; without transverse yellow band on tergite 3; not wasp-like in appearance (Figs 12–16) ..... ***Mesophysa* Macquart, 1838**
- Dorsal profile of abdomen with truncated tergites raised along posterior margins; transverse yellow band on tergite 3; distinctly wasp-like in appearance (Figs 7–11) ..... ***Leucopsina* Westwood, 1876**

### Subfamily Panopinae Schiner, 1868

<http://species-id.net/wiki/Panopinae>

**Type genus.** *Panops* Lamarck, 1804: 263.

**Diagnosis.** Usually large and densely pilose, body shape never arched; antennal flagellum elongate cylindrical to paddle-shaped, sometimes tapered but never stylate, usually lacking terminal setae; postpronotal lobes never meeting medially; wing venation complete to wing margin (rarely reduced), cells  $m_3$ , d, bm and basal  $r_{4+5}$  typically present, closed distally; tibial spines present (rarely absent); larvae exclusively parasitoids of mygalomorph spiders.

### Australasian genera

*Apsona* Westwood, 1876; *Leucopsina* Westwood, 1876; *Mesophysa* Macquart, 1838; *Panops* Lamarck, 1804.


### ***Apsona* Westwood, 1876**

<http://species-id.net/wiki/Apsona>

Figs 2A, 4–6

*Apsona* Westwood, 1876: 510 – Bigot 1890: 317; Hutton 1901: 27; Paramonov 1955: 19; Schlinger 1966: 112; Schlinger and Jefferies 1989: 375. Type species: *Apsona muscaria* Westwood, 1876 by monotypy.

**Diagnosis.** Body length: 7–9 mm. Colouration metallic green; head width slightly smaller than thorax width, hemispherical; postocular ridge and occiput rounded; three ocelli; posterior margin of eye rounded; eye pilose (dense); position of antenna on frons nearer to ocellar tubercle; eyes contiguous above and below antennal base; palpus present; proboscis longer than head length; flagellum shape elongate, tapered apically, apex lacking terminal setae; scapes separate; subscutellum not enlarged, barely visible;



**Figure 2.** Acroceridae wings. Panopinae: **A** *Apsona muscaria* Westwood **B** *Leucopsina odyneroides* Westwood **C** *Mesophysa tenaria* Neboiss **D** *Panops jade* sp. n. Scale line = 0.2 mm.

tibial spines absent; pulvilli present; wing hyaline, markings absent; costa circumambient, costal margin straight apically in both sexes; humeral crossvein present; radial veins curved towards wing anterior margin;  $R_1$  not inflated distally; pterostigma and cell  $r_1$  membranous, not ribbed;  $R_{2+3}$  present;  $R_{4+5}$  present as forked petiolate veins; cell  $r_{4+5}$  bisected by 2r-m, basal cell narrow elongate, closed; 2r-m very short, joining  $M_1$  to stem  $R_{4+5}$ ;  $R_4$  without spur vein; medial vein complement with  $M_1$ ,  $M_2$  and  $M_3$  present ( $M_3$  fused with  $CuA_1$ ); discal cell closed completely;  $M_1$  and  $M_2$  usually not reaching wing margin; cell  $m_3$  present;  $CuA_1$  joining  $M_3$ , petiolate to wing margin;  $CuA_2$  fused to  $A_1$  before wing margin, petiolate; wing microtrichia absent; anal lobe well developed; alula absent; abdominal tergites smooth, rounded; abdomen shape greatly rounded, inflated, conical posteriorly.

**Included species.** *Apsona muscaria* Westwood, 1876.

**Comments.** *Apsona* is a monotypic genus endemic to New Zealand and can be readily differentiated from all other Panopinae based on the lack of tibial spines. *Apsona* shows little relationship to the rest of the Australasian Panopinae and shows remarkable similarity to the New World genus *Eulonchus* Gerstaecker, 1856, sharing numerous characteristics such as metallic green colouration, antennal shape, dense eye pilosity, elongate mouthparts, eyes contiguous below antennal base and absence of an alula (Paramonov 1955).


### ***Leucopsina* Westwood, 1876**

<http://species-id.net/wiki/Leucopsina>

Figs 2B, 7–11

*Leucopsina* Westwood, 1876: 510 – Bigot 1890: 314, 315; Hardy 1921: 78; Paramonov 1957: 524; Neboiss 1971: 219; Schlinger and Jefferies 1989: 375. Type species: *Leucopsina odyneroides* Westwood, 1876 by monotypy.

**Diagnosis.** Body length: 9.0 mm [male], 12.0 mm [female]. Colouration black and yellow [wasp mimic]; head slightly smaller than thorax width, shape hemispherical; postocular ridge and occiput rounded; three ocelli, anterior ocellus reduced in size (female) or absent (male); posterior margin of eye emarginate; eye apilose; position of antennae on head adjacent to ocellar tubercle; male frons width above antennal base not contiguous, eyes contiguous below antennal base; palpus present; proboscis greater than head length; flagellum shape elongate, cylindrical; apex lacking terminal setae; scapes separate; subscutellum enlarged; tibial spines present; pulvilli present; wing markings present (infusate anteriorly); costa circumambient (weaker along anal margin); costal margin straight; humeral crossvein present; radial veins straight;  $R_1$  not inflated distally; pterostigma and cell  $r_1$  membranous, not ribbed;  $R_{2+3}$  present;  $R_{4+5}$  originating separately from cell  $r_{4+5}$  (or at same point); cell  $r_{4+5}$  bisected by 2r-m, basal cell narrow elongate, closed; 2r-m joining  $M_1$  to  $R_5$ ;  $R_4$  with spur vein; medial



**Figure 3.** Acroceridae wings. Philopotinae: **A** *Helle rufescens* Brunetti **B** *Schlingeriella irwini* Gillung & Winterton. Acrocerinae **C** *Ogcodes basalis* Walker **D** *Pterodontia davisii* Paramonov (female). Scale line = 0.2 mm.

vein complement:  $M_1$ ,  $M_2$  and  $M_3$  present ( $M_3$  fused with  $CuA_1$ ); discal cell closed completely; medial veins reaching wing margin; cell  $m_3$  present;  $CuA_1$  joining  $M_3$ , petiolate to margin;  $CuA_2$  fused to  $A_1$  before wing margin, petiolate; wing microtrichia

absent; anal lobe well-developed; alula weakly developed; abdominal tergites smooth, rounded, tergites raised along posterior margins; abdomen constricted anteriorly.

**Included species.** *Leucopsina burnsi* (Paramonov, 1957); *L. odyneroides* Westwood, 1876.

**Comments.** *Leucopsina* is an endemic Australian genus of contrastingly coloured yellow and black flies, with distinct sexual dimorphism between males and females; male having more pronounced constriction of the abdomen anteriorly. The body colouration, darkening of the costal wing margin and abdominal waist allows members of this genus to be convincing wasp mimics (Neboiss 1971). *Leucopsina* can be differentiated from all other acrocerid genera by the wasp mimicking habitus, elongate cylindrical flagellum, apilose eyes and elongate mouthparts. Neboiss (1971) provides a key to species of this genus. *Leucopsina burnsi* was originally described as a variety of *Panops flavipes* (= *Mesophysa flavipes* Latreille, 1811) but subsequently transferred to *Leucopsina* and thoroughly differentiated from *L. odyneroides* by Neboiss (1971).

### ***Mesophysa* Macquart**

<http://species-id.net/wiki/Mesophysa>

Figs 2C, 12–16

*Mesophysa* Macquart, 1838: 166 – Blanchard 1840: 584; Westwood 1876: 517; Brunetti 1926: 580; Edwards 1930: 193; Neboiss 1971: 214; Schlinger and Jefferies 1989: 376. Type species: *Mesophysa scapularis* Macquart, 1838 by subsequent designation of Brunetti 1926: 580 [= *Panops flavipes* Latreille, 1811].

**Diagnosis.** Body length: 8.0–10.0 mm [male], 9.0–11 mm [female]. Colouration non-metallic, usually matte greenish hue; head size slightly smaller than thorax width; shape hemispherical; postocular ridge and occiput rounded; three ocelli; posterior margin of eye emarginate; eye apilose; antennae positioned on head adjacent to ocellar tubercle; eyes not contiguous above antennal base, contiguous below antennal base; palpus present; proboscis greater than head length; flagellum shape elongate, cylindrical (flattened), truncated apically [more pronounced in male]; scapes separate; flagellum apex lacking terminal setae; subscutellum not enlarged, barely visible; tibial spines present; pulvilli present; wing infusate, markings present; costa circumambient (weaker along anal margin); costal margin straight apically; humeral crossvein present; radial veins straight;  $R_1$  not inflated distally; pterostigma and cell  $r_1$  membranous, not ribbed;  $R_{2+3}$  present;  $R_{4+5}$  originating separately from cell  $r_{4+5}$ ; cell  $r_{4+5}$  bisected by 2r-m, basal cell narrow elongate, closed; 2r-m, joining  $M_1$  to  $R_5$ ;  $R_4$  with spur vein; medial vein complement with  $M_1$ ,  $M_2$  and  $M_3$  present ( $M_3$  fused with  $CuA_1$ ); discal cell closed completely; medial veins reaching wing margin; cell  $m_3$  present;  $CuA_1$  joining  $M_3$ , petiolate to margin;  $CuA_2$  fused to  $A_1$  before wing margin, petiolate to margin; wing microtrichia absent; anal lobe well developed; alula well developed; abdominal tergites

smooth, rounded; abdomen shape rounded, cylindrical, similar width to thorax or constricted anteriorly (male), tergites raised along posterior margins.

**Included species.** *Mesophysa flavipes* (Latreille, 1811); *M. ilzei* Neboiss, 1971; *M. tenaria* Neboiss, 1971; *M. ultima* Neboiss, 1971.

**Comments.** *Mesophysa* is an endemic eastern Australian genus closely related to *Leucopsina*. They share a similar habitus with narrowing of the abdomen anteriorly (more pronounced in *Leucopsina*), apilose eyes, infusate wings and flagellum shape, as well as the crossvein 2r-m joining to  $R_5$  rather than to the stem  $R_{4+5}$ . This genus can be differentiated from *Leucopsina* by the lack of black and yellow markings. *Mesophysa* has been considered a synonym of *Panops* by some authors (Erichson 1840; Kertész 1909; Edwards 1930; Hardy 1946; Paramonov 1957) and treated as separate genera by others (e.g. Brunetti 1926; Neboiss 1971). This was complicated by an incorrect synonymy of *Panops* with the distantly related South American genus *Lasia* Wiedemann, 1824 by Kertész (1909) (see discussion in Neboiss 1971). Neboiss (1971) provides a key to species of this genus.

### ***Panops* Lamarck, 1804**

<http://species-id.net/wiki/Panops>

Figs 1, 2D, 17–55

*Panops* Lamarck, 1804: 263 – Latreille 1804: 191, 1809: 316, 1810: 392, 443, 1811: 707, 1816: 608, 1825: 492, 1829: 461; Lamarck 1812: 56; Wiedemann 1830: 18; Macquart 1838: 166; Blanchard 1840: 583; Erichson 1840: 140; Walker 1855: 332; Schiner, 1868: 140; Westwood 1876: 509; Bigot 1890: 314; Hardy 1921: 76, 1946: 66; Brunetti 1926: 580; Paramonov 1957: 525; Neboiss 1971: 208; Schlinger and Jefferies 1989: 376. Type species: *Panops baudini* Lamarck, 1804 by monotypy.

*Epicerina* Macquart, 1850: 97 – Bigot 1890: 316. Synonymy in: Hardy 1921: 79; Hardy 1946: 66; Paramonov 1957: 521. Type species: *Epicerina nigricornis* Macquart, 1850 by original designation.

*Neopanops* Schlinger, 1959: 157 – Schlinger and Jefferies 1989: 376. Type species: *Neopanops boharti*, Schlinger, 1959 by original designation. syn. n.

*Panocalda* Neboiss, 1971: 212 – Schlinger and Jefferies 1989: 376. Type species: *Panocalda grossi*, Neboiss, 1971 by original designation. syn. n.

**Diagnosis.** Body length: 8.0–12.5 mm [male], 9.5–14.5 mm [female]. Colouration non-metallic or metallic; head slightly smaller than thorax width, shape hemispherical; postocular ridge and occiput rounded; three ocelli, anterior ocellus reduced in size or absent; posterior margin of eye emarginate; eye apilose or pilose (sparse) (sometimes localized dorsally); position of antennae on head adjacent to ocellar tubercle; eyes not contiguous above antennal base, contiguous below antennal base; palpus present; proboscis length variable, less than or greater than head length; flagellum



**Figure 4.** *Apsona muscaria* Westwood, male, lateral view [700415]. Body length = 8.0 mm.

shape elongate, slightly tapered (female) or elongate, cylindrical (male); flagellum apex lacking terminal setae; scapes separate; subscutellum not enlarged, barely visible; tibial spines present; pulvilli present; wing hyaline, markings absent; costa circumambient (weaker along anal margin); costal margin at pterostigma straight; humeral crossvein present;  $R_1$  not inflated distally; pterostigma and cell  $r_1$  membranous, not ribbed; vein  $R_{2+3}$  present;  $R_4$  and  $R_5$  present as forked petiolate veins; radial veins straight towards wing apex, slightly angled anteriorly; cell  $r_{4+5}$  bisected by 2r-m, basal cell narrow elongate, closed; 2r-m joining  $M_1$  to stem  $R_{4+5}$ ;  $R_4$  with or without spur vein; medial vein complement with  $M_1$ ,  $M_2$  and  $M_3$  present; discal cell closed completely; medial veins reaching wing margin; cell  $m_3$  present;  $CuA_1$  joining  $M_3$ , petiolate to wing margin;  $CuA_2$  fused to  $A_1$  before wing margin, petiolate to margin; wing microtrichia absent; anal lobe well developed; alula well developed; abdominal tergites smooth, rounded; abdomen shape greatly rounded, inflated (larger in female). Male genitalia (Fig. 17) typical for Panopinae and varying little between species: gonostylus fused with gonocoxite and non-articulated, but with lightly sclerotized areas ventrally indicating flexion of gonostylus with gonocoxite; gonostylus as ventrally curved process with cup-like ventromedial surface; aedeagus consisting of flattened quadrangular, or cylindrical, parameral sheath with ventral rod-like structure with apical gonopore; ejaculatory apodeme poorly developed.



**Figure 5.** *Apsona muscaria* Westwood, male, oblique view [700418]. Body length = 8.0 mm.

**Included species.** *Panops aurum* sp. n.; *P. australae* Neboiss, 1971; *P. baudini* Lamarck, 1804; *P. boharti* (Schlinger, 1959) comb. n.; *P. conspicuus* (Brunetti, 1926); *P. danielsi* sp. n.; *P. grossi* (Neboiss 1971) comb. n.; *P. jade* sp. n.; *P. schlingeri* sp. n.

**Comments.** *Panops* is the type genus for the subfamily Panopinae and includes some large metallic coloured species. The genus is endemic to Australia and neighbouring Papua region of Indonesia. The original concept of the genus was expanded to include species from the New World by some authors, but these have subsequently been placed in the separate and distantly related genus *Lasia* Wiedemann, 1824 (e.g. *Lasia metallica* Rondani, 1863; *Lasia ocelliger* (Wiedemann, 1830)). Bequaert (1931) and later Neboiss (1971), discuss the historically confused and intertwined generic concepts of *Lasia* and *Panops* (sometimes including *Mesophysa*) in previous treatments of the group by various authors. Based on a series of characters, it is clear that those Australasian species are placed in *Panops* or *Mesophysa*, while the New World species are placed in *Lasia*. In his description of *Neopanops*, Schlinger (1959) suggested that the genus was closely related to *Panops* and provided an extensive list of characters distinguishing the two. Similarly, Neboiss (1971) provided a list of characteristics to differentiate *Panocalda* from the closely related *Panops* and *Neopanops*. Both Schlinger (1959) and Neboiss (1971) distinguished their respective genera based on characters such as eye pilosity, length of proboscis, shape of ocellar tubercle, palpi length, head



**Figure 6.** *Apsona muscaria* Westwood, male, anterior view [700419]. Body length = 8.0 mm.

width, parafacial pilosity and wing length. With the inclusion of the four new species described here, and a critical re-examination of the characters used to differentiate *Neopanops* and *Panocalda* from *Panops*, it is clear that all of these characters are variable and that only one genus is warranted. Some species of *Panops* have pilose eyes, either uniformly sparse and minute (i.e. *P. danielsi* sp. n., *P. boharti* comb. n., *P. baudini*) or localized (*P. grossi* comb. n.), with the other species being apilose. In no species of *Panops* are the eyes uniformly dense pilose, as is found in most other panopine genera (e.g. *Apsona*, *Lasia*). This paucity of eye pilosity is shared with only a few other genera, including the Australian *Leucopsina* and *Mesophysa*, as well as the highly derived genus *Corononcodes* Speiser, 1920 from the Palaearctic and Afrotropical regions. Proboscis length is a frequently used character in acrocerid taxonomy, but in *Panops* the length is dramatically variable, with a proboscis much shorter than the head height in some species (e.g. *P. jade* sp. n., *P. schlingeri* sp. n., *P. boharti* comb. n.) while the rest have a proboscis longer than the head height. *Panops* is a variable genus, but can be differentiated from all other Panopinae based on the diagnosis above, and specifically from all other genera in the Australasian region based on tibial spines being present (cf. *Apsona*) and wing crossvein 2r-m joining to  $R_{4+5}$  (cf. *Leucopsina*, *Mesophysa*). Like most acrocerids, species of *Panops* display distinct sexual dimorphism with males often have slightly



**Figure 7.** *Leucopsina odyneroides* Westwood, male, lateral view [700421]. Body length = 9.0 mm.

smaller body size and larger antennae than females. Many Old World panopine genera (e.g. *Apsona*, *Panops*, *Rhysogaster* Aldrich, 1927) have a distinctive unidirectional arrangement of the pile on the head and thorax, giving the individual a dramatic change in appearance when viewed head on (e.g. Figs 20, 23, 40); the biological significance of this is unknown.

### Key to *Panops* species

*Panops baudini* keys to two couplets as the eye pilosity is extremely minute in some individuals and may be overlooked. Females are unknown for *P. boharti* comb. n. and *P. aurum* sp. n., whilst males are unknown for *P. schlingeri* sp. n.

- |   |  |  |
|---|--|--|
| 1 | Eye sparsely pilose (Fig. 39) or pilosity localized dorsally (Fig. 44).....                                    | 2                                      |
| – | Eye completely apilose (Fig. 18) .....   | 5                                      |
| 2 | Proboscis elongate, length greater than head height (Fig. 18).....   | 3                                      |
| – | Proboscis very short, hardly projecting from oral cavity, shorter than head height (Figs 44, 50) .....         | 4                                      |
| 3 | Postpronotal lobe dark, concolourous with rest of pleuron (widely distributed in Australia) (Figs 25–30) ..... | <b><i>P. baudini</i> Lamarck, 1804</b> |



**Figure 8.** *Leucopsina odyneroides* Westwood, male, dorsal view [700423]. Body length = 9.0 mm.

- Postpronotal lobe yellow, pleuron greenish (Queensland) (Figs 39–43) .....  
..... *P. danielsi* sp. n.
- 4 Eye extending posteriorly beyond widest part of head; eye with sparse, minute  
pile of uniform length across eye (length subequal to width of lateral ocellus);  
ocellar tubercle not touching margin of eye; palpus as long or longer than  
proboscis (Papua) (Figs 31–33) ..... *P. bobarti* (Schlinger, 1959), comb. n.
- Eye not extending posteriorly beyond widest part of head; eye pilose on dor-  
sal-lateral region only, pile denser and more elongate (length much greater  
than width of lateral ocellus); ocellar tubercle touching margin of eye; palpus



**Figure 9.** *Leucopsina odyneroides* Westwood, male, anterior view [700426]. Body length = 9.0 mm.

- half as long as proboscis (South Australia) (Figs 44–47).....
- ..... *P. grossi* (Neboiss, 1971), **comb. n.**
- 5 Proboscis short, hardly projecting from oral cavity..... **6**
- Proboscis elongate, length equal to, or greater than head height..... **7**
- 6 Postpronotal lobes dark yellow; femora dark brown, rest of legs cream (North-  
ern Territory) (Figs 53–55)..... *P. schlingeri* **sp. n.**
- Postpronotal lobes and legs dark, concolourous with rest of body (Queens-  
land) (Figs 48–52) ..... *P. jade* **sp. n.**
- 7 Postpronotal lobes pale, contrasting with rest of thorax (Figs 34–38).....
- ..... *P. conspicuus* (Brunetti, 1926)
- Postpronotal lobes dark, concolourous with rest of thorax (Figs 21, 28) ..... **8**





**Figure 11.** *Leucopsina odyneroides* Westwood, female, dorsal view [700447]. Body length = 12.0 mm.

***Panops aurum* sp. n.**

[urn:lsid:zoobank.org:act:3864CACB-368C-4770-88E8-8346544EBED7](http://zoobank.org/urn:lsid:zoobank.org:act:3864CACB-368C-4770-88E8-8346544EBED7)

[http://species-id.net/wiki/Panops\\_aurum](http://species-id.net/wiki/Panops_aurum)

Figs 18–20

**Type material.** **Holotype** male, AUSTRALIA: **Western Australia:** Darlington, 450 ft., E.S. Ross, D.Q. Cavagnaro, 5.ix.1962 [-31.901, 116.081] (CAS).

**Diagnosis.** Eye apilose; proboscis longer than head height; body non-metallic; antennae red-brown; parafacial with yellow marginal pile; postpronotal lobe concoloured with rest of thorax; legs dark yellow, femora brown-black.



**Figure 12.** *Mesophysa tenaria* Neboiss, male, lateral view [700448]. Body length = 10.0 mm.

**Description.** Body length: 11.0 mm (male). Head with eye apilose; ocellar tubercle raised laterally; medial ocellus absent; occiput brown-black, occipital pile yellow, postocular ridge and gena overlain with grey pubescence; clypeus length equal to oral cavity, brown-black; palpus yellow; margin of oral cavity (parafacial) densely pilose (yellow); proboscis longer than head height; flagellum apex of uniform width, truncated apically, flagellum red-brown; scape and pedicel brown. Thorax with postpronotal lobe brown-black; scutum black, scutal vestiture dense yellow-gold pile; scutellum black; pleuron black; coxae black; femora brown-black, apices dark yellow; tibiae dark yellow; tarsi dark yellow; lower calypter white with dark yellow margin; wing hyaline, venation dark; vein  $R_4$  without spur vein. Abdomen shape rounded globose, much larger than thorax, colour orange-red to yellow, dark markings anteriorly and medially, vestiture dense elongate pile, yellow anteriorly, brown posteriorly on tergites 2–5.

**Etymology.** The specific epithet is derived from the Latin, *aurum* – gold; referring to the distinctive golden setal pile on the head and thorax.

**Comments.** *Panops aurum* sp. n. is known only from a single male specimen from Western Australia. The fringing yellow setae around the oral cavity and yellow pile on the thorax are distinctive for the species.



**Figure 13.** *Mesophysa tenaria* Neboiss, male, oblique view [700450]. Body length = 10.0 mm.

***Panops austrae* Neboiss, 1971**

[http://species-id.net/wiki/Panops\\_austrae](http://species-id.net/wiki/Panops_austrae)

Figs 21–24

*Panops austrae* Neboiss, 1971: 209 – Schlinger and Jefferies 1989: 376.

**Type material examined.** **Holotype** female, AUSTRALIA: **Northern Territory:** nr. Mount Olga [-25.3, 130.73], C.A., Paul Genery, ix.1960, picked up dead in sand (Type- T.4177) (NMV).

**Other material examined.** AUSTRALIA: **Western Australia:** male, Wialki [-30.483, 118.117], R. P. McMillan, 12.x.1983 (WAM); male, W of Norseman, *Eucalyptus* woodland, dry gully to salt lake, Malaise trap, C. Lambkin et al., ANIC bulk sample 2184, 1-17.xi.2003 271m [-32.186, 121.721] (ANIC).

**Diagnosis.** Eye apilose; proboscis equal to head height; body metallic green-blue; antennae yellow-brown; parafacial without marginal pile; postpronotal lobe concolourous with rest of thorax; legs black.

**Redescription.** Body length: 8.0–10.0 mm (male), 14.5 mm (female). Head with eye apilose; ocellar tubercle relatively flat, medial ocellus present; occiput metallic green-blue, occipital pile dense, white; postocular ridge and gena overlain with grey



**Figure 14.** *Mesophysa tenaria* Neboiss, male, anterior view [700452]. Body length = 10.0 mm.

pubescence; clypeus length equal to oral cavity, brown-black; palpus white or black; margin of oral cavity (parafacial) glabrous; proboscis equal or slightly longer than head height; flagellum dark yellow-orange, suffused with brown, apex in male tapered, narrow apically; scape and pedicel brown or dark yellow. Thorax postpronotal lobe green; scutum metallic green or metallic blue, scutal vestiture dense white pile; scutellum metallic blue-green; pleuron metallic green or metallic blue; coxae black with metallic blue iridescence; femora black; tibiae black or brown; tarsi black; lower calypter white, with brown margin; wing hyaline (male) or slightly infusate (female), venation dark; vein  $R_4$  with spur vein. Abdomen shape rounded globose, much larger than thorax (female) or rounded to conical, not larger than thorax (male), colour metallic green or metallic blue violet, vestiture as minute setae, dense white-silver elongate setae along anterior margin of tergites 2–5.

**Comments.** *Panops austra*e is a large, metallic coloured species similar to *P. jade* sp. n. and *P. schlingeri* sp. n. It is easily distinguished from these species by the longer proboscis and dense white thoracic pile. This species is known from remote, arid regions of the Northern Territory and Western Australia.



**Figure 15.** *Mesophysa tenaria* Neboiss, female, lateral view [700453]. Body length = 11.0 mm.

***Panops baudini* Lamarck, 1804**

[http://species-id.net/wiki/Panops\\_baudini](http://species-id.net/wiki/Panops_baudini)

Figs 1, 17, 25–30

*Panops baudini* Lamarck, 1804: 265 – Latreille 1809: 316, 1810: 443, 1811: 710; Wiedemann 1830: 19; Erichson 1840: 141; Walker 1855: 333; Kertész 1909: 9; Hardy 1946: 66; Edwards 1930: 193; Paramonov 1957: 526; Neboiss 1971: 208; Schlinger and Jefferies 1989: 376.

*Mesophysa marginata* Macquart, 1838: 168 – Blanchard 1840: 584.

*Epicerina nigricornis* Macquart, 1850: 98 – Kertész 1909: 8; Hardy 1918: 61, 1921: 79, 1946: 66; Brunetti 1926: 578.

*Panops lamarckianus* Westwood, 1876: 508 – Kertész 1909: 9; Paramonov 1957: 526.

*Mesophysa australiae* Thomson, 1869: 475 – Westwood 1876: 517.

*Panops australiae*. Kertész, 1909: 8.

*Mesophysa baudini* Brunetti, 1926: 581.

*Panops nigricornis*. Hardy, 1946: 66.

**Type material.** *Panops baudini* Lamarck. **Neotype** female, AUSTRALIA: **New South Wales:** Asquith (nr, Sydney), 10.x.1962, A.L. Dyce (ANIC) (designated by Neboiss



**Figure 16.** *Mesophysa tenaria* Neboiss, female, oblique view [700454]. Body length = 11.0 mm.

1971). Neboiss (1971) discussed the identity of this species based on the original species description and justification for designating the neotype [examined].


*Mesophysa marginata* Macquart. **Type** female, [no label data] (MHN). See discussion by Neboiss (1971).

*Epicerina nigricornis* Macquart. **Type** male, AUSTRALIA: “2/47 Tasmanie J. Verreaux 1847” (MNHN). See discussion by Paramonov (1957) and Neboiss (1971) regarding synonymy and possible erroneous locality recording.

*Panops lamarckianus* Westwood. **Type** male, AUSTRALIA: **Queensland:** Moreton Bay, 1859 (OUMNH).

*Mesophysa australasiae* Thomson. **Type** male, AUSTRALIA: **New South Wales:** Sydney, Kinb. (NHRS). See discussion by Hardy (1921) and Neboiss (1971) regarding synonymy.

**Other material examined.** AUSTRALIA: **Queensland:** male, female, Isla Gorge National Park, [-25.183, 149.966] 12.ix.1992, 320m, G. Daniels (GDCB); male, Isla Gorge National Park, [-25.183, 149.966] 11.ix.1992, 320m, R. Eastwood (GDCB); 32 km S Theodore, [-25.166, 150.000], 13.ix.1992, 300m, G. Daniels (GDCB); 2 males, female, 43 km WSW Millmerran, [-27.983, 150.933],



**Figure 17.** *Panops baudini* Lamarck. **A** male genitalia, lateral view **B** same, ventral view. Scale line = 0.2 mm. Abbreviations: **c** cercus; **e** epandrium; **g** gonocoxite; **gs** gonostylus; **ps** parameral sheath of aedeagus.



**Figure 18.** *Panops aurum* sp. n., male, lateral view [700495]. Body length = 11.0 mm.

21.ix.1986, G. & A. Daniels (GDCB); 2 females, Lake Broadwater, nr. Dalby, [-27.361, 151.102], site 8, 27.ix.1986, G. & A. Daniels (GDCB); male, Gayndah, Masters (NMV). **New South Wales:** female, Sydney swamps (NMV); male, Sydney, 17.x.1932, G.M. Goldfinch (ANIC); female, Ku-ring-gai Chase National Park [-33.651, 151.201], 2.x.1972, A. & G. Daniels (GDCB); 2 males, Goondera Ridge, Royal National Park [-34.122, 151.063], 24.x.1976, G. & A. Daniels (GDCB). **Victoria:** female, Mitta Mitta River, 8km NW of Dartmouth Dam [-36.566, 147.55], 30.x.1976, A. A. Calder (NMV). **Western Australia:** 3 males, W of Norseman, Eucalyptus woodland, dry gully to salt lake, Malaise trap, C. Lambkin et al., ANIC bulk sample 2184, 1-17.xi.2003 271m [-32.186, 121.721] (ANIC); male, Wongan Hills area [-30.871, 116.771], Greg Guérin, on flowers of *Microcorys* (CAS); female, East Yuna Nature Reserve, 34 km WNW Mullewa [-28.42, 115.42], 23-24.ix.1983, C. & T. Houston, 559-17, on flowers of ?*Helipterum* (WAM); female Australia, Boorabbin Rock National Park [-31.23, 120.16], W Coolgardie, 26.ix.2005, L. Packer (CNC) [not examined but identity confirmed by B. Sinclair].

**Diagnosis.** Eye minutely pilose; proboscis longer than head height; body black (with faint blue iridescence in western population); antennae red-brown to black; parafacial with marginal pile; postpronotal lobe concolourous with rest of thorax; femora black with pale apices, rest of leg dark yellow to white with black on tibiae; abdomen red or yellow laterally; distiphallus broad apically.

**Redescription.** Body length: 9.5–12.5 mm (male), 11.0–14.0 mm (female). Head with eye sparsely pilose with minute setae (appears apilose); ocellar tubercle



**Figure 19.** *Panops aurum* sp. n., male, dorsal view [700496]. Body length = 11.0 mm.

raised laterally or relatively flat; medial ocellus reduced; occiput brown-black, occipital pile white, sparse; postocular ridge and gena overlain with grey pubescence; clypeus length equal to oral cavity, brown-black; palpus white or yellow; margin



**Figure 20.** *Panops aurum* sp. n., male, anterior view [700497]. Body length = 11.0 mm.

of oral cavity (parafacial) pilose; proboscis longer than head height; flagellum red-brown to black; scape and pedicel brown. Thorax with postpronotal lobe brown-black; scutum black, scutal vestiture dense white pile; scutellum black; pleuron black (thorax with slight bluish iridescence in western populations); coxae black; femora black or brown-black, apices dark yellow; tibiae predominantly black with dark yellow to white (apically); tarsi dark yellow to white; lower calypter white, with yellow margin; wing hyaline (male) or slightly infusate (female); venation dark; vein  $R_4$  with spur vein, rarely without. Abdomen shape rounded globose, much larger than thorax, colour highly variable, orange-red to yellow, dark markings anteriorly and medially, or dark yellow, brown anteriorly on tergites 2–6, vestiture as extensive short white-silver pile, longer laterally.



**Figure 21.** *Panops australis* Neboiss, male, lateral view (partially denuded) [700499]. Body length = 8.0 mm.

**Comments.** The type for the genus, *Panops baudini* is the most commonly represented species in collections. This species is distributed in Queensland, New South Wales, Victoria and Western Australia. The apex of the aedeagus is broad and quadrangular in this species (Fig. 17) while in all other species it is much narrower. The record from Tasmania is apparently erroneous (Neboiss 1971). Western Australian individuals have more reddish colouration laterally on the abdomen, particularly in males, and the body has a bluish iridescence (Fig. 1). This bluish iridescence is not seen in specimens from eastern states.

***Panops boharti* (Schlinger, 1959), comb. n.**

[http://species-id.net/wiki/Panops\\_boharti](http://species-id.net/wiki/Panops_boharti)

Figs 31–33

*Neopanops boharti* Schlinger, 1959: 157 – Neboiss 1971: 212; Schlinger and Jefferies 1989: 376.

**Type material examined.** **Holotype** male, INDONESIA: **Papua:** Cyclops Mountains, Sabron, 930 ft. [-2.509, 140.523], iv.1936, L. E. Cheesman, B. M. 1936-271 (BMNH).

**Diagnosis.** Eye pilose; eye extends posteriorly beyond maximum head width; proboscis very short, not extending beyond oral cavity; body brown and yellow; antennae



**Figure 22.** *Panops australae* Neboiss, male, dorsal view [700502]. Body length = 8.0 mm.

yellow; parafacial without marginal pile; postpronotal lobe cream with brown spot; legs yellow, femora brown with yellow apices; lower calypter cream with brown margin.

**Redescription.** Body length: 9.0 mm (male). Head with eye sparsely pilose, slightly denser and elongate laterally; eye extends posteriorly beyond maximum head width; ocellar tubercle relatively flat; medial ocellus present; occiput cream, brown suffusion laterally; occipital pile white, sparse; flagellum yellow, apex uniform width, truncated apically; scape and pedicel dark yellow; clypeus minute, yellow-brown; palpus yellow; margin of oral cavity (parafacial) glabrous; proboscis not extending beyond oral cavity. Thorax with postpronotal lobe cream, brown suffusion dorsally; scutum brown, cream posterolaterally; scutal vestiture dense brown and white, matching respective scutal



**Figure 23.** *Panops australis* Neboiss, male, anterior view [700498]. Body length = 8.0 mm.

markings; scutellum brown with bluish iridescence, cream laterally; pleuron cream with brown markings; coxae cream with brown markings; femora cream with brown on middle half; tibiae dark yellow; tarsi dark yellow; lower calypter white, brown marginally on membrane; wing hyaline, venation brownish, pale yellow distally along costa and radial veins; vein  $R_4$  with spur vein. Abdomen rounded globose, slightly larger than thorax, colour dark yellow, brown on tergites 3–6, vestiture minute setae, dense white-silver elongate setae along anterior margin of tergites 2–5.

**Comments.** *Panops boharti* comb. n. was described by Schlinger (1959) as the sole species in the genus *Neopanops* but is transferred herein to *Panops*. This Indonesian species is the only non-Australian representative of the genus, and is distinctive based on body colouration and markings, very short mouthparts, eye pilosity and eye shape. Only the male is known.



**Figure 24.** *Panops austrae* Neboiss, female, dorsal view [700508]. Body length = 14.5 mm.



**Figure 25.** *Panops baudini* Lamarck (western form), male, lateral view [700505]. Body length = 9.5 mm.



**Figure 26.** *Panops baudini* Lamarck (western form), male, oblique view [700509]. Body length = 9.5 mm.

***Panops conspicuus* (Brunetti, 1926)**

[http://species-id.net/wiki/Panops\\_conspicuus](http://species-id.net/wiki/Panops_conspicuus)

Figs 34–38

*Epicerina conspicua* Brunetti, 1926: 579.

*Panops conspicuus* (Brunetti, 1926) – Edwards 1930: 193; Paramonov 1957: 529; Neboiss 1971: 210; Schlinger and Jefferies 1989: 376.

**Type material examined.** Holotype female, AUSTRALIA: **Western Australia:** Kalamunda [-31.974, 116.058], 14.iii–14.iv.1914, R.E. Turner, 1914-349 (BMNH).

**Other material examined.** AUSTRALIA: **Victoria:** male, female, Kiata [-36.366, 141.791], R. Oldfield, X 4172, captured as copulating pair (NMV). **Western Australia:** female, Boulder Rock [-32.133, 116.166], 15.iii.1981, M.J. Smart, Jarrah Forest, 300m, hovering 2–3 m above ground, taken at rest on leaf (WAM); 4.5 km E Lake Monger on Wanarra Road [-29.544, 116.775], 7.v.2008, T.F. Houston and E. G. Cunningham, 1266-1 (WAM).



**Figure 27.** *Panops baudini* Lamarck (western form), male, anterior view [700510]. Body length = 9.5 mm.

**Diagnosis.** Eye apilose; proboscis longer than head height; body colour and shape sexually dimorphic: male black with slender body, female yellow and brown with globose abdomen; antennae yellow-brown to red-brown with black suffusion; parafacial without marginal pile; postpronotal lobe yellow; legs yellow with brown medially on femora and tibiae.

**Redescription.** Body length: 11.0 mm (male), 12.0–13.0 mm (female). Head with eye apilose; ocellar tubercle raised laterally; medial ocellus present; occiput colour brown-black (male) or brown with dark yellow spot laterally (female); occipital pile yellow; postocular ridge and gena glabrous; clypeus shorter than oral cavity; yellow-brown; palpus yellow; margin of oral cavity (parafacial) glabrous; proboscis longer than head height; flagellum dark yellow, suffused with brown (female) or red with black suffusion (male), apex in male tapered, narrow apically; scape and pedicel brown. Thorax with postpronotal lobe yellow; scutum black (male) or yellow and brown (markings variable) (female); scutal vestiture dense white pile or dense yellow-gold pile; scutellum black or brown; pleuron brown; coxae brown; femora brown-black, apices dark yellow; tibiae dark yellow or dark yellow, suffused with brown; tarsi dark yellow; lower calypter white, with dark yellow margin; wing hyaline (male) or slightly infusate (female), venation dark; vein  $R_4$  with spur vein. Abdomen shape rounded globose, much



**Figure 28.** *Panops baudini* Lamarck (eastern form), female, lateral view [700512]. Body length = 12.0 mm.

larger than thorax (female) or cylindrical along length (male), colour orange-yellow or brown-black, vestiture elongate yellow pile (whitish in male).

**Comments.** *Panops conspicuus* is recorded from arid regions of southwest Western Australia and Western Victoria. There is dramatic sexual dimorphism in both body colouration and shape in this species, with males very similar to species of *Mesophysa*. *Panops conspicuus* can be differentiated from other *Panops* species by the bright yellow postpronotal lobes, elongate mouthparts, yellow and brown colouration (female), and apilose eyes. Females of this species are similarly coloured to females of *P. grossi* comb. n., a species which also displays dramatic sexual dimorphism.

***Panops danielsi* sp. n.**

urn:lsid:zoobank.org:act:3FAB3406-C6A4-42CC-9ABC-B82BCB22FDE8

[http://species-id.net/wiki/Panops\\_danielsi](http://species-id.net/wiki/Panops_danielsi)

Figs 39–43

**Type material.** **Holotype** male, AUSTRALIA: **Queensland:** 3km SW Fox Ck. x-ing [crossing], ‘Wolverton’ [-13.104, 142.970], 13.iv.1989, G. and A. Daniels (AMS).



**Figure 29.** *Panops baudini* Lamarck (eastern form), female, oblique view [700513]. Body length = 12.0 mm.

**Paratypes.** AUSTRALIA: **Queensland:** female, male, same data as holotype (GDCB) (CAS); female, 7 km NNW Coen, [-13.844, 143.163], 17.iv.1989, G. and A. Daniels (GDCB); female, 26 km W 'Fairview', [-15.535, 144.154], 20.iv.1989, G. and A. Daniels (GDCB).

**Diagnosis.** Eye uniformly sparse pilose; proboscis longer than head height; body dark yellow and brown, with metallic green-blue iridescence; antennae red-brown or black; parafacial with marginal pile; postpronotal lobe dark yellow; legs dark yellow and brown.

**Description.** Body length: 11.0 mm (male), 10.5–12.0 mm (female). Head with eye sparsely pilose, uniformly distributed, setae minute; ocellar tubercle raised laterally; medial ocellus absent; occiput metallic green-blue; occipital pile yellow; postocular ridge and gena overlain with grey pubescence; flagellum apex in male uniform width, truncated apically, narrower in female, red-brown (male) or black (female); scape and pedicel dark yellow; clypeus length equal to oral cavity, brown-black; palpus yellow; margin of oral cavity (parafacial) pilose; proboscis longer than head height. Thorax with postpronotal lobe yellow; scutum glossy black (with metallic iridescence), dark yellow marginally; scutal vestiture dense yellow-gold pile; scutellum brown, dark yellow medially; pleuron brown with metallic iridescence; coxae black or brown; femora brown-black, apices dark yellow; tibiae dark yellow, suffused with brown; tarsi dark



**Figure 30.** *Panops baudini* Lamarck (eastern form), female, anterior view [700514]. Body length = 12.0 mm.

yellow; lower calypter white, with yellow margin; wing hyaline, venation dark; vein  $R_4$  with spur vein. Abdomen shape rounded globose, much larger than thorax (female) or rounded to conical, not larger than thorax (male), colour black with metallic green iridescence (female) or dark yellow, brown anteriorly on tergites 2–6 (male), vestiture extensive white-silver elongate setae, brown posteromedially on tergites 3–5 (female) or erect dark pile (male).

**Etymology.** This species is named in honour of the collector of this species, Greg Daniels.

**Comments.** *Panops danielsi* sp.n. is known only from Far Northern Queensland. This species is closely related to *P. baudini* as both species have similar shaped mouth-parts and pilose eyes. *Panops danielsi* sp. n. can be distinguished by the more evident eye pilosity, yellow postpronotal lobes and body colouration.



**Figure 31.** *Panops bobarti* (Schlinger) comb. n., male, lateral view [700515]. Body length = 9.0 mm.

***Panops grossi* (Neboiss, 1971), comb. n.**

[http://species-id.net/wiki/Panops\\_grossi](http://species-id.net/wiki/Panops_grossi)

Figs 44–47

*Panocalda grossi* Neboiss, 1971: 214 – Schlinger and Jefferies 1989: 376.

**Type material examined.** **Holotype** female, AUSTRALIA: **Northern Territory:** Koolpinyah, 21.iv.1916 [-12.331, 131.148] G. F. Hill, (in copula) (SAM).

**‘Allotype’.** AUSTRALIA: **Northern Territory:** same data as holotype (SAM).

**Diagnosis.** Eye pilose dorsally only, relatively dense and elongate; proboscis shorter than head height; body colour and shape sexually dimorphic: male metallic olive green, female yellow and brown, globose; antennae yellow; parafacial without marginal pile; postpronotal lobe and legs concolourous with rest of body.

**Redescription.** Body length: 9.0 mm (male), 12.0 mm (female). Head eye pilose dorsally only, dense and relatively elongate; occiput olive green, occipital pile dense white (male) or yellow (female); postocular ridge and gena overlain with grey pubescence; ocellar tubercle raised laterally or relatively flat; medial ocellus absent; clypeus shorter than oral cavity, yellow-brown; palpus black; margin of oral cavity (parafacial) glabrous; proboscis not extending beyond oral cavity; flagellum yellow, apex in male uniform width, truncated apically; scape and pedicel brown. Thorax with postpronotal lobe yellow (female) or green (male); scutum metallic olive green or yellow-orange; scutal vestiture dense white or yellow-gold pile; scutellum metallic olive green or orange-yellow with brown suffusion; pleuron orange or metallic olive green; coxae brown; femora brown-black, apices dark yellow; tibiae brown; tarsi brown; lower calypter white, brown marginally on membrane



**Figure 32.** *Panops boharti* (Schlinger) comb. n., male, dorsal view [700517]. Body length = 9.0 mm.

or white, with dark yellow margin; wing hyaline or slightly infusate, venation dark; vein  $R_4$  without spur vein. Abdomen shape with male rounded, not larger than thorax, metallic olive green, vestiture dense short pile, longer laterally; female rounded globose, much larger than thorax (female), orange-yellow (female), vestiture elongate yellow pile.

**Comments.** *Panops grossi* comb. n. was described by Neboiss (1971) as the sole species in the genus *Panocalda* but is transferred herein to *Panops*. This species is apparently closely related to *P. boharti* comb. n. based on eye pilosity, and *P. schlingeri* sp. n. and *P. jade* sp. n. based on the short mouthparts. All of these species are northern Australian or Indonesian species. *Panops grossi* comb. n. can be distinguished from all



**Figure 33.** *Panops boharti* (Schlinger) comb. n., male, anterior view [700522]. Body length = 9.0 mm.

other *Panops* based on the dense patch of relatively elongate pile on the dorsal part of the eye. This species displays a dramatic sexual dimorphism similar to that found in *P. conspicuus*, with females being orange-yellow in colour.

***Panops jade* sp. n.**

urn:lsid:zoobank.org:act:96D0BD2A-0C81-4BCE-BB32-671D1C2D901C

[http://species-id.net/wiki/Panops\\_jade](http://species-id.net/wiki/Panops_jade)

Figs 2D, 48–52

**Type material.** **Holotype** male, AUSTRALIA: **Queensland:** Isla Gorge National Park [-25.183, 149.966], 3.x.1991, 320 m, G. Daniels (AMS).

**Paratypes.** AUSTRALIA: **Queensland:** female, Isla Gorge National Park [-25.183, 149.966], 3.x.1991, 320 m, G. Daniels (CAS); female, Isla Gorge National Park [-25.183, 149.966], 14.ix.1992, 320 m, G. Daniels (AMS).



**Figure 34.** *Panops conspicuus* (Brunetti), male, lateral view [700525]. Body length = 11.0 mm.



**Figure 35.** *Panops conspicuus* (Brunetti), male, oblique view [700527]. Body length = 11.0 mm.

**Diagnosis.** Eye apilose; proboscis shorter than head height; body metallic green-blue to violet iridescence; antennae red-brown; parafacial with marginal pile; postpronotal lobe concolourous with rest of thorax; legs black with metallic blue-violet iridescence.



**Figure 36.** *Panops conspicuus* (Brunetti), male, anterior view [700528]. Body length = 11.0 mm.

**Description.** Body length: 11.5 mm (male), 11.5–12.0 mm (female). Head with eye apilose; ocellar tubercle relatively flat; medial ocellus present; occiput metallic green-blue, occipital pile white, sparse; postocular ridge and gena overlain with grey pubescence; clypeus length equal to oral cavity, black with blue-green suffusion; palpus black; margin of oral cavity (parafacial) pilose; proboscis extending beyond oral cavity, but shorter than head height; flagellum apex in male tapered, slightly rounded apically, red-brown; scape and pedicel red-brown. Thorax with postpronotal lobe blue-violet; scutum metallic blue-violet, green posteromedially; scutellum metallic blue-violet; coxae and femora with metallic blue-violet iridescence; tibiae black; tarsi black; lower



**Figure 37.** *Panops conspicuus* (Brunetti), female, lateral view [700529]. Body length = 13.0 mm.

calypter white with brown margin; wing hyaline, venation dark; vein  $R_4$  with spur vein. Abdomen shape rounded globose, much larger than thorax, colour metallic green or blue-violet iridescent, vestiture extensive white-silver short pile, longer laterally.

**Etymology.** This beautifully coloured species is named after my daughter, Jade Tanya Winterton, whose name also describes the deep green colouration found in this species.

**Comments.** *Panops jade* sp. n. is a distinctive species with extensive green to blue-violet iridescence, particularly in the female. It is similar to the western Australian species, *P. austrae*, but is distinguished by the length of the mouthparts, leg colour and different vestiture pattern on the abdomen. *Panops jade* sp. n. is known only from Isla Gorge National Park in southern Queensland. Both males and females are recorded from Spinifex grass (*Triodia* sp.), presumably at rest.

***Panops schlingeri* sp. n.**

urn:lsid:zoobank.org:act:03D163A1-D1DA-4810-8D88-77F76D5CC490

[http://species-id.net/wiki/Panops\\_schlingeri](http://species-id.net/wiki/Panops_schlingeri)

Figs 53–55

**Type material.** **Holotype** female, AUSTRALIA: **Northern Territory:** 9 km NE of Mudginbarry H.S. (on scarp), 10.vi.1973, D. H. Colless [-12.310, 132.579] (ANIC).



**Figure 38.** *Panops conspicuus* (Brunetti), female, oblique view [700530]. Body length = 13.0 mm.



**Figure 39.** *Panops danielsi* sp. n., male, lateral view [700531]. Body length = 11.0 mm.



**Figure 40.** *Panops danielsi* sp. n., male, anterior view [700532]. Body length = 11.0 mm.

**Paratype.** AUSTRALIA: **Northern Territory:** female, 8 km SSW of Oenpelli Mission 7.vi.1973, J. Cardale [-12.381, 133.024] (ANIC).

**Diagnosis.** Eye apilose; proboscis shorter than head height; body metallic green-blue iridescence; antennae orange; parafacial without marginal pile; postpronotal lobe dark yellow; legs dark yellow, femora brown-black with yellow apices.

**Description.** Body length: 9.5–11.0 mm (female only). Head with eye apilose; ocellar tubercle relatively flat; medial ocellus present; occiput metallic green-blue, occipital pile white, dense; postocular ridge and gena overlain with grey pubescence; clypeus shorter than oral cavity, brown-black; palpus black; margin of oral cavity (parafacial) glabrous; proboscis not extending beyond oral cavity; flagellum orange; scape and pedicel dark red-yellow. Thorax with postpronotal lobe yellow; scutum metallic green to blue iridescent; scutal vestiture dense white pile; scutellum metallic blue-green; pleuron metallic green to blue iridescent; coxae brown-black with metal-



**Figure 41.** *Panops danielsi* sp. n., female, lateral view [700533]. Body length = 12.0 mm.



**Figure 42.** *Panops danielsi* sp. n., female, oblique view [700534]. Body length = 12.0 mm.

lic blue iridescence; femora brown-black, apices dark yellow; tibiae dark yellow; tarsi dark yellow; lower calypter white, with dark yellow margin; wing hyaline, venation dark; vein  $R_4$  without spur vein. Abdomen shape rounded globose, much larger than



**Figure 43.** *Panops danielsi* sp. n., female, anterior view [700535]. Body length = 12.0 mm.

thorax, dark with metallic green to blue iridescence, vestiture as dense short pile, longer laterally.

**Etymology.** I am honoured to name this species after the world-renowned Acroceridae taxonomist Dr. Evert Irving Schlinger.

**Comments.** *Panops schlingeri* sp. n. is known only from two female specimens collected in the Northern Territory. This species is differentiated easily by the green-blue iridescence on the body and dark yellow postpronotal lobes.



**Figure 44.** *Panops grossi* (Neboiss) comb. n., male, lateral view [700536]. Body length = 9.0 mm.

### Subfamily Philopotinae Schiner, 1968

<http://species-id.net/wiki/Philopotinae>

**Type genus.** *Philopota* Wiedemann. Schlinger, 1971: 186.

**Diagnosis.** Body shape slightly to strongly arched and never densely pilose; small to medium sized; antennal flagellum stylate; postpronotal lobes enlarged and meeting medially to form collar behind head; tibial spines absent; wing costal vein ending at wing apex, never circumambient; wing venation highly variable, ranging from relatively complete with cells cu-p, bm br, d and basal  $r_{4+5}$  present, to highly reduced with only cell br present; cell  $m_3$  absent; veins  $R_4$  and  $R_5$  always present as single vein  $R_{4+5}$ ; cubital and medial veins not reaching posterior wing margin; larvae exclusively parasitoids of araneomorph spiders.

### Australasian genera

*Helle* Osten Sacken, 1896; *Schlingeriella* Gillung & Winterton, 2011.

#### *Helle* Osten Sacken, 1896

<http://species-id.net/wiki/Helle>

Figs 3A, 56–59

*Helle* Osten Sacken, 1896: 16 – Hutton 1901: 28; Paramonov 1955: 21; Schlinger and Jefferies 1989: 376. Type species: *Acrocera longirostris* Hudson, 1892: 56 by monotypy.



**Figure 45.** *Panops grossi* (Neboiss) comb. n., male, dorsal view [700537]. Body length = 9.0 mm.

**Diagnosis.** Body length: 4.0–6.0 mm [male], 6.0–7.0 mm [female]. Body shape strongly arched; colouration non-metallic (brown or black); head size slightly narrower than thorax width, shape sub-spherical; postocular ridge and occiput rounded; three ocelli, anterior ocellus reduced in size; posterior margin of eye rounded; eye apilose; position of antennae on head near middle of frons; eyes contiguous above



**Figure 46.** *Panops grossi* (Neboiss) comb. n., male, anterior view [700538]. Body length = 9.0 mm.

antennal base, not contiguous below antennal base; palpus present; proboscis greater than head length; flagellum stylate, apex with terminal seta; postpronotal lobes enlarged, medially contiguous to form collar; subscutellum enlarged; legs not elongated; wing markings absent; costa ending near wing apex, costal margin straight; humeral crossvein absent; radial veins straight or curved towards wing anterior margin;  $R_1$  inflated distally at pterostigma; pterostigma and cell  $r_1$  membranous, not ribbed;  $R_{2+3}$  present;  $R_{4+5}$  angled anteriorly approximately midway; cell  $r_{4+5}$  bisected by 2r-m, basal cell very narrow elongate, closed; 2r-m joining  $M_1$  to  $R_{4+5}$ ; cell  $r_{4+5}$  present, narrow elongate, closed (open apically when 2r-m rarely absent); crossvein 2r-m present (rarely absent);  $R_4$  without spur vein; medial vein complement with  $M_1$ ,  $M_2$  and  $M_3$  present ( $M_3$  fused with  $CuA_1$ ); discal cell closed completely; medial veins not reaching wing margin;  $CuA_1$  joining  $M_3$ , petiolate to margin;  $CuA_2$  fused to  $A_1$  before wing margin, petiolate; wing microtrichia absent; anal lobe well developed; alula well developed; abdominal tergites smooth, rounded; abdomen shape elongate, narrow cylindrical or conical (male), or rounded and inflated (female).



**Figure 47.** *Panops grossi* (Neboiss) comb. n., female, oblique view [700539]. Body length = 12.0 mm.



**Figure 48.** *Panops jade* sp. n., male, lateral view [700540]. Body length = 11.5 mm.

**Included species.** *Helle longirostris* (Hudson, 1892); *H. rufescens* Brunetti, 1926.

**Comments.** *Helle* is an endemic genus to New Zealand that is closely related to *Schlingeriella*, the only other philopotine genus in the region (Gillung and Winterton 2011; Winterton et al. 2007). Characteristics supporting this close relationship include thickening of wing vein  $R_1$  at the pterostigma, elongate mouthparts, apilose eyes, 2r-m absent (rarely in *Helle*) and  $R_{4+5}$  angled anteriorly approximately half way along vein.



**Figure 49.** *Panops jade* sp. n., male, dorsal view [700541]. Body length = 11.5 mm.

*Helle* can be differentiated from all other philopotine genera based on the relatively complete wing venation, inflated  $R_1$  at pterostigma, palpi present and apilose eyes.

#### Key to *Helle* species

- 1      Body colour brown-black, sometimes with metallic iridescence, scutum without dark markings (Figs 56–57) ..... *H. longirostris* (Hudson, 1892)
- Body colour yellowish-orange, scutum with dark longitudinal stripes, narrower anteriorly (Figs 58–59) ..... *H. rufescens* Brunetti, 1926

#### *Schlingeriella* Gillung & Winterton

urn:lsid:zoobank.org:act:99EAC1BE-4A6F-43E0-B61A-6460BF68694E

<http://species-id.net/wiki/Schlingeriella>

Figs 3B, 60–62

*Schlingeriella* Gillung & Winterton, 2011: 22. Type species: *Schlingeriella irwini* Gillung & Winterton, 2011: 23.



**Figure 50.** *Panops jade* sp. n., female, lateral view [700542]. Body length = 12.0 mm.



**Figure 51.** *Panops jade* sp. n., female, dorsal view [700543]. Body length = 12.0 mm.



**Figure 52.** *Panops jade* sp. n., female, anterior view [700545]. Body length = 12.0 mm.

**Diagnosis.** Body length: 2.4–4.0 mm [male], 4.4–6.0 mm [female]. Body shape arched; body colouration non-metallic dark brown; head width much smaller than thorax (female) or slightly smaller than thorax (male); head spherical; postocular ridge and occiput extended posteriorly into slight ridge; posterior margin of eye rounded; eyes bare; position of antennae on head near middle of frons, slightly nearer to mouth-parts; eyes contiguous above antennal base, not contiguous below; palpus present; proboscis longer than head; antennal flagellum stylate, apex with terminal seta; thorax with postpronotal lobes enlarged, medially contiguous to form collar; subscutellum



**Figure 53.** *Panops schlingeri* sp. n., female, lateral view [700546]. Body length = 11.0 mm.



**Figure 54.** *Panops schlingeri* sp. n., female, oblique view [700547]. Body length = 11.0 mm.



**Figure 55.** *Panops schlingeri* sp. n., female, anterior view [700548]. Body length = 11.0 mm.

enlarged; legs not greatly elongated; pulvilli present; wing hyaline, markings absent; costa ending in radial field; costal margin straight in both sexes; humeral crossvein absent; radial veins meeting wing margin before wing apex;  $R_1$  inflated distally at pterostigma;  $R_{2+3}$  present;  $R_{4+5}$  slightly curved anteriorly midway; veins  $M_1$ ,  $M_2$  and  $M_3$  present; discal cell absent; medial veins reaching wing margin (or nearly so); crossvein 2r-m absent; Cu reduced, not reaching wing margin; anal lobe not enlarged; alula well developed; abdomen smooth, rounded, cylindrical in shape, similar width to thorax (male) or greatly rounded, inflated (female).



**Figure 56.** *Helle longirostris* (Hudson), male, lateral view [700556]. Body length = 5.0 mm.

**Included species.** *Schlingeriella irwini* Gillung & Winterton, 2011.

**Comments.** *Schlingeriella* is differentiated from other Philopotinae by medial veins mostly reaching the wing margin,  $R_1$  inflated apically, reduced wing venation (i.e. absence of all wing cells except cell *br*), elongate mouthparts and apilose eyes. See results of Winterton et al. (2007) for phylogenetic placement and divergence times. This genus is represented by only a single species (*S. irwini* sp. n.) from New Caledonia (France). There is dramatic sexual dimorphism in body size, with females considerably larger than the males. This genus was described by Gillung and Winterton (2011) to honour the decades of work by Evert I. Schlinger on world Acroceridae taxonomy. Evert Schlinger not only collected many of the specimens in New Caledonia, he also recognized that it represented a completely new genus of endemic spider flies.



**Figure 57.** *Helle longirostris* (Hudson), female, lateral view [700557]. Body length = 5.5 mm.

**Subfamily Acrocerinae Zetterstedt, 1837**

<http://species-id.net/wiki/Acrocerinae>

**Type genus.** *Acrocera* Meigen 1803: 266.

**Diagnosis.** Small to medium sized, densely pilose to apilose, body rarely arched; antennal flagellum stylate; postpronotal lobes widely separated, never medially contiguous; wing venation highly variable, ranging from complete with cells cu-p, bm br, d, m<sub>3</sub> and basal r<sub>4+5</sub> present, to highly reduced with few closed cells; humeral cross-vein rarely well developed; tibial apical spines absent (rarely present); larvae exclusively parasitoids of araneomorph spiders.



**Figure 58.** *Helle rufescens* Brunetti, male, lateral view [700558]. Body length = 8.5 mm.

### Australasian genera

*Ogcodes* Latreille, 1797; *Pterodontia* Gray, 1832

### *Ogcodes* Latreille, 1797

<http://species-id.net/wiki/Ogcodes>

Figs 3C, 63–64

*Ogcodes* Latreille, 1797: 154 – Schlinger 1960: 245; Schlinger & Jefferies 1989: 377. Type species, *Musca gibbosa* Linnaeus, by subsequent monotypy (Latreille 1802: 432).

*Oncodes* Meigen 1822: 99 [emendation of *Ogcodes* Latreille] – White 1914: 69; Hardy 1921: 77, 1946: 66; Paramonov 1955: 23, 1957: 531.

**Note.** Synonymy and usage restricted to Australasian region fauna only; see Schlinger (1960) for more exhaustive list.

**Diagnosis.** Body length: 3.0–5.0 mm [male], 4.0–8.0 mm [female]. Body shape not arched, colouration black, yellow or white, non-metallic; head much smaller than



**Figure 59.** *Helle rufescens* Brunetti, female, lateral view [700559]. Body length = 7.0 mm.

thorax width, shape sub-spherical; postocular ridge and occiput rounded; two or three ocelli, anterior ocellus sometimes absent; posterior margin of eye rounded; eye apilose; position of antennae on head adjacent to mouthparts; eyes contiguous above antennal base, not contiguous below antennal base; palpus absent; proboscis apparently absent; flagellum shape stylate; apex with terminal setae (or single seta); antenotum not collar-like behind head; subscutellum enlarged; tibial spines absent; pulvilli present; wing hyaline, markings absent; costa ending near wing apex, costal margin straight; humeral crossvein absent; radial veins straight;  $R_1$  inflated or not inflated distally; pterostigma and cell  $r_1$  membranous, not ribbed; only two radial veins present,  $R_{2+3}$  absent,  $R_{4+5}$  not reaching wing margin; medial vein complement with  $M_1$ ,  $M_2$  and  $M_3$  present, or two  $M$  veins present; discal cell weakly formed or absent; medial veins not reaching wing margin; cell  $m_3$  absent;  $CuA_1$  absent;  $CuA_2$  separate from  $A_1$ , ending just before wing margin; crossvein  $2r-m$  absent; wing microtrichia absent; anal lobe well developed; alula well developed; abdominal tergites smooth, rounded (rarely with tubercles in fossil species); abdomen shape greatly rounded, inflated.

**Included species.** *Ogcodes* is a distinctive and cosmopolitan genus and the most species-rich in the family. Thirty-four species in two subgenera (*Ogcodes* and *Protogcodes* Schlinger, 1960) are listed by Schlinger and Jefferies (1989) for the Australasian region.



**Figure 60.** *Schlingeriella irwini* Gillung & Winterton, male, lateral view [700560, 693079]. Body length = 2.4 mm.

**Comments.** *Ogcodes* is in need of revision and no recent keys to species have been published for the region. The most recent revision of the genus was by Schlinger (1960), but there are many undescribed species in collections and a world revision of the genus is needed. *Ogcodes* is a derived genus with a typical globose body, relatively small head and reduced wing venation. Characters which differentiate *Ogcodes* from all other Acroceridae genera include antennae proximal to mouthparts, palpi absent, proboscis very short, almost all wing cells absent or poorly formed, eyes apilose and  $R_{2+3}$  absent.

### ***Pterodontia* Gray, 1832**

<http://species-id.net/wiki/Pterodontia>

Figs 3D, 65–66

*Pterodontia* Gray, 1832: 779 – Macquart 1838: 174; Erichson 1840: 161; Walker 1855: 346; Westwood 1876: 513; White 1914: 68; Hardy 1921: 76, 1946: 66;



**Figure 61.** *Schlingeriella irwini* Gillung & Winterton, female, lateral view [700561, 693080]. Body length = 4.4 mm.

Paramonov 1957: 529; Schlinger 1959: 158. Type species: *Pterodontia flavipes* Gray, 1832: 779 by monotypy.

*Nothra* Westwood, 1876: 514 – Hardy 1921: 77, 1946: 66. Type species: *Nothra bicolor* Westwood, 1876: 514 by monotypy.

**Note.** Synonymy and usage list restricted to Australasian region fauna only.

**Diagnosis.** Body length: 3.0–7.0 mm [male], 4.0–10.0 mm [female]. Body shape not arched. Body colouration non-metallic; head much narrower than thorax width; shape nearly spherical; postocular ridge and occiput rounded; three ocelli; posterior margin of eye rounded; eye pilose (dense); antennae located adjacent to mouthparts; eyes contiguous above antennal base, not contiguous below antennal



**Figure 62.** *Schlingerella irwini* Gillung & Winterton, female, anterior view [700562]. Body length = 4.4 mm.

base; palpus absent; proboscis greatly reduced; flagellum stylate, apex with terminal setae (multiple); antenotum shape not collar-like behind head; subscutellum not enlarged, barely visible; tibial spines present; pulvilli present; wing markings absent; costa circumambient; wing costal margin straight or with anterior projection



**Figure 63.** *Ogcodes* sp., male, lateral view [700563]. Body length = 9.0 mm.

(males); humeral crossvein present or reduced; radial veins curved or angled towards wing anterior margin;  $R_1$  inflated distally at pterostigma (especially in male); pterostigma and cell  $r_1$  membranous, not ribbed;  $R_{2+3}$  present;  $R_{4+5}$  present as single vein; basal cell  $r_{4+5}$  (portion basal to bisecting  $2r-m$ ) merged with discal cell to form composite cell comprising  $d+r_{4+5}$ ; cell  $m_3$  absent; medial vein complement usually a single M vein fused with  $CuA_1$ , petiolate to margin, sometimes with second medial vein originating from cell  $d+r_{4+5}$ ;  $CuA_2$  fused to  $A_1$  before wing margin, petiolate, rarely open to wing margin; wing microtrichia absent; anal lobe well developed; alula present or absent, rarely well developed; abdominal tergites smooth, rounded; abdomen shape greatly rounded, inflated.

**Included species.** *Pterodontia davisi* Paramonov, 1957; *P. longisquama* Sabrosky, 1947; *P. mellii* Erichson 1840 (= *P. variegata* White, 1914 syn. n.).



**Figure 64.** *Ogcodes* sp., female, lateral view [700564]. Body length = 5.0 mm.

### Key to Australasian *Pterodontia* species

- 1 Thorax black, or yellow suffused with dark brown to black ventrally, abdomen yellow to red laterally on segments 2–4; mid and hind femora brown to black; lower calypter hyaline medially, relatively small ( $< \frac{1}{2}$  length of wing) (Western Australia, Tasmania, New South Wales, Queensland) (Fig. 66) ..... *P. melli* Erichson 1840
- Thorax and abdomen completely brown to black; all legs uniformly yellow to white; lower calypter relatively large ( $> \frac{1}{2}$  length of wing), uniformly brown...2
- 2 Wing brown infusate (Papua New Guinea)... *P. longisquama* Sabrosky, 1947
- Wing hyaline (Queensland, New South Wales) (Fig. 65)..... *P. davisi* Paramonov 1957



**Figure 65.** *Pterodontia davisii* Paramonov, male, dorsal view [700565]. Body length = 7.0 mm.

**Comments.** *Pterodontia* is a cosmopolitan genus containing 19 valid species, three of which are recorded from the Australasian region (Schlinger and Jefferies 1989). *Pterodontia variegata* was described by White (1914) and differentiated from *P. melli* (as *P. macquarti* Westwood, 1848) based on colouration of the fore femur, scutellum and abdomen. Paramonov (1957) examined a range of specimens from various localities and suggested that the former was likely a synonym of the latter. Based on examinations of these and additional specimens this synonymy is supported herein.

Some species of *Pterodontia* have greatly enlarged and sclerotized lower calypters, appearing somewhat like a second pair of wings (e.g. *P. davisii*). Males in this genus typically have sclerotized projections on the costal margin of the wing. Characteristics which diagnose this genus from other acrocerids include head very small relative to thorax width, tibial spines present, cells  $m_3$ , d and basal  $r_{4+5}$  fused to form a single cell, eyes densely pilose, antennae adjacent to the ocellar tubercle and mouthparts reduced.



**Figure 66.** *Pterodontia mellii* Erichson, female, lateral view [700566]. Body length = 11.0 mm.

Contrary to other authors, *Pterodontia* has been placed previously in Panopinae by Schlinger (1981, 1987, 1989) based on the presence of tibial spines. The wing venation of *Pterodontia* is unique among acrocerids.

## Acknowledgements

Thank you to Greg Daniels (Brisbane, Australia), Chris Grinter and Norman Penny (CAS), Erica McAllister (BMNH), Terry Houston (WAM), Ken Walker (NMV), Peter Hudson (South Australian Museum) and David Yeates (Australian National Insect Collection) for loan of specimens. Thank you to Evert Schlinger and F. Christian Thompson for access to unpublished bibliographical and nomenclatural information on Acroceridae that were used in part to compile lists of nomenclatural acts and usage. Much of this research was based on years of dedicated work, both published and unpublished, by E.I. Schlinger on Acroceridae of the world; I am grateful for the opportunity for consolidate some of this work and publish some of the results herein. Thank you also to Dan Schoknecht (WAM) and Terry Houston for providing the image of the living *Panops baudini* and identification of the host plant, respectively. Thank you to Jéssica Gillung, Torsten Dikow and Bradley Sinclair for their comments on the draft manuscript. Thank

you also to F. Christian Thompson for providing his expertise on the etiquette of not publishing. This research was supported by the Australian Biological Resource Study (ABRS-209-48). Statements and viewpoints expressed herein do not necessarily reflect the opinion of ABRS.

## References

- Bequaert J (1931) The genus *Lasia* (Diptera, Cyrtidae) in North America, with descriptions of two new species. *American Museum Novitates*. 455: 1–11.
- Bigot JMF (1890) Dipteres nouveaux ou peu connus. 35e partie (1). XLIII: Cyrtidi (J. Bigot). *Ann. Soc. Ent. Fr.* 9: 313–320.
- Blanchard CE (1840) Histoire naturelle des insectes. Orthoptères, névroptères, hémiptères, hyménoptères, lépidoptères et diptères. Vol. 3, 672 pp., 67 pls. In: Laporte FL de. Histoire naturelle des animaux articulés. Paris. [Acroceridae, p. 584]
- Brunetti E (1926) New and little-known Cyrtidae (Diptera). *Ann. Mag. Nat. Hist.* 18: 561–606. doi: 10.1080/00222932608633552
- Edwards FW (1930) Bombyliidae, Nemenstrinidae, Cyrtidae. *Dipt. Patagonia S. Chile* 5: 166–197.
- Erichson WF (1840) Die Henopier, Eine familie aus der Ordnung der Dipteren, 135–180, pl. 1, Figs 7–10.
- Gillung J, Winterton S (2011) New genera of philopotine spider flies (Diptera, Acroceridae) with a key to living and fossil genera. *ZooKeys* 127: 15–27. doi: 10.3897/zookeys.127.1824
- Gray GR (1832) [New genera and species.] In: Griffith E, Pidgeon E. The class Insecta arranged by Baron Cuvier, with supplementary additions to each order, and notices of new genera and species by George Gray, Esq. Vol. 2. Whittaker, Treacher & Co., London, 796 pp.
- Hardy GH (1918) Notes on Tasmanian Diptera and descriptions of new species. *Pap. Proc. R. Soc. Tasmania* 1917: 60–66.
- Hardy GH (1921) Australian Bombyliidae and Cyrtidae. *Pap. Proc. R. Soc. Tasmania* 1921, 41–83.
- Hardy GH (1946) Miscellaneous notes on Australian Diptera. XII. Cyrtidae, Dolichopodidae and Phoridae. *Proc. Linn. Soc. New South Wales* 71: 65–71.
- Hudson GV (1892) An elementary manual of New Zealand Entomology. Being an introduction to the study of our native insects. West, Newman & Co., London.
- Hutton FW (1901) Synopsis of the Diptera Brachyera of New Zealand. *Trans. Proc. N. Z. Inst.* 33: 1–95.
- Kerr PH, Winterton SL (2008) Do parasitic flies attack mites? Evidence in Baltic amber. *Biological Journal of the Linnean Society* 93: 9–13. doi: 10.1111/j.1095-8312.2007.00935.x
- Kertész K (1909) *Catalogus Dipteriorum hucusque descriptorum*. Vol. 4, Oncodidae, Nemenstrinidae, Mydidae, Apioceridae, Asilidae. *Museum Nationale Hungaricum, Budapestini* [=Budepest], 349 pp.
- Lamarck JBPA (1804) Sur deux nouveaux genres d'insectes de la Nouvelle-Hollande. *Ann. Mus. Hist. Nat. Paris* 3: 260–265.

- Lamarck JBPA (1812) Extrait du cours de Zoologie du Muséum d'Histoire naturelle, sur les Animaux sans vertèbres; présentant la distribution et la classification de ces animaux, les caractères des principales divisions, et une simple liste des genres; à l'usage de ceux qui suivent ce cours. d'Hautel & Gabon, Paris, 127 pp.
- Latreille PA (1802) Histoire naturelle, générale et particulière, des crustacés et des insectes. Tome troisième. Familles naturelles des genres. Dufart, Paris, xii + 13–467 pp.
- Latreille PA (1804) Tableau méthodique des insectes. In: Société de Naturalistes de d'Agriculteurs, Nouveau dictionnaire d'histoire naturelle ... Tome 24 [Section 3]: Tableaux méthodique d'histoire naturelle. 84 + 4 + 85 + 238 + 18 + 34 pp. Déterville, Paris, 129–200.
- Latreille PA (1809) Genera crustaceorum et insectorum ... Tomus quartus et ultimas. A. Koenig, Parisiis et Argentorati [= Paris & Strasbourg], 399 pp.
- Latreille PA (1810) Considerations generales sur l'ordre naturel des animaux ... F. Schoell, Paris, 444 pp.
- Latreille PA (1811) Panops. In: Encyclopedie methodique. Historie naturelle, Insects. Vol. 8, 722 pp. H. Agasse, Paris, 707–710.
- Latreille PA (1816) Les crustacés, les arachnides et les insectes. In: Cuvier, [G.C.L.F.D.], Le règne animal. Tome III. Déterville, Paris, xxix + 653 pp.
- Latreille PA (1825) Familles naturelles du règne animal ... J.-B. Baillière, Paris, 570 pp.
- Latreille PA (1829) Suite et fin des insectes. In: Cuvier, [G. L. C. F. D.], Le règne animal. Tome V. Déterville et Crochard, Paris, xxiv + 556 pp.
- Macquart J (1838) Diptères exotiques nouveaux ou peu connus. Tome premier. 2e partie. Roret, Paris, 207 pp., 14 pls.
- Macquart J (1850) Diptères exotiques nouveaux ou peu connus. Supplement. [4]. Mem. Soc. R. Sci. Agric. Arts, Lille 1850: 5–160, 14 pls.
- McAlpine JF (1981) Morphology and terminology-Adults, In: McAlpine JF, Peterson BV, Shewell GE, Teskey HJ, Vockeroth JR, Wood DM (Eds) Manual of Nearctic Diptera. Research Branch, Agriculture Canada Monograph 1: 9–63.
- Meigen JW (1822) Systematische Beschreibung der bekannten europäischen zweiflügeligen Insekten. Dritter Theil. Schulz-Wundermann, Hamm. x + 416 pp
- Neboiss A (1971) Australian Panopinae (Diptera: Acroceridae). Journal of the Australian Entomological Society 10: 205–222. doi: 10.1111/j.1440-6055.1971.tb00031.x
- Osten Sacken CR (1896) A new genus of Cyrtidae (Dipt.) from New Zealand. Ent. mon. mag. 7: 16–18.
- Pape T, Thompson FC [editors] (2010) Systema Dipterorum, Version 1.0. <http://www.diptera.org/> [accessed on 23 August 2010]
- Paramonov SJ (1955) New Zealand Cyrtidae (Diptera) and the problem of the Pacific island fauna. Pacific Science 9: 16–25.
- Paramonov SJ (1957) A Review of the Australian Acroceridae (Diptera). Australian Journal of Zoology 5: 521–546. doi: 10.1071/ZO9570521
- Pyle RL, Michel E (2008) Zoobank: Developing and nomenclatural tool for unifying 250 years of biological information. Zootaxa 1950: 39–50.

- Rondani C (1863) *Diptera exotica revisa et annotata. Novis non nullis descriptis*. E. Soliani, Modena, 99 pp.
- Sabrosky CW (1947) A new Pterodontia from New Guinea (Diptera, Acroceridae). *Pan-Pacific Entomologist* 23: 74–75.
- Schiner IR (1868) Diptera. In: Wullerstorff-Urbair B von (Ed) *Reise der oesterreichischen Fregatte Novara*. Zool. 2(1) B. K. Gerold's Sohn, Wien, vi + 388 pp., 4 pls.
- Schlenger EI (1959) A review of the genus *Rhysogaster* Aldrich, with descriptions of new genera and new species of Oriental, Ethiopian and Australian Acroceridae (Diptera). *Annals of the Entomological Society of America* 52: 150–159.
- Schlenger EI (1960) A revision of the genus *Ogcodes* Latreille, with particular reference to species of the Western Hemisphere. *Proc. U. S. Nat. Mus.* 111: 227–336.
- Schlenger EI (1966) Distributional patterns of selected western North American Insects. An analysis of the distribution of *Eulonchus* Gerstaecker (Diptera: Acroceridae). *Bull. Ent. Soc. Amer.* 12: 112–113.
- Schlenger EI (1981) Acroceridae. In: McAlpine JF, Peterson BV, Shewell GE, Teskey HJ, Vockeroth JR, Wood DEM (Eds) *Manual of Nearctic Diptera*. vol. I. Research Branch, Agriculture Canada. Monograph 27: 575–584.
- Schlenger EI (1987) The biology of Acroceridae (Diptera): True endoparasitoids of spiders. Pp. 319–327. In Nentwig W (Ed) *Ecophysiology of spiders*. Springer Verlag, Berlin, xi + 448 pp.
- Schlenger EI (2009) Acroceridae (spider flies, small-headed flies). pp. 551–556 In Brown B.V. et al. (Eds) *Manual of Central American Diptera: Volume 1*. NRC Research Press, Ottawa, Ontario, Canada, 714 pp.
- Schlenger EI, Jefferies MG (1989) Family Acroceridae. In: Evenhuis NL (Ed) *Catalog of Diptera of the Australasian and Oceanian regions*. Bishop Museum Special Publication. Bishop Museum Press, 86: 375–37.
- Thomson CG (1869) Diptera. Species nova descriptis. In: *Kongliga svenska fregatten Eugenies resa omkring jorden under befäl af C. A. Virgin, åren 1851–1853*. 2 (Zoologi) 1, Insecta. P. A. Norstedt & Söner, Stockholm, 443–614.
- Walker F (1855) List of the specimens of dipterous insects in the collection of the British Museum. Part VI. Supplement II. British Museum, London, 331–506.
- Westwood JO (1876) *Notae Dipterologicae*. No. 3. – Descriptions of new genera and species of the family Acroceridae. *Trans. Ent. Soc. London* 1876, 507–518.
- White A (1914) The Diptera-Brachycera of Tasmania. Part I. Families Leptidae, Stratiomyidae, Nemestrinidae, & Cyrtidae. *Pap. Proc. R. Soc. Tasmania* 1914, 35–74.
- Wiedemann CRW (1830) *Aussereuropäische zweiflügelige Insekten*. Zweiter Theil. Schulz, Hamm. xii + 684 pp.
- Winterton SL, Wiegmann BM, Schlenger EI (2007) Phylogeny and Bayesian divergence time estimations of small-headed flies (Diptera: Acroceridae) using multiple molecular markers. *Molecular Phylogenetics and Evolution* 43: 808–832.