

Two new spider species of the genus *Chryso* O. P.- Cambridge, 1882 (Araneae, Theridiidae) in Hainan Island, China

Bao-Shi Zhang^{1†}, Feng Zhang^{2‡}

1 Baoding University, Baoding, 071051, P. R. China **2** College of Life Sciences, Hebei University, Baoding, Hebei 071002, P. R. China

† [urn:lsid:zoobank.org:author:484F2B50-D862-476D-9651-B09A81534ECC](https://doi.org/urn:lsid:zoobank.org:author:484F2B50-D862-476D-9651-B09A81534ECC)

‡ [urn:lsid:zoobank.org:author:904C4445-AAEA-40BC-B418-502053EF39B8](https://doi.org/urn:lsid:zoobank.org:author:904C4445-AAEA-40BC-B418-502053EF39B8)

Corresponding author: Feng Zhang (dudu06042001@163.com)

Academic editor: D. Dimitrov | Received 12 february 2012 | Accepted 27 April 2012 | Published 4 May 2012

[urn:lsid:zoobank.org:pub:DE2AD1BC-7A09-4B30-BED0-D5A868EC1EA8](https://doi.org/urn:lsid:zoobank.org:pub:DE2AD1BC-7A09-4B30-BED0-D5A868EC1EA8)

Citation: Zhang B-S, Zhang F (2012) Two new spider species of the genus *Chryso* O. P.-Cambridge, 1882 (Araneae, Theridiidae) in Hainan Island, China. ZooKeys 190: 21–31. doi: 10.3897/zookeys.190.2924

Abstract

Two new spider species of the genus *Chryso* O. P.-Cambridge, 1882 are reported from Hainan Island, China, *Chryso bifurca* **sp. n.** (male, female) and *C. bicuspadata* **sp. n.** (male, female). *C. bimaculata* Yoshida, 1998 is recorded from China for the first time.

Keywords

Chryso, taxonomy, new species, newly recorded, China

Introduction

The genus *Chryso* was erected by O. P.-Cambridge (1882). It was regarded as a junior synonymy of *Theridion* Walckenaer, 1805, and removed by Bryant (1940). Levi and Levi (1962) considered this genus as a senior synonym of *Arctachaea* Levi, 1957, *Argyroaster* Yaginuma, 1958 and *Meotipa* Simon, 1894, but Deeleman-Reinhold (2009) removed the genus *Meotipa* from the synonymy of *Chryso*. Until now no large revision of this genus has been done. However, Agnarsson (2004) listed 10 autapomorphies of the *Chryso* in his phylogeny, these characteristics are putative *Chryso* synapomorphies.

Currently 62 *Chryso* species are reported, mostly from America and Asia (Levi 1957; Levi and Levi 1962; Barrion and Litsinger 1995; Miller and Agnarsson 2005; Gonzaga et al. 2006; Yoshida 2009; Siliwal 2009; Platnick 2012), among them, 22 species are known from China. 94 species out of 24 genera are known from Hainan Island, China (Zhu 1998; Song et al. 1999; Yoshida et al. 2000; Tang et al. 2003; Song et al. 2006), including six *Chryso* species: *C. cyclocera* Zhu, 1998, *C. trispinula* Zhu, 1998, *C. trimaculata* Zhu, Zhang & Xu, 1991, *C. scintillans* (Thorell, 1895), *C. pulcherrima* (Mello-Leitão, 1917) and *C. nigra* (O. P.-Cambridge, 1880).

During the examination of spider specimens collected from 2007 to 2009 in Hainan Island, China, two new species, *Chryso bifurca* sp. n. and *C. bicuspidata* sp. n. were recognized and are described here. *C. bimaculata* Yoshida, 1998, known from Japan previously, is newly recorded from Hainan, China.

Material and methods

All specimens were kept in 75% ethanol and examined, drawn and measured under a Tech XTL-II stereomicroscope equipped with an Abbe drawing device. Carapace length was measured medially from the anterior margin to the rear margin of the carapace. Eye sizes were measured as the maximum diameter of the lens in dorsal or frontal view. MOA length was measured medially from the anterior margin to the rear margin of MOA. Leg measurements are given as: total length (femur, patella, tibia, metatarsus, tarsus). Epigynes were cleared in warm potassium hydroxide (KOH) and transferred to 75% ethanol for drawing. The labeling of the palpal sclerites is adopted following Agnarsson (2004). All measurements are in millimeters. All specimens studied are deposited in the Museum of Hebei University (MHBUS), Baoding, China.

Abbreviations

AER	anterior eye row
ALE	anterior lateral eyes
AME	anterior median eyes
C	conductor
CD	copulatory ducts
E	embolus
FD	fertilization ducts
MA	median apophysis
MOA	median ocular area
PER	posterior eye row
PLE	posterior lateral eyes
PME	posterior median eyes
S	spermathecae

ST subtegulum
T tegulum
TTA theridiid tegular apophysis

Taxonomy

Chryso bifurca sp. n.

urn:lsid:zoobank.org:act:5B93397C-1884-4E05-9738-5B5A003645A6

http://species-id.net/wiki/Chryso_bifurca

Figs 1–6

Type material. **Holotype** ♂, **CHINA, Hainan Island:** Jianfengling Mountain (19°07'N, 109°13'E), 29 May 2009, C. Zhang leg. **Paratypes:** 2 ♀, same data as holotype; 1 ♀, Limu Mountain (19°10'N, 109°39'E), 20 August 2007, C. Zhang leg.; 3 ♂, 12 ♀, Limu Mountain, 20 November 2008, G.X. Han leg.


Diagnosis. Males can be distinguished from other *Chryso* species by the following characters: apex of cymbium bifurcate; embolus short and thin; theridiid tegular apophysis erect, with obtuse apex (Figs 3–4). Females differ from other *Chryso* species by the bigger and oval spermathecae, and the longer and winding copulatory ducts, differ from *C. octomaculata* (Bösenberg & Strand, 1906) by the process of copulatory ducts (Fig. 6).

Etymology. The species name refers to bifurcate apex of cymbium.

Description. Male (holotype). Total length 2.70: cephalothorax 1.26 long, 1.08 wide; abdomen 1.44 long, 0.99 wide, 0.90 high. Carapace yellow, longer than wide, lateral margin with thin black striations. Only anterior part with several gray setae. Cervical groove distinct. Median furrow almost triangle, yellowish. Each eye with a black ring. Both the eye rows recurved from dorsal view (Fig. 1). Diameters of eyes: AME 0.12, ALE 0.10, PME 0.10, PLE 0.10. Interdistances of eyes: AME–AME 0.10, AME–ALE 0.08, ALE–ALE 0.64, PME–PME 0.10, PME–PLE 0.14, PLE–PLE 0.66, ALE and PLE closed to each other. MOA long 0.29, front width 0.34, back width 0.31. Clypeus 0.32 high, yellow and furnished with few short gray setae. Chelicerae armed with sparse gray setae, promargin with 2 teeth, fangs short and thin. Anterior margin of endites armed with gray scopula. Labium armed with sparse long black setae. Sternum furnished with sparse black setae, its anterior margin lightly procurved. Legs yellow, patella, metatarsus and the base of femur and tibia reddish-brown. Measurements of legs: leg I 9.44 (2.52, 0.59, 2.55, 2.88, 0.90), II 4.38 (1.49, 0.45, 1.13, 0.90, 0.41), III 3.16 (0.90, 0.36, 0.59, 0.90, 0.41), IV 5.09 (1.58, 0.36, 1.17, 1.44, 0.54). Leg formula: 1423.

Abdomen oval, longer than wide and armed with brown setae. Dorsum yellowish, armed with six pairs of irregular black patches, posterior with a median black patch (Figs 1–2). Venter yellowish, with a black median patch. Anal tubercle yellow. Spinnerets yellow.

Male palp with long cymbium, apical part of cymbium bifurcate; embolus short, base with a lunate process, distal part thin; theridiid tegular apophysis erect, with wide base, apex thin and obtuse; conductor lying behind theridiid tegular apophysis (Figs 3–4).



Figures 1–6. *Chryso bifurca* sp. n., 1–4 male holotype 1 body, dorsal view 2 abdomen, lateral view 3 male left palp, ventral view 4 same, prolateral view 5–6 female paratype 5 epigynum, ventral view 6 vulva, dorsal view. Scale bars: 0.5 mm (1–2); 0.1 mm (3–6).

Female (one paratype from Jianfengling Mountain) total length 3.33: cephalothorax 0.95 long, 0.94 wide; abdomen 2.34 long, 1.62 wide, 1.58 high. Diameters of eyes: AME 0.12, ALE 0.10, PME 0.10, PLE 0.10. Interdistances of eyes: AME–AME 0.13, AME–ALE 0.09, ALE–ALE 0.59, PME–PME 0.14, PME–PLE 0.16, PLE–PLE 0.61, ALE and PLE closed to each other. MOA long 0.27, front width 0.30, back width 0.31. Clypeus 0.32 high. Measurements of legs: leg I 10.81 (3.24, 0.63, 2.70, 3.29, 0.95), II 6.44 (2.03, 0.54, 1.44, 1.80, 0.63), III 3.52 (1.13, 0.41, 0.63, 0.90, 0.45), IV 7.08 (1.94, 0.54, 2.21, 1.80, 0.59). Leg formula: 1423. Other characters as in holotype.

Female genitalia lightly sclerotized, posterior part with a kidney-shaped atrium; spermathecae big, oval; copulatory ducts long, thick, winding and connected with spermathecae from posterior part (Figs 5–6).

The new species with putative *Chryso* synapomorphies as follow: (1) carapace pars stridens irregular; (2) abdomen extending beyond spinnerets; (3) cymbial hood of male palp broad; (4) median apophysis of male palp with distinct apophysis; (5) palpal claw of female dentition sparse.

Variation. The lateral part of dorsal abdomen of some females examined with eight to eleven pairs of irregular black patches. Males total body length from 2.64–2.80, female total length from 3.28–3.35.

Distribution. China (Hainan).

***Chryso bicuspadata* sp. n.**

urn:lsid:zoobank.org:act:660AA46B-0F72-4257-A1FC-8B734804876B


http://species-id.net/wiki/Chryso_bicuspadata

Figs 7–12

Type material. Holotype ♂, CHINA, Hainan Island: Jianfengling Mountain, 29 May 2009, C. Zhang leg. **Paratypes:** 2 ♀, same data as holotype; 3 ♀, Jianfengling Mountain, 12 November 2008, G.X. Han leg.; 3 ♂, 1 ♀, Bawangling Mountain (19°07'N, 109°04'E), 25 May 2009, C. Zhang leg.; 5 ♀, Bawangling Mountain, 7 November 2008, G.X. Han leg.; 2 ♂, 1 ♀, Diaoluo Mountain (18°45'N, 109°45'E), 6 June 2009, C. Zhang leg.; 1 ♂, 2 ♀, Limu Mountain, 19 August 2007, F. Zhang leg.

Diagnosis. Males can be distinguished from most *Chryso* species by membranous conductor, wider embolus and acute apex of cymbium. And it resembles *C. cyclocera* Zhu, 1998 and *C. oxycera* Zhu & Song, 1993 in the general shape of the palpal organ, but differs from them by the wider embolus (Figs 8–9). Females differ from all other *Chryso* species except *C. viridiventris* Yoshida, 1996 by the epigynum with a big atrium and a large posterior lobe. It differs from *C. viridiventris* by the bigger spermathecae and shorter copulatory ducts (Figs 10–12).

Etymology. The species name refers to the shape of embolic tip.



Figures 7–12. *Chryso bicuspidata* sp. n., **7–9** female holotype **7** body, dorsal view **8** male left palp, ventral view **9** same, prolateral view **10–12** female paratype **10** epigynum, lateral view **11** same, ventral view **12** vulva, dorsal view. Scale bars: 0.5 mm (7); 0.1 mm (8–12).

Description. Male (holotype). Total length 1.84: cephalothorax 0.92 long, 0.71 wide; abdomen 0.92 long, 0.82 wide, 0.92 high. Carapace longer than wide, yellowish, lateral margin with thin black striations, anterior median with a triangular black patch. Only anterior part with several gray setae. Cervical groove yellowish. Median furrow yellowish, almost triangle. Each eye with a black ring. Both the eye rows recurved from dorsal view (Fig. 7). Diameters of eyes: AME 0.05, ALE 0.04, PME 0.04, PLE 0.04. Interdistances of eyes: AME–AME 0.10, AME–ALE 0.03, ALE–ALE 0.33, PME–PME 0.10, PME–PLE 0.08, PLE–PLE 0.38, ALE and PLE closed to each other. MOA long 0.13, front width 0.18, back width 0.18. Clypeus 0.20 high and furnished with few short gray setae. Chelicerae armed with sparse gray setae, promargin with 2 teeth, fangs short and thin. Endites, labium and sternum yellowish. Anterior margin of endites armed with gray scopula. Sternum furnished with sparse gray setae. Legs yellowish, the end of tibia with gray spots. Measurements of legs: leg I 7.14 (1.94, 0.41, 1.94, 1.63, 1.22), II 3.69 (1.33, 0.31, 0.82, 0.82, 0.41), III 2.14 (0.71, 0.20, 0.31, 0.61, 0.31), IV 3.86 (1.12, 0.31, 0.82, 1.20, 0.41). Leg formula: 1423.

Abdomen oval, longer than wide and armed with brown setae. Dorsum yellowish, anterior part with a pair of black patches (Fig. 7). Venter yellowish. Spinnerets yellowish.

Apical cymbium of male palp acute; embolus big, thick, and end with a thin ramus; conductor membranous, falciform from ventral view; apex of median apophysis with some small tubers (Figs 8–9).

Female (one paratype from Limu Mountain) total length 2.32: cephalothorax 0.92 long, 0.71 wide; abdomen 1.43 long, 1.22 wide, 1.43 high. Diameters of eyes: AME 0.05, ALE 0.05, PME 0.03, PLE 0.05. Interdistances of eyes: AME–AME 0.05, AME–ALE 0.02, ALE–ALE 0.30, PME–PME 0.09, PME–PLE 0.08, PLE–PLE 0.33, ALE and PLE closed to each other. MOA long 0.13, front width 0.15, back width 0.15. Clypeus 0.30 high. Measurements of legs: leg I 7.75 (2.24, 0.51, 1.94, 2.35, 0.71), II 3.59 (1.24, 0.31, 0.71, 0.92, 0.41), III 2.04 (0.71, 0.20, 0.31, 0.51, 0.31), IV 3.79 (1.24, 0.31, 0.81, 1.02, 0.41). Leg formula: 1423. Dorsal abdomen yellowish. Other characters as in holotype.

Female genitalia lightly sclerotized, and with a circular atrium and a large posterior lobe; spermathecae big, oval; copulatory ducts short, thick and connected with spermathecae from anterior part; each fertilization duct with a global head (Figs 10–12).

The new species with putative *Chryso* synapomorphies as follow: (1) carapace pars stridens smooth; (2) abdomen extending beyond spinnerets; (3) cymbial hood of male palp broad; (4) subconductor of male palp present; (5) median apophysis of male palp with apophysis; (6) palpal claw of female dentition sparse; (7) anterior margin of female genital atrium medially acute.

Variation. The anterior part of dorsal abdomen of some species examined with a pair of black patches, some species without. Males total body length from 1.73–1.88, females total length from 2.22–2.35.

Distribution. China (Hainan).

***Chryso bimaculata* Yoshida, 1998**

http://species-id.net/wiki/Chryso_bimaculata

Figs 13–17

Chryso bimaculata Yoshida, 1998: 105, f. 1–6; Yoshida 2003: 125, f. 330–335; Yoshida 2009: 378, f. 203–204.

Material examined. CHINA, Hainan Island: 1 ♂, 1 ♀, Jianfengling Mountain, 31 May 2009, C. Zhang leg.; 1 ♀, Bawangling Mountain, 6 November 2008, G.X. Han leg.; 1 ♀, Bawangling Mountain, 25 May 2009, C. Zhang leg.; 5 ♀, Limu Mountain, 29 August 2007, G.X. Han leg.; 1 ♂, 3 ♀, Limu Mountain, 21 November 2008, G.X. Han leg.

Description. Male (one specimen from Jianfengling Mountain) total length 1.40: cephalothorax 0.59 long, 0.54 wide; abdomen 0.81 long, 0.59 wide, 0.54 high. Carapace longer than wide, yellowish. Cervical groove yellowish. Median furrow yellowish and almost triangle. Each eye with a red ring. AER recurved and PER procurved from dorsal view (Fig. 13). Diameters of eyes: AME 0.08, ALE 0.07, PME 0.07, PLE 0.07. Interdistances of eyes: AME–AME 0.08, AME–ALE 0.07, ALE–ALE 0.34, PME–PME 0.08, PME–PLE 0.06, PLE–PLE 0.36, ALE and PLE closed to each other. MOA long 0.18, front width 0.18, back width 0.17. Clypeus 0.16 high and furnished with few short gray setae. Chelicerae armed with sparse black setae, promargin with 2 teeth. Endites, labium and sternum yellowish. Anterior margin of endites armed with gray scopula. Sternum furnished with sparse black setae. Legs yellow, the end of tibia yellow brown. Measurements of legs: leg I 5.10 (1.44, 0.36, 1.22, 1.58, 0.50), II 3.07 (0.99, 0.32, 0.63, 0.77, 0.36), III 1.81 (0.50, 0.23, 0.45, 0.36, 0.27), IV 3.20 (1.17, 0.27, 0.63, 0.77, 0.36). Leg formula: 1423.

Abdomen oval, armed with gray setae. Dorsum yellowish, posterior part with two pairs of white patches, lateral part armed with three pairs of black patches and several black stripes (Fig. 13). Venter yellowish. Spinnerets yellowish.


Embolus of male palp small, thin and wind; conductor wind, membranous; median apophysis big, posterior part with a small tuber from prolateral view (Figs 14–15).

Female (one specimen from Jianfengling Mountain) total length 1.80: cephalothorax 0.63 long, 0.54 wide; abdomen 1.08 long, 0.95 wide, 1.04 high. Diameters of eyes: AME 0.08, ALE 0.07, PME 0.07, PLE 0.07. Interdistances of eyes: AME–AME 0.09, AME–ALE 0.03, ALE–ALE 0.35, PME–PME 0.10, PME–PLE 0.07, PLE–PLE 0.38, ALE and PLE closed to each other. MOA long 0.17, front width 0.17, back width 0.20. Clypeus 0.16 high. Measurements of legs: leg I 4.73 (1.35, 0.27, 1.17, 1.44, 0.50), II 2.81 (0.90, 0.23, 0.59, 0.77, 0.32), III 1.95 (0.63, 0.18, 0.32, 0.50, 0.32), IV 2.94 (0.95, 0.27, 0.59, 0.77, 0.36). Leg formula: 1423. Other characters as in holotype.

Female genitalia lightly sclerotized, median part with a circular atrium; spermathecae oval; the copulatory ducts very short (Figs 16–17).

Variation. The amount of white patches and black patches of dorsal abdomen varied from two pairs to three pairs. Males total body length of some species examined from 1.40–1.55, females total length from 1.68–1.82.

Distribution. China (Hainan), Japan.



Figures 13–17. *Chryso bimaculata* Yoshida, 1998 **13–15** male **13** body, dorsal view **14** male left palp, ventral view **15** same, prolateral view **16–17** female **16** epigynum, ventral view **17** vulva, dorsal view. Scale bars: 0.5 mm (**13**); 0.1 mm (**14–17**).

Acknowledgements

We are grateful to Dr. Dimitar Dimitrov, Dr. Ingi Agnarsson and an anonymous referee for valuable suggestions. We are also grateful to Chao Zhang and Guangxin Han for the collection of specimens. This study was supported by the Ministry of Science and Technology of the People's Republic of China (2006FY110500), the National Natural Science Foundation of China (No. 31071885, 30970325, 31093430), and partly by the Education Office of Hebei Province, China (No. Z2009101), and the Natural Science Foundation of Baoding University (No. 2010Z01).

References

- Agnarsson I (2004) Morphological phylogeny of cobweb spiders and their relatives (Araneae, Araneioidea, Theridiidae). *Zoological Journal of the Linnean Society* 141: 447–626. doi: 10.1111/j.1096-3642.2004.00120.x
- Barrion AT, Litsinger JA (1995) *Riceland Spiders of South and Southeast Asia*. CAB International, Wallingford, UK, 700 pp.
- Bryant EB (1940) Cuban spiders in the Museum of Comparative Zoology. *Bulletin of the Museum of Comparative Zoology* 86: 247–554.
- Cambridge OP (1882) On new genera and species of Araneidea. *Proceedings of the Zoological Society of London*, 423–442.
- Deeleman-Reinhold CL (2009) Spiny theridiids in the Asian tropics. Systematics, notes on behaviour and species richness (Araneae: Theridiidae: *Chryso*, *Meotipa*). *Contributions to Natural History* 12: 403–436.
- Gonzaga MO, Leiner NO, Santos AJ (2006) On the sticky cobwebs of two theriid spiders (Araneae: Theridiidae). *Journal of Natural History* 40: 293–306. doi: 10.1080/00222930600703433
- Levi HW (1957) The spider genera *Chryso* and *Tidarren* in America. *Journal of the New York Entomological Society* 63: 59–81.
- Levi HW, Levi LR (1962) The genera of the spider family Theridiidae. *Bulletin of the Museum of Comparative Zoology at Harvard University* 127: 1–71.
- Miller J, Agnarsson I (2005) A redescription of *Chryso nigriceps* (Araneae, Theridiidae) with evidence for maternal care. *The Journal of Arachnology* 33: 711–714. doi: 10.1636/H04-10.1
- Platnick NI (2012) The world spider catalog, version 12.5. American Museum of Natural History, online at <http://research.amnh.org/entomology/spiders/catalog/index.html> [accessed 31 December, 2011]
- Siliwal M (2009) A new species of the genus *Chryso* O. P.-Cambridge (Araneae: Theridiidae) from Arunachal Pradesh, India. *Biosystematica* 3: 5–9.
- Song DX, Zhang JX, Zhu MS (2006) Araneae. In: Li ZZ, Jin DC (Eds) *Insects from Fanjingshan Landscape*. Guizhou Science and Technology Publishing House, Guiyang, 656–690.
- Song DX, Zhu MS, Chen J (1999) *The Spiders of China*. Hebei Science & Technology Publishing House, Shijiazhuang, 640 pp.

- Tang G, Yin CM, Peng XJ (2003) Two new species of the genus *Chryso* from Hunan Province, China (Araneae, Theridiidae). *Acta zootaxonomica Sinica* 28: 53–55.
- Yoshida H (2009) Uloboridae, Theridiidae, Ctenidae. In: Ono H (Ed) *The Spiders of Japan with keys to the families and genera and illustrations of the species*. Tokai University Press, Kanagawa, 142–147, 356–393, 467–468.
- Yoshida H, Tso IM, Severinghaus LL (2000) The spider family Theridiidae (Arachnida: Araneae) from Orchid Island, Taiwan: Descriptions of six new and one newly recorded species. *Zoological Studies* 39: 123–132.
- Zhu MS (1998) *Fauna Sinica: Arachnida: Araneae: Theridiidae*. Science Press, Beijing, 436 pp.