

Two new species of the genus *Abrus* Dai & Zhang, 2002 (Hemiptera, Cicadellidae, Deltocephalinae) from China

Jichun Xing¹, Zizhong Li¹

¹ Institute of Entomology, Guizhou University; Guizhou Key Laboratory for Plant Pest Management of Mountainous Region, Guizhou University, Guiyang, Guizhou Province, 550025, China.

Corresponding author: Zizhong Li (lizizhong38@163.com)

Academic editor: A. Sanborn | Received 12 March 2014 | Accepted 10 June 2014 | Published 24 June 2014

<http://zoobank.org/B6F17820-8C84-47F0-9D3F-96627F11B8F0>

Citation: Xing J, Li Z (2014) Two new species of the genus *Abrus* Dai & Zhang, 2002 (Hemiptera, Cicadellidae, Deltocephalinae) from China. ZooKeys 419: 103–109. doi: 10.3897/zookeys.419.7481

Abstract

Two new species of leafhoppers: *Abrus damingshanensis* **sp. n.** (from Guangxi) and *A. expansivus* **sp. n.** (from Guizhou) are described and illustrated from China. A map showing the geographic distribution of the two new species is given. Taxonomic notes on species of the genus *Abrus* is also provided.

Keywords

Homoptera, leafhopper, morphology, taxonomy, distribution

Introduction

The genus *Abrus* belonging to the tribe Athysanini of subfamily Deltocephalinae, was established by Dai and Zhang (2002) with six species: *A. hengshanensis*, *A. brevis*, *A. huangi*, *A. wuyiensis*, *A. bifurcatus* and *A. coneus* from China and with *A. hengshanensis* as its type species. It belongs to the tribe Athysanini of subfamily Deltocephalinae (Hemiptera: Cicadellidae). Later, Li and Wang (2006) described two new species: *A. concavelus* and *A. leigongshanensis*. Dai and Zhang (2008) reviewed this genus

and added a new species *A. breviolus*. Recently, Li (in Li et al. 2011) described two new species: *A. biprocessus* and *A. graciaedeagus*, and recorded *A. brevis*, *A. coneus* and *A. leigongshanensis* feeding on bamboo. Chen, Yang and Li (2012) described four new species, namely, *A. anlongensis*, *A. bambusanus*, *A. daozenensis* and *A. yunshanensis*. Yang and Chen (2013) described two new species: *A. xishuiensis* and *A. langshanensis*, and provided a key to 13 known species. Moreover, *A. brevis*, *A. coneus*, *A. leigongshanensis*, *A. anlongensis*, *A. bambusanus*, *A. daozenensis*, *A. yunshanensis*, *A. xishuiensis* and *A. langshanensis* were recorded that they feed on bamboo (Li et al. 2011; Chen et al. 2012; Yang and Chen 2013). So far, 17 species of this genus were known from China, of them, all species are distributed in the Oriental Region (China: Guizhou, Sichuan, Hunan, Hubei, Guangxi, Guangdong, Fujian and Zhejiang), and only *A. coneus* is also distributed in the Palaearctic Region (China: Gansu).

This genus is distinguished by its crown with two pairs of similar black spots on anterior margin, clypellus expanded apically, male pygophore with a long membranous process from its inner apex, and aedeagus with a well-developed basal projection dorsally (except *A. breviolus* and *A. langshanensis*).

In the present paper, two new species: *A. damingshanensis* sp. n. and *A. expansivus* sp. n. are described and illustrated from China (Oriental Region, Fig. 23). The type specimens of the new species are deposited in the Institute of Entomology, Guizhou University, Guiyang, China (GUGC).

Material and methods

Terminology of morphological and genital characters follow Dai and Zhang (2008). Male specimens were used for the description and illustration. External morphology was observed under a stereoscopic microscope and characters were measured with an ocular micrometer. Color pictures for adult habitus were obtained by KEYENCE VHX-1000 system. The genital segments of the examined specimens were macerated in 10% NaOH and drawn from preparations in glycerin jelly using a Leica MZ 12.5 stereomicroscope. Illustrations were scanned with Canon CanoScan LiDE 200 and imported into Adobe Photoshop CS8 for labeling and plate composition.


Descriptions of species

Abrus damingshanensis Xing & Li, sp. n.

<http://zoobank.org/211F15BE-9FD5-4A84-9D74-E3A16BA498DD>

Figs 1–11

Description. Yellowish brown species. Crown with two pairs of similar blackish brown spots on anterior margin. Eyes brown. Ocelli pale yellow. Pronotum with yellowish-brown stripe on anterior part. Inner and central anteapical cells at apex, third and



Figures 1–11. *Abrus damingshanensis* sp. n., **1** ♂, dorsal view **2** ♂, lateral view **3** ♂, head and thorax, dorsal view **4** ♂, face **5** Male pygofer side, lateral view **6** Valve, ventral view **7** Subgenital plate, ventral view **8** Aedeagus, lateral view **9** Aedeagus, caudal view **10** Connective, ventral view **11** Style, dorsal view.

fourth apical cells at base each with a dark brown spot. Face brown, frontoclypeus yellowish brown. Forewings yellowish. Legs marked with brown.

Body elongate, robust. Head including eyes narrower than greatest width of pronotum. Vertex with fore margin produced roundly, median length shorter than width between eyes. Eyes fairly large. Ocelli on anterior margin, separated from corresponding eye by approximately their own diameter. Frontoclypeus distinctly longer than wide, anteclypeus expanded apically. Antennae arising near lower corner of eye. Pronotum with anterior margin strongly and roundly produced, posterior margin slightly concave. Scutellum triangular, slightly shorter than pronotum, with transverse suture curved and depressed. Forewing with 3 subapical and 4 apical cells, 4 times as long as wide, appendix wide. Hind wings with three apical cells and two anteapical cells. Profemur with 2 dorsoapical setae. Hind femur apical setal formula 2+2+1. Hind tibia flattened and nearly straight, with PD setae very long. Metabasitarsomere with three platellae and two setae on apical transverse row.

Male genitalia. Male pygofer side longer than high, with many macrosetae posteriorly and some at midventral margin; posterior margin lobe alongate and with a long membranous process at inner apex (Fig. 5). Valve subtriangular with anterior margin concaved and posterior margin strongly produced medially (Fig. 6). Subgenital plate narrowing to rounded apex, outer margin rounded, with uniseriate row of macrosetae along lateral margin (Fig. 7). Aedeagus with well-developed basal projection on dorsal margin, tapered to acute apex, with pair of dorsal quadrilateral flange at midlength on dorsal margin; aedeagal shaft about half length of basal projection, expanded medially, apically branched in the caudal view, gonopore apical (Figs 8, 9). Connective Y-shaped, stem robust, arms well developed, articulated with the aedeagus (Fig. 10). Style long, broad at base, narrow at middle, apical margin expanded (Fig. 11).

Measurement. Length (including tegmen): ♂, 9.1–9.2 mm.

Type material. Holotype ♂, **China: Guangxi** Autonomous Region, Nanning City, Mt. Damingshan, 13 August 2011, coll. Zaihua Yang (GUGC); paratypes 2♂♂, same data as holotype (GUGC).

Diagnosis. The new species is similar to *A. leigongshanensis* Li & Wang, 2006, but can be distinguished from the latter by the aedeagal shaft broad and short (about half length of basal projection); the basal projection tapered to acute apex, with pair of quadrilateral flange at midlength; and the apical process of style expanded.


Etymology. This new species is named after the type locality, Damingshan, Guangxi Autonomous Region in China.

***Abrus expansivus* Xing & Li, sp. n.**

<http://zoobank.org/9C25F88C-D706-481F-B6B1-089C0EE2D4D8>

Figs 12–22

Description. External features resemble as *Abrus damingshanensis* sp. n., but color light yellow and body slightly small.



Figures 12–22. *Abrus expansivus* sp. n., 12 ♂, dorsal view 13 ♂, lateral view 14 ♂, head and thorax, dorsal view 15 ♂, face 16 Male pygofer side, lateral view 17 Valve, ventral view 18 Subgenital plate, ventral view 19 Aedeagus, lateral view 20 Aedeagus, caudal view 21 Connective, ventral view 22 Style, dorsal view.




Figure 23. Geographic distribution of two new *Abrus* species in China: *A. damingshanensis* sp. n. (■); *A. expansivus* sp. n. (●).

Male genitalia. Male pygofer side elongate with many macrosetae posteriorly and a few at midventral margin; posterior margin lobe elongate and with a long membranous process at inner apex (Fig. 16). Valve subtriangular with anterior margin concave and posterior margin strongly produced medially (Fig. 17). Subgenital plate broad and short, outer margin rounded, with many macrosetae on lateral margin (Fig. 18). Aedeagus with broad and flat basal projection from dorsal margin, deeply concave at dorsal margin, the basal projection without processes; aedeagal shaft slightly shorter than basal projection, with a pair of slender apical processes and its length equal to aedeagal shaft, gonopore apical (Figs 19, 20). Connective Y-shaped, stem robust, arms well developed, articulated with the aedeagus (Fig. 21). Style long, broad at base, narrow at middle, apex slightly widening (Fig. 22).

Measurement. Length (including tegmen): ♂, 8.1 mm.

Type material. Holotype ♂, **China: Guizhou** Province, Dushan County, 16 July 2012, coll. Qiongzhang Song (GUGC).

Diagnosis. This new species is very similar to *A. brevis* Dai & Zhang, 2002 in aedeagal shaft with a pair of long apical appendages, but can be distinguished from the latter by the aedeagal shaft longer than half length of basal projection; aedeagal shaft with apical processes located medially in lateral view and its length equal to aedeagal shaft; the apical process of style wide and flat; and the subgenital plate narrow apically.

Etymology. The new species name is derived from the Latin word “*expansivus*”, referring to the apical process of style wide and expand.

Taxonomic notes on *Abrus* species

Species of *Abrus* are all very similar in coloration and difficult to distinguish externally, but the structure of aedeagus are markedly different. This genus now contains 19 species, they can be divided into 3 types based on the structure of aedeagus: 1) basal projection of aedeagus very small or absent (*A. breviolus* and *A. langshanensis*); 2) aedeagal shaft obviously shorter than basal projection (*A. brevis*, *A. leigongshanensis*, *A. damingshanensis* sp. n. and *A. expansivus* sp. n.); 3) aedeagal shaft as long as or longer than basal projection (other 13 species).

Above mentioned the second type structure of aedeagus, of them, two species (*A. brevis* and *A. damingshanensis* sp. n.) with aedeagal shaft about half length of basal projection, two species (*A. brevis* and *A. expansivus* sp. n.) with aedeagal shaft apically have a pair of slender processes, and two species (*A. damingshanensis* sp. n. and *A. expansivus* sp. n.) with the apical process of style expand.

Acknowledgements

We are grateful to Dr. Yang Zaihua and Ms. Song Qiongzhang for collecting specimens, and also grateful to two anonymous referees for reading the manuscript and making some suggestions. This work was supported by China Postdoctoral Science Foundation funded project (2013T60864, 2012M521719) and the National Natural Science Foundation of China (31301909).

References

- Chen XS, Yang L, Li ZZ (2012) Bamboo-feeding leafhoppers in China. China Forestry Publishing House, Beijing. [In Chinese with English summary]
- Dai W, Zhang YL (2002) A new genus and six new species of Deltocephalinae from China (Homoptera: Cicadellidae). *Acta Zootaxonomica Sinica* 27: 313–322. [In Chinese with English summary]
- Dai W, Zhang YL (2008) A review of the genus *Abrus* Dai & Zhang (Hemiptera: Cicadellidae: Deltocephalinae) from China with description of one new species. *Zootaxa* 1688: 37–53.
- Li ZZ, Dai RH, Xing JC (2011) Deltocephalinae from China (Hemiptera: Cicadellidae). Popular Science Press, Beijing. [In Chinese with English summary]
- Li ZZ, Wang LM (2006) Descriptions of two new species of the genus *Abrus* from China (Hemiptera, Cicadellidae). *Acta Zootaxonomica Sinica* 31: 840–842. [In Chinese with English summary]
- Yang L, Chen XS (2013) Two new species of the bamboo-feeding leafhopper genus *Abrus* Dai & Zhang (Hemiptera, Cicadellidae, Deltocephalinae) from China. *ZooKeys* 318: 81–89. doi: 10.3897/zookeys.318.5799