

# The distribution and taxonomy of *Lissotriton* newts in Turkey (Amphibia, Salamandridae)

Ben Wielstra<sup>1,2</sup>, Emin Bozkurt<sup>3</sup>, Kurtuluş Olgun<sup>3</sup>

**1** Department of Animal and Plant Sciences, University of Sheffield, S10 2TN Sheffield, UK **2** Naturalis Biodiversity Center, P.O. Box 9517, 2300 RA Leiden, The Netherlands **3** Adnan Menderes University, Faculty of Science and Art, Department of Biology, 09010, Efeler, Aydın, Turkey

Corresponding author: Ben Wielstra (b.wielstra@sheffield.ac.uk)

Academic editor: F. Andreone | Received 31 October 2014 | Accepted 16 February 2015 | Published 26 February 2015

<http://zoobank.org/A5F4CFEC-78B3-46E2-9999-D22ACC45840A>

**Citation:** Wielstra B, Bozkurt E, Olgun K (2015) The distribution and taxonomy of *Lissotriton* newts in Turkey (Amphibia, Salamandridae). ZooKeys 484: 11–23. doi: 10.3897/zookeys.484.8869

## Abstract

Two and perhaps three taxa of *Lissotriton* newt occur in Turkey. Their species status is controversial. The distribution of these taxa and the taxonomic status of each are reviewed and discussed. A database of 128 Turkish *Lissotriton* localities was compiled and species distribution models were constructed. We reiterate that the presence of *L. (v.) lantzi* in Turkey is disputed and needs confirmation. The range of *L. (v.) kosswigi* is restricted to north-western Anatolia – given the small global range of this Turkey endemic, a closer look at its conservation status is warranted. The distribution of *L. v. schmidleri* covers western Asiatic and European Turkey. The findings support an allopatric distribution of the Turkish *Lissotriton* species. We reflect on the biological significance of previously reported morphological intermediates between *L. (v.) kosswigi* and *L. v. schmidleri* in the light of the recent proposal to recognize *kosswigi* at the species level. The available data are in line with species status for *L. (v.) lantzi* and *L. (v.) kosswigi*. Although *L. v. schmidleri* is a genetically diverged taxon as well, the extent of gene flow with parapatric European *Lissotriton* taxa is as yet unknown.

## Keywords

Anatolia, Bosphorus, historical biogeography, *Lissotriton kosswigi*, *Lissotriton lantzi*, *Lissotriton vulgaris schmidleri*, Smooth newt

## Introduction

The Smooth newt *Lissotriton vulgaris* group (Amphibia: Salamandridae) is distributed in Europe and adjacent Asia (Schmidtler and Franzen 2004). The taxonomy of the group is a matter of dispute, with the inclusive taxa usually referred to as subspecies, although some of these have been occasionally regarded as specifically distinct (see Dubois and Raffaëlli 2009, Speybroeck et al. 2010). Based on the taxonomy of Babik et al. (2005) the *Lissotriton vulgaris* group consists of seven taxa, namely *ampelensis* (Fuhn 1951), *graecus* (Wolterstorff 1906), *kosswigi* (Freytag 1955), *lantzi* (Wolterstorff 1914), *meridionalis* (Boulenger 1882), *schmidtleri* (Raxworthy 1988) [following the rationale of Dubois (2007), Dubois and Raffaëlli (2009) make the case that the original name *schmidtleri* as in Raxworthy (1988) is correct, rather than the name *schmidtlerorum* introduced in Raxworthy (1990)], and the nominal species *vulgaris* (Linnaeus 1758).

In Turkey, two and perhaps three *Lissotriton* occur (Fig. 1; Schmidtler and Franzen 2004). The range of *lantzi* covers the Caucasus region and the taxon might occur in the extreme north-east of Turkey, near the border with Georgia (Schmidtler and Franzen 2004, Skorinov et al. 2014). The taxon *kosswigi* is restricted to north-western Anatolia (Schmidtler and Franzen 2004). The taxon *schmidtleri* was originally considered to be restricted to western Asiatic Turkey, but Raxworthy (1988, 1990) suggested it might extend into Europe. Genetic data have subsequently confirmed that this taxon's range encompasses European Turkey (Nadachowska and Babik 2009; Pabijan et al. 2014). The range of *schmidtleri* protrudes further into the Balkan Peninsula, but its range limit is as yet unclear; newts with mitochondrial DNA typical of *schmidtleri* have been recorded as far north-west as easternmost Greece and central Bulgaria (Pabijan et al. 2014). Previous records of *vulgaris* from Turkey reflect incomplete taxonomy and can be referred to the other taxa (cf. Dubois and Raffaëlli 2012, Olgun et al. 1999).

The *Lissotriton vulgaris* group comprises two main morphological types: one with a smooth crest and flappy feet and another with a ragged crest and limited fringing on the feet (Fig. 2). Distinguishing the taxa within the two main groups is less straightforward and this topic is beyond the scope of the present paper (we refer to Raxworthy (1990) and Schmidtler and Franzen (2004) for a detailed treatment). Relevant for the current paper is that *kosswigi* belongs to the 'smooth-crested with flappy feet' type and *schmidtleri* to the 'ragged-crested with limited feet-fringing' type and that morphological intergradation has been reported between these two taxa (e.g. Freytag 1955, 1957, Tabrizi 1980, Yılmaz 1983). In Fig. 2 typical males of *kosswigi* and *schmidtleri* are depicted. Next to the smooth crest and flappy feet, *kosswigi* possesses a tail filament and its crest starts at a more posterior position than in *schmidtleri*. Although *lantzi* belongs to the 'ragged-crested with limited feet-fringing' type as well, confusion with *schmidtleri* is ruled out based on geography.

An overview of the distribution of the Turkish *Lissotriton* taxa is provided by composing a database of localities and constructing species distribution models. The focus is mainly on the taxa *kosswigi* and *schmidtleri* and particularly the supposed genetic admixture between the two. Finally, we reflect on the as yet controversial proposal to treat the Turkish *Lissotriton* taxa as distinct species.


## Material and methods

The distribution of *Lissotriton* in Turkey has been reviewed and a database compiled of localities based on: 1) the collection of the Zoology Laboratory of the Department of Biology at Science and Arts Faculty, Adnan Menderes University, 2) extensive personal field observations, and 3) a review of the literature (Bozkurt et al. in press, Çevik et al. 1997, Çiçek and Ayaz 2011, Demirsoy 1996, Eiselt 1966, Freytag 1955, Freytag 1957, Mulder 1995, Olgun et al. 1999, Raxworthy 1988, Schmidtler and Schmidtler 1967, Skorinov et al. 2014, Sparreboom and Arntzen 1987, Tabrizi 1980, Taşkın and Olgun 2003, Yılmaz 1983, 1989). In this paper we particularly focused on *kosswigi*, this being the rarest and most restricted taxon globally. The aim was not to be exhaustive for *schmidtleri*, which is common were not included, and widely distributed in western Turkey. Localities within one kilometre of one another and in such cases the locality with the most accurate information available was chosen. We particularly focused on records of presumed transitional forms between *kosswigi* and *schmidtleri* reported in the literature, considering their relevance in the taxonomic treatment of the different *Lissotriton* taxa occurring in Turkey.

For a species distribution modelling exercise for *lantzi* (and a comprehensive overview of the distribution of this taxon outside of Turkey) we refer to Skorinov et al. (2014). Species distribution models were constructed for *kosswigi* and *schmidtleri* using Maxent 3.3.3k (Phillips et al. 2006). For climate layers bioclimatic variables were used, at 2.5 arcminute resolution (c. 5 × 5 km) available from the WorldClim database 1.4 (Hijmans et al. 2005; <http://www.worldclim.org>). We trimmed these layers to an extent that broadly encompasses the distribution of the genus *Lissotriton*: the area between -15 and 65 degrees longitude and between 30 and 75 degrees latitude. Following Guisan and Thuiller (2005) and Peterson (2011) a subset considered to reflect physiological limitations of the study species (in this case seasonality) was selected while showing little multicollinearity (a Pearson's correlation of  $r < 0.7$ ): bio10 = mean temperature of warmest quarter, bio11 = mean temperature of coldest quarter, bio15 = precipitation seasonality, bio16 = precipitation of wettest quarter, and bio17 = precipitation of driest quarter. To determine whether our species distribution model performs better than random expectation, we tested its AUC value against a null model based on 99 models for random localities (see Raes and ter Steege 2007 for details). Random point data were created with ENMTtools 1.3 (Warren et al. 2010). To more thoroughly cover the range of environmental conditions experienced by *schmidtleri* the only four confirmed populations from outside the Turkish range (noted on Fig. 1; details in Pabijan et al. 2014) were included.

## Results

A database of 128 distribution records of Turkish *Lissotriton* newts (49 *kosswigi*, 78 *schmidtleri* and one *lantzi*) is provided in Suppl. material 1. Fig. 1 shows these records



**Figure 1.** Map showing the distribution of the taxa of the *Lissotriton vulgaris* group that occur in Turkey. The inset shows the rough outlines of the ranges of *lantzi* (in green), *kosswigi* (in red) and *schmidleri* (in blue). Type localities are marked with a black star. The blue hatched area reflects the unclear range of *schmidleri* outside of Turkey (see discussion), with four confirmed records denoted with white dots. The green hatched area reflects the potential occurrence of *lantzi* in the extreme northeast of Turkey, with a black dot depicting the single historical record for Turkey (see discussion); question marks denote historical records in Armenia and Azerbaijan. The cut-out shows Turkish localities for *kosswigi* and *schmidleri* as black dots. Localities supposedly showing intergradation between *kosswigi* and *schmidleri* are marked with a white star. The arrow highlights a poorly documented locality attributed to *kosswigi* (see discussion). Details on Turkish localities are provided in Suppl. material 1.

plotted on a map. The map also shows the type localities of *kosswigi* and *schmidtleri*, as well as populations reported to contain morphological intermediates between the two taxa. Fig. 3 shows the species distribution models for *kosswigi* and *schmidtleri*. The AUC values of these models (0.991 for both *kosswigi* and *schmidtleri*) rank above the 99 AUC values based on random points, meaning our species distribution models perform significantly better than random expectation ( $P < 0.05$ ).

## Discussion

### Distribution

The taxon *lantzi* is widely distributed in the Caucasus region (Schmidtler and Franzen 2004, Skorinov et al. 2014). A species distribution modelling exercise (Skorinov et al. 2014) revealed that suitable environmental conditions protrude into the extreme north-east of Turkey, near the border with Georgia. However, the continued occurrence of *lantzi* in Turkey needs confirmation; there is only a single record, dating from the beginning of the twentieth century (Schmidtler and Franzen 2004, Skorinov et al. 2014). Intriguingly, there are also old reports of the Crested newt *Triturus karelinii* (Strauch 1870) from this part of Turkey (Wielstra et al. 2010). Just as *lantzi*, *T. karelinii* is widely distributed in the Caucasus and, although its occurrence in Turkey is suggested by species distribution modelling (Wielstra et al. 2013c), its actual presence requires further scrutiny. In any case, *lantzi* is allopatric from the other Turkish taxa: *Lissotriton* newts are absent from north-east Anatolia (Schmidtler and Franzen 2004; Fig. 1).

The distribution of the Turkey endemic *kosswigi* is restricted to north-western Anatolia (our exhaustive survey revealed 49 localities; Fig. 1). The species distribution model suggests that suitable environmental conditions extend further to the east along most of the Turkish Black Sea coast (Fig. 3). However, this area appears to be devoid of *Lissotriton* newts (Fig. 1). Over-prediction is a well-known problem in species distribution modelling (Elith et al. 2011). This could suggest that the climate layers used to create the species distribution model do not properly reflect the factors limiting the distribution of the species, but it could also suggest that not all suitable area could be colonized due to dispersal constraints.

The taxon *schmidtleri* occurs in the west of Asiatic Turkey and is now known to extend into Europe, across the marine corridor connecting the Aegean and Black Seas (Nadachowska and Babik 2009, Pabijan et al. 2014). The permeability of this apparent barrier can be ascribed to sea level fluctuations related to glacial cycles and the disjunct distribution pattern of *schmidtleri* is mirrored by the co-distributed crested newt species *T. ivanbureschi* Arntzen & Wielstra, 2013 in Wielstra et al. (2013a) (Wielstra and Arntzen 2012). Although the Balkan range of *schmidtleri* outside of Turkey is poorly understood, the taxon appears to occur well into Bulgaria (Pabijan et al. 2014). This is in conflict with the species distribution model (Fig. 3). It could be that mitochondrial DNA does not properly reflect the range of *schmidtleri* and overestimates



**Figure 2.** Example of the two morphological types comprising the *Lissotriton vulgaris* group of newts. Shown (not to scale) are a typical *kosswigi* male (A) and a typical *schmidtleri* male (B). Notice the shape of the dorsal fin (smooth in *kosswigi* and ragged in *schmidtleri*), the position where the dorsal fin starts (approximately above the forelimbs in *kosswigi* and at the back of the head in *schmidtleri*), the presence of a thread-like tail filament (found in *kosswigi* but not in *schmidtleri*) and the extensiveness of the fringing on the feet (with *kosswigi* having much more flappy feet than *schmidtleri*).

its occurrence in Bulgaria. However, we consider it more likely that, due to the lack of confirmed *schmidtleri* localities from Bulgaria (whereas the taxon might well be abundant there), the species distribution model underestimates the environmental space inhabited by *schmidtleri*.

The taxa *kosswigi* and *schmidtleri* currently appear allopatric. We have particularly surveyed the area for *Lissotriton* (pers. obs.) and no localities are known between the *schmidtleri* locality Gemlik (Olgun et al. 1999; locality 62 in Suppl. material 1) and *kosswigi* locality Yalova (Demirsoy 1996; locality 18 in Suppl. material 1). The Yalova locality lacks documentation and needs confirmation (note that the locality was not included in Schmidtler and Franzen 2004) and it is suggested that there is probably a larger distribution gap, with the next closest *kosswigi* locality from the perspective of *schmidtleri* being Kocaeli (museum record; locality 3 in Suppl. material 1). This apparent distribution gap disagrees with the species distribution models, which suggest suitable environmental conditions for both *kosswigi* and *schmidtleri* occur south of the Marmara Sea (Fig. 3).


Based on introgression of *schmidtleri* mitochondrial DNA into *kosswigi* (very similar to mitochondrial DNA found in *schmidtleri* today) it has been hypothesized that

*kosswigi* displaced *schmidtleri* on the Istanbul Peninsula as the waterway between the Black and Marmara Seas rerouted within the last 10,000 years (Nadachowska and Babik 2009, Wielstra et al. 2013b). Similarly, an as yet undescribed *Triturus* species was proposed to have displaced *T. ivanbureschi* in this region (Wielstra et al. 2013a, 2013b). The species distribution models suggest suitable environmental conditions here for both *kosswigi* and *schmidtleri* and hence do not provide further insight into how *kosswigi* was able to locally outcompete *schmidtleri* (Fig. 3).

### Genetic admixture

In light of the current allopatric distribution pattern of *kosswigi* and *schmidtleri*, previous reports of transitional forms are curious. Following up on a possible intermediate specimen from Sapanca, Eiselt (1966) could only confirm the presence of pure *kosswigi* there. Freytag (1955) indicated that in a *Lissotriton* population from Kanlıca (locality 37 in Suppl. material 1), on the eastern side of the Bosphorus, some males showed characteristics of *schmidtleri*, namely the dorsal fin being ragged and starting at the back of the head and the lack of a tail filament (cf. Fig. 2). Tabrizi (1980) studied a larger sample of newts from populations throughout the range of *kosswigi*. He found that four out of 70 newts in Kanlıca showed a *schmidtleri*-like, relatively anterior starting position of the dorsal fins; all other newts were classified as typical *kosswigi*. Considering the biogeographical scenario outlined above, a relict *schmidtleri* population in the process of being replaced by *kosswigi* via genetic swamping is a possibility. A study on historical gene flow between the two taxa unfortunately did not include samples from the potentially admixed populations, but did suggest ancient gene flow from *schmidtleri* into *kosswigi* (Nadachowska and Babik 2009).

Furthermore, Freytag (1957) mentioned that in a *Lissotriton* population from Baltalimanı (locality 104 in Suppl. material 1), on the western side of the Bosphorus, some males shared similarities with *kosswigi*, in terms of possessing tail filaments and smooth dorsal fins that started relatively posteriorly (Fig. 2). Yılmaz (1983) studied a larger sample encompassing more populations from European Turkey. He noted newts with *kosswigi* characteristics at Habibler and Küçükçekmece (localities 121 and 125 in Suppl. material 1). Out of 80 studied newts, 20 had dorsal fins that began at the forelimbs rather than at the back of the head, 41 had smooth dorsal fins and 37 had tail filaments to varying degree (17 with 0–2 mm, 10 with 2–4 mm, 5 with 4–6 mm, and 5 with over 6 mm). Schmidtler and Franzen (2004) state that in *schmidtleri* males can show *kosswigi*-like characteristics, but do not provide further details. The presence of *kosswigi* west of the Bosphorus would not make sense in light of the biogeographical scenario outlined above, unless it could be proven that the Bosphorus on initial formation had a more westward position or formed only after *kosswigi* reached European Turkey. The rerouting of the marine connection between the Marmara and Black Seas is not yet fully understood and a matter of debate in the paleogeological literature (e.g. Nazik et al. 2011, 2012, Yalıtırak et al. 2012). We suggest that historical



**Figure 3.** Species distribution models for two Turkish *Lissotriton* taxa. Shown are species distribution models for *kosswigi* (A) and *schmidtleri* (B). The maps depict predicted suitability, which ranges from 0 to 1, in ten equal intervals, with higher values expressed by warmer colours.

biogeographical patterns such as shown by *Lissotriton* (and *Triturus*) newts might assist paleogeological reconstruction.

Genetic data from the potentially admixed *kosswigi* and *schmidtleri* populations are as yet lacking, but would provide more insight in the matter. However, considering the expansion of the Istanbul agglomeration it should be taken into account that these populations might well have gone extinct. We conclude that potential *kosswigi-schmidtleri* admixture represents, at most, the remnants of a former contact zone. The main ranges of the two taxa are currently isolated in the region by the Bosphorus and hence the influence of potentially admixed populations on the genetic integrity of the two taxa can be expected to be negligible. In this light we make some remarks on the not (yet) generally accepted treatment of the Turkish *Lissotriton* taxa as distinct species (Dubois and Raffaelli 2009, Frost 2014).

## Taxonomy

Following the taxonomy of Babik et al. (2005), the *Lissotriton vulgaris* group consists of seven taxa, namely *ampelensis*, *graecus*, *kosswigi*, *lantzi*, *meridionalis*, *schmidtleri* and the nominal *vulgaris*. Four of these taxa, *graecus*, *kosswigi*, *lantzi* and *meridionalis*, are sometimes regarded as specifically distinct (Dubois and Raffaelli 2009, Frost 2014). The split of *graecus* and *meridionalis* has been criticised (Speybroeck et al. 2010) as a misinterpretation of the phylogenetic position of the congener *L. montandoni* which, due to mitochondrial DNA introgression, is nested within the *L. vulgaris* group from the perspective of mitochondrial DNA (Babik et al. 2005, Zieliński et al. 2013). However, the taxa *kosswigi* and *lantzi* are genuinely genetically diverged for mitochondrial DNA (Babik et al. 2005).

Within the *Lissotriton vulgaris* group mitochondrial DNA suggests a basal split between *lantzi* and the rest (Babik et al. 2005). Although the distinction of *lantzi* from the perspective of the nuclear genome has as yet not been determined, the divergence in the mitochondrial genome and the at least currently disjunct distribution support a scenario of long-term disrupted gene flow with other *Lissotriton newts*.

The next split in the *Lissotriton vulgaris* group is between *kosswigi* and the remaining taxa (Babik et al. 2005). The distinction of *kosswigi* from its geographical neighbour *schmidtleri* has been supported in a study exploring gene flow based on eight nuclear DNA markers (Nadachowska and Babik 2009). Given that *kosswigi* is genetically distinct and currently allopatric from other *Lissotriton* taxa, its treatment at the species level seems justified. From the conservation perspective it is important whether this geographically restricted, Turkish endemic is treated as a 'unique species' or 'merely a subspecies'.

Although *schmidtleri* represents a distinct mitochondrial DNA clade as well, it is genetically nested within the European *Lissotriton* taxa (Babik et al. 2005, Pabijan et al. in prep.). The phylogeography of *Lissotriton* on the Balkan Peninsula is highly complex, with morphologically distinct subspecies being highly polyphyletic from the mitochondrial DNA perspective (Babik et al. 2005, Pabijan et al. in prep.). Furthermore, no doubt in part because of its turbulent taxonomical history, the morphological distinctiveness of *schmidtleri* is not well understood (Schmidtler and Franzen 2004). Hence, we refrain from making further comments on the taxonomic status of *schmidtleri* and rather await further research on nuclear gene flow between *schmidtleri* and the other *Lissotriton* taxa on the Balkan Peninsula.

## Acknowledgements

Wieslaw Babik, Sergé Bogaerts and Maciej Pabijan provided helpful comments during the preparation of this paper. BW is a Newton International Fellow.

## References

- Babik W, Branicki W, Crnobrnja-Isailovic J, Cogalniceanu D, Sas I, Olgun K, Poyarkov NA, Garcia-Paris M, Arntzen JW (2005) Phylogeography of two European newt species discordance between mtDNA and morphology. *Molecular Ecology* 14(8): 2475–2491. doi: 10.1111/j.1365-294X.2005.02605.x
- Boulenger GA (1882) Catalogue of the Batrachia Gradientia s. Caudata and Batrachia Apoda in the Collection of the British Museum. 2nd Edition. Taylor and Francis, London, UK.
- Bozkurt E, Olgun K, Wielstra B (in press) First record of facultative paedomorphism in the Kosswig's newt *Lissotriton (vulgaris) kosswigi* (Freytag, 1955) (Urodela; Salamandridae), endemic to northwestern Turkey. doi: 10.3906/zoo-1408-53
- Çevik İE, Atatür MK, Arikan H, Akyurtlaklı N (1997) Occurrence of neotenic *Triturus vulgaris* (Urodela: Salamandridae) larvae in western Anatolia. *Israel Journal of Zoology* 43(3): 301–304. doi: 10.1080/00212210.1997.10688916
- Çiçek K, Ayaz D (2011) New data on facultative paedomorphism of the smooth newt, *Lissotriton vulgaris*, in Western Anatolia, Turkey. *Journal of Freshwater Ecology* 26(1): 99–103. doi: 10.1080/02705060.2011.553924
- Demirsoy A (1996) Türkiye Omurglıları, Amfibiler. Meteksan, Ankara, Turkey.
- Dubois A (2007) Genitives of species and subspecies nomina derived from personal names should not be emended. *Zootaxa* 1550: 49–68.
- Dubois A, Raffaëlli J (2009) A new ergotaxonomy of the family Salamandridae Goldfuss, 1820 (Amphibia, Urodela). *Alytes* 26(1-4): 1–85.
- Dubois A, Raffaëlli J (2012) A new ergotaxonomy of the order Urodela Duméril, 1805 (Amphibia, Batrachia). *Alytes* 28(3-4): 77–161.
- Eiselt J (1966) Ergebnisse zoologischer Sammelreisen in der Türkei: Amphibia caudata. *Annalen des Naturhistorischen Museums in Wien* 69: 427–445.
- Elith J, Phillips SJ, Hastie T, Dudík M, Chee YE, Yates CJ (2011) A statistical explanation of MaxEnt for ecologists. *Diversity and Distributions* 17(1): 43–57. doi: 10.1111/j.1472-4642.2010.00725.x
- Freytag GE (1955) Ein neuerer Teichmolch aus der Türkei. *Zoologischer Anzeiger* 154: 195–200.
- Freytag GE (1957) Über *Triturus vulgaris* aus dem Gebiet von Istanbul. *Zoologischer Anzeiger* 158: 49–53.
- Frost DR (2014) Amphibian Species of the World: an Online Reference. Version 6.0 (31 October 2014). American Museum of Natural History, New York, USA. Electronic Database accessible at <http://research.amnh.org/herpetology/amphibia/index.html>
- Fuhn IE (1951) Contributiuni la sistematica Salamazdrelor din Republica Populara Romana. I. Studiul catorva populatii de *Triturus vulgaris* L. *Buletinul Stiintific. Sectiunea de Stiinte Biologice, Agronomice, Geologice si Geografice*. Bucuresti, Romania, 501.
- Guisan A, Thuiller W (2005) Predicting species distribution: offering more than simple habitat models. *Ecology Letters* 8(9): 993–1009. doi: 10.1111/j.1461-0248.2005.00792.x
- Hijmans RJ, Cameron SE, Parra JL, Jones PG, Jarvis A (2005) Very high resolution interpolated climate surfaces for global land areas. *International Journal of Climatology* 25(15): 1965–1978. doi: 10.1002/joc.1276

- Linnaeus C (1758) *Systema Naturae per Regna Tria Naturae, Secundum Classes, Ordines, Genera, Species, cum Characteribus, Differentiis, Synonymis, Locis*. 10th Edition. Volume 1. L. Salvii, Stockholm, Sweden.
- Mulder J (1995) Herpetological observations in Turkey (1987–1995). *Deinsea* 2: 51–66.
- Nadachowska K, Babik W (2009) Divergence in the face of gene flow: the case of two newts (Amphibia: Salamandridae). *Molecular Biology and Evolution* 26(4): 829–841. doi: 10.1093/molbev/msp004
- Nazik A, Meriç E, Avşar N, Ünlü S, Esenli V, Gökaşan E (2011) Possible waterways between the Marmara Sea and the Black Sea in the late Quaternary: evidence from ostracod and foraminifer assemblages in lakes İznik and Sapanca, Turkey. *Geo-Marine Letters* 31(2): 75–86. doi: 10.1007/s00367-010-0216-9
- Nazik A, Meriç E, Avşar N (2012) Reply to Discussion: a critique of Possible waterways between the Marmara Sea and the Black Sea in the late Quaternary: evidence from ostracod and foraminifer assemblages in lakes İznik and Sapanca, Turkey, *Geo-Marine Letters*, 2011. *Geo-Marine Letters* 32(3): 275–277. doi: 10.1007/s00367-012-0281-3
- Olgun K, Baran İ, Tok CV (1999) The Taxonomic Status of *Triturus vulgaris* (Linnaeus, 1758) Populations in Western Anatolia, Turkey. *Turkish Journal of Zoology* 23: 133–140.
- Pabijan M, Zieliński P, Dudek K, Chloupek M, Sotiropoulos K, Liana M, Babik W (2014) The dissection of a Pleistocene refugium – phylogeography of the smooth newt, *Lissotriton vulgaris*, in the Balkans. *Journal of Biogeography*. doi: 10.1111/jbi.12449
- Peterson AT (2011) Ecological niche conservatism: a time-structured review of evidence. *Journal of Biogeography* 38(5): 817–827. doi: 10.1111/j.1365-2699.2010.02456.x
- Phillips SJ, Anderson RP, Schapire RE (2006) Maximum entropy modeling of species geographic distributions. *Ecological Modelling* 190(3–4): 231–259. doi: 10.1016/j.ecolmodel.2005.03.026
- Raes N, ter Steege H (2007) A null-model for significance testing of presence-only species distribution models. *Ecography* 30(5): 727–736. doi: 10.1111/j.2007.0906-7590.05041.x
- Raxworthy CJ (1988) A description and study of a new dwarf sub-species of smooth newt, *Triturus vulgaris*, from western Anatolia, Turkey. *Journal of Zoology* 215(4): 753–763. doi: 10.1111/j.1469-7998.1988.tb02409.x
- Raxworthy CJ (1990) A review of the smooth newt (*Triturus vulgaris*) subspecies, including an identification key. *Herpetological Journal* 1 (11): 481–490.
- Schmidler J, Schmidler F (1967) Über die Verbreitung der Molchgattung *Triturus* in Kleinasien. *Salamandra* 3: 15–36.
- Schmidler JF, Franzen M (2004) *Triturus vulgaris* Linnaeus, 1758 – Teichmolch. In: Grosenbacher K, Thiesmeier B (Eds) *Handbuch der Reptilien und Amphibien Europas*. Aula Verlag, Wiesbaden, Germany, 847–967.
- Skorinov DV, Doronin IV, Kidov AA, Tuniev BS, Litvinchuk SN (2014) Distribution and conservation status of the Caucasian newt, *Lissotriton lantzi* (Wolterstorff, 1914). *Russian Journal of Herpetology* 21(4): 251–268.
- Sparreboom M, Arntzen P (1987) Über die Amphibien in der Umgebung von Adapazari, Türkei. *Herpetofauna* 9: 27–34.

- Speybroeck J, Beukema W, Crochet PA (2010) A tentative species list of the European herpetofauna (Amphibia and Reptilia). *Zootaxa* 2492: 1–27.
- Strauch A (1870) Revision der Salamandriden-Gattungen nebst Beschreibung einiger neuen oder weniger bekannten Arten dieser Familie. *Mémoires de l'Académie impériale des sciences de St.-Petersbourg Serie 7*, 16: 17–426.
- Tabrizi FH (1980) On some new material of *Triturus vulgaris* (L.) from the area between Abant (Bolu) and Bosphorus in NW Anatolia. *E. Ü. Fen Fakültesi Dergisi*, 1, 2, 3, 4: 115–140.
- Taşkın N, Olgun K (2003) Morphological comparison of two populations of the Smooth Newt, *Triturus vulgaris*, from western Anatolia. *Zoology in the Middle East* 28 (1): 61–66. doi: 10.1080/09397140.2003.10637956
- Warren DL, Glor RE, Turelli M (2010) ENMTTools: a toolbox for comparative studies of environmental niche models. *Ecography* 33(3): 607–611. doi: 10.1111/j.1600-0587.2009.06142.x
- Wielstra B, Espregueira Themudo G, Güçlü Ö, Olgun K, Poyarkov NA, Arntzen JW (2010) Cryptic crested newt diversity at the Eurasian transition: the mitochondrial DNA phylogeography of Near Eastern *Triturus* newts. *Molecular Phylogenetics and Evolution* 56(3): 888–896. doi: 10.1016/j.ympev.2010.04.030
- Wielstra B, Arntzen JW (2012) Postglacial species displacement in *Triturus* newts deduced from asymmetrically introgressed mitochondrial DNA and ecological niche models. *BMC Evolutionary Biology* 12: 161. doi: 10.1186/1471-2148-12-161
- Wielstra B, Litvinchuk SN, Naumov B, Tzankov N, Arntzen JW (2013a) A revised taxonomy of crested newts in the *Triturus karelinii* group (Amphibia: Caudata: Salamandridae), with the description of a new species. *Zootaxa* 3682(3): 441–453. doi: 10.11646/zootaxa.3682.3.5
- Wielstra B, Baird AB, Arntzen JW (2013b) A multimarker phylogeography of crested newts (*Triturus cristatus* superspecies) reveals cryptic species. *Molecular Phylogenetics and Evolution* 67(1): 167–175. doi: 10.1016/j.ympev.2013.01.009
- Wielstra B, Crnobrnja-Isailović J, Litvinchuk SN, Reijnen BT, Skidmore AK, Sotiropoulis K, Toxopeus AG, Tzankov N, Vukov T, Arntzen JW (2013c) Tracing glacial refugia of *Triturus* newts based on mitochondrial DNA phylogeography and species distribution modeling. *Frontiers in Zoology* 10: 13. doi: 10.1186/1742-9994-10-13
- Wolterstorff W (1906) Über *Triton vulgaris* L. subsp. *graeca* Wolt. n. subsp. *Zoologischer Anzeiger* 29: 137–139.
- Wolterstorff W (1914) Zwei neue Tritonformen der paläarktischen Region. *Abhandlungen und Berichte aus dem Museum für Natur- und Heimatkunde zu Magdeburg* 2: 371–381.
- Yalıtırak C, Ülgen U, Zabcı C, Franz S, Ön S, Sakınç M, Çağatay MN, Alpar B, Öztürk K, Tunoğlu C, Ünlü S (2012) Discussion: a critique of Possible waterways between the Marmara Sea and the Black Sea in the late Quaternary: evidence from ostracod and foraminifer assemblages in lakes İznik and Sapanca, Turkey, *Geo-Marine Letters*, 2011. *Geo-Marine Letters* 32(3): 267–274. doi: 10.1007/s00367-011-0270-y
- Yılmaz İ (1983) Trakya Kuyruklu Kurbağaları Üzerine Morfolojik ve Taksonomik Bir araştırma. *Doğa Bilim Dergisi* 7: 119–130.

- Yılmaz İ (1989) Kuzey Anadolu Amfibilerinin Yayılışı Üzerine Bir Çalışma (Amphibia: Urodela, Anura). *Doğa TU Zooloji* 13(2): 130–140.
- Zieliński P, Nadachowska-Brzyska K, Wielstra B, Szkotak R, Covaciu-Marcov S, Cogălniceanu D, Babik W (2013) No evidence for nuclear introgression despite complete mtDNA replacement in the Carpathian newt (*Lissotriton montandoni*). *Molecular Ecology* 22(7): 1884–1903. doi: 10.1111/mec.12225

## **Supplementary material I**

### **Locality data for Turkish *Lissotriton* newts**

Authors: Ben Wielstra, Emin Bozkurt, Kurtuluş Olgun

Data type: ZIP archive

Explanation note: The ZIP archive contains the locality database as an Excel file and a shape file.

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.