

Caribbean Amphipoda (Crustacea) of Panama. Part I: parvorder Oedicerotidira

Elizabeth L. Durham¹, Kristine N. White¹

¹ Georgia College & State University, Department of Biological and Environmental Sciences, Milledgeville, GA 31061, USA

Corresponding author: Kristine N. White (kristine.white@gcsu.edu)

Academic editor: Alan Myers | Received 15 February 2023 | Accepted 5 April 2023 | Published 24 April 2023

<https://zoobank.org/8FEFD7CD-4AC1-484F-ACB8-42A440ADB866>

Citation: Durham EL, White KN (2023) Caribbean Amphipoda (Crustacea) of Panama. Part I: parvorder Oedicerotidira. ZooKeys 1159: 37–50. <https://doi.org/10.3897/zookeys.1159.102034>

Abstract

Amphipods in the parvorder Oedicerotidira are burrowers, furrowers, or surface skimmers. Members of the parvorder share a well-developed posteroventral lobe on coxa 4, an equilobate coxa 5, an immensely elongate pereopod 7 that differs in structure from pereopod 6, and an entire telson. Within the parvorder, only the family Oedicerotidae has been documented from Bocas del Toro, Panama, represented by two species. This research documents a range extension for *Hartmanodes nyei* (Shoemaker, 1933) and describes a new species of *Synchelidium* Sars, 1892. An identification key to the species of Caribbean Oedicerotidae of Panama is provided.

Keywords

Bocas del Toro, Caribbean, *Hartmanodes nyei*, identification key, new species, Oedicerotidae, Panama, *Synchelidium purpurivitellum* sp. nov.

Introduction

Parvorder Oedicerotidira Lilljeborg, 1865 (Lilljeborg 1865b) is comprised of 302 species, with representative species documented around the world. Members of Oedicerotidira share a well-developed posteroventral lobe on coxa 4, an equilobate coxa 5, an immensely elongate pereopod 7 that differs in structure from pereopod

6, and an entire telson (Lowry and Myers 2017). The parvorder contains three families of burrowing, furrowing, or surface skimming amphipods: Exoedicerotidae Barnard & Drummond, 1982 (20 spp.), Oedicerotidae Lilljeborg, 1865 (262 spp.), and Paracalliopiidae Barnard & Karaman, 1982 (20 spp.). Only one of these families, the Oedicerotidae, is known to occur in the Caribbean Sea and, to date, only six species within that family have been reported from the Caribbean: *Aceroides (Patoides) synparis* (Barnard, 1964); *Americhelidium americanum* (Bousfield, 1973); *Hartmanodes nyei* (Shoemaker, 1933); *Kroyera carinata* Bate, 1857 (as *Monoculodes* cf. *carinatus*); *Periocolodes cerasinus* Thomas & Barnard, 1985; *Westwoodilla longimana* Shoemaker, 1934 (LeCroy et. al. 2009; Martín et. al. 2013). Of these Caribbean species, only *A. synparis* has been documented from Panamanian waters, occurring at a depth of 850 m (Barnard 1964).

Defining characteristics of amphipods in the family Oedicerotidae include having a well-developed antenna 2, reaching at least half the length of antenna 1; a strong down-curved rostrum; well-developed dorsolateral eyes; coxae 1–3 well-developed, each longer than the previous coxa; a subchelate gnathopod 1; article 3 of gnathopod 2 less than $2 \times$ as long as wide; and a distally attenuate pereopod 7 that is longer and more slender than pereopod 6. Amphipods in the family Oedicerotidae differ from those in the families Exoedicerotidae and Paracalliopiidae in having separate urosomite segments and lacking the oblique setal row on the maxilla 2 inner plate (Lowry and Myers 2017). Most species of the parvorder burrow into sediment, but little else is known about their ecology.

Two species of oedicerotid amphipods were collected from Bocas del Toro, Panama, one of which is new to science. Both species are diagnosed and the new species is described herein, and an identification key is provided to distinguish between the three species known from the Caribbean waters of Panama.

Materials and methods

Amphipods were collected by hand using a Ziploc bag to scoop up fine sand from Crawl Cay, Bocas del Toro, Panama at depths of 1.5–5.0 m. The sand was elutriated with freshwater to remove amphipods. Live specimens were sorted to morphospecies, placed in clove oil for imaging, and preserved in 99.5% EtOH for later examination. Preserved specimens were transferred to glycerol, measured from the tip of the rostrum to the base of the telson, and dissected under a stereomicroscope. Specimens were illustrated using a Meiji MT5900L phase contrast microscope with an Olympus U-DA drawing tube attached. Illustrations were digitally inked following Coleman (2003) in Adobe Illustrator 2020 using a Wacom Intuos Pro Pen Tablet. Specimens are deposited in the Smithsonian Institution, U.S. National Museum of Natural History (USNM).

Results

Descriptions

Parvorder Oedicerotidira Lilljeborg, 1865 (Lilljeborg, 1865b)

Superfamily Oedicerotoidea Lilljeborg, 1865 (Lilljeborg, 1865a)

Family Oedicerotidae Lilljeborg, 1865 (Lilljeborg, 1865b)

Genus *Hartmanodes* Bousfield & Chevrier, 1996

Generic diagnosis. Antenna 1 shorter than antenna 2; male antenna 2 much longer than that of female; head, rostrum large, apex deflexed. Gnathopods 1 and 2 not sexually dimorphic; gnathopod 1 carpus broad, propodus long, ovate; Gnathopod 2 subchelate, carpus narrow, propodus elongate, narrowing distally. Pereopod 5 coxa large, deep, equilobate. Pereopod 7 basis with small posterodistal lobe. Telson short, apex truncate or emarginate.

***Hartmanodes nyei* (Shoemaker, 1933)**

Figs 1, 2, 6A

Monoculodes nyei Shoemaker, 1933: 9, fig. 5.

Hartmanodes nyei: Bousfield and Chevrier 1996: 92; LeCroy 2000: 169, fig. 205.

Material examined. PANAMA • 1 ♀, 4.0 mm; Bocas del Toro, Crawl Cay; 9.2475°N, 82.1290°W; depth 5 m, in sand; 12 Aug 2021; K.N. White leg; USNM 1522425 • 1 ♂, 3.5 mm; same station data as for preceding; USNM 1522426 • 4 ♀; same station data as for preceding; USNM 1522427 • 1 ♀, 1 ♂, 2 juvenile; Bocas del Toro, Crawl Caye; 9.2376°N, 82.1438°W; depth 2 m, in sand, 11 Aug 2021; K.N. White leg.; USNM 1522428.

Diagnosis. Antenna 1 of female subequal to peduncle of antenna 2; antenna 2 of male much longer than that of female; head, anterodorsal angle broadly subquadrate. Pereopods 3 and 4 propodus subrectangular, dactylus elongate, slender; pereopods 5 and 6 dactylus elongate, subequal to propodus in length; pereopod 7 basis posterior margin with several short setae, carpus and propodus, posterior margin with several spine groups, dactylus elongate, slender.

Distribution. U.S.A.: Gulf of Mexico (Rakocinski et al. 1993), South Florida (Shoemaker 1933; Thomas 1993), Pacific California (Barnard 1962; Bousfield and Chevrier 1996); South America: Brazil (Shoemaker 1933); Central America: Belize (Thomas 1993), Panama (present study).

Ecology. These amphipods burrow into sand in shallow subtidal habitats.



Figure 1. *Hartmanodes nyei*, female, 4.0 mm, pereopod 3, pereopod 3 dactyl with setae removed, pereopod 6 and 7, antennae 1 and 2; male, 3.5 mm, antenna 1 and antenna 2 (broken). Scale bars: 0.1 mm.




Figure 2. *Hartmanodes nyei*, female, 4.0 mm, gnathopods 1 and 2 lateral. Scale bar 0.1 mm.

Genus *Synchelidium* Sars, 1892

Generic diagnosis. Antenna 1 shorter than antenna 2; male antenna 2 much longer than that of female; head, rostrum strong, distally deflexed. Gnathopods 1 and 2 not sexually dimorphic; gnathopod 1 carpus elongate, slender, propodus broad; Gnathopod 2 chelate, carpal lobe slender, propodus elongate. Pereopod 5 coxa medium, deep, equilobate. Pereopod 7 basis lacking or with weak posterior lobe. Telson short, apex emarginate or rounded.

Synchelidium purpurivitellum sp. nov.

<https://zoobank.org/117F5F75-7DE0-403D-8C9E-34DCB11C8EC1>

Figs 3–5, 6B

Type locality. Bocas del Toro, Panama: Crawl Cay, 9.2475°N, 82.1290°W, depth 5 m, in sand.

Material examined. Holotype: PANAMA • 1 ♀, 2.3 mm; Bocas del Toro, Crawl Cay; 9.2475°N, 82.1290°W; depth 5 m, in sand; 12 Aug 2021; K.N. White leg; USNM 1522429.

Paratypes: PANAMA • 1 ♂, 2.0 mm; same station data as for preceding; USNM 1522430 • 1 ♀, 2.0 mm; same station data as for preceding; USNM 1522431.




Figure 3. *Synchelidium purpurivitellum* sp. nov. holotype female, 2.3 mm, habitus; paratype male, 2.0 mm: head, antennae broken. Scale bars: 0.5 mm.

Other material: PANAMA • 2 ♀, 1 ♂, 3 juvenile, same station data as for preceding; USNM 1522432.

Diagnosis. Gnathopod 1 propodus, palm regularly toothed. Gnathopod 2 propodus slender, $6 \times$ length of dactyl. Pereopod 3 propodus with anteroproximal margin longer than anterodistal margin, dactylus short, stubby. Coxa 4 posteroventral angle slightly produced. Coxa 6 posteroventral angle narrowly rounded. Pereopod 7 merus with spines on posterior margin slightly shorter than width of article. Epimeron 3 anteroventral margin narrowly produced. Telson thickened dorsoventrally, narrowing distally, apex subtruncate with two medium setae dorsolaterally, two short setae medially.

Description. Female (holotype, 2.3 mm). **Head.** Rostrum deflexed, reaching ventral margin of head, not surpassing article 1 of antenna 1. Eyes large, covering entire anterior portion of head. Antenna 1 length surpassing peduncle of antenna 2, moderately setose, peduncle segments subequal; flagellum 5-articulate. Antenna 2 is $1.2 \times$ length of antenna 1, flagellum 5-articulate. Maxilliped, inner plate with four apical setae, outer plate with four or five distomedial, marginal setae. Lower lip, inner lobes rounded, outer lobes with large gape, apically setose. Maxilla 1 outer lobe with five apical plumose setae; palp bi-articulate, with three apical setae. Maxilla 2 inner plate with two apical setae, outer plate with three apical setae. Mandibles similar, incisors dentate; left mandible lacinia mobilis 6-dentate, right mandible lacinia mobilis 4-dentate; four




Figure 4. *Synchelidium purpurivitellum* sp. nov. holotype female, 2.3 mm, gnathopods 1 and 2 lateral, G1 medial palm, left mandible, maxilla 1 and 2; paratype male, 2.0 mm: pereopod 7 (dactyl broken), telson. Scale bars: 0.1 mm.

accessory spines; molar process small; palp tri-articulate, article 2 with three setae, article 3 subequal in length with article 1, with two or three setae. Upper lip asymmetrical, apically setose.

Pereon. Coxae weakly setose on distal margin; coxae 1–3 subrectangular; coxa 4 subquadrate, slightly produced posterodistally. Gnathopod 1 subchelate; basis with few short setae on anterior margin; merus not expanded; carpal lobe reaching palmar angle, with two distal setae; propodus ovate, palm oblique, regularly toothed, defined by blunt tipped spine; dactylus reaching palmar angle. Gnathopod 2 chelate; basis slender, with one short seta on anterodistal margin; propodus minutely toothed on cutting edge of fixed finger, sparsely setose; total length of dactylus one-sixth of propodus, smooth. Pereopods 3 and 4 bases slender, with long plumose setae on anterodistal and posterodistal margins; propodus with anteroproximal margin longer than anterodistal margin; dactylus short, stout, one-fourth length of propodus. Pereopod 5 basis anterior margin with distally located plumose setae and a single row of medial plumose setae along the midline; propodus subequal to carpus. Pereopod 6 moderately setose; propodus subequal to carpus. Pereopod 7 basis subrectangular, lacking posterodistal lobe, posterior margin lined with short setae; dactylus styliform, at least as long as propodus (broken).

Pleon. Epimera 1–3 margins smooth, bare; epimeron 3, posteroventral margin evenly rounded. Uropod 1, peduncle $1.2 \times$ length of outer ramus, inner and outer rami subequal in length, inner ramus with one robust marginal seta, outer ramus with four robust setae, inner and outer rami lined with fine setae. Uropod 2 peduncle subequal in length to outer ramus, inner ramus broken, inner ramus with one robust marginal seta, outer ramus with three robust setae, inner and outer rami lined with fine setae. Uropod 3 peduncle $0.3 \times$ length of outer ramus, inner ramus $1.2 \times$ length of outer ramus, each ramus with one robust seta, lined with fine setae. Telson thickened dorsoventrally, narrowing distally, apex subtruncate with two medium setae dorsolaterally, two short setae medially.

Male (paratype, 2.0 mm). Similar in all aspects to the female with the exception of the following: Eye slightly larger; antenna 2 flagellum elongate, at least $0.5 \times$ body length (broken); gnathopods 1 and 2 bases slightly wider than in female.

Etymology. After the Latin *purpur*, meaning purple and *vitellum*, meaning yolk and referring to the striking purple color of the eggs in the brood pouch of females of this species.

Distribution. Panama: Bocas del Toro (present study).

Ecology. These amphipods burrow into sand in shallow subtidal habitats.

Remarks. *Synchelidium purpurivitellum* sp. nov. is similar to the geographically close species *Americhelidium americanum* in many aspects, but differs in having a strongly toothed gnathopod 1 propodal palm (smooth in *A. americanum*), a broadly rounded posteroventral corner of epimeron 2 (produced in *A. americanum*), and lacking the posterodistal lobe found on the basis of pereopod 7 in *A. americanum*.

Within the genus, *Synchelidium purpurivitellum* sp. nov. is most similar to *Synchelidium maculatum* Stebbing, 1906, sharing the long gnathopod 1 carpus and toothed




Figure 5. *Synchelidium purpurivitellum* sp. nov. holotype female, 2.3 mm, antennae 1 and 2, pereopods 3 and 4, uropods 1–3; paratype male, 2.0 mm: maxilliped. Scale bars: 0.1 mm.



Figure 6. Photographs of live specimens **A** *Hartmanodes nyei*, male **B** *Synchelidium purpurivitellum* sp. nov. female. Scale bar: 1.0 mm.

propodus palm, minute pereopod 3 and 4 dactyls, and rounded posteroventral corner of epimeron 2. It differs from *S. maculatum* in having shorter antenna 1 flagellum, subrectangular gnathopod 1 propodus (ovate in *S. maculatum*), and subtruncate telson (rounded in *S. maculatum*). The subtruncate telson distinguishes *S. purpurivitellum* sp. nov. from all other *Synchelidium* species, but this character most resembles *Synchelidium intermedium* Sars, 1892, which has a truncate telson.

Identification Key to the Caribbean Oedicerotae of Panama

- 1 Gnathopod 1, carpal lobe slender; gnathopod 2 chelate; pereopods 3 and 4, dactylus short..... ***Synchelidium purpurivitellum* sp. nov**
- Gnathopod 1, carpal lobe broad; gnathopod 2 subchelate; pereopods 3 and 4, dactylus long..... **2**
- 2 Eye well-developed; pereopods 3 and 4, carpus subquadrate, not produced, subequal in width to propodus; pereopod 7, basis without facial setae; telson apically truncate..... ***Hartmanodes nyei***
- Eye absent; pereopods 3 and 4, carpus posteroventrally produced, 3 × as wide as propodus; pereopod 7, basis with facial setae; telson apically convex.....
..... ***Aceroides (Patoides) synparis***

Discussion

The results of this study increase the number of Caribbean oedicerotid amphipods known from Panama to three, with the documentation of a new species and a range extension to include Panama for *H. nyei*. *Hartmanodes nyei* has previously been reported from the western Atlantic Ocean, including the Caribbean Sea (Shoemaker 1933; Rakocinski et al. 1993; Thomas 1993; LeCroy 2000), with two questionable records in the Pacific Gulf of California (Barnard 1962; Bousfield and Chevrier 1996).

The genus *Synchelidium* now contains eight species worldwide, with *Synchelidium purpurivitellum* sp. nov. being the first Caribbean species of the genus. *Americhelidium americanum* was reported from the Caribbean as *Synchelidium americanum* before the closely related genus *Americhelidium* Bousfield & Chevrier, 1996 was erected (Thomas 1993). Bousfield and Chevrier (1996) designated *Americhelidium* as a North Pacific coastal marine genus, including *A. americanum*, and *Synchelidium* as a European Atlantic genus. Despite this designation, *Synchelidium purpurivitellum* sp. nov. is placed in the genus based on the strongly toothed gnathopod 1 propodus, broadly rounded epimeron 2, and the lack of a posterodistal lobe on the basis of pereopod 7, all of which are the diagnostic characters selected by Bousfield and Chevrier (1996) to distinguish this genus from *Americhelidium*. It is likely that the genera have a wider distribution, but have yet to be documented properly, given the similarities among the species, which would explain its presence in Caribbean Panama. Future investigation

of additional material, including type material will aid in understanding the relationships between species of the *Americhelidium* and *Synchelidium*. Documenting these sand burrowing species from Bocas del Toro may allow their inclusion in future applied studies, including studies on trophic interactions, habitat-use, and population or community analyses.

Funding

Funding for this study was provided by a National Science Foundation grant: Collaborative Research: ARTS: Understanding Tropical Invertebrate Diversity Through Integrative Revisionary Systematics and Training (1856421). Publication costs were provided by the Georgia College & State University GC Journeys Program.

Acknowledgements

Logistical support and facilities were provided by Georgia College & State University Department of Biological and Environmental Sciences and the Smithsonian Tropical Research Institute (STRI). Special thanks to Carolina Cesar for assistance with diving and collecting in Bocas del Toro. Comments and suggestions from Sara LeCroy and Azman Abdul Rahim greatly improved the manuscript.

References

- Barnard JL (1962) Benthic marine Amphipoda of southern California: Family Oedicerotidae. *Pacific Naturalist* 3(12): 349–371. <http://hdl.handle.net/1834/27427>
- Barnard JL (1964) Deep-sea Amphipoda (Crustacea) collected by R.V. “Vema” in the eastern Pacific Ocean and the Caribbean and Mediterranean Seas. *Bulletin of the American Museum of Natural History* 127(1): 3–46. <http://hdl.handle.net/2246/1119>
- Barnard JL, Drummond MM (1982) Redescription of *Exoediceros fossor* (Stimpson, 1856), an Australian marine fossorial Amphipod, the type-genus of the new family Exoedicerotidae. *Proceedings of the Biological Society of Washington* 95(3): 610–620. <http://www.biodiversitylibrary.org/page/1694499>
- Barnard JL, Karaman GS (1982) Classificatory revisions in gammaridean Amphipoda (Crustacea), Part 2. *Proceedings of the Biological Society of Washington* 95(1): 167–187. <https://doi.org/10.5479/si.00810282.360>
- Bate CS (1857) A synopsis of the British edriophthalmous Crustacea Part I. Amphipoda. *The Annals and Magazine of Natural History*, series 2 19(110): 135–152. <https://doi.org/10.1080/00222935708697715>
- Bousfield EL (1973) Shallow-water Gammaridean Amphipoda of New England. Cornell University Press, New York, 312 pp.

- Bousfield EL, Chevrier A (1996) The Amphipod Family Oedicerotidae on the Pacific Coast of North America. 1. The *Monoculodes* & *Synchelidium* Generic Complexes. Systematics and Distributional Ecology: Amphipacifica 2(2): 75–148.
- Coleman CO (2003) “Digital inking”: How to make perfect line drawings on computers. Organisms, Diversity & Evolution 3(14): 1–14. <https://doi.org/10.1078/1439-6092-00081>
- LeCroy SE (2000) An Illustrated Identification Guide to The Nearshore Marine and Estuarine Gammaridean Amphipoda of Florida: Families Gammaridae, Hadziidae, Isaeidae, Melitidae and Oedicerotidae. Florida Department of Environmental Protection Annual Report Contract No. WM724 1: 1–195.
- LeCroy SE, Gasca R, Winfield I, Ortiz M, Escobar-Briones E (2009) Amphipoda (Crustacea) of the Gulf of Mexico. In: Felder DL, Camp DK (Eds) Gulf of Mexico: Origin, Waters, and Biota. Texas A&M University Press, Texas, 941–972.
- Lilljeborg W (1865a) Bidrag till kannedomen om underfamiljen Lysianassina inom underordningen Amphipoda bland kraftdjuren. Nova Acta Regiae Societatis Scientiarum Upsaliensis 3: 1–25.
- Lilljeborg W (1865b) On the *Lysianassa magellanica* H. Milne Edwards, and on the Crustacea of the suborder Amphipoda and the subfamily Lysianassina found on the coast of Sweden and Norway. Nova Acta Regiae Societatis Scientiarum Upsaliensis 6(1): 1–38. <https://doi.org/10.5962/bhl.title.6806>
- Lowry JK, Myers AA (2017) A Phylogeny and Classification of the Amphipoda with the establishment of the new order Ingolfiellida (Crustacea: Peracarida). Zootaxa 4265(1): 1–89. <https://doi.org/10.11646/zootaxa.4265.1.1>
- Martín A, Díaz Y, Miloslavich P, Escobar-Briones E, Guerra-García JM, Ortiz M, Valencia B, Giraldo A, Klein E (2013) Regional diversity of Amphipoda in the Caribbean Sea. Revista de Biología Tropical 61(4): 1681–1720. <https://doi.org/10.15517/rbt.v61i4.12816>
- Rakocinski CF, Heard RW, LeCroy SE, McLelland JA, Simons T (1993) Seaward change and zonation of the sandy shore macrofauna at Perdido Key, Florida, U.S.A. Estuarine, Coastal and Shelf Science 36(1): 81–104. <https://doi.org/10.1006/ecss.1993.1007>
- Sars GO (1892) Amphipoda. Part XIV. Oediceridae (continued). An account of the Crustacea of Norway, with short descriptions and figures of all the species 1: 297–320[, pls 105–112]. <https://doi.org/10.5962/bhl.title.1164>
- Shoemaker CR (1933) Amphipoda from Florida and the West Indies. American Museum Novitates 598: 1–24.
- Shoemaker CR (1934) Three new amphipods. Reports on the collections obtained by the first Johnson-Smithsonian deep-sea expedition to the Puerto Rican Deep. Smithsonian Miscellaneous Collections 91(12): 1–6.
- Stebbing TRR (1906) Amphipoda I. Gammaridea. Das Tierreich 21: 1–806. <https://doi.org/10.5962/bhl.title.1224>
- Thomas JD (1993) Identification manual for marine Amphipoda (Gammaridea): I. Common coral reef and rocky bottom amphipods of South Florida. Florida Department of Environmental Protection Final Report Contract No. SP290: 1–83.
- Thomas JD, Barnard JL (1985) *Periculodes cerasinus*, n. sp., the first record of the genus from the Caribbean Sea (Amphipoda: Oedicerotidae). Proceedings of the Biological Society of Washington 98(1): 98–106.

Supplementary material I

Collection data for Caribbean Oedicerotidira of Panama

Authors: Elizabeth L. Durham, Kristine N. White

Data type: COL (.xlsx file)

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/zookeys.1159.102034.suppl1>