

Research Article

# Two new species of *Amaurobius* C.L. Koch, 1837 from China (Araneae, Amaurobiidae)

Lu-Yu Wang<sup>10</sup>, Muhammad Irfan<sup>10</sup>, Huan Zhou<sup>10</sup>, Zhi-Sheng Zhang<sup>10</sup>

1 Key Laboratory of Eco-environments in Three Gorges Reservoir Region (Ministry of Education), School of Life Sciences, Southwest University, Chongqing 400715, China

Corresponding author: Zhi-Sheng Zhang (zhangzs327@qq.com)

#### Abstract

Two new species of the genus *Amaurobius* are described from China: *A. guangwushanensis* **sp. nov.** ( $3^{\circ}$ ) from Sichuan and *A. wulongdongensis* **sp. nov.** ( $3^{\circ}$ ) from Shaanxi. With the addition of two new species, the number of *Amaurobius* species endemic to China now reaches four. Morphological descriptions, photos, and illustrations of copulatory organs, as well as a map of distribution records, are provided.

**Key words:** *Amaurobius guangwushanensis* sp. nov., *Amaurobius wulongdongensis* sp. nov., description, illustration, morphology, taxonomy

# Introduction

*Amaurobius* is the most species-rich genus in the spider family Amaurobiidae Thorell, 1869, comprising 65 species mainly distributed in North America and Europe (WSC 2023). Three European species [*A. erberi* (Keyserling, 1863), *A. fenestralis* (Ström, 1768) and *A. jugorum* L. Koch, 1868] were spread or introduced into west, central, or south Asia. Two Asian endemic species (*A. songi* Zhang, Wang & Zhang, 2018 and *A. spinatus* Zhang, Wang & Zhang, 2018) were described from the northern parts of Chongqing and Sichuan, China (Zhang et al. 2018), suggesting that there should be more local undescribed *Amaurobius* species from China.

Our recent exploration revealed two more new *Amaurobius* species from Sichuan and Shaanxi provinces of China. The four currently known endemic species of *Amaurobius* in China are distributed on the south slope of the Qinling Mountains and the eastern extension of the Hengduan Mountains. Most likely, the uplift of the Qinghai-Tibet Plateau influenced the origin and expansion of this spider group.

## Materials and methods

All specimens were examined, illustrated, photographed and measured using a Leica M205A stereomicroscope equipped with a drawing tube, a Leica DFC420 camera, and LAS (Leica Application Suite) software (version 4.6). Epigynes were cleared immersing them in pancreatin (Álvarez-Padilla and Hormiga 2007).



Academic editor: Ingi Agnarsson Received: 23 February 2023 Accepted: 26 June 2023 Published: 18 July 2023

ZooBank: https://zoobank. org/0D298638-F6F1-47C6-817E-D31B80C42C36

Citation: Wang L-Y, Irfan M, Zhou H, Zhang Z-S (2023) Two new species of *Amaurobius* C.L. Koch, 1837 from China (Araneae, Amaurobiidae). ZooKeys 1169: 307–315. https://doi. org/10.3897/zookeys.1169.102581

Copyright: © Lu-Yu Wang et al. This is an open access article distributed under terms of the Creative Commons Attribution License (Attribution 4.0 International – CC BY 4.0). Leg measurements are rendered here as: total length (femur, patella and tibia, metatarsus, tarsus). All measurements are in millimeters. Morphological terminology follows Zhang et al. (2008). All specimens examined here are deposited in the School of Life Sciences, Southwest University, Chongqing, China (SWUC). The following abbreviations are used in the text:

Somatic characters: ALE-anterior lateral eye; AME-anterior median eye; DTA-dorsal tibial apophysis; MOA-median ocular area; PLE-posterior lateral eye; PME-posterior median eye; RTA-retrolateral tibial apophysis.

Male palp: **Co**-conductor; **DTA**-dorsal tibial apophysis; **eDTA**-exterior branch of DTA; **E**-embolus; **iDTA**-interior branch of DTA; **MA**-median apophysis; **RTA**-retrolateral tibial apophysis; **TA**-tegular apophysis.

Epigyne: CD-copulatory ducts; FD-fertilization duct; LT-lateral teeth; ML-median lobe; S-spermathecae.

## Taxonomy

Family Amaurobiidae Thorell,1869 Genus Amaurobius C. L. Koch, 1837 (暗蛛属)

# Amaurobius guangwushanensis sp. nov. (光雾山暗蛛)

https://zoobank.org/73040AE8-D0AB-47FE-8390-23A719420954 Figs 1, 2, 5

**Type material.** *Holotype*  $\mathcal{J}$ : CHINA, **Sichuan Province**: Nanjiang County, Guangwu Mountain, Taoyuan, Sandaoguan, 32°41'15"N, 106°47'45"E, elev. 1377 m, 3.XI.2018, Z.S. Zhang, L.Y. Wang, T. Yuan, L. Yuan & P. Liu leg.; *Paratypes* ( $2\mathcal{J}\mathcal{J}12\mathcal{P}\mathcal{P}$ ):  $2\mathcal{P}\mathcal{P}$ , same data as holotype;  $2\mathcal{J}\mathcal{J}3\mathcal{P}\mathcal{P}$ , Guangwu Mountain, Taoyuan, Lianghekou,  $32^\circ40'31$ "N, 106°46'08"E, elev. 1002 m, 3.XI.2018, Z.S. Zhang, L.Y. Wang, T. Yuan, L. Yuan & P. Liu leg.;  $7\mathcal{P}\mathcal{P}$ , Guangwu Mountain, Taoyuan, 32°41'24"N, 106°47'39"E, elev. 1703 m, 16.V.2013, X.K. Jiang & D. Wang leg.

**Etymology.** The epithet refers to the type locality (Mt. Guangwu is read as Guangwushan in Chinese); adjective.

Diagnosis. Amaurobius guangwushanensis sp. nov. resembles A. wulongdongensis sp. nov. and A. spinatus in having a similar embolus in the male [Figs 1A, 2C; see Zhang et al. (2018), figs 4C, 5E, 6A], but can be differentiated by the thumb-shaped retrolateral tibial apophysis in ventral view (Figs 1B, 2D) [vs. somewhat rectangular in A. spinatus (see Zhang et al. 2018, figs 4D, 5F); semicircular in A. wulongdongensis sp. nov. (Figs 3B, C, 4D, E)]; interior branch of dorsal tibial apophysis somewhat triangular with a sharp pointed end in prolateral view in A. guangwushanensis sp. nov. (Figs 1D, 2C) and A. wulongdongensis sp. nov. (Figs 3A, 4C) [vs. about thumb-shaped with blunt tip in A. spinatus (see Zhang et al. 2018, figs 4C, 5E, 6C); tegular apophysis apex bifurcated in ventral view in A. guangwushanensis sp. nov. (Figs 1B, 2D) [vs. broad with small depression at the center in A. spinatus (see Zhang et al. 2018, figs 4D, 5F; somewhat rectangular in A. wulongdongensis sp. nov. (Figs 3B, 4D)]; median apophysis doorknob-shaped, tip slightly curved in ventral view in A. guangwushanensis sp. nov. (Figs 1B, 2D) [vs. hook-shaped in A. spinatus (see Zhang et al. 2018, figs 4D, 5F); doorknob-like, tip strongly curved in A. wulongdongensis sp. nov. (Figs 3B,



Figure 1. Amaurobius guangwushanensis sp. nov., male holotype (A–C) and female paratype (D–E). A left male palp, prolateral view B same, ventral view C same, retrolateral view D epigyne, ventral view E same, dorsal view.

4D)]; conductor apex narrow, apical margin about half the length of the embolus in retrolateral view in *A. guangwushanensis* sp. nov. (Figs 1C, 2E) [vs. broad, apical margin as long as embolus both in *A. spinatus* (see Zhang et al. 2018, figs 4E, 5G) and *A. wulongdongensis* sp. nov. (Figs 3C, 4E)]. The epigyne can be differentiated from the related species by the lateral teeth wider than long with round margin (Figs 1A, 2F) (vs. longer than wide with pointed tip in *A. spinatus* Zhang, Wang & Zhang, 2018, figs 4A, 5C, 6B); median lobe somewhat sheet-like in *A. guangwushanensis* sp. nov. (Figs 1A, 2F) [vs. oval in *A. spinatus* (see Zhang et al. 2018, figs 4A, 5C, 6B); round in *A. wulongdongensis* sp. nov. (Figs 3D, 4F)], lateral teeth c. 1/4 length of median lobe with round tip (Figs 1A, 2F) [vs. about half the length of median lobe, horn-like with tapering tip in *A. spinatus* (see Zhang et al. 2018, figs 4A, 5C, 6B); broad, semicircular, almost two times larger than median lobe in *A. wulongdongensis* sp. nov. (Figs 3D, 4F)].

**Description. Male** (holotype, Fig. 2A). Total length 3.12–3.52. Holotype (Fig. 2A) total length 3.52. Carapace 1.77 long, 1.26 wide; opisthosoma 1.59 long, 1.14 wide. Carapace yellowish. Cervical groove indistinct, radial furrows



Figure 2. Amaurobius guangwushanensis sp. nov., male holotype (A, C–E) and female paratype (B, F–G). A male habitus, dorsal view B female habitus, dorsal view C left male palp, prolateral view D same, ventral view E same, retrolateral view F epigyne, ventral view; G same, dorsal view.

distinct. Eye measurements and interdistances: AME 0.06, ALE 0.11, PME 0.10, PLE 0.11; AME-AME 0.04, AME-ALE 0.05, PME-PME 0.06, PME-PLE 0.09, ALE-PLE 0.04. MOA 0.31 long, front width 0.14, back width 0.29. Clypeus height 0.11. Chelicerae brown, with 4 promarginal and 4 retromarginal teeth. Endites

and labium yellowish brown, longer than wide. Sternum yellowish brown, with brown setae. Legs yellowish. Leg measurements: I 5.88 (1.62, 2.04, 1.35, 0.87); II 4.55 (1.30, 1.50, 1.05, 0.70); III 3.83 (1.11, 1.23, 0.93, 0.56); IV 5.00 (1.42, 1.66, 1.29, 0.63). Opisthosoma oval, dorsum yellowish, with six brown chevrons extending posteriorly, venter yellowish brown.

Palp (Figs 1A–C, 2C–E). Femur almost as long as cymbium. Patella with strong dorso-apical short spine. Tibia with large thumb-shaped retrolateral tibial apophysis, originating near the base of tibia. Dorsal tibial apophysis large, exterior branch of dorsal tibial apophysis longer than wide, c. 1/3 length of tibia, with wavey tip in retrolateral view, interior branch of dorsal tibial apophysis short, somewhat triangular with pointed end in prolateral view. Cymbium longer than both tibia and patellae, with retrolateral angular projection. Bulb oval, slightly longer than wide. Tegulum widest in middle part, tegular apophysis sclerotized, doorknob-like, about more than half-length of embolus, present at the center of bulb. Sperm duct visible in prolateral and retrolateral view. Embolus originating prolaterally, short, flat, with a round tip.

**Female** (Fig. 2B). Total length 4.29–4.36. One of paratypes (Fig. 2B) total length 3.84. Carapace 1.52 long, 1.05 wide; opisthosoma 2.40 long, 1.67 wide. Eye measurements and interdistances: AME 0.05, ALE 0.10, PME 0.09, PLE 0.10; AME-AME 0.05, AME-ALE 0.06, PME-PME 0.10, PME-PLE 0.12, ALE-PLE 0.04. MOA 0.30 long, front width 0.15, back width 0.28. Clypeus height 0.12. Legs yellowish brown. Leg measurements: I 4.14 (1.20, 1.46, 0.88, 0.60); II 3.90 (1.05, 1.14, 0.76, 0.50); III 3.09 (0.93, 1.07, 0.68, 0.41); IV 4.06 (1.21, 1.38, 0.97, 0.50). Opisthosoma color pattern as in male.

Epigyne (Figs 1D, E, 2F, G). Median lobe wider than long; lateral teeth with round margin. Copulatory ducts transverse, located between spermathecae. Spermethecae globular, spaced by about half of radius. Fertilization ducts originating posteriorly, slightly curved, directed laterally.

Distribution. Known only from the type locality, Sichuan, China (Fig. 5).

## Amaurobius wulongdongensis sp. nov. (五龙洞暗蛛) https://zoobank.org/A9F7F1B9-0D48-4895-85A6-32603A937A21 Figs 3-5

**Type material.** *Holotype* ♂: CHINA, **Shaanxi Province:** Lueyang County, Wulongdong Forest Park, 33°36'17"N, 106°18'34"E, elev. 1786 m, 17 October 2018, L.Y. Wang leg.; *Paratypes*: 3♀♀, same data as holotype.

**Etymology.** The specific name refers to the type locality; adjective.

Diagnosis. See diagnosis of Amaurobius guangwushanensis sp. nov.

**Description. Male** (holotype, Fig. 4A) total length 3.99. Carapace 2.12 long, 1.43 wide; opisthosoma 1.83 long, 1.34 wide. Carapace yellowish brown. Cervical groove and radial furrows distinct. Eye measurements and interdistances: AME 0.05, ALE 0.12, PME 0.11, PLE 0.11; AME-AME 0.05, AME-ALE 0.06, PME-PME 0.12, PME-PLE 0.13, ALE-PLE 0.05. MOA 0.36 long, front width 0.17, back width 0.34. Clypeus height 0.11. Chelicerae dark, with 4 promarginal and 4 or 5 retromarginal teeth. Endites and labium yellowish brown, longer than wide. Sternum yellowish, with brown setae. Legs yellow brown. Leg measurements:



Figure 3. Amaurobius wulongdongensis sp. nov., male holotype (A–C) and female paratype (D–E). A left male palp, prolateral view B same, ventral view C. same, retrolateral view D epigyne, ventral view E same, dorsal view.

I 7.40 (2.01, 2.50,1.87, 1.02); II 5.19 (1.37, 1.85, 1.23, 0.74); III 4.70 (1.40, 1.55, 1.16, 0.59); IV 5.83 (1.74, 2.02, 1.49,0.58). Opisthosoma oval, dorsum yellowish brown, with six brown chevrons, extending posteriorly, venter yellowish brown.

Palp (Figs 3A–C, 4C–E). Femur almost as long as cymbium. Patella with strong dorso-apically short spine. Tibia with large semicircular retrolateral tibial apophysis, originating near the base of tibia. Dorsal tibial apophysis large, exterior branch of dorsal tibial apophysis large, longer than wide, interior branch of dorsal tibial apophysis short, somewhat triangular. Cymbium longer than both tibia and patellae, with retrolateral angular projection. Bulb oval, slightly longer than wide. Tegulum widest in middle part. Conductor membranous, sheet-like. Median apophysis sclerotized, doorknob-like, about more than half-length of embolus, present at the center of bulb. Sperm duct visible in prolateral and retrolateral view. Embolus originating prolaterally, short, flat, with bifurcate tip.

**Female** (Fig. 4B). Total length 4.29–4.36. One of paratypes (Fig. 4B) total length 4.36. Carapace 2.05 long, 1.36 wide; opisthosoma 2.40 long, 1.75 wide. Eye measurements and interdistances: AME 0.07, ALE 0.13, PME 0.12, PLE 0.11; AME–AME



Figure 4. Amaurobius wulongdongensis sp. nov., male holotype (A, C–E) and female paratype (B, F–G). A male habitus, dorsal view B female habitus, dorsal view C left male palp, prolateral view D same, ventral view E same, retrolateral view F epigyne, ventral view G same, dorsal view.

0.07, AME-ALE 0.09, PME-PME 0.12, PME-PLE 0.14, ALE-PLE 0.06. MOA 0.37 long, front width 0.20, back width 0.35. Clypeus height 0.14. Leg measurements: I 5.39 (1.55, 1.84, 1.20, 0.80); II 4.23 (1.24, 1.45, 0.95, 0.59); III 3.60 (1.13, 1.03, 0.91, 0.53); IV 4.79 (1.46, 1.59, 1.19, 0.55). Opisthosoma color pattern as in male.



Figure 5. Distribution records of four Amaurobius species in China.

Epigyne (Figs 3D, E, 4F, G). Median lobe reduced, somewhat round, with lateral teeth large, semicircular. Copulatory ducts V-shaped, located between spermathecae. Spermathecae semiglobular, almost touching each other. Fertilization ducts originating postero-laterally.

Distribution. Known only from the type locality, Shaanxi, China (Fig. 5).

## Acknowledgements

The manuscript benefited greatly from comments by Ingi Agnarsson, Feng Zhang, Lina Maria Almeida-Silva, and Yuri Marusik. We thank Dong Wang, Xuan-Kong Jiang, Li Yuan, Tao Yuan and Piao Liu (SWUC formerly) for their assistance during the field work.

# **Additional information**

## **Conflict of interest**

The authors have declared that no competing interests exist.

#### **Ethical statement**

No ethical statement was reported.

#### Funding

This study was supported by the Science & Technology Fundamental Resources Investigation Program (Grant No. 2022FY202100), the Fundamental Research Funds for the Central Universities (SWU120051), the Chongqing Provincial Funding for Postdoc (cstc2021jcyj-bsh0196), Foreign Youth Talent Program Funding (QN2022168002L), the Investigation Project of Basic Science and Technology (2018FY100305), the fund on survey of spiders and insects from Yintiaoling Nature Reserve and the Fundamental Research Funds for the Central Universities to Zhou Huan (XDJK2017D097).

#### **Author contributions**

LY Wang and ZS Zhang collected spider individuals. LY Wang and M Irfan performed the experiments. LY Wang, M Irfan, and ZS Zhang wrote the paper.

### **Author ORCIDs**

Lu-Yu Wang <sup>©</sup> https://orcid.org/0000-0002-5250-3473 Muhammad Irfan <sup>©</sup> https://orcid.org/0000-0003-0445-9612 Huan Zhou <sup>©</sup> https://orcid.org/0009-0004-4507-6241 Zhi-Sheng Zhang <sup>©</sup> https://orcid.org/0000-0002-9304-1789

#### Data availability

All of the data that support the findings of this study are available in the main text.

# References

- Álvarez-Padilla F, Hormiga G (2007) A protocol for digesting internal soft tissues and mounting spiders for scanning electron microscopy. The Journal of Arachnology 35(3): 538–542. https://doi.org/10.1636/Sh06-55.1
- WSC (2023) World Spider Catalog. Natural History Museum Bern. http://wsc.nmbe.ch, [version 19.5] [Accessed on 23 February 2023]
- Zhang ZS, Zhu MS, Song DX (2008) Revision of the spider genus *Taira* (Araneae, Amaurobiidae, Amaurobiinae). The Journal of Arachnology 36(3): 502–512. https://doi. org/10.1636/H07-49.1 [N.B.: reprinted in Korean Arachnol. 24: 127–148]
- Zhang L, Wang LY, Zhang ZS (2018) The first record of *Amaurobius* C.L. Koch, 1837 (Araneae, Amaurobiidae) from China, with description of two new species. Zootaxa 4402(2): 363–372. https://doi.org/10.11646/zootaxa.4402.2.8