

Research Article

## Architecture, construction, retention, and repair of faecal shields in three tribes of tortoise beetles (Coleoptera, Chrysomelidae, Cassidinae: Cassidini, Mesomphaliini, Spilophorini)

Caroline Simmrita Chaboo<sup>10</sup>, Sally Adam<sup>20</sup>, Kenji Nishida<sup>30</sup>, Luke Schletzbaum<sup>40</sup>

- 1 University of Nebraska State Museum, Systematics Research Collections, W436 Nebraska Hall, University of Nebraska, Lincoln, NE, 68588-0514, USA
- 2 Blommekloof, Leeukloof, Mossel Bay, South Africa
- 3 Museo de Zoología, Universidad de Costa Rica, San José, & Estación Biológica Monteverde, Monteverde, Puntarenas, Costa Rica
- 4 USGS National Wildlife Health Center, 6006 Schroeder Rd, Madison, WI, 53711, USA

Corresponding author: Caroline Simmrita Chaboo (cchaboo2@unl.edu)



Academic editor: Michael Schmitt Received: 24 February 2023 Accepted: 10 May 2023 Published: 30 August 2023

ZooBank: https://zoobank.org/ C4092EE1-1D66-447F-8419-6AE3839B7756

**Citation:** Chaboo CS, Adam S, Nishida K, Schletzbaum L (2023) Architecture, construction, retention, and repair of faecal shields in three tribes of tortoise beetles (Coleoptera, Chrysomelidae, Cassidinae: Cassidini, Mesomphaliini, Spilophorini). In: Chaboo CS, Schmitt M (Eds) Research on Chrysomelidae 9. ZooKeys 1177: 87–146. https://doi.org/10.3897/ zookeys.1177.102600

**Copyright:** © Caroline Simmrita Chaboo et al. This is an open access article distributed under terms of the Creative Commons Attribution License (Attribution 4.0 International – CC BY 4.0).

#### Abstract

Animal constructions are the outcomes of complex evolutionary, behavioural, and ecological forces. A brief review of diverse animal builders, the materials used, and the functions they provide their builders is provided to develop approaches to studying faecal-based constructions and faecal-carrying in leaf beetles (Coleoptera: Chrysomelidae). Field studies, rearing, dissections, photography, and films document shields constructed by larvae in two species in two tribes of the subfamily Cassidinae, Calyptocephala attenuata (Spaeth, 1919) (Spilophorini), and Cassida sphaerula Boheman, 1853 (Cassidini). Natural history notes on an undetermined Cassidini species and Stolas cucullata (Boheman, 1862) (Tribe Mesomphaliini) outline the life cycle of tortoise beetles and explain terms. Commonly, the cassidine shield comprises exuviae onto which faeces are daubed, producing a pyramidal-shaped shield that can cover most of the body (up to the pronotum). In Cal. attenuata the larval shield comprises only exuviae, while in Cass. sphaerula, instar 1 initiates the shield by extending its telescopic anus to apply its own faeces onto its paired caudal processes; at each moult the exuvia is pushed to the caudal process base but remains attached, then more faeces are applied over it. The larva's telescopic anus is the only tool used to build and repair the shield, not mouthparts or legs, and it also applies chemicals to the shield. Pupae in Cal. attenuata retain part of the exuviae-only shield of instar VI, while pupae in Cass. sphaerula retain either the entire 5th instar larval shield (faeces + all exuviae) or only the 5th larval exuvia. The caudal processes are crucial to shield construction, shield retention on the body, and as materials of the central scaffold of the structure. They also move the shield, though the muscular mechanism is not known. Altogether the faecal + exuviae shields may represent a unique morpho-behavioural synapomorphy for the crown-clade Cassidinae (10 tribes, ~ 2669 species) and may have been a key innovation in subsequent radiation. Defensive shields and domiciles may help explain the uneven radiation of chrysomelid subfamilial and tribal clades.

Key words: Behaviour, Calyptocephala, camouflage, Cassida, debris-carrying, exuviae, faeces, pupae

#### Introduction

Animal constructions have fascinated humans for centuries (Smeathman 1781), perhaps as building is one hallmark for our own genus, Homo L. (Hominidae). Coral reefs, beaver dams, bird nests, and spider webs are familiar structures, long attracting research attention (von Frisch 1974). The size of animal constructions ranges from microscopic diatoms to coral reef formations visible from space; between their dams and lodges, beavers (Rodentia: Castor L.) construct the largest mammalian constructions (Larsen et al. 2021). Animals build with many endogenous and/or exogenous materials secreted or excreted by the maker, taken from other animals, or gathered from the environment. For example, silk is the most renowned animal fibre and is produced only by arthropods; it is very versatile, in cocoons, webs, and for knitting other materials together. Silk is even secondarily co-opted by other animals, including by humans. A bird's nest may be constructed from exogenous materials (e.g., plants, spider webs), lined with feathers (endogenous), or comprise salivary secretions (endogenous) as in nests of swifts (Aves: Apodidae) which humans eat as the birds' nest soup delicacy (Hobbs 2004; Marcone 2005). Constructions may be fashioned by an individual or a community to serve diverse purposes-nurseries and homes, traps, pantry, defences, dispersal devices, to mark territory, to aid communication (e.g., sexual and courtship displays), as physical and chemical barriers to deter predation and parasitism, or as camouflage to sneak up on prey (Hansell 2005). Constructions may be built to withstand wind, tide, and rain and some provide thermoregulation with air-conditioning. In the marine environment, decorator crabs (Hultgren and Stachowicz 2009), sea urchins (Ziegenhorn 2017) and sand mason worms (Carey 1987) build structures for camouflage, defence, and dwelling. A few books offer a primer into the diversity, roles, and engineering skills of animal architects (e.g., McCook 1907; von Frisch 1974; Hansell 1984, 2005, 2007; Turner 2000; Gould and Gould 2007; Arndt 2013). There are also children's book on this topic (Hutchins 1959; Dewey 1991; Nicholson 2003; Nassar and Blasco 2015; Butterfield and Hutchinson 2017). Building behaviours overlap with self-decoration behaviours where animals accumulate diverse debris on their body (see review of Ruxton and Stevens 2015).

This paper concerns building behaviours and structures of certain beetles (Coleoptera: Chrysomelidae). As context for our study, we briefly review animal builders to understand the range of study, research approaches, and implications of materials and architecture. Constructions are the outcomes of complex evolutionary, behavioural, and ecological forces. In his chapter on "Instinct", Darwin (1859: 247–256) discussed these elements in his experiments and analyses of "cell-making insect in the Hive-bee". His approach remains valid today: observe building repertoires, design elements, materials, and purposes. Comparative multi-level analyses of physiology, ecology, ontogeny, and history are required to understand these remarkable morpho-behavioural complexes. Constructions are rich opportunities to investigate the "extended phenotypes" of their builders (Dawkins 1982).

The study of constructions is well-developed in birds, mammals, spiders, and Hymenoptera, as evidenced by documentation of specimens (i.e., in museum collections), construction behaviours, materials, terminology, and functions. The best-known insect architects are those social insects where the entire colony builds a communal "city", Hymenoptera (ants, bees, wasps; Fabre 1915; Wheeler 1928; Sakagami and Michener 1962; Wilson 1971) and Isoptera (termites; Lüscher 1961; Krishna and Weesner 1969; Jeanne 1975; Mathews and Mathews 1978). Constructions can be prominent surface features or extend over a wide expanse and deep underground, where specialised chambers and corridors support different activities of members and enable precise control of ventilation, heat, humidity, and responses to invasions.

Many insects are solitary architects (Figs 1-15), but they are far less known, likely due to few collections of these builders, their constructions, scant study of building repertoires, and limited evolutionary analyses. Their constructions serve most commonly to protect the vulnerable egg, larval, and pupal stages that cannot easily escape an attack. These insect mothers invest in protective devices around eggs, including elaborate oothecae (e.g., Dictyoptera: Legendre et al. 2015) and nests (e.g., mud and clay cells of some Carabidae beetles: Claassen 1919; Brandmayr and Brandmayr Zetto 1974). In Scolytidae beetles, females oviposit on or under the bark and the larvae tunnel through the wood by eating the wood and creating galleries under bark. Many insects build protections for their sedentary pupae (e.g., golden cages in Curculionidae: Hyperinae: Hoffman 1954; Scherf 1964; Janzen 1979, 1983; Aiello and Stockwell 1996). Constructions may serve as nutritional shelters, protecting the individual and providing a food source; for example, in "cigar" weevils (Curculionidae: Rhynchitinae), females roll leaves into a dual-purpose nest that serves later as a paedotrophic chamber where larvae feed on the inner walls (Brandmayr 1992).

Building materials are as diverse as the builders. Materials may be secreted by the body (endogenous), extracted from the environment (exogenous), or a combination. Endogenous secretions can create colonial structures (e.g., a coral reef) or be carried by a single individual (e.g., molluscs in their secreted shells; McDougal and Degnan 2018). Integumental secretions of slime and wax occur in sawfly larvae (Hymenoptera; Eisner 1994). Homoptera species exhibit diverse constructions: wax tail filaments (Smith 2010), sugary 'lerp' domiciles of scale insects (Gilby et al. 1976), and liquid marbles in aphids (Kasahara et al. 2019). Salivary secretions can serve as a glue or a building material (e.g., salivary foam moulded into pupation chambers for Criocerinae leaf beetles (Tishechkin et al. 2011). Anal secretions form the elaborate oothecae in Dictyoptera (Legendre et al. 2015).

Exogenous building materials of insects are difficult to catalogue, being so diverse, and include both organic and inorganic materials. Soil is a readily available building resource; tiger beetle larvae (Coleoptera: Cicindelidae) build burrows in the ground from where they can grab prey; some add a mud turret to raise the entrance above possible flooding (Kinsley and Pearson 1981; Shivashankar et al. 1988). Mobile residences include ornate cases by Trichoptera larvae with small pebbles or leaves (Figs 1, 2), a behaviour even co-opted for insect-built jewellery (Fig. 3; Duprat 2020). Leaves are an abundant resource; simple leaf constructions can be achieved by targeted cutting to bend over the leaf (e.g., some Lepidoptera, Loefler 1996; some cassidine beetles, Prathapan et al. 2009). Complex leaf constructions require more time (e.g., rolled leaves of Attelabidae weevils, Vanin and Bená 2020; glued leaves of some Thysanoptera (thrips), Mound and Morris 1999). Many Lepidoptera caterpillars use their



Figures 1–15. Insects with backpacks. 1 Trichoptera: Caddisfly larvae in case (photograph: S. Marshall) 2 Trichoptera: Caddisfly larvae in case (photograph: S. Marshall) 3 Trichoptera: larva with its case, 1980–1994, gold, opal, pearls (case length = 1 inch; photograph: H. Del Olmo (from Hubert Duprat exhibition, ADAGP)) 4 Neuroptera: Chrysopidae: larva with exuvial debris (photograph: Masayuki Hayashi) 5 Hemiptera: Reduviidae: assassin bug, Singapore (photograph: Nicky Bay) 6 Hemiptera: Reduviidae, assassin bug, Costa Rica (photograph: Dieter Mahsberg) 7 Lepidoptera: Psychidae: caterpillar with its bag 8 Lepidoptera: Geometridae: Wavy Emerald Moth caterpillar, *Synchlora aerata* (Fabricius, 1798), covering itself with petals of its host, *Liatris* Gaertn. ex Schreb. sp. (Asteraceae) (photograph: Hope Abrams) 9 Lepidoptera: Nolidae: caterpillar of *Uraba lugens* Walker, 1866 with stack of their exuvial head capsules, Australia (photograph: Alan Henderson) 10 Coleoptera: Curculionidae, *Gymnopholus* Heller, 1901 weevil carrying lichen garden, Papua New Guinea (photograph: Adrian Tejedor) 11 Phasmida: stick insect, *Trychopeplus laciniatus* (Westwood, 1874), with exoskeleton modified to appear like moss, Costa Rica (photograph: Kenji Nishida) 13 Coleoptera: Curculionidae: weevil larva retains moist faecal coat (photograph: Filip Trnka) 14 Coleoptera: Erotylidae: larva of *Toramus* Grouvelle, 1916 with shield of exuviae held on setae (photograph: Takahiro Yoshida) 15 Coleoptera: Cassidinae: Cassidini: larva of *Microctenochira* Spaeth, 1926 undetermined species with shield of exuviae only (photograph: Kenji Nishida).

silk to sew twigs (Fig. 7) or leaves into tunnels, tubes, and portable cases (e.g., Psychidae bagworms, Sharp 1899; Frowhawk 1913; Bucheli 2002). Embioptera make silken galleries where they live (Büsse et al. 2015). Exogenous materials may be harvested from the droppings of other animals; for example, mollusc shells adopted or robbed by hermit crabs (Rodrigues and Rodrigo 2009) or homopteran wax stolen by Neuroptera (Eisner and Silberglied 1988). Some constructions are compound combinations of exogenous and endogenous materials (e.g., a bird's nest of twigs and spider silk, Hansell 2005).

Many solitary insect builders carry a 'backpack' with simple or compound 'debris' (endogenous, exogenous, environmentally acquired, organic or inorganic). Debris backpacks provide the builder with a mobile cloak that is usually assumed as a camouflage to avoid predators or a disguise for hunting (Cardé and Bell 1995; Tauber et al. 2014; Wang et al. 2016). Inorganic 'debris cloaks' of soil dust and small sand grains are found in insects (Odihiambo 1958; McMahan 1982, 1983a, b; Cardé and Bell 1995; Eisner 2003). In Trichoptera (caddisflies) constructed cases of silk may be decorated with sand, stones or shells and are used as retreats, homes, and to seine water for food (Wallace 1975; Wallace and Sherberger 1975; Otto and Svenson 1980; Ferry et al. 2013). The plaster bagworm (Lepidoptera: Tineidae) similarly makes a silken case that traps soil, lint, and even paint chip (Aiello 1979; Villanueva-Jimenez and Fasulo 1996). Organic debris cloaks can comprise small plant fragments such as twigs, leaves, trichomes, and wood fibres (Eisner et al. 2001). Nymphs of Reduvius personatus (L.) (Hemiptera: Reduviidae) are called "masked hunters" because of their debris of dust and soil (Fig. 6; Harz 1952; Dispons 1955; Cardé and Bell 1995; Cai et al. 2002; Weirauch 2006; Ramírez et al. 2013). Some Chrysopidae (Neuroptera) retain trichome debris covers (Smith 1922, 1923; Stitz 1931; Eisner et al. 1978, 2001; Eisner and Eisner 2000b; Anderson et al. 2003; Nakahira and Arakawa 2006; Haruyama et al. 2012). Remarkably, some insects grow a living backpack, a garden of lichens, algae, mosses, and fungi (Fig. 10; Gressitt et al. 1965, 1968). Gressitt (1977) used the term "epizoic symbiosis"; this camouflage resembles those insects that truly are morphologically adapted with a moss-like appearance that matches their lichen + moss covered habitat (e.g., Figs 11, 12).

Organic debris backpacks comprising insect exoskeletons (exuviae, cast skins) appear in diverse insects (Figs 4, 5, 9, 14, 15). These exuviae can be the builder's own castoffs or, more macabre, from their prey. Examples of the first type, retaining their own exuviae, are exhibited in some Lepidoptera and Coleoptera larvae. An Australian caterpillar retains a stack of its previous head capsules, giving it the nickname "mad hatterpillar" (Fig. 9; Lepidoptera: Nolidae; McFarland 1980; Pearson 2013). In Coleoptera, exuvial retention by larvae is known in some Erotylidae (Figs 14; Leschen 2003; Yoshida and Leschen 2020) and in Cassidinae (Fig. 15; Chaboo 2007). The second type of exuvial retention uses those of prey and has been described as a "corpse cover" (Brandt and Mahsberg 2002), a "corpse camouflage" (Stromberg 2012), and a "wolf in sheep's clothing" strategy (Eisner et al. 1978, after ancient rhetorical Greek and Italian fables, e.g., Basilakis in the 12th century; Beneker and Gibson 2016; Abstemius 14<sup>th</sup> century; the Bible (King James Bible Online 2023)). Some Chrysopidae larvae (Fig. 4; Neuroptera) carry the exuviae of their aphid prey, to fool aphid-tending ants (Hayashi and Nomura 2011). Many Hemiptera adults and nymphs retain corpse backpacks (Fig. 5; Odihiambo 1958; McMahan

1982, 1983a, b; Zeledón et al. 1973; Weirauch 2006), some adding dust and soil, for a mix of organic and inorganic debris. Corpse covers and debris cloaks may provide mechanical protection, from weather or predators (e.g., spiders, lizards), or permit aggressive mimicry towards their prey (e.g., ants, termites). Olfactory cues can mask the predator (Odihiambo 1958; Brandt and Mahsberg 2002; Jackson and Pollard 2007; Stromberg 2012) or may become a secondary signal that attracts enemies (Agelopoulus et al. 1995; Benelli et al. 2013; Huang et al. 2022).

Dung (faeces, frass, fecula) is an unconventional organic debris as faeces are typically considered unappetising and unhygienic waste products, vectors of pathogens, and an offensive by-product of animal metabolism. Most animals simply eliminate and avoid their waste, even finding creative ways to dispose of their faeces (e.g., mining insects, Frost 1942). Yet, faeces are a cost-free and readily available benefit of regular feeding. In Mammals, faecal piles function as territory markers (e.g., Stewart et al. 2001) and latrine sites (e.g., meerkats, Jordan et al. 2007). Counter-intuitively, faeces are a resource; indeed, humans have been using dung (Henry et al. 2016; Arranz-Otaegui et al. 2017; Smith et al. 2022) as a fertiliser since early agriculture, to burn as fuel, for plastering adobe walls and floors (faeces mixed with mud and twigs), in beauty facials ("Uguisu no fun", Moore 2001) and even in ancient (Ge 2000 [4<sup>th</sup> century]) and contemporary medical faecal transplants and enemas (e.g., Fecal Microbiota Transplantation or FMT; Eiseman et al. 1958; Zhang et al. 2012).

Dung beetles (Scarabaeidae) may be the most famous insects associated with faeces. Both dung beetles and burying beetles (Silphidae) use vertebrate dung for brood balls (Waterhouse 1974; Scholtz and Grebennikov 2004). Many fly groups are also renowned to use faecal habitats.

Many terms for insect faeces appear in the literature. Frost (1942) used 'faeces, fecula and frass' which have become widely used. Other terms are excrement (Hislop 1872; Scudder 1891; Muir and Sharp 1904; Blatchley 1924; Flinte and Valverde de Macédo 2004), excreta (Wood 1966; LeSage 1982; Jolivet and Verma 2002), and scat (Lécaillon 1896; Hinton 1981). Faeces are produced mainly by immature insects since most adult insects produce little wastes. Insect faeces can serve various purposes, such as adult aggregation, finding mates, brooding, or oviposition deterrent; they can signal pest issues. They can also recycle faeces in multiple ways; the process is sophisticated in social insects where faeces are used as a structural component of the nest and hive walls and as a substrate for growing fungi (Hansell 2005; Weiss 2006). In Coleoptera, faeces can serve for adult aggregation (Tenebrionidae: flour beetles; Suzuki 1985), to find mates (e.g., Cerambycidae: *Hylotrupes bajulus* (L.); Fettköther et al. 2000), brooding, or as an oviposition deterrent (e.g., weevils and cerambycids; Anbutsu and Togashi 2002; Addesso et al. 2007).

Insects in Coleoptera, Diptera, and Lepidoptera have evolved dung-carrying behaviours. Some Lepidoptera caterpillars retain a dry crust of their excreta (e.g., Noctuidae; Preston-Mafham and Preston-Mafham 2003: 406); others use their silk to knit their faeces into "frass chains" (resembling sticks) to build a retreat (e.g., Nymphalidae; Freitas and Oliveira 1992; Caldas 1994; Machado and Freitas 2001). Excremental cases are known in Diptera (e.g., Mycetophilidae; Holmgren 1907; Knab and van Zwaluwenburg 1918) and in Lepidoptera (e.g., Hesperiidae; Sharp 1899). Weevils (Curculionidae) exhibit diverse constructions: leaf-rollers (e.g., Attelabidae, Daanje 1975; Mathews and Mathews 1978), lichen-carriers (e.g., Fig. 13, Gressitt et al. 1965; Gressitt 1977; Jolivet 1988a), solid dung (e.g., Ceutorhynchini, Knab 1915), and liquid excremental covers (e.g., Cionini, Gonipterini; Knab 1915; Arzone and Meotto 1978; Janzen 1979, 1983; Crowson 1981; Aiello and Stockwell 1996). Other beetles may construct a faecal or faecal-fungal canopy or retreat (Leschen and Carlton 1993; Leschen 1994; Hanley 1996). It is important to note that faecal retention is most often exhibited by insect larvae and the behaviour has been interpreted mostly as armour, camouflage, or physical barrier to enemies (Weiss 2006).

Debris-carrying, including dung-carrying, is not simply just 'carrying' since individuals often exhibit specialised morphology associated with handling faeces (e.g., anal comb in some Lepidoptera, Frost 1919) or with retaining materials (special setation; Weirauch 2006; Skuhrovec et al. 2017) to build, carry, wear and even repair structures. Enhanced survivorship is often assumed, and in cases where tested, the adaptive value of debris such as frass and faeces has been demonstrated.

In this paper, we focus on faecal-recycling behaviours in Chrysomelidae (leaf beetles), one of the largest clades of beetles with > 40,000 species (Leschen and Beutel 2014). Chrysomelids use their faeces as a biomaterial for constructions and self-decoration behaviours that serve as defensive coats, mobile debris shields, and protective domiciles. Such faecal-based constructions appear as a striking pattern within Chrysomelidae, diagnosing some large subfamilies and appearing also in some small clades.

In general, leaf beetles exhibit diverse building behaviours, including oothecae with multi-layered colleterial secretions (e.g., some Cassidinae), faecal covers (Kalaichelvan and Verma 2000), or with stomach regurgitate (Jolivet and Verma 2002), larval galls (e.g., Sagrinae, Reid and Beatson 2019), and pupation chambers of soil, sand (e.g., some Galerucinae, Prathapan and Chaboo 2011), faeces (Cryptocephalinae; Brown and Funk 2005), or salivary 'foam' (e.g., some Criocerinae, Tishechkin et al. 2011). Bruchine adults build walls within seeds to inhibit fighting (Mano and Toquenaga 2008). Simple leaf shelters are made by larvae and adults of *Leptispa* Baly, 1858 (Cassidinae: Leptispini; Prathapan et al. 2009). Many chrysomelid mothers coat eggs with glandular and excremental applications, often mixed with anal and buccal secretions, and then may cover eggs further with plant pieces or oothecal membranes or faecal cases (Muir and Sharp 1904; Fiebrig 1910; Prevett 1966; Hinton 1981; Jolivet and Verma 2002; Müller and Hilker 2004).

The faecal-based constructions of Chrysomelidae are not a diffuse pattern but are taxonomically focused, are ancient, dated at least 45 million years ago (Chaboo and Engel 2008; Chaboo et al. 2009), and may have three or four independent origins given simple mapping on recent phylogenetic hypotheses of the family (Figs 16–18): within the subfamily Cassidinae; the *Blepharida*-group within the subfamily Galerucinae; Criocerinae; and in the sister subfamilies Cryptocephalinae + Lamprosomatinae. Within Cassidinae (6,320 species in 37 tribes), faecal constructions diagnose a derived monophyletic clade of ten tribes (= the tortoise beetle tribes) where most larvae use their exuviae and/or faeces to build shields over the body (Figs 15, 19–26; Chaboo 2007); these shields may be retained in pupae of some species (Fig. 25). Cryptocephalinae + Lamprosomatinae (~ 6000 species) form a well-accepted clade, called Camptosomata,



**Figures 16–18.** Recent phylogenetic hypotheses of Chrysomelidae subfamily relationships, redrawn by L. Schletzbaum from original sources **16** Reid (2000) (morphology-based) **17** Gómez-Zurita et al. (2007, 2008) (sequence-based) **18** Nie et al. (2020) (sequence-based). Other chrysomelid hypotheses to compare are Farrell (1998), Hunt et al. (2007), and Zhang et al. (2018, 2022). These available hypotheses are based on less than 1% taxon sampling of clade diversity. Subfamilies in bold font exhibit major patterns of faecal-based constructions. Alticinae (flea-beetles) is now regarded as the tribe Alticini within Galerucinae, so the faecal-retaining *Blepharida*-group is recognized now within Galerucinae. Only a single species in *Phola* Weise, 1890 (Chrysomelinae) has been reported to retain faeces (Chen 1964, 1985) so it is not a major pattern.

that is distinguished by a complex behaviour where females construct a faecal case around the single eggs and the natal larva keeps that egg case as a rigid portable home (Figs 27-34; Lawson 1976). This faecal case (= faecal bag, scatoshell) becomes the nucleus that the larva continues expanding with their own faeces; eventually the pupa inherits this construction as their pupation chamber (Brown and Funk 2005; Chaboo et al. 2016). Criocerinae is a smaller subfamily of ~ 1400 species whose larvae maintain a wet or semi-solid mass of their faeces directly on their back (Figs 35-37; Vencl et al. 2004). The Blepharida-group comprises ~ 21 genera (D'Alessandro and Biondi 2023) within the hyperdiverse Galerucinae (7145 species: Lingafelter and Konstantinov 2000; Nie et al. 2017); this group is distinguished by larvae that keep a single faecal strand held over the body (Fig. 38; Furth 1982, 2004; Furth and Lee 2000) or many faecal pellets directly on the dorsum (Figs 39, 40; Prathapan and Chaboo 2011; Calcetas et al. 2023). In Chrysomelinae (~ 3000 species), larval faecal tubes have been reported only in Phola octodecimguttata (Fabricius, 1775) (Chen 1964, 1985) and is a minor building pattern within this large subfamily. It is unclear at present what could be trends in innovations and maternal investments in oviposition site selection, and oothecal and egg-case construction. These chrysomelid constructions and body coats appear to be composites of endogenous and exogenous materials (Table 1), with their own faeces, exuviae, plant materials (trichomes, bark, twigs, decomposing fragments), chemical (plant or animal made), and even fungi. The endogenous materials can include faeces, anal, buccal, and other glandular products, and exuviae. The roles of each material are unknown.

Comparative surveys of the architectures of leaf-beetle constructions, detailed study of morphology associated with construction, retention and repair, and study of constructing behaviours are all needed to elucidate the apparent multiple origins and diversification of these structures. Experimental studies



Figures 19–26. Shields of larvae and pupae in four tribes of Cassidinae (Coleoptera: Chrysomelidae) 19 Hemisphaerotini: *Hemisphaerota* Chevrolat, 1836 20 Ischyrosonychini: *Physonota* Boheman, 1854 21 Cassidini: *Agroiconota bivittata* (Say, 1827) 22 Aspidimorpha sanctaecrucis (Fabricius, 1792) 23 Cassidini: undetermined sp. 1 24 Cassidini: undetermined sp. 2 from Africa, collected by C.S. Chaboo 25 Cassidini: undetermined sp. 3 pupa from Brazil, collected by D. Yanega 26 Cassidinae: Undetermined sp. 4 Costa Rica, collected by K. Nishida. Darkened sections = faeces. Redrawn by L. Schletzbaum from original sources or from specimens.



Figures 27–34. Faecal structures of larvae and pupae in Cryptocephalinae (Coleoptera: Chrysomelidae) 27 Adiscus taiwanus 28 Chlamisus sp. 1 29 Chlamisus sp. 30 Coenobius taiwanus 31 Cryptocephalus trifasciatus 32 Fulcidax 33 Neochlamisus 34 Lamprosomatinae. Redrawn by L. Schletzbaum from original sources.

are needed to test proposed hypotheses about the adaptive significance of faecal-based constructions. Such data can explain if and how these unusual faecal constructions could have influenced chrysomelid diversification, producing such uneven subfamilial species diversities. Caroline Simmrita Chaboo et al.: Construction behavior in Cassidinae beetles



Figures 35–37. Faecal-based coats of larvae in Criocerinae (shining leafbeetles). **35** Criocerinae sp. 1 **36** Criocerinae sp. 2 **37** *Lema hexastigma.* Redrawn by L. Schletzbaum from original sources.



Figures 38–40. Faecal-based structures and coats maintained by larvae in the *Blepharida*-group (Galerucinae: Alticini; flea beetles). **38** *Blepharida* sacra **39** *Diamphidia* sp. **40** *Podontia* sp. Redrawn by L. Schletzbaum from original sources.

In Cassidinae (~ 6000 species), faecal-based construction behaviour is a significant macroevolutionary event with a radiation of ~ 2700 species after its origin (Chaboo 2007), assuming a single evolutionary origin of faecal shield construction. This crown-clade is called "tortoise beetles" and is distinguished by a unique combination of morphology and behaviours: the larvae are exophagous (or ectophagous) and have paired caudal processes (= urogomphi) onto which they build and carry a debris shield (Figs 15, 19–26) moulded from their own exuviae and faeces. These larvae use a telescopic anus to apply faeces to the shield. The shield can be moved over the body like an umbrella or parasol (Fiebrig 1910; Takizawa 1980; Chaboo 2007; Flinte et al. 2009; Świętojańska 2009). Some exceptions, absence of shield retention, are also known but these appear to be secondary losses given the current phylogenetic views. Tortoise beetles comprise ~ 2700 species classified into ten tribes: Basiprionotini Hincks, 1952; Cassidini Gyllenhal, 1813 (now includes Aspidimorphini and Charidotini); Dorynotini Monrós & Viana, 1949; Eugenysini Hincks, 1952;

Table 1. Architects and materials used for faecal-based shields in subfamilies of Chrysomelidae: Cassidinae (Chaboo 2007; Świętojańska 2009), Criocerinae (Vencl et al. 2004), Cryptocephalinae and Lamprosomatinae (Chaboo et al. 2016), and Galerucinae (Prathapan and Chaboo 2011). Comparison of life stages, materials of larval/pupal faecal-based cases and shields, and larval body parts for construction. Key: + = present; - = absent; ? = unknown.

Feature		Cassidinae: 10 tribes, tortoise beetles	Chrysomelinae: Phola sp. <sup>8</sup>	Criocerinae	Galerucinae: Alticini: Blepharida-group	"Camptosomata"	
						Cryptocephalinae	Lamprosomatinae
Stage	Mother	-	?	-	+	+	+
	Egg	+/-	?	-	+/-	+	+
	Larvae	+/-	+	+	+	+	+
	Pupae	+/-	?	-	?	+	+
Morphology Larval/pupal material	Endogenous						
	Faeces	+/-	+	+	+	+	+
	Exuviae	+/-	?	+1/-	-	_	-
	Chemicals	+/-	?	+/-	+/-	?	?
	Waxes	?	?	?	?	?	?
	Saliva	?	?	?	?	?	?
	Regurgitates	?	?	?	?	?	?
	Exogenous						
	Soil	-	?	-	-	+/-	+/-
	Debris	_	?	+	_	+/-	+/-
	Trichomes	_	?	-	_	+/-	+9/-
	Leaf fragments, fresh	_	_	_	-	+/-	?
	Leaf fragments, decomposed	-	_	-	-	+/-	?
	Bark, twigs	_	_	-	-	+5/-	?
	Chemicals	+/-	?	+/-	+/-	?	?
	Fungi	+7/-	?	-	-	+/-	?
	Micro-organisms	?	?	?	?	?	?
	Abdomen	+	?	-	-	+	+
	Caudal Process	+	?	-	_	-	-
	Setation	?	?	?	?	?	?
	Anus	+	?	+	+	+	+

<sup>1</sup>Lema jacobiana Linell includes exuviae in faecal coat (Kaufmann 1967). <sup>2</sup>Waxes were reported in *Saxinis saucia* LeConte, 1857 (Spruyt 1925) and in *Fulcidax bacca* (Bokerman 1964). <sup>3, 4</sup>Neochlamisus use saliva mixed with faeces in cases (Briggs 1905; Brown and Funk 2005). <sup>5</sup>*Fulcidax cuprea* (Klug, 1824) females integrate bark in egg-cases (Bokerman 1964). <sup>6</sup>*Podontia lutea* (Olivier, 1790) include exuviae in faecal coat (Takizawa 1978). <sup>7</sup>*Fungi was* found in larval shields of *Laccoptera* (*Sindia*) *sulcata* (Olivier, 1808) (Rane and Ghate 2005) and *Cyrtonota sericinus* (Erichson, 1847) (Cedeño-Loja and Chaboo 2020); mycelia can be seen in other shields (e.g., *Canistra*, Flinte et al. 2009). <sup>6</sup>*Phola* Weise, 1890 (Chrysomelinae) reported by Chen (1964, 1985). <sup>9</sup>Described in Chaboo et al. (2008). Questions about Lamprosomatine cases arise due to their close relationship to Cryptocephaline cases that suggests possibly similar materials and constructions.

Goniocheniini Spaeth, 1942; Hemisphaerotini Monrós & Viana, 1951; Ischyrosonychini Chapuis, 1875; Omocerini Hincks, 1952; Mesomphaliini Chapuis, 1875; and Spilophorini Chapuis, 1875. The systematics of Cassidinae has been dynamic in the last 15 years and there are some disagreements on classification; we briefly discuss some issues relevant to our taxon focus in 'Materials and methods' below.

An obvious question is "How do tortoise beetles build their shields?" We address this specifically in three tribes Cassidini, Mesomphaliini, and Spilophorini. We aim to understand how the architecture is achieved and what morphological equipment is involved. We examine the materials, building processes, retention and repair of faecal constructions, and their inheritance from one instar to the next. Still images and short films document building behaviours and dissections help puzzle out how the materials are fitted together. We briefly review explanatory hypotheses for possible functions of cassidine shields. To date, the only study of chrysomelid faecal-constructing behaviour has been in *Neochlamisus* Karren, 1972 in the hyperdiverse subfamily Cryptocephalinae (~ 6000 spp.) by Brown and Funk (2005). Our study complements that work. Finally, we discuss the evolutionary-phylogenetic context to frame future research on chrysomelid faecal-based constructions.

## Materials and methods

We compare architectures and study construction behaviours in four species in three tortoise beetle tribes (derived Cassidinae, *sensu* Chaboo 2007) based on fieldwork in Costa Rica (2011–2021) and South Africa (2021–2022). To minimise confusion of species, we use these genus-name abbreviations: *S. cucullata* for *Stolas cucullata* (Boheman, 1862) (tribe Mesomphaliini), *Cassidini* undet. sp. 4 for an unidentified species (tribe Cassidini), *Cal. attenuata* for *Calyptocephala attenuata* (Spaeth, 1919) (tribe Spilophorini), *Cass. sphaerula* for *Cassida sphaerula* Boheman, 1854 (tribe Cassidini).

## **Research approach**

First, we introduce concepts of life stages, structures and morphology involved in Cassidinae construction by reporting the natural history of *S. cucullata* and Cassidini undet. sp. 4 (three undet. species of Cassidini are illustrated in Figs 23–25). Second, we report on shield construction in two focal species, *Cal. attenuata* (Spilophorini) and *Cass. sphaerula* (Cassidini). Third, we compare and contrast the construction behaviours and resultant architectures, contextualising our findings within Cassidinae and Chrysomelidae. Our focal taxa here are:

1. Tribe Mesomphaliini: Stolas cucullata (Boheman, 1862) (Figs 41-44). Observations, photographs, and specimen collection were made at COS-TA RICA: Cartago Province, Orosi, Tapantí National Park, 9°45'38.63"N, 83°47'3.98"W, 1280 m ele., 24-vii-2011, coll. Kenji Nishida. Oviposition was observed and photographs were taken also on 31-vii-2011 by KN. The live beetles were observed in a cloud forest habitat, along an open trail. The host plant was not determined initially as the female was flying then and landed on vegetation. Later, oviposition was observed, the host plant could be identified, and the hatched larvae were followed in the field on that host plant. Identifications: there are only five or six Stolas Billberg, 1820 species in Costa Rica. The red marginal spot on the black elytra is found in adults of three species: one spot in Stolas cucullata (Boheman, 1862), two spots in Stolas costaricensis (Champion, 1893), and two spots in Stolas lebasii (Boheman, 1850). Świętojańska (2009) indicated that juvenile stages are known for just five of the 187 recognised species of Stolas: Stolas chalybea (Germar, 1824), Stolas festiva (Klug, 1829), Stolas implexa (Boheman, 1850), Stolas lacardairei (Boheman, 1850), and Stolas lineaticollis (Boheman, 1850). Vouchers are deposited in the Museo de Zoología (MZUCR), Universidad de Costa Rica, San Pedro de Montes de Oca, Costa Rica. *Stolas cucullata* was identified by CSC using the online catalogue of Borowiec and Świętojańska (2002–present). The latter indicates that the type specimen was collected by Warszewicz in Panama: Veraguas, and that Bolivia is an inaccurate locality; the type is supposed to be in the J. Weise collection, Zoologisches Museum, Humboldt Universitat, Berlin, Germany, but it cannot be located (Bernd Jaeger, pers commun.). This species is distributed in Costa Rica and Panama (Chaboo 2003). Plant: this was identified as *Neomirandea angularis* (B.L.Rob.) (Asteraceae) by B. Haber, Monteverde. This is a new host record; Windsor et al. (1992) previously recorded *Neomirandea homogama* (Hieron.) Rob & Brett. as a host of *S. cucullata* in Panama.

- Tribe Cassidini: Cassidini undet. sp. 4 (Figs 45–50). All life stages have been documented on the host plant by KN in COSTA RICA: Puntarenas Province, Monteverde, 2016. Identifications: We await further study for more conclusive species determination. Plant: *Chione sylvicola* (Standl.) W. C. Burger (Rubiaceae) was identified by B. Haber, Monteverde. This is a new host record for Cassidinae; only six species of Cassidinae (4 Cassidini, 2 Notosacanthini) have been reported on Rubiaceae hosts (Borowiec and Świętojańska 2002–present; Monteith et al. 2021).
- 3. Tribe Spilophorini: Calyptocephala attenuata (Spaeth, 1919 (Figs 51-58). Live populations were studied on four Smilax spp. (Smilacaceae) at COSTA RICA: Puntarenas Province, Monteverde, 1530m, 10°19'08.5"N, 84°48'32.0"W, periodically over 2014–2020, Author KN led field studies and published some natural history reports (Nishida 2014, 2015; Nishida et al. 2020). Beetles were identified by CSC. Vouchers are deposited in the Museo de Zoología (MZUCR), Universidad de Costa Rica, San Pedro de Montes de Oca, Costa Rica. The four species of Smilax host plants were identified by L. Ferrufino-Acosta. The life cycle of Cal. attenuata includes six larval instars and the pupa; all carry exuvio-faecal shields on paired caudal processes (Figs 54-57; = urogomphi). The shield is composed solely of exuviae of previous instars and no faeces. Adults exit the pupal exuvia by splitting the anterior margin of the pupa (Figs 57, 58). Interestingly, adults eclose partly but stay in situ for 2-3 days, hardening up, before exiting completely from the pupal exuvia. Photographs of juveniles (Figs 59, 60) of an unidentified third species from Ecuador were sent by photographer Eerika Schulz to author CSC in 2018 who identified the species as belonging to Spilophorini. Pedro Ríos Guayasamín and students, Universidad Estatal Amazónica, are studying this population on an Orchidaceae host, and will send specimens to CSC for identification.
- 4. Tribe Cassidini: Cassida sphaerula (Figs 65–89). Author SA conducted fieldwork in 2021–2022, observing populations of an endemic beetle on its host, Arctotheca prostrata (Salisb.) Britten (Asteraceae) in various locations around Mossel Bay, South Africa, 33°57'58"S, 22°5'24"E. Adam et al. (2022) reported on natural history. The life cycle has five larval stages, all with exuvio-faecal shields, and the pupa that may carry shields of exuviae only or shields of exuviae and faeces. Identifications. Beetles were identified by CSC and confirmed by E. Grobbelaar. Vouchers. These are deposited at South Africa National Insects Collection (SANBI) and loaned to CSC.

## Permits

Resolutions # 039-2013-SINAC; # 080-2013-SINAC; SINAC-SE-GASP-PI-R-058-2014 (3 total) were issued by Ministerio de Ambiente y Energia (MINAE), Costa Rica. These allowed research/collecting and specimen export. Permits were issued by Sistema Nacional de Áreas de Conservación, Ministerio de Ambiente y Energía (MINAE), San José, Costa Rica, with assistance of Lourdes Vargas-Fallas and Javier Guevara-Sequeira.

## Photography and film

Various digital cameras were used for photography and filming KN used Nikon Coolpix E4500, Canon EOS 7D, Olympus STYLUS TG-4 Tough, and Sony  $\alpha$ 7S. The movie of *Calyptocephala* moulting was filmed with at 4K movie resolution using Sony's digital camera " $\alpha$ 7S" with Canon MP-E65mm F2.8 1–5× Macro Photo lens. SA used a Panasonic DMC-FZ200 camera plus a Raynox macroscopic lens M-150 and live individuals were observed with a Zeiss stereoscopic microscope plus a Dino-Lite eyepiece digital microscope/camera. CSC used a Basler camera attachment on a Nikon SMZ800 microscope. Photo editing was done in Paint.net or Photoshop. LS did the illustrations in Adobe Photoshop and Adobe Illustrator.

## Taxonomic names, morphological terms, phylogenetic characters

We follow the Cassidinae classification and taxonomic names of Staines (2015) and Borowiec and Świętojańska (2002–present). We follow Chaboo (2007) for morphological terms and phylogenetic character numbers discussed herein (see more discussion under Phylogeny below). Other group-taxon names for beetles follow Bouchard et al. (2011).

## Terminology

This section provides definitions of entomology and cassidine larvae terms that are used to describe the shield construction process. In addition to our illustrative plates, shields can be found in these other synthetic sources: Takizawa (1980), Chaboo (2007), Flinte et al. (2009), and Świętojańska (2009).

In holometabolous insects, larvae instars are demarcated by ecdysis events. Since the process of ecdysis lasts a few seconds (Hemimetabola juveniles are called nymphs), in practice, entomologists recognise the new instar starting when the previous instar's exuvia separates from the epidermal cells of the new instar's exoskeleton (a process called *"apolysis"*). The section aims to help readers understand the interactions between processes and parts involved in shield formation, described in the 'Results' section.

## Exuviae

We use "exuvia" (singular) and "exuviae" (plural) for the exoskeletons ("skins") shed at ecdysis following Snodgrass (1935) and Chapman (1982: 519). Entomologists have co-opted the Latin terms that translates as "things stripped off"

(Latin is Simple 2023). Schuh (1989) recognises "exuviae" only. Exuvium is linguistically incorrect and hardly used. "Pharate" is used to describe when the exuvia is retained and encloses the teneral insect (Chapman 1982: 518); in tortoise beetles, the exuvia is retained without enclosing the emerging larvae, so the latter is not pharate. In *Aproida* Pascoe (tribe Aproidini), the pupa is suspended from the larval exuvia (Monteith 1970), probably by everted forgut cuticle lining as in some other beetles (Frania 2011); this is unlike the exuvial retention of tortoise beetles. We describe some shields below as exuvia-only (single exoskeleton of Instar II larvae and pupae?) or exuviae-only (with more than one exoskeleton).

## Caudal process

'Urogomphus' (singular) and 'urogomphi' (plural) are used widely in insects, referring to the paired spine-like dorsal projections originating from the 9th abdominal tergite of many larvae (Duporte 1977; Schuh 1989). They are not homologues of cerci, projections of the 11th abdominal segment, nor are they universally homologous across Insecta. Within Chrysomelidae, the ten tribes of tortoise beetles (= the crown clade) in the Cassidinae share a character of larvae having paired projections (a few species secondarily exhibit a single process, Chaboo 2007: char. 11). Plesiomorphic 'hispine' larvae lack these dorso-caudal processes but some mining and cryptic feeders have their 9th abdominal tergite modified, heavily sclerotised and concave, into a "urogomphal plate" (Maulik 1931; Chaboo 2007). The tortoise beetle processes are also not morphologically homologous with such processes in juveniles of other chrysomelid subfamilies (e.g., urogomphi in Chrysomelinae larvae sensu Reid 1992a, b), other beetle families, or other insects. In the chrysomelid literature, the cassidine caudal processes have been called many terms: posterior spikes (Kershaw and Muir 1907), anal furca (Buzzi and Miyazaki 1992), supra-anal furca (Heron 2007), and supra-anal processes (Borowiec and Świętojańska 2014). Some labelled figures of Cassidine structures are in Chaboo (2007: fig. 18F of larva and 19C of pupa) and Adam et al. (2022: figs 21-25). We use the term caudal processes here for Cassidinae but indicate "(= urogomphi)" in discussions below to remind readers who may be more familiar with that term.

## Scolus, scoli

We follow the Torre-Bueno Glossary of Entomology (Schuh 1989) using both singular and plural terms for lateral projections from the thorax and abdomen of the larval and pupal body. Scoli are not homologous with tergal-originating caudal processes (= urogomphi). Cassidinae larvae and pupae may have scoli on the pronotum, metathorax and abdomen; these are unbranched and can be simple, spinose or have short setae.

**Anus** (Figs 45–50). Tortoise beetle larvae have a unique anus, sub-terminally-opening, muscular, extensible, and highly manoeuvrable unlike other chrysomelid larvae, which have a simple pore-like anus. The telescopic anus of Cassidinae likely represents the plesiomorphic abdominal segments X–XI. The anus is moved by peristaltic movements (Gómez et al. 1999).

**Shield** (Figs 19–26). This is attached to the caudal processes and held over the cassidine larval and pupal body, sometimes reaching over the pronotum.

Annex, parasol, shield, and umbrella (Jolivet and Verma 2002) have been used to describe the structure. Tortoise beetle shields have been called other names: larval clothing (Muir and Sharp 1904), 'kotanhang''' (= faecal appendage; Fiebrig 1910), faecal mask (Engel 1935), faecal shield (Eisner et al. 1967), faecal pad (Hawkeswood 1982), and exuvio-faecal annex (Buzzi 1988). Buzzi's (1988) term is precise about materials (i.e., exuvio-faecal) and does not imply function (i.e., annex is neutral compared to shield). "Annex" is probably the best term, however at this time, shield has become so widely used in the literature and concurs with the experimental work demonstrating its functions, thus we will retain this term.

#### Faeces, frass, fecula

Many terms for insect excrement appear in the literature: excrement (Hislop 1872; Scudder 1891; Blatchley 1924; Muir and Sharp 1904; Flinte and Valverde de Macédo 2004), faeces (Snodgrass 1935), 'faeces, fecula and frass' (Frost 1942), excreta (Wood 1966; LeSage 1982; Jolivet and Verma 2002), scat (Lécaillon 1896; Hinton 1981), and fecula (Gómez et al. 1999). In this context, the term "faeces" is used to refer to waste substances emerging at the anus (Snodgrass 1935; Chapman 1982; Schuh 1989), which should not be confused with other exudations (honey dew, spittle froth, glandular and salivary secretions, etc.). Terms like merdigery (Jones 1994) and psammophory (Bameul 1989) refer to faeces and sand, respectively.

#### Experiments to unravel shield architecture

Chaboo's (2007) subfamily phylogenetic study of Cassidinae determined that the exuvio-faecal shield represents a unique morpho-behavioural complex supporting monophyly of tortoise beetles (10 tribes, ~ 2700 species). The majority of these species has exophagous larvae that retain the cast exuviae and apply their own faeces to build the distinct globular or pyramidal structure. This is held on their caudal processes and can be moved about. Within this crown clade, a few species do not retain a shield and we will discuss this pattern in our evolutionary discussion below.

Typically, a tortoise beetle female may deposit faecal pellets onto eggs or oothecae, but it is the instar 1 that initiates the shield with its faecal material. Instar II retains the exuvia of instar I on its own caudal processes and attaches its own faeces. For *Cass. sphaerula*, we dissected shields to understand how it is fitted and held together.

*Calyptocephala attenuata* (Spaeth, 1919) (Figs 51–58). Observations and imaging were made over a 2-yr period by KN; specimens collected by KN were studied by CSC by dissection and imaging to determine how the shields are held together. The moulting process was filmed in Costa Rica for some Japanese television nature documentaries (Yamamoto 2018, 2020), assisted by KN; KN also photographed published nature notes (Nishida 2014, 2015; Nishida et al. 2020). We describe the moulting process and shield architecture under 'Results'. The moulting process exhibits active and quieter periods; to ease description, we use 'phases' and timing to describe the sequence filmed.

*Cassida sphaerula* (Figs 65–89). Given access to a large population, we were able to access many live specimens for various manipulations indoor to document

the construction, enlargement, and transfer of the exuvio-faecal shield from one instar to the next, then to the pupa. We observed multiple larvae of various instars indoors, maintained in plastic containers at ambient temperatures and light and supplied daily with fresh host leaves. We followed these larvae until the emergence of adults. We studied the effects of shield removal in three experiments as follows:

- Experiments 1–2: remove the shield entirely, sliding the structure off the caudal processes and leaving the live larva naked.
- Experiment 3: abrade only the faecal part of the shield, leaving the exuviae *in situ* on the caudal processes.

We photographed and filmed these individuals at 2-hr time (T) intervals to capture the initiation, expansion, and maintenance of the exuvio-faecal shield. We paid attention to larval movements and pupation. Based on these observations, dissections, and imagery, we describe the shield architecture, shield construction and reconstruction, and the moulting process under 'Results' below.

#### **Phylogeny relations**

Faecal constructions are considered here at two levels, first in Chrysomelidae and second in Cassidinae. For Chrysomelidae (Figs 16–18), we only present the broad pattern of faecal constructions and their possible role in sub-familial diversification, so we simplify the original sampled taxa to subfamily names to show the overall topology of recent major analyses (Reid 1995, 2000; Gómez-Zurita et al. 2007, 2008; Nie et al. 2020). We do not discuss the underlying evidence and premises supporting these topologies.

For Cassidinae, subfamilial monophyly is well-supported in hypotheses of chrysomelid evolutionary relationships (Farrell 1998; Reid 1995, 2000; Gómez-Zurita et al. 2007, 2008; Hunt et al. 2007; Haddad and McKenna 2016; Song et al. 2017; Nie et al. 2020). The internal relations are not fully settled. Cassidinae were historically treated as two subfamilies, Hispinae ("hispines") and Cassidinae ("tortoise beetles"), but are now recognised as a single subfamily, Cassidinae sensu lato, based on life history, morphological, and molecular evidence (Borowiec 1995; Hsiao and Windsor 1999; Chaboo 2007); other phylogenetic studies target subsets of tribes. Two online catalogues are available for "hispines" (plesiomorphic Cassidinae, 3,371 species in 24 tribes; Staines 2015) and for "tortoise beetles" (2,948 species in 12 tribes; Borowiec and Świętojańska 2002-present). Opinions differ about the status of certain tribes, arising largely from lack of natural history data, and are reflected in the catalogues (Staines 2015; Borowiec and Świętojańska 2002-present) and in higher-level phylogenies. For example, the catalogues overlap regarding Imatidiini and Spilophorini. These catalogues are still valuable and allow us to extract information on faecal-building behaviours from the documented life cycles. Chaboo and Engel (2009) examined the phylogenetic positions of two crucial fossils, Denaeaspis chelonopsis Chaboo and Engel 2008 (tribe Imatidiini) and Eosacantha delocranioides Chaboo and Engel 2008 (tribe Notosacanthini) at the transition zone between basal Cassidinae ("hispiforms") and tortoise beetles (derived Cassidinae or Cassidinae sensu stricto) so these topologies are also pertinent to discussing the origins and timing of the shield-constructing behaviour.

## Results

We report on four tortoise beetle species from three tribes: Cassidini, Mesomphaliini, and Spilophorini. We outline the basic life cycle of tortoise beetles with two models, *S. cucullata* and Cassidini undet. sp. 4, and introduce special terminology and morphology used for tortoise beetle shields. Then we describe shield architecture, shield retention, and shield construction and reconstruction in *Cal. attenuata* and *Cass. sphaerula*, based on field observations and laboratory manipulations and dissections. We pay particular attention to the caudal processes and the telescopic anus in the two latter species to understand their roles.

#### Natural history of Stolas cucullata (Tribe Mesomphaliini) (Figs 41-44)

This species serves to outline the general life cycle of tortoise beetles and to explain special terms and definitions in Cassidinae. The female (Fig. 41) was captured and provided with a dry twig on which she deposited three eggs (Fig. 42). Egg. Cassidinae eggs may be solitary or grouped, and some are even guarded by mothers (i.e., subsocial); in S. cucullata, the female oviposits a group but each egg is separated. Cassidine eggs may be covered with plant debris, oothecal membranes, or faecal depositions; in S. cucullata, the eggs are naked. They are initially white, then turn grey within a few minutes, then reddish brown with a black apical disc (Fig. 42). Egg size (n = 2: 2.4 mm long; 1.0 mm wide. Larvae (Fig. 44). The neonate larvae have a yellow body with yellowish cream scoli and are densely setose. They wandered away after hours/days, living a solitary life which contrasts with many tortoise beetles that maintain a gregarious group that can additionally be guarded by the mother (subsociality; Chaboo et al. 2014). Comments. The host plant, Neomirandia, has 56 known species and may host other Stolas species; its interesting chemistry (Tamayo-Castillo et al. 1989) is suggestive of a possible role in the beetle's biology and its exuvio-faecal shield.

**Shield construction behaviour.** The natal larva (Fig. 44) has many scoli and paired caudal processes, all with long setation. As the larva feeds, faeces accumulate on these paired caudal processes and, it appears, are held additionally by the long hairs. We have not yet observed the other life stages of this species, but we note how the shield is initiated in Instar I.

## Natural history of Cassidini undet. sp. 4 (Tribe Cassidini) (Figs 45-50)

These larvae build a wide fan-like shield. **Shield construction behaviour** (Figs 45–50). At each moult, the exuvia is shed, from the head to the hind end, but is not cast off. Instead, the exuvia remains attached to the caudal processes. Faeces are added all over, enlarging the shield structure which becomes dry and black-brown in colour. We observed the long, telescopic anus extend and deposit faeces; the anus is highly manoeuvrable and can extend nearly  $^{2}/_{3}$  of the body length (note different positions of the anus in Figs 45–50). The shield becomes a large triangular structure with the exuviae stacked internally but not apparent externally, being so daubed over with thick faeces. *Materials*. The instar I initiates the faeces-only shield but later instars have a shield of all larval exuviae and faeces. This is inherited by the pupa (Figs 49, 50); note the fungal



**Figures 41–44.** Life history in *Stolas cucullata* (Boheman, 1862) (tribe Mesomphaliini) in Costa Rica **41** adult **42** female laying eggs **43** eggs, grouped but not in contact **44** neonate larvae resting on egg shell (photographs: K. Nishida).

hyphae growing upon the shield. **Associated morphology.** The extensible anus builds the shield, placing wet faeces on the caudal process (instar 1) or on the exuviae + faeces of older instars. In older instars, the chaetotaxy is much smaller, raising a question if long chaetotaxy on the instar I caudal processes help hold on to moist faeces, until a hardened structure forms; older instars do not have such long chaetotaxy. The caudal processes in both larva and pupa provide the scaffold of construction (internally, the exuviae become inter-nested at their caudal processes, giving stability). In the larva, caudal processes also rotate the shield vertically, forward and lowered onto the dorsum, backward and extending flat behind the body, and side to side. This raises a question of stability of the larva's body while moving such a relatively large structure; certainly, the feet must be firmly anchored, temporarily glued perhaps, on the leaf and stem substrate. The two caudal processes move but we do not know if each Caroline Simmrita Chaboo et al.: Construction behavior in Cassidinae beetles



**Figures 45–50.** Telescopic anus and shield of larva, Cassidini undetermined sp. 4 on *Chione sylvicola* (Standl.) W. C. Burger (Rubiaceae) in Costa Rica **45–48** anus at different positions **49** pupa, postero-dorsal view **50** pupa, dorsal view (photographs: K. Nishida).

process can move independently of the other. In the solitary pupa (Figs 49, 50), we noted shields held in different positions, directly on the dorsum (Fig. 49) or backwards (Fig. 50). The pupa's abdomen is firmly glued and anchored to the leaf substrate.

# Natural history of *Calyptocephala attenuata* (Spaeth, 1919) (Spilophorini) (Figs 51–56)

Illustrated natural history notes have been reported (Nishida 2014, 2015; Yamamoto 2018, 2020; Nishida et al. 2020). Shield construction behaviour: The larvae retain a shield comprised solely of exuviae of previous instars on the paired caudal processes ("urogomphi") (Figs 51–54). The mature larva carries five exuviae (Figs 51, 52), thus indicating that larva as Instar VI and this is an atypical life cycle for Cassidinae (Chaboo 2007).

The process of shield-building in *Cal. attenuata* begins at the end of Instar I. We describe this process, based on field data and photographs of KN (Nishida 2014, 2015; Nishida et al. 2020), including his assistance on staging the beetle to film the behaviour for two nature documentaries (Yamamoto 2018, 2020; we indicate time (T) in minutes and seconds below based on the film, but readers must access film).

Phase 1 (Fig. 61). Larva, instar I (~ 4.1 mm long), naked, lacking a shield. The larva becomes quiescent as it prepares to moult (T 0-1 min). The six legs are firmly anchored on the leaf and the claw tips appear to be a little embedded on the leaf surface. With a few large inspirations, air fills the gap between the old instar I exuvia and the new instar II; the former seems to lift away. Then the old thoracic nota split medially (T 1 min 35 secs). The abdomen and caudal process move slowly and gently forward and back. The larva inspires air again, inflates a little, and the new prothorax pushes out of the old skin (T 2 mins 5 secs), further widening the breach along the notum. The head capsule splits along the epicranial suture (T 2 mins 28 secs); the new prothorax pushes out further (T 2 mins 40 secs), freeing the lateral scoli (T 2 mins 38 secs), and pulling the head out (Time 3 mins 5 secs). The head and thorax are lifted and freed of the exuvia I, then the new legs are lifted free of their old exoskeleton (T 3 mins 16 secs - 3 mins 25 secs); the instar II abdomen is still encased in instar I abdominal exoskeleton that has not yet split open (Fig. 62).

There is a pause as the head, thorax and legs are lifted vertically, with only slight movements of new legs. The instar II integument is white; yellow haemolymph is apparent internally at the coxal bases. The six pairs of stemmata are black.

Phase 2 (T 6 mins 30 secs - 6 mins 44 secs). Exuvia II legs drop to the surface, then position on the leaf rib and surface, perhaps anchoring claws into the substrate. The entire body heaves a little, gently, then faster, pulling the instar II abdomen free of the Instar I exuvia. The instar I legs lift free of the substrate. Instar II does not walk forward, but pro- and meso-legs stay fixed on the vein as at the start of Phase II. The body is now lifted and rotated, in 360°, extending the abdomen which pushes the anterior section of the old exoskeleton further posteriad (T 7 mins 10 secs). The larva heaves the body anteriad and posteriad, pushing the instar I exuvia backwards (T 6 mins 53 secs). The metaleg positions and re-positions during this phase. The abdomen is held close to the substrate allowing the old head capsule to be dragged against the substrate and pushed further posteriad. At T 7 mins 36 secs, abdominal segments I–II become liberated of the old exoskeleton;



Figures 51–58. *Calyptocephala attenuata* on the host, *Smilax domingensis* Willd. (Smilacaceae), Monteverde, Costa Rica 51 larva with shield of five exuviae, dating this as instar VI 52 dorsal view 53 showing exuviae folded to expose head capsule and caudal processes 54 teneral instar II larva has just exited exuvia I and is retaining it on elaborate paired caudal processes (photographs: K. Nishida) 55 instar I (~ 42 mm long), showing caudal processes 56 instar I caudal processes (photographs: CS Chaboo) 57 adult partially exiting pupal exuvia, fronto-lateral view 58 adult partially exiting pupal exuvia, frontal view (photographs: K. Nishida).

Caroline Simmrita Chaboo et al.: Construction behavior in Cassidinae beetles



**Figures 59, 60.** Unidentified genus, 5<sup>th</sup> instar larvae of Spilophorini on orchid host in Ecuador **59** mature larvae feeding in a group; note color contrast which may be aposematic and the leaf fragment on shield of one larva **60** single larva, dorsal view, with shield of four exuviae. Note exuvial folding exposes the anus and head capsule. Bases of caudal processes are also exposed (photographs: E. Schulz).



**Figures 61–64.** Timing of moulting process and exuviae retention in *Calyptocephala attentuata*. **61** at 17 seconds. Instar I larva lacks the shield **62** at 7 mins, instar II exiting from instar I exuvia **63** at 8 mins, the old head capsule is folded caudad, the instar II pulls forward, pushing the exuvia posteriad **64** at 13 mins, instar II larva with exuvia of instar I on urogomphi. Other instars with additional exuviae (drawn by L. Schletzbaum; timing follows films (Yamamoto 2018, 2020).

shortly after, most of the larval abdomen is extracted from the old exuvia (Fig. 63). By T 9 mins 31 secs (Fig. 64), the old exuvia has been pushed to the posterior half of the new caudal processes (In the sped-up film, the process looks violent). The entire process takes about 17 mins in real time.

- Phase 3. T 11 mins, it appears that abdominal sternites I and II may anchor to the substrate. The anus appears protuberant. The larva sits for another 6 mins before it slightly repositions all its legs. The exuvia of instar I is now positioned on the posterior half of the new caudal processes, with the body folded over and its caudal processes free. The abdominal section of the exuvia is oriented anteriad; the legs, thorax and head sections are folded over and oriented posteriad. Only the posterior half of instar II's caudal processes are inserted into exuvia I, holding it together.
- **Phase 4.** T 14 mins 3 secs, the instar II larvae changes position and we gain a posterior view of its abdomen and caudal processes. The curvature and width of the new caudal processes retains the exuvia I firmly, with some tension.
  - **Instars II–V.** These instars were not observed, but ecdysis at the end of each instar probably follows a similar process as above, with the preceding shield retained on the posterior section of the caudal processes.
  - Pupa. The mature larva attaches the abdomen to the leaf and undergoes pupation. Of the pupae collected, all retained a shield; some shields comprised of two or three older exuviae, but not the younger exuviae that would be most apical in the stacked structure. These shields extended only up to the pronotum, so it is possible that the younger exuviae fell off during ecdysis or were subsequently abraded. Figs 57 and 58 show a pupa with five exuviae, suggesting that exuvia I is detached (these pupae could have six larval exuviae); thus, the pupa inherits shields with varying numbers of exuviae. Figs. 57–58 also show the teneral adult partly exiting this pupal exoskeleton.

## Materials

The larval shield of *Cal. attenuata* is comprised only of exuviae; there are no faecal deposits, secretions, nor plant materials.

## Morphology

Roles of caudal processes in larva and in pupa. These are critical to retaining the shield on the body and to connecting all the previous exuviae together in a single structure. The posterior sections of each caudal process are entirely enclosed within the previous exuvia. *Repair*. It seems obvious that the larvae have no way to repair these exuviae-only shields; if one or more exuviae are removed, the larva must wait until the next moult to add a new exuvia. The movements of the abdomen and caudal processes are responsible for moving the shield in various directions, forwards, laterally and backwards, including above the head. The pupae lack the processes; instead, the final larval exuvia is wrapped around the pupa's caudal region and retains the larval exuvial shield. Some shields (Fig. 59) in our unidentified Ecuadorean Spilophorini have leaf fragments attached; these are possibly accidental.

## Natural history of Cassida sphaerula (Cassidini) (Figs 65-89)

*Cassida* Linnaeus, 1758 comprises 484 species (Borowiec and Świętojańska 2002–present). Immatures have been described for 64 species and exuvio-faecal shields have been noted in most documented larvae to date (Świętojańska and Borowiec 2007: Table 1; Świętojańska et al. 2013). Natural history of *Cass. sphaerula* was reported by Adam et al. (2022) and we summarise in Figs 65–70. Females oviposit small clusters of eggs with oothecal membranes, there are five larval instars (Figs 65, 67), all solitary, and pupae are solitary (Figs 68, 69).

#### Shield construction behavior

Soon after the natal larva (Fig. 65) begins feeding, it begins accumulating its faeces on its paired caudal processes (Fig. 66). At each moult, the cast exuvia is pushed to the base of the caudal processes. The shield becomes a rough triangular-shape, with dark brown-black faeces obscuring the lighter-brown exuviae slightly visible at the base (Fig. 68). The pupa retains the entire larval shield of exuviae + faeces (Fig. 68) or retains only the 5<sup>th</sup> larval exuvia (Fig. 69). The faeces are dense, at different times appearing wet, moist, or desiccated.

#### Incorporation of exuviae into shield

At ecdysis, the old exuvia splits along the ecdysial line of the head and is peeled and pushed backwards, as the teneral instar pulls forward to free its legs. It fixes the legs to the leaf surface, then wriggles its abdomen forward to free itself of the old exuvia. In this way, the previous exuvia becomes positioned at the base of the caudal processes of the teneral larva, beneath the existing exuvio-shield structure. Since all the caudal processes are nested (all previous exuviae atop the living caudal processes of the current instar), the former exuvia becomes crumpled at the base of the existing shield. Soon this recently added exuvia becomes daubed with faeces, and so becomes indistinguishable within the entire shield structure (unless the latter is dissected). No exuviae are omitted from the central scaffold. Apart from the shield structure, excess faeces may be left on the leaf.

We address the question "Will larvae rebuild the shield" with several shield-removal experiments to observe responses of larvae. We present results of three experiments below.

## Experiment 1, instar I (Figs 71-76)

T 0 mins, (Fig. 71): Shield removed completely, exposing the living paired caudal processes. T 2 hours, (Fig. 72): a small quantity of faeces accumulates on the anus. T 4 hours (Fig. 73): moist faecal material has accumulated on the urogomphi, covering it up to the apices. T 6 hours (Fig. 74): faecal material almost the same as at T 4 hours. T 23 hours (Fig. 75): The faecal shield is almost twice as large. T 48 hours (Fig. 76): The faecal shield is about three times larger than it was at T 2 hours.



**Figures 65–70.** Life stages of *Cassida sphaerula* Boheman, 1853 (Cassidini) **65** instar I, neonate **66** instar II, with faeces on caudal processes **67** mature larva with faeces + exuviae shield **68** pupa with entire larval shield (faeces + exuviae) **69** pupa with shield comprised of only 5<sup>th</sup> instar exuvia **70** adult (photographs: S. Adam, September 2021).

Experiment 2, instar I (Figs 77–82)

Time 0 (Fig. 77): We removed the shield entirely, both exuvia I and faeces. T 2 hours (Fig. 78): a small amount of fresh faeces accrue on the caudal processes. T 4 hours (Fig. 79): more new faeces accumulate, obscuring the caudal processes. T 6 hours (Fig. 80): more new faeces accumulate. T 23 hours (Fig. 81): the faeces have grown into a small, dimensional inverted pyramid. T 48 hours (Fig. 82): the inverted pyramidal shield is larger, held together on the caudal processes. This shape seems unstable, that lateral sections should break off yet hold together.

Experiment 3, instar II (Figs 83-86)

T 0 mins (Fig. 83): we scraped away only the faeces to expose the Instar I exuvia still attached to the caudal processes. T 23 hours (Fig. 85): faeces have been applied to the sides of the old exuvia, so the overall shield width is almost as wide as the larva. T 48 hours (Fig. 86): More faeces have been applied to the



Figures 71–76. Re-construction of faeces on exuvio-faecal shield in Experiment 1, starting with instar I larva (so no prior exuvia), *Cassida sphaerula* Boheman, 1853 (Cassidini; photos: S. Adam, September 2021) 71 instar 1 (~ 2 mm long) at time 0 when faecal shield is removed, exposing urogomphi 72 larva at two hours, small faecal blob at anus 73 larva at four hours, urogomphi encased in faeces 74 larva at six hours, urogomphi encased in faeces 75 larva at 23 hours, lateral view. 76 larva at 48 hours, dorso-ventral view (photographs: S. Adam, September 2021).



**Figures 77–82.** Re-construction of faeces on exuvio-faecal shield in Experiment 2 with *Cassida sphaerula* Boheman, 1853 (Cassidini) **77** instar I (~ 2 mm long) before shield construction **78** instar II at time 0 with faeces removed (scraped off) **79** after 2 hours, dorsal view **80** after four hours, dorsal view **81** after 23 hours **82** after 48 hours (photographs: S. Adam, September 2021).



**Figures 83–86.** Faecal re-construction in experiment 3 with instar II larva, *Cassida sphaerula* Boheman, 1853 (Cassidini) **83** time 0 when faecal shield is removed, exposing instar I exuvia **84** larva at two hours, exuvia I still exposed **85** larva at four hours, faeces attached to lateral projections (scoli) of exuvia I **86** larva at six hours, exuvia I with a lot of faeces (photographs: S. Adam, September 2021).

lateral margins, so the shield is now a little wider than the larva. The old exuvia is in the centre, exposed, and the moist black faeces hang on to the sides.

The entire exuvio-faecal structure was gently eased off the living caudal processes using forceps and these intact larvae continued feeding. In each case, the larva soon produced a faeces-only shield, small at 2 hours after removal, then bigger and bigger at hours 4 and hours 6 after removal. By hours 23 and 48, 1–2 days after the earlier removal, the new shield was larger and club-shaped. In the three experiments of shield manipulation, the timing, and responses to reconstruct a new shield were similar. The experimental larvae of *Cass. sphaerula* moulted normally and retained the exuvia into the inherited shield.

The larva can rotate the shield in a circular plane over the body, forward up to the mesothorax, and backward almost 180°, and in a horizontal plane with the body (Suppl. material 1). Films of the acrobatic movements of the larva's



**Figures 87–89.** Shield of pupa of *Cassida sphaerula* Boheman, 1853 (Cassidini) **87** host leaf chewed by beetles, with one larva and two pupae (dorsal views; upper one with exuvio-faecal shield; lower one with exuviae-only shield) **88** posterior view showing exuvial-faecal shield (of instars I–IV) attached to caudal processes of instar V exuviae **89** ventral view showing complete instar V exuvia and exuvio-faecal shield (photographs: S. Adam, September 2021).

extensible anus reveal that it applies faeces to the existing shield and, also periodically exudes large, mostly clear, droplets that are applied to and absorbed into the shield (Suppl. material 2). We found no trichomes in dissected shields even though we observed consumption of trichomes in *Cass. sphaerula* (Adam et al. 2022).

#### Shield retention in pupae (Figs 87–89)

In *Cass. sphaerula*, we observed pupae can have either an exuvia-only shield (Figs 69, 87) or the entire inherited exuvio-faecal shield structure of the earlier larvae (Figs 68, 88, 89). The faeces of the latter are dry since new faeces are not being applied. We found several discarded exuvio-faecal shields next to pupae. Given the observation of the moulting process (from 6<sup>th</sup> instar to pupa) in *Calyptocephala* (described above), we infer that pupation is similar, with splitting of the ecdysial sutures on the cranium and thorax of the 5<sup>th</sup> instar split and the pupa pulls forward and out. In the larval moults, the new legs and the old legs serve to anchor the emerging larva at different times in the process.

## **Discussion**

Faecal-based constructions and faecal debris-carrying are widespread behaviours in Chrysomelidae. Chrysomelid faecal-based constructions have been studied in terms of ecological function (Olmstead and Denno 1992; Gómez 1997, 2004; Morton 1997; Vencl and Morton 1998a, b, 1999; Morton and Vencl 1998; Gómez et al. 1999; Müller and Hilker 1999, 2001a, b, 2003, 2004; Vencl et al. 1999, 2005, 2011; Nogueira-de-Sá and Trigo 2002, 2005; Bacher and Luder 2005; Bottcher et al. 2009; Huang et al. 2022). Until now, this faecal building behaviour has been studied in only one chrysomelid, *Neochlamisus* by Brown and Funk (2005).

## Materials in cassidine shields

The macro-materials in shields of our observed species comprised exuviae only or faeces + exuviae. These two materials are side effects of metabolism and moulting respectively. Additional analyses may identify other possible components (Table 1) and their functions. The construction processes we documented allow us to now analyse how the two primary materials originate, are manipulated into the construction, and are held to the body. We briefly discuss evolutionary insights as we compare these aspects with other Cassidinae and other Chrysomelidae.

#### **Building stages in Cassidinae?**

Larvae are the builders in our four studied species and building begins in two possible ways: 1) during instar I when faeces are deposited and held on the caudal processes as the larva feeds (in *S. cucullata*, Cassidini undet. sp. 4, *Cass. sphaerula*) or, 2) in the transition moult from instar I to instar II when the cast exoskeleton is retained on the caudal processes (*Cal. attenuata*). Cassidinae pupae in tortoise beetle tribes are not active builders; they receive their shields as an inheritance from the final larval instar and their shield is either the entire exuvio-faecal shield or only the final instar exuvia. Given the life cycle of Spilophorini, the final instar could be the 5<sup>th</sup> or 6<sup>th</sup> for tortoise beetles. For pupation, the pre-pupa anchors itself by gluing the abdomen to the host surface. Then the larval exoskeleton splits along the head and thoracic midlines and the pupa pushes out as the larval exuvia is propelled caudad. The shield is retained passively, attached on the pupa's own caudal processes. This pupal inheritance recalls that of camptosomate chrysomelids where the final instar seals the larval faecal case to the substrate and so provides a pupation chamber (Chaboo et al. 2008).

## **Cassidinae shield architectures**

The common pattern is the exuvio-faecal shield built by larvae, retained in all instars, and which may be inherited by pupae. The faeces are variable in moisture, from desiccated (Figs 19, 22, 23, 25, 26) to wet (e.g., *Plagiometriona flavescens* (Boheman, 1855): Flinte et al. 2009); this is certainly tied to the excretion of water, retention by Malpighian tubules, and rectal resorption. Dried faeces can be in long strands; these strands are arranged in a circular heap in Hemisphaerotini (Fig. 19; Chaboo and Nguyen 2004) or are held as vertical strands (Figs 22, 23). Within the tribe Ischyrosonychini, larval shields are varied: desiccated stacked exuvio-faecal shields (e.g., *Cistudinella obducta* (Boheman, 1894) (Fiebrig 1910; Buzzi 1988), wet faeces (e.g., *Physonota unipunctata* (Say, 1823); Keefover-Ring 2013, 2015), or older larvae that lack shields altogether (e.g., some *Physonota* Boheman, 1854). Larvae of *Eurypepla* Boheman, 1854 have a unique tapered body that is curved verticad, allowing wet faeces to slide down and coat the body (Chaboo 2004).

Architectural elements of cassidine faecal structures may be diagnostic for species-, genus- or tribal-level diagnoses. Shield architecture is determined by how exuviae are compressed and how faeces are arranged (long vertical strands, a dense clump, or a fan). Basket-like shields are diagnostic of

Hemisphaerotini (Fig. 19; Eisner et al. 1967; Beshear 1969; Chaboo and Nguyen 2004) and appear to have some limited mobility, particularly in younger stages (note its position in Fig. 19). As the larval shield enlarges, it becomes less mobile, suggesting that this shield is relatively heavy and/or the caudal processes may not be as freely mobile. Although exuviae are retained in Hemisphaerotini, these are so compressed that only torn remnants remain at the base of the caudal processes, and it seems impossible to determine how many distinct exuviae are held. This shield architecture has been demonstrated to be protective (Eisner and Eisner 2000a).

We propose here that the particular exuviae-only shield architecture described herein is diagnostic for Spilophorini (Figs 51-54, 57, 59, 60). Life cycles of two species of Calyptocephala Chevrolat, 1836 (Buzzi and Miyazaki 1992; Córdova-Ballona and Sánchez-Soto 2008) on palm hosts reveal larvae with paired caudal processes and exuviae-only shields. Maulik (1932) described the larva of an Oediopalpa Baly, 1858 species with paired caudal processes and an exuvial shield; Chaboo (2007: 184) examined larvae in this genus and noted the unique pattern of exuviae compression. Hsiao and Windsor (1999) determined Oediopalpa as most closely related to Calyptocephala and Spilophora Boheman 1850 and Staines (2002) re-classified it in Spilophorini. Sekerka et al. (2014) reported an Orchidaceae host, the larval form, and exuviae-only shields for one species of Cladispa Baly, 1858 (Spilophorini). Monophyly of Spilophorini has been supported by adult characters (Chaboo 2007) and molecular data (Sekerka et al. 2014). The documented larvae exhibit exuviae-only materials arranged in a similar architecture, with a stable exuvial stack, a distinct spatial arrangement, and large partly exposed caudal processes. The exuviae are compressed and curved so the head capsule and the anus are exposed in posterior view. The shape of the caudal processes, like the yoke of a lyre, is unique in Cassidinae; the exposure (Fig. 57) of the large basal section of each process is also unique. These features altogether support monophyly of Spilophorini.

Other tortoise beetles exhibit exuviae-only shields (Figs 15, 21, 24) but the spatial arrangement of those exuviae and the underlying caudal process morphologies are unlike those in Spilophorini. In other documented species, exuviae are compressed differently, more closely at caudal processes, and head capsules are exposed in distinct ways (see examples: *Stolas implexa* (Boheman, 1850), Flinte et al. 2009: pl. 18K; *Chiridopsis undecimnotata* (Boheman, 1855), Świętojańska 2009: fig. 128). As species and their constructions are documented, it may be possible to diagnose more groups based on more shield and process features.

Shields may be present or absent in pupae of tortoise beetles. We found that pupae of *Cass. sphaerula* retain different shields (the entire structure or only the final exuvia). Some pupae retain only the 5<sup>th</sup> instar exuvia and their caudal processes are a dominant exposed feature (e.g., *Anacassis* Spaeth, 1913, Buzzi 1975; *Discomorpha* Chevrolat, 1836; Flowers and Chaboo 2015). In the Cassidini, pupal shields are known in species of *Charidotis* Boheman, 1854, *Drepanocassis* Spaeth, 1936, *Metriona* Weise, 1896, and *Syngambria* Spaeth, 1911 (Buzzi 1988). In some cassidines, the 5<sup>th</sup> exuvia is retained by the pupa, encircling the terminal abdominal segments, e.g., *Anacassis* Spaeth, 1913 (Buzzi 1975, 1996). In *Eugenysa columbiana* (Boheman, 1850) (Chaboo 2002), *Dorynota pugionota* (Germar, 1824) (Buzzi 1976), and *Chelymorpha* Chevrolat,

1836 (Buzzi 1998) this exuvia becomes part of the pupal attachment to the substrate. Shield removal is required to determine if this exuvia is wrapped around the base of the abdomen only or if it is attached to a pupal caudal process.

#### What is the building equipment in Cassidinae?

We documented the anus moving freely over the posterior surface of the shield (Figs 45–48). We observed anal droplets excreted and quickly absorbed into the shield (Suppl. material 2). We also documented the application of fresh moist faecal deposits to the intact shield and, in our experiments, application to the exposed exuviae to rebuild the shield (Figs 71–86). Gómez (1997) reported the repair of damaged shields with precise deposits of faeces. Thus, the anus is the applicator for constructing and repairing the shield and appears to replenish the shield with moist droplets. Certainly, the cassidine anus has manipulative skill for these distinct roles (applying, building, repair, replenishment). Such replenishment may involve chemicals that sustain the shield's chemo-barrier functioning. If pupal shields are not being replenished, this raises a question about their chemistry and functional effectiveness versus larval shields.

The musculated extensible anus of larvae is a second synapomorphy of the ten tortoise beetle tribes. Plesiomorphic Cassidinae larvae which do not exhibit shield-retaining behaviours have the typical posterior or ventrally opening simple anus pore and also lack caudal processes. As far as we know currently, no other chrysomelid larvae have an extensible anus. One question we have is the status of the anus in those Cassidinae with exuviae-only shields; we were unable to determine this in *Cal. attenuata*. Pinpointing the first appearance of the telescopic anus on phylogenetic topologies is one crucial element in the assembly of shield building traits.

Cassidinae larvae do not use their legs or mouthparts as building tools. Females may defecate on their eggs, but their genitalia lack rectal plates (as in Camptosomata: Erber 1968, 1969, 1988). In Camptosomata, the larva's arrangement within its case positions the anus near the mouthparts and legs. Brown and Funk (2005) reported that faeces are mixed with a regurgitated yellow fluid and then applied to the margin of the case to continue building it or to repair holes, so the larva's position with the mouth, anus and legs in proximity allows the faecal mixing and manipulation. Camptosomate larvae use their mouthparts to cut a longitudinal section which is then filled with faeces; this expands the girth of the case to accommodate the growing larva (Brown and Funk 2005; Chaboo et al. 2008). Calcetas et al. (2023) reports that *Podontia* larvae use legs and mouthparts to manipulate soil and faeces to build the pupation chamber.

#### **Building routines in Cassidinae**

Cassidinae larvae use simple materials in simple building routines. Each shield has a distinct appearance due to the compression pattern of individual exuviae (Figs 15, 21, 24, 51, 52, 57, 59, 60) and due to the arrangements of faeces (strands, blobs, fan, bird nests, etc.). The shield enlarges at each transformation to the next instar as another exuvia is added basally to the mass. The extensible anus deposits faeces precisely on various parts of the exuviae to give the distinct appearance of shields. Faeces are extruded moist or wet, allowing

attachment to the existing structure, before drying. Our simple experiments allowed us to understand the repair of the shields. If a portion of faeces is removed or broken off on one side of the structure, the anus can repair the faecal part to apply fresh faeces to recover a more balanced shield. Our study demonstrates that the shield-constructing behaviour is intrinsic and is probably not requiring any external activator to elicit the building response.

#### Role of caudal processes (= urogomphi)

Many animals that retain debris covers possess fastening structures, frequently specialised chaetotaxy (e.g., Weirauch 2006). We determined here several roles of the paired caudal processes— the anchorage or fastener for cast exuviae and faeces to the body, part of the shield materials, the crucial central scaffold by their inter-nesting, and movement of the shield. During instar I, faeces are applied directly to the caudal proceses; in *S. cucullata* dense chaetotaxy around the anal area may enhance faecal retention.

The exuvia is added to the shield with each moult, expanding the area for faecal attachment. In some species, exuviae alone make up the shield. The caudal processes become inter-nested from instar to instar, further strengthening the central scaffold of the exuvio-faecal shield and provide mobility, allowing it to be moved as needed to startle or hit an attacker or be the distasteful barrier. The caudal processes move the shield for a more active defence.

In pupae, we have no reports of cassidine pupae moving their shields, although there are reports of pupa jerking reflexively when disturbed (even in unison in gregarious pupae). It appears the entire pupal body jerks so pupal caudal processes may not be mobile.

## Role of chaetotaxy

In one unidentified species and in *Cass. sphaerula* we observed that dense chaetotaxy in the caudal area of the neonate larva appears to aid initial faecal build-up. Specialised chaetotaxy may aid faecal retention in the faecal retaining chrysomelid clades. Specialised setae to hold on to debris have been described in unrelated beetles (Leschen and Carlton 1993; Leschen 1994; Yoshida and Leschen 2020), in other insects (e.g., Reduviidae, Weirauch 2006), and in other animals (e.g., spiders, Duncan et al. 2007; Gawryszewski 2014). In *Uraba* caterpillars (Fig. 9), it is a question how the old head capsules become stacked on the living caterpillar's head, since the head capsule typically splits first during the moulting process, then becomes distorted as it is pushed posteriad, and the larva propels forward to exit its old exoskeleton. We suspect that specialised chaetotaxy on the caudal processes of tortoise beetle larvae and on the dorsum of larvae in Criocerinae and in the *Blepharida*-group may hold onto the faecal debris. Each debris-retaining animal has different strategies for attaching and retaining debris.

#### Materials of coats, cases, and shields across Chrysomelidae (Table 1)

Chrysomelid constructions are composed mainly of endogenous faeces and, in Cassidinae, of exuviae. Documented exogenous materials are soil, fungi, leaf fragments (fresh, undigested, decayed), plant extracts, and trichomes (Table 1). We will not review here how exactly these materials may be mixed or intermingled with the other structural materials. In Cassidinae, fungal elements have been noted but not identified taxonomically (Figs 49, 50; Rane and Ghate 2005; Flinte et al. 2009; Cedeño-Loja and Chaboo 2020). Fig. 59 shows a larva with plant fragment on the exuvial shield but this may be accidental. It has been well-established that animal guts are rich with microbiota that can be passed to the next generation via the egg surface; Stammer (1935) established such transmission in Cassidinae. Faeces are also rich with microbiota (thus, Faecal Transplant technique); we can presume that the cassidine shield is harbouring microbiota that await discovery and study. The exuviae are a low-cost material that add substantial structural value to the shield (like straw added to dung) but we do not know yet their chemical contributions. All debris materials have pros and cons, depending on how they originate (time to produce or assemble) and their consequences (e.g., weight, odour, chemistry) so every chrysomelid material likely has a functional role simply because of the cost in carrying the weight and bulk of a structure; it is unlikely that unnecessary materials are selected. Most of these chrysomelid materials are actively manipulated, although it is possible that some (e.g., blown soil) may be passively integrated.

## Chrysomelid construction behaviours: ecological implications

Chrysomelid larval and pupal shields are hypothesised to serve multiple functions, including protection from extreme temperature (Réaumur 1737), humidity, precipitation and desiccation (Weise 1893), camouflage or mimicry (e.g., bird or caterpillar droppings: Briggs 1905; Blatchley 1910; Jenks 1940; Balsbaugh 1988; plant detritus: Lee and Morimoto 1991a, b), as a distasteful physical barrier deterring predators and parasitoids (Réaumur 1737; Eisner et al. 1967; Olmstead and Denno 1992; Olmstead 1994, 1996; Eisner and Eisner 2000a; Bacher and Luder 2005), or as chemical deterrents from exocrine glands of retained exuviae (Olmstead 1994). They can also be used as a mobile club to hit intruders or as a protective umbrella (CSC, pers. obs.). The term 'shield' implies passive protection, which may lower the body temperature or decrease wind shear (Olmstead and Denno 1992). The material and consistency (including cementing and chemistry) must ease accumulation and attachment. It appears that chrysomelid shields are generally resistant to rain as they do not absorb water and fall apart.

## **Testing of function hypotheses**

The hypothesis of a mechanical defence against predators has been tested experimentally and found to be supported (Eisner et al. 1967; Wallace 1970; Olmstead and Denno 1992; Eisner and Eisner 2000a; Schaffner and Müller 2001; Müller 2002). Blum's (1994) hypothesis of defensive chemicals in shields has led to some analytical studies, usually of single chrysomelid species, aimed at comparing compounds in the faecal shields and the host plants (Mummery and Valadon 1974; Morton and Vencl 1998; Gómez et al. 1999; Vencl et al. 1999; Aregullín and Rodríguez 2003; Bacher and Luder 2005; Nagasawa and Matsuda 2005 Nogueira-de-Sá and Trigo 2005; Vencl et al. 2005, 2009, 2011; Bottcher et al. 2009; Keefover-Ring 2013, 2015; Vencl and Srygley 2013). Maybe a chemical barrier is achieved by integrating plant tissues and trichomes or by applying secretions (plant-sequestered or de novo chemicals) that volatilise around the animal, maybe creating a small chemosphere. Exuvial glands may have residual chemicals that may disguise the wearer or deter enemies.

In testing Ehrlich and Raven's (1964) "escape and radiate" hypothesis, Vencl et al. (2011) compared differential functioning in defence of shields with/without faeces, larval solitary/gregarious living, and maternal care and deduced a sequence of trait accumulation correlated with enhanced defences and, likely, species diversification. Such creative experiments can assess the contribution of each trait within the defense array.

Others have determined the shields to have mixed effects, deterring some predators yet attracting others (Müller and Hilker 1999; Bacher and Luder 2005; Huang et al. 2022). Certainly, faeces can have chemical signatures that attract enemies (Van Leerdam et al. 1985; Agelopoulus et al. 1995).

Chemical deterrents in exocrine glands of retained exuviae (Hinton 1951; Olmstead 1994) have not been investigated. Furthermore, the traits accumulated in the defence arsenal must now include the morphological features that accompany the structures; for example, caudal processes enhance shield mobility in tortoise beetles and may enhance defence success. Our research here highlights the morphological features used by tortoise beetle larvae within their arsenal of weapons.

#### **Construction behaviours: evolutionary implications**

The primary hypotheses proposed to explain chrysomelid hyperdiversity have been their ancient age (Farrell et al. 1992), herbivory and the rise of angiosperms (Farrell 1998), adaptive radiation with plants (Gómez-Zurita et al. 2007), and chemical adaptation to plants (Farrell et al. 1992). However, the great unevenness in subfamilial diversity begs for additional explanations. Transitions to new habitats within Chrysomelidae (e.g., aquatic, seeds, subterranean, mosses) and to the jumping escape mechanism (in ~8000 flea-beetle species, Begossi and Benson 1988; Furth 1988) await finer-scale study of correlated adaptations in morphology, physiology, and behaviour. Behaviours such as cycloalexy (larval defence formations; Jolivet 1988b), sound production (Schmitt 1994), myrmecophily (Agrain et al. 2015), and subsociality with maternal care (Chaboo et al. 2014) probably impact speciation in more restricted clades of Chrysomelidae. On the available phylogenetic hypotheses of Chrysomelidae (Figs 16-18), faecal armours appear as independent macroevolutionary events in speciose clades (part of Cassidinae; Cryptocephalinae; Lamprosomatinae) and in minor lineages (Blepharida-group within Galerucinae; Criocerinae; Phola within Chrysomelinae). Systematic analyses of these nodes of transitions, from no faecal recycling to faecal recycling, are needed to understand possible speciation impacts after the origin of constructions. We can surmise a shared genetic history for faecal constructions, that they have value in the survival of their builders, and they could be considered adaptive. To understand evolutionary relevance and even possible character information for phylogeny reconstruction, many more species-level studies are needed to document the life stages and to compare roles of different building materials and building repertoires.

Chrysomelid faecal-based constructions are not homologous, being formed in different ways and are held to the body by different structural modifications. Interesting points emerge when subfamily comparisons are made (Table 1). The common material of faeces points to its cheapness and ready availability. Some architectures may be convergent. Dorsal coats of faecal pellets and similar anus position in Criocerinae and in the Blepharida-group suggest similar neuro-physiological mechanisms (a "conveyor belt") to move faecal pellets from anus towards the head and similar purposes. The cassidine Eurypepla Boheman, 1854 (Chaboo 2004) also has a wet shield, but this is built differently - the upwardly held abdomen permits the flow of viscous faeces (not pellets) down the body to coat it. It is highly likely, given findings in other non-chrysomelid debris-carriers, that specialised chaetotaxy hold the pellets onto the dorsum. The case architecture of Camptosomata-similar architecture, similar construction behaviours, and similar correlated morphologies (i.e., maternal abdominal fovea and genital 'kotpresse'; larval flattened head, swollen abdomen, long legs, and curved claws)-support the close relationship of Cryptocephalinae and Lamprosomatinae. Comparing these aspects in the arboreal, terrestrial, and myrmecophilous species of this clade might reveal additional informative characters for taxonomy and phylogeny.

Cassidinae (e.g., Chaboo 2007) and Criocerinae (Vencl et al. 2004) both exhibit mining, cryptic and exposed larval feeders but faecal shields are made only in exposed forms; this pattern suggests these larvae use shields to protect themselves against a range of abiotic and biotic dangers that are different from those faced by their mining relatives. A bulky structure like a shield is unlikely in the constrained space of a mine.

A question in Cassidinae now is "Which tribe is the sister for the ten tortoise beetle tribes?" Borowiec (1995: fig. 2) proposed two major lineages of tortoise beetles, without identifying a particular basal tribe. Hsiao and Windsor's topology (1995: fig. 1) resolved Spilophorini + *Oediopalpa* as phylogenetically distant from other tortoise beetles; their topology suggests either two origins of shield construction or a single origin with some losses. Chaboo (2007) found *Oediopalpa* among "hispines" and Spilophorini and Hemisphaerotini at the base of the tortoise beetle clade; this also suggests a minimum of two origins of the shield construction, yet the shields and caudal processes in these two tribes are very different. A few tortoise beetle species lack a shield, but our current phylogenetic hypotheses suggest these are secondary losses. We also know now that exuviae-only shields appear scattered over the tortoise beetle clade, suggesting multiple origins.

Two Cassidinae fossils (Chaboo and Engel 2008) support a close relationship between Notosacanthini which have mining larvae (Monteith et al. 2021) and Delocraniini which have cryptic exophagous larvae but no shield (CSC, pers. obs.). These fossils suggest that the typical tortoise beetle larval shields probably originated once and during the latest Paleocene or earliest Eocene (Chaboo and Engel 2008).

Recent field observations of *Aproida* (Aproidini) in Australia reveal that the larvae have a single caudal process and that faeces can pile up from time to time but falls off quickly: there is no fixed stable faecal shield and exuviae are not retained by larvae except at the pre-pupation stage (Chaboo, Sandoval, Campos, and Monteith unpubl. data). Leptispini have exophagous larvae that live in a cryptic leaf shelter they construct; these larvae also exhibit a single

caudal process, but no shield (Prathapan et al. 2009). Species of *Eurispa* Baly, 1858 (Eurispini) have exophagous sheath-feeding larvae but the illustrations of Hawkeswood and Takizawa (1997) are unclear if they have typical caudal processes (tergal) or marginal extensions of an urogomphal plate (not homologous with caudal processes). The single caudal process appears as multiple independent origins within Cassidinae. *Discomorpha* (Omocerini) larvae exhibit a functionally single process but this appears to be a fusion of two and it retains the exuvio-faecal shield (Flowers and Chaboo 2015).

## Conclusions

We demonstrate general and widespread models of shield construction in tortoise beetles. We indicate variations in shields over the tortoise beetle clade that raise new challenges to study odd species. Many characters of shields can be defined to benefit phylogeny reconstruction, including construction repertoire, architecture, materials, and associated morphology. Natural history studies and specimen collections can integrate more species to achieve finer-scaled phylogenies of Cassidinae, particularly around nodes of transitions (e.g., mining to exophagy; presence/absence of caudal processes; presence/ absence of shields). Clarifying these nodes will help us understand how life history and shields affected diversification within Cassidinae.

Defecation ecology is an important yet under-researched area that is intertwined with the building behaviours and morphology of chrysomelid beetles. Their constructions are crucial for their survival and represent adaptive macro-evolutionary events. Comparative and inter-disciplinary studies of construction behaviours are needed to better understand the evolution of chrysomelids. Until now, explanations of chrysomelid hyperdiversity have relied on the association and radiation with plants. Yet, constructions are a pervasive feature that may help explain the great subfamilial unevenness in Chrysomelidae. The major challenge is fieldwork and specimen assembly of juvenile stages and their constructions, as they are poorly represented in museum collections.

## Acknowledgements

In Costa Rica fieldwork, we thank Mauricio Fernandez, Bill Haber, Paul Hanson, Alan Pounds, and Angel Solís for help and Sistema Nacional de Áreas de Conservación, Ministerio de Ambiente y Energía (MINAE) for the research permits. We thank Michael Thomas and Paul Skelley for specimen Ioans of *S. cucullata*. In South Africa fieldwork, we thank Beth Grobbelaar (SANBI), Hugh Heron, and Maureen Coetzee, University of Witwatersrand, for help and discussion. This paper had a long gestation with valuable discussions, citations, and specific examples from: John Wenzel; colleagues at Cornell University—Thomas Eisner (deceased), Maurice (deceased) and Catherine Tauber, Karen Sime, Nico Franz, Kelly Miller, and Quentin Wheeler; many beetle colleagues—Cheryl Barr, Mary Liz Jameson, Elizabeth Grobbelaar, Hugh Heron, Alfred Newton, Karen Olmstead, Margaret Thayer, William Shepard, and Martha Weiss; and many chrysomelid specialists— Federico Agrain, Christopher Brown, Lourdes Chamorro, Cibele Rebeiro Costa, Vivian Flinte, R. Wills Flowers, Daniel Funk, David Furth, Ken Keefover-Ring, Karen Olmstead, Divarkaran Prathapan, Chris Reid, Paula Alex Trillo, Fred Vencl, and Rob Westerduijn. We thank the following for sending photographs and permission to use: Hope Abrams, Nicky Bay, H. Del Olmo, Hubert Duprat, Dimitri Forero, Jan Hamryski, Masayuki Hayashi, Alan Henderson, Steve Marshall, Dieter Mahsberg, Nick Porch, Filip Trnka, Eerika Schulz, Adrian Tejedor, Gil Wizen, and Takahiro Yoshida. We also thank Ichiro Yamamoto (Director), KAZE Co., Japan for allowing us to view his film of Cal. attenuata. We thank Federico Agrain, Jesús Gómez-Zurita, Divarkaran Prathapan, and Michael Schmitt for discussing chrysomelid phylogenies, and particularly thank Sara López-Pérez for collaborating on phylogenetic characters. During this long study some results were presented at scientific meetings: Coleopterists Society, Entomological Collections Network, Entomological Society of America, European Congress of Entomology, Kansas Entomological Society, and International Congress in Entomology. We are very grateful to our manuscript readers, Orly Calcetas, Charles Staines, and Nick Upton, to our two valued reviewers, and to Editor Michael Schmitt for all comments and suggestions that improved the final text. Finally, CSC dedicates this paper to her husband, Fernando Merino, and their daughter Teresa Merino.

## **Additional information**

## **Conflict of interest**

No conflict of interest was declared.

## **Ethical statement**

No ethical statement was reported.

## Funding

This work was supported by the U.S.A. National Science Foundation grant EAGER 1663680 (PI: CS Chaboo).

## **Author contributions**

Conceptualization: CSC. Data curation: CSC, SA, KN. Formal analysis: CSC. Investigation: CSC, SA, KN. Methodology: CSC, SA, KN. Project administration: CSC. Visualization: CSC, SA, KN, LS. Writing – original draft: CSC, SA, LS. Writing – review and editing: CSC, SA, KN, LS.

## Author ORCIDs

Caroline Simmrita Chaboo ID https://orcid.org/0000-0002-6983-8042 Sally Adam ID https://orcid.org/0009-0002-4971-4488 Kenji Nishida ID https://orcid.org/0000-0002-1846-9195 Luke Schletzbaum ID https://orcid.org/0000-0001-7995-7136

## Data availability

All of the data that support the findings of this study are available in the main text or Supplementary Information.

## References

Abstemius L (14<sup>th</sup> century) Hecatomythium Secundium. In: Desbillons FJT (Ed. [translator]) Francisci-Josephi Desbillons Fabulae Aesopiae, curis posterioribus omnes feré emendatae: quibis accesserunt plus quam CLXX novae. Typis J. Barbou, Paris, 504 pp.

- Adam S, Campos M, Heron HDC, Staines C, Westerduijn R, Chaboo CS (2022) Natural history notes of *Cassida sphaerula* Boheman, 1853 (Coleoptera: Chrysomelidae: Cassidini) on *Arctotheca prostrata* (Salisb.) Britten (Asteraceae) in South Africa. Insecta Mundi 0945: 1–23. https://journals.flvc.org/mundi/article/view/131550
- Addesso KM, McAuslane HJ, Stansly PA, Schuster DJ (2007) Host-marking by female pepper weevils, *Anthonomus eugenii*. Entomologia Experimentalis Applicata 125(3): 269–276. https://doi.org/10.1111/j.1570-7458.2007.00626.x
- Agelopoulus NG, Dicke M, Posthumus MA (1995) Role of volatile infochemicals emitted by faeces of larvae in host-searching behaviour of parasitoid *Cotesia rubecula* (Hymenoptera: Braconidae): A behavioural and chemical study. Journal of Chemical Ecology 21(11): 1789–1811. https://doi.org/10.1007/BF02033677
- Agrain FA, Buffington M, Chaboo CS, Chamorro ML, Schöller ME (2015) Leaf beetles are ant-nest beetles: the curious life histories of the juvenile stages of case-bearers (Chrysomelidae: Cryptocephalinae). ZooKeys 547: 133–164. https://doi. org/10.3897/zookeys.547.6098
- Aiello A (1979) Life history and behaviour of the case-bearer *Phereoeca allutella* (Lepidoptera: Tineidae). Psyche 86(2–3): 125–136. https://doi.org/10.1155/1979/84541
- Aiello A, Stockwell HP (1996) The golden-cage weevil, *Isorhinus undatus* (Champion) (Coleoptera: Curculionidae). The Coleopterists Bulletin 50(2): 192–194.
- Anbutsu H, Togashi K (2002) Oviposition deterrence associated with larval frass of the Japanese pine sawyer, *Monochamus alternatus* (Coleoptera: Cerambycidae). Journal of Insect Physiology 48(4): 459–465. https://doi.org/10.1016/S0022-1910(02)00067-7
- Anderson KL, Seymour JE, Rowe R (2003) Influence of a dorsal trash-package on interactions between larvae of *Mallada signata* (Schneider) (Neuroptera: Chrysopidae).
  Australian Journal of Entomology 42(4): 363–366. https://doi.org/10.1046/j.1440-6055.2003.00373.x
- Arranz-Otaegui A, López-Sáez JA, Araus JL, Portillo M, Balbo A, Iriarte E, Gourichon L, Braemer F, Zapata L, Ibañez JJ (2017) Landscape transformations at the dawn of agriculture in southern Syria (10.7e9.9 ka cal. BP): Plant-specific responses to the impact of human activities and climate change. Quaternary Science Reviews 158: 145–163. https://doi.org/10.1016/j.quascirev.2017.01.001
- Aregullín M, Rodríguez E (2003) Sesquiterpene lactone sequestration by the tortoise beetle *Physonota arizonae* (Cassidinae). Revista de la Sociedad Química de México 47: 143–145.
- Arndt I (2013) Animal Architecture. Abrams, New York, 160 pp. https://www.abramsbooks.com/product/animal-architecture\_9781419711657/
- Arzone A, Meotto F (1978) Reperti biologici su *Goniopterus scutellatus* Gyll. (Col. Curculionidae) infestante gli eucalypti della riviera ligure. Redia (Firenze) 61: 205–222.
- Bacher S, Luder S (2005) Picky predators and the function of a faecal shield in a Cassidinae larva. Functional Ecology 19(2): 263–272. https://doi.org/10.1111/j.1365-2435.2005.00954.x
- Balsbaugh Jr EU (1988) Mimicry and the Chrysomelidae. In: Jolivet P, Petitpierre E, Hsiao TH (Eds) Biology of the Chrysomelidae. Kluwer Academic Publisher, Dordrecht, 261–284. https://doi.org/10.1007/978-94-009-3105-3\_16
- Baly JS (1858) Catalogue of Hispidae in the Collection of the British Museum. Printed by order of the Trustees, London, 172 pp. https://doi.org/10.5962/bhl.title.9686

- Bameul F (1989) Description du comportement de camouflage d'un Coléoptère: Le déguisement actif de *Georissus crenulatus* (Coleoptera Georissidae), et proposition d'une nouvelle classification des déguisements chez les Invertébrés. Compte Rendus de l'Academie des Sciences 309(III): 351–356.
- Begossi A, Benson WW (1988) Host plants and defense mechanisms in Oedionychina (Alticinae). In: Jolivet PH, Petitpierre E, Hsiao TH (Eds) Biology of Chrysomelidae. Series Entomologica (Vol. 42). Kluwer Academic Publishers, Dordrecht, 57–72. https:// doi.org/10.1007/978-94-009-3105-3\_4
- Beneker J, Gibson CA [Eds and translators] (2016) The Rhetorical Exercises of Nikephoros Basilakes. *Progymnasmata* from Twelfth Century Byzantium. Dumbarton Oaks Medieval Library, Harvard University Press, Boston, 416 pp. https://www.hup.harvard. edu/catalog.php?isbn=9780674660243
- Benelli G, Pacini N, Conti B, Canale A (2013) Following a scented beetle: larval faeces as a key olfactory cue in host location of *Stegobium paniceum* (Coleoptera: Anobiidae) by *Lariophagus distinguendus* (Hymenoptera: Pteromalidae). Chemoecology 23(2): 129–136. https://doi.org/10.1007/s00049-012-0125-7
- Beshear RJ (1969) Observations on the life cycle of *Hemisphaerota cyanea* in Georgia (Coleoptera: Chrysomelidae). Journal of the Georgia Entomological Society 4: 168–170.
- Blatchley WS (1910) The Coleoptera or Beetles Known to Occur in Indiana, pt. 2. Nature Publishing Co, Indianapolis, 1386 pp. https://doi.org/10.5962/bhl.title.79314
- Blatchley WS (1924) The Chrysomelidae of Florida. The Florida Entomologist 7: 49–57. https://doi.org/10.2307/3491997
- Blum MS (1994) Antipredator devices in larvae of Chrysomelidae: a unified synthesis for defensive ecleticism. In: Jolivet PH, Cox ML, Petitpierre E (Eds) Novel Aspects of the Biology of Chrysomelidae. Series Entomologica 50. Kluwer Academic Publishers, Dordrecht, 277–288. https://doi.org/10.1007/978-94-011-1781-4\_21
- Boheman CH (1850) Monographia Cassididarum. Holmiae (Vol. 1). 425 pp. https://doi. org/10.5962/bhl.title.8973
- Boheman CH (1854) Monographia Cassididarum. Holmiae (Vol. 2). 506 pp. https://doi. org/10.5962/bhl.title.8973
- Boheman CH (1855) Monographia Cassididarum. Tomus tertius. Holmiae, 543 pp. https://doi.org/10.5962/bhl.title.8973
- Boheman CH (1862) Monographia Cassididarum. Holmiae (Vol. 4). 504 pp. https://doi. org/10.5962/bhl.title.8973
- Bokerman WC (1964) Notes on the biology of *Poropleura cuprea* (Klug 1928) (Coleoptera, Chrysomelidae, Chlamisinae (14<sup>th</sup> contribution). Anais da Academia Brasileira de Ciências 36: 165–172.
- Borowiec L (1995) Tribal classification of the cassidoid Hispinae (Coleoptera: Chrysomelidae), In: Pakaluk J, Ślipiński SA (Eds) Biology, Phylogeny and Classification of Coleoptera. Muzeum I Instytut Zoologii, Warszawa, 451–558.
- Borowiec L, Świętojańska J (2002–present) World Catalog of Cassidinae, Wrocław, Poland. http://www.cassidae.uni.wroc.pl/katalog%20internetowy/index.htm [Accessed 12 February 2023]
- Borowiec L, Świętojańska J (2014) 2.7.2. Cassidinae Gyllenhall, 1813. In: Leschen RAB, RG Beutel (Eds) Coleoptera, Beetles (Vol. 3): Handbook of Zoology. De Gruyter, Berlin/ Boston, 198–217.
- Bottcher A, Zolin JP, Nogueira-de-Sá F, Trigo JR (2009) Faecal shield chemical defence is not important in larvae of the tortoise beetle *Chelymorpha reimoseri* (Chrysomelidae:

Cassidinae: Stolaini). Chemoecology 19(1): 63–66. https://doi.org/10.1007/s00049-009-0006-x

- Bouchard P, Bousquet Y, Davies A, Alonso-Zarazaga M, Lawrence J, Lyal C, Newton A, Reid C, Schmitt M, Ślipiński A, Smith A (2011) Family-group names in Coleoptera (Insecta). ZooKeys 88: 1–972. https://doi.org/10.3897/zookeys.88.807
- Brandmayr P (1992) Short review of the presocial evolution in Coleoptera. Ethology Ecology and Evolution 2(Special Issue): 7–16. https://doi.org/10.1080/03949370.1992.1 0721939
- Brandmayr P, Brandmayr Zetto T (1974) Sulle cure parentali e su altri aspetti della biologia de *Carterus (Sabienus) calydonius* Rossi, con alcune considerazioni sui fenomeni di cura della prole sino ad oggi riscontrati in Carabidi. Redia (Firenze) 20: 143–175.
- Brandt M, Mahsberg D (2002) Bugs with a backpack: The function of nymphal camouflage in the West African assassin bugs *Paredocla* and *Acanthmaspis* spp. Animal Behaviour 63(2): 277–284. https://doi.org/10.1006/anbe.2001.1910
- Briggs EM (1905) The Life History of Case Bearers: 1. *Chlamys plicata*. Cold Harbor Spring Monographs IV, 13 pp.
- Brown CG, Funk DJ (2005) Aspects of the natural history of *Neochlamisus* (Coleoptera: Chrysomelidae): faecal-case associated life history and behaviour, with a method for studying the construction of insect defensive structures. Annals of the Entomological Society of America 98(5): 711–725. https://doi.org/10.1603/0013-8746(2005)09 8[0711:AOTNHO]2.0.CO;2
- Bucheli S (2002) Larval case architecture and implications of host-plant associations for North American *Coleophora* (Lepidoptera: Coleophoridae). Cladistics 18(1): 71–93. https://doi.org/10.1006/clad.2001.0181
- Büsse S, Hörnschemeyer T, Hohu K, McMillan D, Edgerly JS (2015) The spinning apparatus of webspinners—Functional-morphology, morphometrics and spinning behaviour. Scientific Reports 4(1): e9986. https://doi.org/10.1038/srep09986
- Butterfield M, Hutchinson T (2017) How Animals Build (1<sup>st</sup> edn.). Lonely Planet Global Limited, Malaysia, 24 pp. https://shop.lonelyplanet.com/products/how-animals-build-us-1
- Buzzi ZJ (1975) Contribuçao ao conhecimento da biologia de *Anacassis fuscata* (Klug 1829) (Coleoptera, Chrysomelidae). Revista Brasileira de Biologia 35(4): 767–774.
- Buzzi ZJ (1976) Uma nova especie de *Anacassis* do sul do Brasil (Coleoptera, Chrysomelidae). Dusenia 9(3): 17–111. [ISSN 0418-1867]
- Buzzi ZJ (1988) Biology of Neotropical Cassidinae. In: Jolivet P, Petitpierre E, Hsiao TH (Eds) Biology of the Chrysomelidae. Kluwer Academic Publisher, Dordrecht, 559– 580. https://doi.org/10.1007/978-94-009-3105-3\_32
- Buzzi ZJ (1996) Morfologia dos imaturos e ciclo evolutivo de Anacassis dubia (Boheman), A. fuscata (Klug), A. languida (Boheman), A. phaeopoda Buzzi e A. punctulata (Klug) (Coleoptera: Chrysomelidae: Cassidinae). Revista Brasileira de Zoologia 13(1): 211–214. https://doi.org/10.1590/S0101-81751996000100020
- Buzzi ZJ, Miyazaki RD (1992) Calyptocephala paralutea sp. n. e descrição de larva e pupa (Coleoptera, Chrysomelidae, Cassidinae). Revista Brasileira Zoologica 9: 157– 166. https://doi.org/10.1590/S0101-81751992000100017
- Cai W, Zhao P, Mi Q (2002) Camouflaging in assassin bugs. Entomological Knowledge 39: 317–319. [In Chinese, English summary]
- Calcetas OA, Adorada JL, Staines CL, Chaboo CS, Anabo RAA (2023) Biology of the sineguelas leaf beetle, *Podontia quatuordecimpunctata* (L.) (Chrysomelidae: Galerucinae: Alticini), on *Spondias purpurea* L. (Anacardiaceae) in the Philippines. Insecta Mundi 1001: 1–39.

- Caldas A (1994) Biology of *Anaeaa ryphea* (Nymphalidaae) in Campinas, Brazil. Journal of the Lepidopterists Society 48(3): 248–257. https://www.biodiversitylibrary.org/part/80704
- Cardé RT, Bell WJ (1995) Chemical Ecology of Insects 2. Chapman & Hall, New York, 433 pp. https://doi.org/10.1007/978-1-4615-1765-8
- Carey D (1987) Sedimentological effects and palaeoecological implications of the tube-building polychaete *Lanice conchilega* Pallas. Sedimentology 34(1): 49–66. https://doi.org/10.1111/j.1365-3091.1987.tb00559.x
- Cedeño-Loja PE, Chaboo CS (2020) Natural history notes on *Cyrtonota sericinus* (Erichson, 1847) in Ecuador (Coleoptera: Chrysomelidae: Cassidinae: Mesomphaliini). Revista Peruana de Biología 27(2): 127–130. https://doi.org/10.15381/rpb.v27i2.17873
- Chaboo CS (2002) First report of immatures, genitalia and maternal care in *Eugenysa columbiana* (Boheman) (Coleoptera: Chrysomelidae: Cassidinae: Eugenysini). The Coleopterists Bulletin 56(1): 50–67. https://doi.org/10.1649/0010-065X(2002)056[0 050:FROIGA]2.0.CO;2
- Chaboo CS (2003) Tortoise beetles of Costa Rica: new records and localities (Coleoptera: Chrysomelidae: Cassidinae). Genus 14(1): 109–120. https://doi.org/10.1649/0 010-065X(2003)057[0071:ACOTBO]2.0.CO;2
- Chaboo CS (2004) Natural history observations in *Eurypepla calochroma* Blake (Chrysomelidae: Cassidinae: Physonotini). The Coleopterists Bulletin 58(1): 142–143. https://doi.org/10.1649/658
- Chaboo CS (2007) Biology and phylogeny of the Cassidinae (tortoise and leaf-mining beetles) (Coleoptera: Chrysomelidae). Bulletin of the American Museum of Natural History 305: e250. https://doi.org/10.1206/0003-0090(2007)305[1:BAPO TC]2.0.CO;2
- Chaboo CS, Engel MS (2008) Eocene tortoise beetles from the Green River Formation in Colorado, U.S.A. (Coleoptera: Chrysomelidae: Cassidinae). Systematic Entomology 34(2): 202–209. https://doi.org/10.1111/j.1365-3113.2008.00456.x
- Chaboo CS, Nguyen TC (2004) Immatures of *Hemisphaerota palmarum* (Boheman), with discussion of the caudal processes and shield architecture in the tribe Hemisphaerotini (Chrysomelidae: Cassidinae). In: Jolivet P, Santiago-Blay J, Schmitt M (Eds) New Developments in the Biology of Chrysomelidae. Kluwer Academic Press, 171–184. https://doi.org/10.1163/9789004475335\_019
- Chaboo CS, Brown CG, Funk D (2008) Faecal case architecture in the *gibbosus* species group of *Neochlamisus* Karren 1972 (Coleoptera: Chrysomelidae: Cryptocephalinae: Chlamisini): instar, host plant, species, generic and tribal variations. Zoological Journal of the Linnean Society 152(2): 315–351. https://doi.org/10.1111/j.1096-3642.2007.00343.x
- Chaboo CS, Engel MS, Chamorro-Lacayo ML (2009) Maternally-inherited architecture in Tertiary leaf beetles: Paleoichnology of cryptocephaline faecal cases in Dominican and Baltic amber. Naturwissenschaften 96(9): 1121–1126. https://doi.org/10.1007/ s00114-009-0573-2
- Chaboo CS, Frieiro-Costa FA, Gómez-Zurita J, Westerduijn R (2014) Subsociality in leaf beetles (Coleoptera: Chrysomelidae: Cassidinae, Chrysomelinae). Journal of Natural History 48(37–38): 1–44. https://doi.org/10.1080/00222933.2014.909060
- Chaboo CS, Chamorro-Lacayo L, Schöller M (2016) Catalog of known immature stages of Camptosomate leaf beetles (Coleoptera: Chrysomelidae: Cryptocephalinae and Lamprosomatinae). Proceedings of the Washington Entomological Society 118(2): 150–217. https://doi.org/10.4289/0013-8797.118.1.150

- Chapman RF (1982) The Insects Structure and Function (3<sup>rd</sup> edn.). Harvard University Press, Cambridge, 919 pp. [ISBN-10: 052111389X]
- Chapuis F (1875) In: Lacordaire JT (Ed.) Histoire Naturelle des Insectes. Genera des Coléoptères (Vol. 11), Famille des Phytophages. Encylopédique de Roret, Paris, 420 pp. https://doi.org/10.5962/bhl.title.8864
- Chen S (1964) Evolution and classification of the Chrysomelid beetles. Acta Entomologica Sinica 13(4): 469–483.
- Chen S (1985) Phylogeny and classification of the Chrysomeloidea. In: Furth DG, Seeno TN (Eds) First international symposium on the Chrysomelidae. Entomography 3: 465–475.
- Chevrolat LAA (1836) In: Dejean PFMA (Ed.) Catalogue des Coléoptères de la Collection de M. le compte Dejean. Ed. 3, revue, corrigée et augmentée. Paris. Livr. 5: 385–503. https://doi.org/10.5962/bhl.title.8771
- Claassen PW (1919) Life history and biological notes on *Chlaenius impunctifrons* Say (Coleoptera, Carabidae). Annals of the Entomological Society of America 12: 95–100. https://doi.org/10.1093/aesa/12.2.95
- Córdova-Ballona L, Sánchez-Soto S (2008) Bionomics data and descriptions of the immatures of *Calyptocephala gerstaeckeri* Boheman (Coleoptera: Chrysomelidae), pest of the oil palm (*Elaeis guineensis* J.) and camedor palm (*Chamaedorea elegans* Mart.) (Arecaceae) in Tabasco, Mexico. Neotropical Entomology 37(6): 674–680. https://doi.org/10.1590/S1519-566X2008000600008
- Crowson RA (1981) The Biology of the Coleoptera. Academic Press, London, 802 pp.
- Daanje A (1975) Some special features of the leaf-rolling technique of Byctiscus populi L. (Coleoptera: Rhynchitini). Behaviour 53: 285–316. https://doi. org/10.1163/156853975X00236
- D'Alessandro P, Biondi M (2023) Generic placement of the African flea beetle *Polyclada maculipennis* Bryant and the possible occurrence of the genus *Procalus* Clark in sub-Saharan Africa (Coleoptera, Chrysomelidae, Galerucinae, Alticini). ZooKeys 1145: 181–189. https://doi.org/10.3897/zookeys.1145.90667
- Darwin C (1859) The Origin of Species by Means of Natural Selection. John Murray, London, 477 pp. [reprinted 1985 as "The Origin of Species" in Penguin Classics, Clays Ltd., London] http://darwin-online.org.uk/content/frameset?itemID=F373&viewtype=text&pageseg=1
- Dawkins R (1982) The Extended Phenotype. W.H. Freeman, Oxford.
- Dewey JO (1991) Animal Architecture. Orchard Books, New York, 72 pp.
- Dispons P (1955) Les Réduvidés de l'Afrique nord-occidentale. Biologie et biogeographie. Mémoires du Muséum National d'Histoire Naturelle, Ser. A. Zoologie 10(2): 93-240.
- Duncan RP, Autumn K, Binford GJ (2007) Convergent setal morphology in sand-covering spiders suggests a design principle for particle capture. Proceedings of the Royal Society B, Biological Sciences 274(1629): 3049–3056. https://doi.org/10.1098/ rspb.2007.1039
- Duporte EM (1977) Manual of Insect Morphology. Robert E. Krieger Publishing Company, Huntington, 224 pp.
- Duprat H (2020) Miroir du Trichoptère/ The Caddisfly's Mirror (French/English). Fage Editor Lyon, France, 628 pp.
- Ehrlich PR, Raven PH (1964) Butterflies and plants: A study in coevolution. Evolution; International Journal of Organic Evolution 18(4): 586–608. https://doi. org/10.2307/2406212

- Eiseman B, Silen W, Bascom GS, Kauvar AJ (1958) Faecal enema as an adjunct in the treatment of pseudomembranous enterocolitis. Surgery 44(5): 854–859.
- Eisner T (1994) Integumental slime and wax secretion: Defensive adaptations of sawfly larvae. Journal of Chemical Ecology 20(10): 2789–2492. https://doi.org/10.1007/ BF02036205
- Eisner T (2003) For Love of Insects. Belknap Press, Cambridge, 448 pp.
- Eisner T, Eisner M (2000a) Defensive use of a faecal thatch by a beetle larva (*Hemisphaerota cyanea*). Proceedings of the National Academy of Sciences of the United States of America 97(6): 2632–2636. https://doi.org/10.1073/pnas.050002197
- Eisner T, Eisner M (2000b) Coiling into a sphere: Defensive behaviour of a trash-carrying chrysopid larva *Leucochrysa* (*Nodita*) *pavida* (Neuroptera: Chrysopidae). Entomological News 113: 6–10. https://biostor.org/reference/66885
- Eisner T, Silberglied RE (1988) A chrysopid larva that cloaks itself in mealybug wax. Psyche 95(1-2): 15-19. https://doi.org/10.1155/1988/97301
- Eisner T, van Tassel E, Carrel JE (1967) Defensive use of a 'faecal shield' by a beetle larva. Science 158(3807): 1471–1473. https://doi.org/10.1126/science.158.3807.1471
- Eisner T, Hicks K, Eisner M, Robson DS (1978) "Wolf-in-sheep's clothing" strategy of a predaceous insect larva. Science 199(4330): 790–794. https://doi.org/10.1126/science.199.4330.790
- Eisner T, Carrel JE, van Tassel E, Hoebeke ER, Eisner M (2001) Construction of a defensive thrash packet from sycamore leaf trichomes by a chrysopid larva (Neuroptera: Chrysopidae). Proceedings of the Entomological Society of Washington 104: 437– 446. https://biostor.org/reference/57175
- Engel H (1935) Biologie und ökologie von *Cassida viridis* L. Zeitschrift für Morphologie und Ökologie der Tiere 30: 1: 41–96. [ISSN: 2365-0087] https://doi.org/10.1007/ BF00418068
- Erber D (1968) Bau, Funktion und Bildung der Kotpresse mittleuropäischer Clytrinen und Cryptocephalinen (Coleoptera: Chrysomelidae). Zeitschrift für Morphologie der Tiere 62(3): 245–306. https://doi.org/10.1007/BF00401486
- Erber D (1969) Beitrag zur Entwicklungs-Biologie mitteleuropäischer Clytrinae und Cryptocephalinen (Coleoptera, Chrysomelidae). Zoologische Jahrbucher. Abteilung fur Systematik, Ökologie und Geographie der Tiere 96: 453–477.
- Erber D (1988) Biology of Camptostomata Clytrinae-Cryptocephalinae-Chlamisinae-Lamprosomatinae. In: Jolivet P, Petitpierre E, Hsiao TH (Eds) Biology of the Chrysomelidae. Kluwer Academic Publisher, Dordrecht, 513–552. https://doi.org/10.1007/978-94-009-3105-3\_30
- Erichson WF (1847) Conspectus insectorum coleopterorum, quae in Republica Peruana observata sunt. Archiv für Naturgeschichte 13: 67–185. https://biostor.org/reference/61389
- Fabre JH (1915) The Hunting Wasps. Dodd Mead, New York, 393 pp. https://www.gutenberg.org/files/67110/67110-h/67110-h.htm
- Fabricius JCh (1775) Systema Entomologiae, Sistens Insectorum Classes, Ordines, Genera, Species, Adiectis Synonymis, Locis, Descriptionibus, Observationibus. Officina Libraria Kortii, Flensburgi et Lipsiae, [32 +] 832 pp. https://doi.org/10.5962/bhl. title.36510
- Fabricius JCh (1798) Supplementum entomologiae systematicae. Hafniae, [II +] 572 pp. https://www.biodiversitylibrary.org/item/132638#page/5/mode/1up
- Farrell BD (1998) "Inordinate Fondness" explained: Why are there so many beetles? Science 281(5376): 555–559. https://doi.org/10.1126/science.281.5376.555

- Farrell BD, Mitter C, Futuyma DJ (1992) Diversification at the insect-plant interface. Bioscience 42(1): 34–42. https://doi.org/10.2307/1311626
- Ferry EE, Hopkins GR, Stocks AN, Mohammadi S, Brodie ED, Gall BG (2013) Do all portable cases constructed by larvae function in defense? Journal of Insect Science 13(5): 1–5. https://doi.org/10.1673/031.013.0501
- Fettköther R, Reddy GVP, Noldt U, Dettner K (2000) Effect of host and larval frass volatiles on behavioural response of the old-house borer *Hylotrupes bajulus* (L) (Coleoptera: Cerambycidae), in a wind tunnel bioassay. Chemoecology 10(1): 1–10. https://doi. org/10.1007/s000490050001
- Fiebrig K (1910) Cassiden und Cryptocephaliden Paraguays. Ihre Entwicklungsstadien und Schutzvorrichtungen. Zoologische Jahrbucher 2(Supplement 12): 161–264. https://doi.org/10.5962/bhl.title.60191
- Flinte V, Valverde de Macédo M (2004) Biology and seasonality of *Fulcidax monstrosa* (F.) (Chrysomelidae: Chlamisinae). The Coleopterists Bulletin 58(4): 457–465. https://doi.org/10.1649/629
- Flinte V, Borowiec L, de Freitas S, Viana JH, Fernandes F, Nogueira-de-Sá F, de Valverde de Macédo M, Monteiro RF (2009) Tortoise beetles of the State of Rio de Janeiro, Brazil (Coleoptera: Chrysomelidae: Cassidinae). Genus 20: 571–614.
- Flowers RW, Chaboo CS (2015) Natural history of the tortoise beetle, *Discomorpha* (*Discomorpha*) *biplagiata* (Guérin, 1844) (Chrysomelidae: Cassidinae: Omocerini). Insecta Mundi 439: 1–10. https://doi.org/10.5281/zenodo.5182476
- Frania HE (2011) Ecdysis and other aspects of metamorphosis in rove beetles with exarate or obtect pupae (Coleoptera: Staphylinidae). Canadian Journal of Zoology 70(12): 2326–2332. https://doi.org/10.1139/z92-312
- Freitas AVL, Oliveira PS (1992) Biology and behaviour of the neotropical butterfly *Eunica bechina* (Nymphalidae) with special reference to larval defense against ant predation. Journal of Research on the Lepidoptera 31(1–2): 1–11. https://doi. org/10.5962/p.266580
- Frost SW (1919) The function of the anal comb of certain lepidopterous larvae. Journal of Economic Entomology 12(6): 446–447. https://doi.org/10.1093/jee/12.6.446
- Frost SW (1942) Insect life and insect natural history. Dover Publications, New York, 526 pp.
- Frowhawk FW (1913) Faeces ejector of lepidopterous larvae. Entomologist 46(602): 201–202.
- Furth DG (1982) *Blepharida* biology, as demonstrated by the Sacred Sumac Flea Beetle (*B. sacra* Weise). Spixiana, Supplement 7: 43–52. [Corpus ID: 52065429]
- Furth DG (1988) The jumping apparatus of flea beetles (Alticinae) The metafemoral spring.
  In: Jolivet P, Petitpierre E, Hsiao TH (Eds) Biology of Chrysomelidae. Series Entomologica (Vol. 42). Springer, Dordrecht, 285–297. https://doi.org/10.1007/978-94-009-3105-3\_17
- Furth DG (2004) Fun with flea beetle faeces. Chrysomela newsletter 43: 1–10.
- Furth DG, Lee JE (2000) Similarity of the *Blepharida*-group genera using larval and adult characters (Coleoptera: Chrysomelidae: Alticinae). Journal of the New York Entomological Society 108(1): 26–51. https://doi.org/10.1664/0028-7199(2000)108[0026: SOTBGG]2.0.CO;2
- Gawryszewski FM (2014) Evidence suggests that modified setae of the crab spiders *Stephanopis* spp. fasten debris from the background. Zoomorphology 133(2): 205–215. https://doi.org/10.1007/s00435-013-0213-4
- Ge H (2000 [4<sup>th</sup> century]) The first use of donor faeces as a therapeutic agent for food poisoning and diarrhea was recorded in the [= Handbook of Emergency Medicine (Chinese)]. Tiangin Science & Technology Press, Tianjin.

- Gilby AR, McKellar JW, Beaton CD (1976) The structure of lerps: Carbohydrates, lipid and protein components. Journal of Insect Physiology 22(5): 689–696. https://doi. org/10.1016/0022-1910(76)90234-1
- Gómez NE (1997) The faecal shields of larva of tortoise beetles (Cassidinae: Chrysomelidae): a role in chemical defense using plant-derived secondary compounds. Ph.D. dissertation, Technische Universität Carolo-Wilhemina zu Braunschweig, 124 pp.
- Gómez NE (2004) Survivorship of immature stages of *Eurypedus nigrosignatus* Boheman (Chrysomelidae: Cassidinae: Physonotini) in Central Panama. The Coleopterists Bulletin 58(3): 73–84. [ISSN: 0010-065X] https://doi.org/10.1649/634
- Gómez NE, Witte L, Hartmann T (1999) Chemical defense in larval tortoise beetles: Essential oil composition of faecal shields of *Eurypedus nigrosignata* and foliage of its host plant, *Cordia curassavica*. Journal of Chemical Ecology 25(5): 1007–1027. https://doi.org/10.1023/A:1020821507014
- Gómez-Zurita J, Hunt T, Kopliku F, Vogler AP (2007) Recalibrated tree of leaf beetles (Chrysomelidae) indicates independent diversification of angiosperms and their insect herbivores. PLoS ONE 2(4): e360. https://doi.org/10.1371/journal.pone.0000360
- Gómez-Zurita J, Hunt T, Vogler AP (2008) Multilocus ribosomal RNA phylogeny of the leaf beetles (Chrysomelidae). Cladistics 24(1): 34–50. https://doi.org/10.1111/j.1096-0031.2007.00167.x
- Gould JR, Gould CG (2007) Animal Architects: Building and the Evolution of Intelligence. Basic Books, New York, 324 pp.
- Gressitt JL (1977) Papuan weevil genus *Gymnopholus*: Third supplement with studies in epizoic symbiosis. Pacific Insects 17(2–3): 179–195. https://eurekamag.com/research/006/060/006060248.php
- Gressitt JL, Samuelson GA, Vitt DH (1968) Moss growing on living Papua moss-forest weevils. Nature 217(5130): 765–767. https://doi.org/10.1038/217765a0
- Gressitt JL, Sedlacek J, Szent-Ivany JJH (1965) Fauna and flora on the backs of large Papuan moss-forest weevils. Science 150(3705): 1833–1835. https://doi.org/10.1126/science.150.3705.1833
- Gyllenhal L (1813) Insecta Suecica. Classis 1. Coleoptera sive Eleuterata, Tomus I, pars III. Scaris, 734 pp. https://doi.org/10.5962/bhl.title.8767
- Haddad S, McKenna DD (2016) Phylogeny and evolution of the superfamily Chrysomeloidea (Coleoptera: Cucujiformia). Systematic Entomology 41(4): 697–716. https:// doi.org/10.1111/syen.12179
- Hanley RS (1996) Immature stages of *Scaphisoma castaneum* Motschulsky (Coleoptera: Staphylinidae: Scaphidiinae), with observations of natural history, fungal hosts and development. Proceedings of the Entomological Society of Washington 98(1): 36–43. [biostor-56733]
- Hansell MH (1984) Animal Architecture and Building Behaviour. Longman, New York, 324 pp. [ISBN-13: 978-0582468153]
- Hansell M (2005) Animal Architecture. Oxford Animal Biology Series. Oxford University Press, Oxford, 322 pp. [ISBN: 0-19-850751-8 (hc); 0-19-850752-6 ISBN-13: 978-0198507529]
- Hansell M (2007) Built by Animals: the Natural History of Animal Architecture. Oxford University Press, Oxford, 268 pp. [ISBN-13: 978-0199205578]
- Haruyama N, Miyazaki Y, Nakahira K, Mochizuki A, Nomura M (2012) Developmental time and survival of trash-carrying versus naked green lacewings, with implications for their utility as augmentative biological control agents. Annals of the Entomological Society of America 105(6): 46–851. https://doi.org/10.1603/AN12003

- Harz K (1952) Ein Beitrag zur Biologie von *Reduvius personatus* L. (Rynchita/ Heteroptera). Nachrichtenblatt der Bayerischen Entomologen 1(10): 73–75.
- Hawkeswood TJ (1982) Notes on the life history of *Aspidomorpha maculatissima* Boheman (Coleoptera: Chrysomelidae: Cassidinae) at Townsville, North Queensland. Victorian Naturalist 99(3): 92–101.
- Hawkeswood TJ, Takizawa H (1997) Taxonomy, ecology and descriptions of the larva, pupa and adult of the Australian hispine beetle, *Eurispa vitatta* Baly (Insecta, Coleoptera, Chrysomelidae). Spixiana 20(3): 245–253. [BHL: 28280564]
- Hayashi M, Nomura M (2011) Larvae of the green lacewing *Malladades jardinsi* (Neuroptera: Chrysopidae) protect themselves against aphid-tending ants by carrying dead aphids on their backs. Applied Entomology and Zoology 46(3): 407–413. https://doi. org/10.1007/s13355-011-0053-y
- Heller KM (1901) Dritter Beitrag zur Papuanischen Käferfauna. Abhandlungen und Berichte des Königlichen Zoologischen und Anthropologisch-Ethnographischen Museums zu Dresden 10(2): 1–20.
- Henry DO, Cordova CE, Portillo M, Albert RM, DeWitt R, Emery-Barbier A (2016) Blame it on the goats? Desertification in the Near East during the Holocene. The Holocene 27(5): 625–637. [ISSN 0959-6836. SAGE Publications] https://doi. org/10.1177/0959683616670470
- Heron HDC (2007) The life history of *Aspidimorpha areata* (Coleoptera: Chrysomelidae: Cassidinae). African Entomology 15(1): 75–87. https://doi.org/10.4001/1021-3589-15.1.75
- Hincks WD (1952) The genera of the Cassidinae. Transactions of the Royal Entomological Society of London 103(10): 327–358. https://doi.org/10.1111/j.1365-2311.1952. tb01061.x
- Hinton HE (1951) On a little-known protective device of some chrysomelid pupae (Coleoptera). Proceedings of the Royal Entomological Society of London, Series A 26: 67–73. https://doi.org/10.1111/j.1365-3032.1951.tb00123.x
- Hinton HE (1981) Biology of Insect Eggs. Pergamon Press, Oxford, 1125 pp. [ISBN 0080215394, 9780080215396]
- Hislop R (1872) Note on the larva-case, etc., of *Clythra 4-punctata*. Entomologist's Monthly Magazine 8: 269–270. https://www.biodiversitylibrary.org/item/36494#page/7/ mode/1up
- Hobbs JJ (2004) Problems in the harvest of edible birds' nests in Sarawak and Sabah, Malaysian Borneo. Biodiversity & Conservation 13: 2209–2226. https://doi. org/10.1023/B:BIOC.0000047905.79709.7f
- Hoffman A (1954) Fauna de France, 59. Coléoptères Curculionidae (Deuxieme Partie). Lechevalier, Paris, 722 pp.
- Holmgren N (1907) Monographische Bearbeitung einer schalentragenden Mycetophilidenlarve (*Mycetophila ancyliformans* n.sp.). Zeitschrift für Wissenschaftliche Zoologie 38: 1–77.
- Hsiao TH, Windsor DM (1999) Historical and biological relationships among Hispinae inferred from 12S mtDNA sequence data. In: Cox ML (Ed.) Advances in Chrysomelidae biology 1: 39–50. Backhuys, Leiden, 671 pp.
- Huang Z-Z, Dong Z-Q, Liang Z-L, Zhang B, Xue H-J, Ge S-Q (2022) The fecal shield is a double-edged sword for larvae of a leaf beetle. Current Zoology 69(2): 173–180. https://doi.org/10.1093/cz/zoac026
- Hultgren KM, Stachowicz JJ (2009) Evolution of decoration in majoid crabs: A comparative phylogenetic analysis of the role of body size and alternative defensive strategies. American Naturalist 173(5): 566–578. https://doi.org/10.1086/597797

- Hunt T, Bergsten J, Levkanicova Z, Papadopoulou A, St. John O, Wild R, Hammond PM, Ahrens D, Balke M, Caterino MS, Gómez-Zurita J, Ribera I, Barraclough TG, Bocakova M, Bocak L, Vogler AP (2007) A Comprehensive Phylogeny of Beetles Reveals the Evolutionary Origins of a Superradiation. Science 318: 1913–1916. https://doi.org/10.1126/science.1146954
- Hutchins RE (1959) Insect Builders and Craftsmen. Rand McNally and Company, Chicago, 96 pp. [ASIN: B0007DXJDK]
- Jackson RR, Pollard SD (2007) Bugs with backpacks deter vision-guided predation by jumping spiders. Journal of Zoology 273(4): 358–363. https://doi.org/10.1111/ j.1469-7998.2007.00335.x
- Janzen DH (1979) Natural history of *Phelypera distigma* (Boheman), Curculionidae, a Costa Rican defoliator of *Guazuma ulmifolia* Lam. (Sterculiaceae). Brenesia 16: 213– 219. [ISSN: 0304-3711]
- Janzen DH (1983) *Guazuma ulmifolia* (Guácimo, Guácima, Caulote, Tapaculot). In: Janzen DH (Ed.) Costa Rican Natural History. University of Chicago Press, 246–248. https://doi.org/10.7208/chicago/9780226161204.001.0001
- Jeanne RL (1975) The adaptiveness of social wasp architecture. The Quarterly Review of Biology 50(3): 267–287. https://doi.org/10.1086/408564
- Jenks GE (1940) Dwarfs that live in their hats. Nature Magazine 33: 337-340.
- Jolivet P (1988a) Interrelationships Between Insects and Plants. CRC Press, Baton Rouge, 309 pp. [ISBN 9780367400422]
- Jolivet P (1988b) Une nouvelle stratégie de defense: La stratégie de defense annulaire (cycloalexie) chez quelques larves de Chrysomelides brésiliens. Bulletin de la Société Entomologique de France 92: 291–299. https://doi.org/10.3406/bsef.1987.17510
- Jolivet P, Verma KK (2002) Biology of Leaf Beetles. Intercept, Andover, 332 pp. [ISBN: 1898298866]
- Jones RA (1994) Merdigery and maternal care in a leaf beetle. Entomologist's Record and Journal of Variation 106: 7–12.
- Jordan N, Cherry M, Manser MB (2007) Latrine distribution and patterns of use by wild meerkats: implications for territory and mate defence. Animal Behaviour 73(4): 613– 622. https://doi.org/10.1016/j.anbehav.2006.06.010
- Kalaichelvan T, Verma KK (2000) Faecal cover of eggs of Indian Cassidinae (Coleoptera: Chrysomelidae). Insect Environment 6(1): 41–42.
- Karren JB (1972) A revision of the subfamily Chlamisinae of America north of Mexico (Coleoptera: Chrysomelidae). The University of Kansas Science Bulletin 49(12): 875–988. https://www.jstor.org/stable/3999451
- Kasahara M, Akimotoa S-I, Hariyama T, Takaku Y, Yusa S-I, Okada S, Nakajima K, Hirai T, Mayama H, Okada S, Deguchi S, Nakamura Y, Fugii S (2019) Liquid Marbles in Nature: Craft of Aphids for Survival. Langmuir 35(18): 6169–6178. https://doi.org/10.1021/ acs.langmuir.9b00771
- Kaufmann DL (1967) Notes on the biology of three species of *Lema* (Coleoptera: Chrysomelidae) with larval descriptions and key to described United States species. Journal of the Kansas Entomological Society 40(3): 361–372. https://www.jstor.org/ stable/25083643
- Keefover-Ring K (2013) Making scents of defense: Do faecal shields and herbivore-caused volatiles match host plant chemical profiles? Chemoecology 23(1): 1–13. https://doi.org/10.1007/s00049-012-0117-7
- Keefover-Ring K (2015) Bergamot versus beetle:evidence for intraspecific chemical specialization. AoB PLANTS7 7: plv132. [1–10] https://doi.org/10.1093/aobpla/plv132

- Kershaw JC, Muir F (1907) On the egg-cases and early stages of some South China Cassididae. The Transactions of the Entomological Society of London 1907(2): 249–252. https://doi.org/10.1111/j.1365-2311.1907.tb01762.x
- King James Bible Online (2023) Matthew 7:15. https://www.kingjamesbibleonline.org/ Matthew-7–15/ [Accessed 21 Jan 2023]
- Kinsley CB, Pearson DL (1981) The function of turret building behaviour in the larval tiger beetle *Cicindela willistoni* (Coleoptera: Cicindelidae). Ecological Entomology 6: 401–410. [ISSN: 0307-6946] https://doi.org/10.1111/j.1365-2311.1981.tb00631.x
- Klug JCF (1824) Entomologische Monographien. G. Reimer, Berlin, [10 pls +] 242 pp. https://doi.org/10.5962/bhl.title.3346
- Knab F (1915) Dung-bearing weevil larvae. Proceedings of the Entomological Society of Washington 17(4): 193–194.
- Knab F, van Zwaluwenburg RH (1918) A second *Mycetophila* with dungbearing larva (Diptera; Mycetophilidae). Entomological News 29(4): 138–142.
- Krishna K, Weesner FM (1969) Biology of Termites (Vol. 1). Academic Press, New York, 612 pp. https://doi.org/10.1007/BF02040996
- Larsen A, Larsen J, Lane S (2021) Dam builders and their works: Beaver influences on the structure and function of river corridor hydrology, geomorphology, biogeochemistry and ecosystems. Earth-Science Reviews 218(5): e103623. https://doi.org/10.1016/j. earscirev.2021.103623
- Latin is Simple (2023) Erhalt und Digitalisierung indoeuorpäischer Sprachen, Graz, Austria. https://www.latin-is-simple.com/en/ [Accessed 8 May 2023]
- Lawson FA (1976) Egg and larval case formation by *Pachybrachis bivittatus*. Annals of the Entomological Society of America 69(5): 942–944. https://doi.org/10.1093/ aesa/69.5.942
- Lécaillon A (1896) Note relative a la coque excrémentielle des oeufs et des larves de certaines insectes en particulier du *Clytra quadripunctata*. Comptes Rendus Hebdomadaires des Séances et Mémoires de la Société de Biologie 3: 506–510.
- Lee JE, Morimoto K (1991a) Descriptions of the egg and first-instar larva of *Clytra arida* (Weise) (Coleoptera: Chrysomelidae). Journal of the Faculty of Agriculture, Kyushu University 35(3–4): 93–101. [Corpus ID: 83245949] https://doi.org/10.5109/23954
- Lee JE, Morimoto K (1991b) The egg and first-instar larva of *Oomorphoides cupreatus* (Baly) from Japan, with notes on the systematic position of Lamprosomatinae (Coleoptera: Chrysomelidae). Journal of the Faculty of Agriculture, Kyushu University 35(3-4): 101–107. [ISSN: 0023-6152] https://doi.org/10.5109/23955
- Legendre F, Nel A, Svenson GJ, Robillard T, Pellens R, Grandcolas P (2015) Phylogeny of Dictyoptera: Dating the origin of cockroaches, praying mantises and termites with molecular data and controlled fossil evidence. PLoS ONE 10(7): e0130127. https://doi.org/10.1371/journal.pone.0130127
- LeSage L (1982) The immature stages of *Exema canadensis* Pierce (Coleoptera: Chrysomelidae). The Coleopterists Bulletin 36(2): 318–327. https://www.jstor.org/stable/4008075
- Leschen RAB (1994) Retreat building by larval Scaphidiinae (Staphylinidae). Mola 4: 3–5.
- Leschen RAB (2003) Erotylidae (Insecta: Coleoptera: Cucujoidea): Phylogeny and review. Fauna of New Zealand 47: 1–108.
- Leschen RAB, Beutel RG (2014) Volume 3: Morphology and Systematics (Phytophaga). Handbook of Zoology. Walter de Gruyter, Berlin. [ISBN 978-3-11-027370-0] https:// doi.org/10.1515/9783110274462

- Leschen RAB, Carlton CE (1993) Debris cloaking in Endomychidae: A new species from Peru (Coleoptera). Zoological Journal of the Linnean Society 109(1): 35–51. https:// doi.org/10.1111/j.1096-3642.1993.tb01258.x
- Lingafelter SW, Konstantinov AS (2000) The monophyly and relative rank of alticine and galerucine leaf beetles: a cladistic analysis using adult morphological characters (Coleoptera: Chrysomelidae. Entomologica Scandinavica 30(4): 397–416. https://doi.org/10.1163/187631200X00525
- Linnaeus C (1758) Systema Naturae, Sive Regna Tria Naturae, Secundum Classes, Ordines, Genera, Species, cum Characteribus, Differentiis, Synonymis, Locis. Editio Decima, reformata. I. Holmiae, [IV +] 824 pp. https://doi.org/10.5962/bhl.title.542
- Loefler CC (1996) Adaptive trade-offs of leaf folding in *Dichomeris* caterpillars on goldenrods. Ecological Entomology 21(1):34–40. https://doi.org/10.1111/j.1365-2311.1996. tb00263.x
- Lüscher M (1961) Air-conditioned termite nests. Scientific American 205(1): 138–145. https://doi.org/10.1038/scientificamerican0761-138
- Machado G, Freitas AVL (2001) Larval defence against ant predation in the butterfly *Smyrna blomfildia*. Ecological Entomology 26(4): 436–439. https://doi.org/10.1046/ j.1365-2311.2001.00328.x
- Mano H, Toquenaga Y (2008) Wall-making behaviour in *Callosobruchus maculatus* (Coleoptera: Bruchidae). Annals of the Entomological Society of America 101(2): 449–455. https://doi.org/10.1603/0013-8746(2008)101[449:WBICMC]2.0.C0;2
- Marcone MF (2005) Characterization of the edible bird's nest the 'Caviar of the East'. Food Research International 38(10): 1125–1134. https://doi.org/10.1016/j. foodres.2005.02.008
- Mathews RW, Mathews JR (1978) Insect Behaviour. John Wiley & Sons, New York, 507 pp. [ISBN 13: 9780471576853]
- Maulik S (1931) On the structure of larvae of Hispine beetles I. Proceedings of the Zoological Society of London 1931(3): 1137–1162. https://doi. org/10.1111/j.1096-3642.1931.tb01055.x
- Maulik S (1932) On the structure of larvae of Hispine beetles II. Proceedings of the Zoological Society of London 1932: 293–322. https://doi.org/10.1111/j.1096-3642.1932. tb01078.x
- McCook HC (1907) Nature's Craftsmen: Popular Studies of Ants and Other Insects. Harper & Bros., New York, 317 pp. [ISBN-13: 978-1167000881]
- McDougal C, Degnan BM (2018) The evolution of mollusc shells. Wiley Interdisciplinary Reviews. Developmental Biology 7(3): e313. https://doi.org/10.1002/wdev.313
- McFarland N (1980) Retention of cast head capsules by some nolid immatures in four Old World countries. Journal of Research on the Lepidoptera 17(4): 209–217. [ISSN: 0022-4324] https://doi.org/10.5962/p.333757
- McMahan EA (1982) Bait-and-capture strategy of a termite-eating assassin bug. Insectes Sociaux 29(2): 346–351. https://doi.org/10.1007/BF02228761
- McMahan EA (1983a) Bugs angle for termites. Natural History, New York 83: 40–46.
- McMahan EA (1983b) Adaptations, feeding preferences, and biometrics of a termite-baiting assassin bug (Hemiptera: Reduviidae). Annals of the Entomological Society of America 76(3): 483–486. https://doi.org/10.1093/aesa/76.3.483
- Monrós F, Viana MJ (1949) Revisión de las especies Argentinas de Dorynotini (Col., Cassidinae). 1 Contribución al conocimiento de Cassidinae. Acta Zoológica Lilloana 8: 391–426.
- Monrós F, Viana MJ (1951) Las Cassidinae de la sección Hemisphaerotina con revisión de las especies argentinas. (Col., Cassidinae). Acta Zoológica Lilloana 11: 367–395.

- Monteith G (1970) Life history of the chrysomelid, *Aproidea bayli* Pascoe. News Bulletin Entomological Society of Queensland 72: 9–10.
- Monteith GB, Sandoval Gomez VE, Chaboo CS (2021) Natural history of the Australian tortoise beetle *Notosacantha dorsalis* (Waterhouse, 1877) (Coleoptera: Chrysomelidae: Cassidinae: Notosacanthini) with summary of the genus in Australia. Australian Entomologist 48(4): 329–354.
- Moore JH (2001) The Nightingale Facial. Asian Wall Street Journal. [Dec 14 2001] https://www.wsj.com/articles/SB1008292800778950720
- Morton TC (1997) Sequestration of host-plant-chemistry into frass-based defenses of Chrysomelidae: *Lema trilineata, Neolema sexpunctata* (Criocerinae), *Plagiometriona clavata* (Cassidinae) and *Blepharida rhois* (Alticinae). Pennsylvania State University. Ph.D. Thesis, 88 pp.
- Morton TC, Vencl FV (1998) Larval beetles form a defense from recycled host-plant chemicals discharged as faecal wastes. Journal of Chemical Ecology 24(5): 765–785. https://doi.org/10.1023/A:1022382931766
- Mound LA, Morris DC (1999) *Carcinothrips*: A genus of *Acacia* phyllode-glueing thrips with grossly enlarged fore legs (Thysanoptera: Phlaeothripidae). Australian Journal of Entomology 38(1): 10–14. https://doi.org/10.1046/j.1440-6055.1999.00076.x
- Muir F, Sharp D (1904) On the egg cases and early stages of some Cassididae. The Transactions of the Entomological Society of London 1904(1): 1–23. https://doi.org/10.1111/j.1365-2311.1904.tb02735.x
- Müller C (2002) Variation in the effectiveness of abdominal shields of cassidine larvae against predators. Entomologia Experimentalis et Applicata 102: 191–198. https://doi.org/10.1046/j.1570-7458.2002.00939.x
- Müller C, Hilker M (1999) Unexpected reactions of a generalist predator towards defensive devices of cassidine larva (Coleoptera: Chrysomelidae). Oecologia 118(2): 166–172. https://doi.org/10.1007/s004420050715
- Müller C, Hilker M (2001a) Exploitation of the faecal shield of lily leaf beetle, *Lilioceris lilii* (Coleoptera: Chrysomelidae), by the specialist parasitoid *Lemophagus pulcher* (Hymenoptera: Ichneumonidae). Journal of Insect Behavior 14(6): 739–757. https:// doi.org/10.1023/A:1013085316606
- Müller C, Hilker M (2001b) Host finding and oviposition behaviour in a chrysomelid specialist – the importance of host plant surface waxes. Journal of Chemical Ecology 27(5): 985–994. https://doi.org/10.1023/A:1010343205114
- Müller C, Hilker M (2003) The advantages and disadvantages of larval abdominal shields on the Chrysomelidae: a mini-review. In: Furth DG (Ed.) Special Topics in Leaf Beetle Biology. Pensoft Publishers, Sofia-Moscow, 243–259. [ISBN-13: 978-9546421708]
- Müller C, Hilker M (2004) Ecological relevance of faecal matter in Chrysomelidae. In: Jolivet P, Santiago-Blay J, Schmitt M (Eds) New Developments in the Biology of the Chrysomelidae. Kugler Publications, The Hague, 693–705. https://doi.org/10.1163/9789004475335\_059
- Mummery RS, Valadon LRG (1974) Carotenoids of the lily beetle (*Lilioceris lilii*) and of its food plant (*Lilium hansonii*). Journal of Insect Physiology 20(3): 429–433. https://doi.org/10.1016/0022-1910(74)90073-0
- Nagasawa A, Matsuda K (2005) Effects of feeding experience on feeding responses to spinach in *Cassida nebulosa* L. (Coleoptera: Chrysomelidae). Applied Entomology and Zoology 40(1): 83–89. https://doi.org/10.1303/aez.2005.83
- Nakahira K, Arakawa R (2006) Defensive functions of the trash-package of a green lacewing, Mallada desjardinsi (Neuroptera: Chrysopidae), against a ladybird, Harmonia axyridis

(Coleoptera: Coccinellidae). Applied Entomology and Zoology 41: 111–115. https:// doi.org/10.1303/aez.2006.111

- Nassar D, Blasco JA (2015) Animal Architects: Amazing Animals Who Build Their Homes. Laurence King Publishing, London, 32 pp. [ISBN-13: 978-1780676548]
- Nicholson J (2003) Animal Architects. Allen Unwin, Crow's Nest. 32 pp. [ISBN-13: 978-1865089553]
- Nie R-E, Bezděk J, Yang X-K (2017) How many genera and species of Galerucinae s. str. do we know? Updated statistics (Coleoptera, Chrysomelidae). In: Chaboo CS, Schmitt M (Eds) Research on Chrysomelidae 7. ZooKeys 720: 91–102. https://doi. org/10.3897/zookeys.720.13517
- Nie RE, Andújar C, Gómez-Rodríguez C, Bai M, Xue HJ, Tang M, Yang CT, Tang P, Yang XK, Vogler AP (2020) The phylogeny of leaf beetles (Chrysomelidae) inferred from mitochondrial genomes. Systematic Entomology 45(1): 188–204. https://doi.org/10.1111/syen.12387
- Nishida K (2014) Discovered a Strange Insect with Several Heads Attached (Vol. 73). Insect-Centered Life, Web National Geographic Japan. https://natgeo.nikkeibp.co.jp/ nng/article/20140421/393881/ [In Japanese] [Accessed 15 Feb 2023]
- Nishida K (2015) The Miracle World of Insects in Costa Rica. National Geographic Japan, Tokyo, 128 pp.
- Nishida K, Ferrufino-Acosta L, Chaboo CS (2020) A new host plant family for Cassidinae sensu lato: *Calyptocephala attenuata* (Spaeth, 1919) (Coleoptera: Chrysomelidae: Cassidinae: Spilophorini) on *Smilax* (Smilacaceae) in Costa Rica. The Pan-Pacific Entomologist 96(4): 263–267. https://doi.org/10.3956/2020-96.4.263
- Nogueira-de-Sá F, Trigo JR (2002) Do faecal shields provide physical protection to larvae of the tortoise beetles *Plagiometriona flavescens* and *Stolas chalybea* against natural enemies. Entomologia Experimentalis et Applicata 104(1): 203–206. https://doi.org/10.1046/j.1570-7458.2002.01007.x
- Nogueira-de-Sá F, Trigo JR (2005) Faecal shield of the tortoise beetle *Plagiometriona* aff. *flavescens* (Chrysomelidae: Cassidinae) as chemically mediated defence against predators. Journal of Tropical Ecology 21(2): 189–194. https://doi.org/10.1017/S0266467404002147
- Odihiambo T (1958) Some observations on the natural history of *Acanthaspis petax* Stål (Hemiptera: Reduviidae) living in termite mounds in Uganda. Physiological Entomology 33(10–12): 167–175. https://doi.org/10.1111/j.1365-3032.1958.tb00449.x
- Olivier AG (1790) Encyclopédie Méthodique, Histoire Naturelle. Insectes. Tome cinquiéme. Paris, 368 pp. https://doi.org/10.5962/bhl.title.82248
- Olivier AG (1808) Entomologie, ou histoire naturelle des Insectes, avec leur caracteres géneriques et spécifiques, leur description, leur synonymie, et leur figure enluminée. Coléopteres, Vol. VI. Paris. 613–1104 pp. https://doi.org/10.5962/bhl.title.61905
- Olmstead KL (1994) Waste products as chrysomelid defenses. In: Jolivet PH, Cox ML, Petitpierre E (Eds) Novel Aspects of the Biology of Chrysomelidae. Series Entomologica 50. Kluwer Academic Publishers, Dordrecht, 311–318. https://doi.org/10.1007/978-94-011-1781-4\_24
- Olmstead KL (1996) Cassidine defences and natural enemies. In: Jolivet PHA, Cox ML (Eds) Chrysomelidae Biology: Ecological Studies (Vol. 2). SPB Academic Publishers, Amsterdam, 25 pp.
- Olmstead KL, Denno RF (1992) Cost of shield defence for tortoise beetles (Coleoptera: Chrysomelidae). Ecological Entomology 17(3): 237–243. https://doi. org/10.1111/j.1365-2311.1992.tb01053.x

- Otto C, Svenson BS (1980) The significance of case material selection for the survival of caddis larvae. Journal of Animal Ecology 49(3): 855–865. https://doi.org/10.2307/4231
- Pearson G (2013) The Mad Hatterpillar. WIRED magazine. https://www.wired. com/2014/01/the-mad-hatterpillar/ [Retrieved 11 Nov 2022]
- Prathapan KD, Chaboo CS (2011) Natural history of *Podontia congregata* Baly 1865, an endemic flea beetle from southern India (Coleoptera: Chrysomelidae: Galerucinae: Alticini). ZooKeys 157: 95–130. https://doi.org/10.3897/zookeys.157.1472
- Prathapan KD, Karthikeyan K, Chaboo CS (2009) Natural history and leaf shelter constructions of the Asian rice leptispa beetle, *Leptispa pygmaea* Baly (Coleoptera: Chrysomelidae: Cassidinae: Leptispini). Zoological Studies (Taipei, Taiwan) 48(5): 625–631.
- Preston-Mafham K, Preston-Mafham R (2003) The Natural World of Bugs & Insects. PRC Publishing Ltd., London, 512 pp. [ISBN-13: 978-1571452887]
- Prevett PF (1966) Observations on biology in the genus *Caryedon* Schoenherr in Northern Nigeria, with a list of parasitic Hymenoptera. Proceedings of the Entomological Society of London 41: 9–16. https://doi.org/10.1111/j.1365-3032.1966.tb01127.x
- Ramírez PA, González A, Botto-Mahan C (2013) Masking behaviour by Mepraia spinolai (Hemiptera: Reduviidae): Anti-predator defense and life history trade-offs. Journal of Insect Behavior 26(4): 592–602. https://doi.org/10.1007/s10905-012-9371-3
- Rane N, Ghate HV (2005) Notes on the life history of *Laccoptera* (*Sindia*) *sulcata* (Olivier) (Coleoptera: Chrysomelidae: Cassidinae). Journal of the Bombay Natural History Society 102(3): 217–222.
- Réaumur R-AFde (1737) Mémoires pour server à l'histoire des insects. Tome 3. Imprimie Royale, Paris. https://doi.org/10.5962/bhl.title.14146
- Reid CAM (1992a) Descriptions of pupae of nine genera of Australian paropsine Chrysomelinae (Coleoptera: Chrysomelidae). Proceedings of the Linnean Society of New South Wales 113(4): 311–337.
- Reid CAM (1992b) The leaf-beetle genus *Microdonacia* Blackburn (Coleoptera, Chrysomelidae, Galerucinae): Revision and systematic placement. Systematic Entomology 17(4): 359–387. https://doi.org/10.1111/j.1365-3113.1992.tb00557.x
- Reid CAM (1995) A cladistic analysis of subfamilial relationships of the Chrysomelidae sensu lato (Chrysomeloidea). In: Pakaluk J, Ślipiński, SA (Eds) Biology, Phylogeny and Classification of Coleoptera: Papers Celebrating the 80<sup>th</sup> Birthday of Roy A. Crowson.
  2. Muzeum I Instytut Zoologii PAN, Warsaw, 559–631.
- Reid CAM (2000) Spilopyrinae Chapuis: A new subfamily in the Chrysomelidae and its systematic placement (Coleoptera). Invertebrate Taxonomy 14(6): 837–862. https://doi.org/10.1071/IT00042
- Reid CAM, Beatson M (2019) Descriptions of the larva and pupa of *Mecynodera balyi* Clark, 1864, with notes on its life history (Coleoptera: Chrysomelidae: Sagrinae). Zootaxa 4686(4): 551–570. https://doi.org/10.11646/zootaxa.4686.4.5
- Rodrigues P, Rodrigo RK (2009) Shell occupancy by the hermit crab *Clibanarius erythropus* (Crustacea) on the south coast of São Miguel, Açores. Acoreana (Suplemento 6): 211–216.
- Ruxton GD, Stevens M (2015) The evolutionary ecology of decorating behaviour. Biology Letters 11(6): e20150325. https://doi.org/10.1098/rsbl.2015.0325
- Sakagami SF, Michener CD (1962) The Nest Architecture of the Sweat Bees (Halictinae). University of Kansas Press, Lawrence, 135 pp.
- Schaffner U, Müller C (2001) Exploitation of the faecal shield of the lily leaf beetle, *Lilioceris lilii* (Coleoptera: Chrysomelidae), by the specialist parasitoid *Lemophagus pulcher*

(Hymenoptera: Ichneumonidae). Journal of Insect Behavior 14(6): 740–757. https:// doi.org/10.1023/A:1013085316606

- Scherf H (1964) Die Entwicklungsstadien der mitteleuropäischen Curculioniden (Morfologie, Bionomie, Ökologie). Abhandlungen der Senckenbergischen Naturforschenden Gesellshcaft, 506. Verlag Waldemar Kramer, Frankfurt am Main, 335 pp.
- Schmitt M (1994) Stridulation in leaf beetles (Coleoptera, Chrysomelidae). In: Jolivet P (Ed.) Novel Aspects of the Biology of Chrysomelidae. Series Entomologica 50. Kluwer Academic Publishers, Dordrecht, 319–325. https://doi.org/10.1007/978-94-011-1781-4\_24
- Scholtz CH, Grebennikov VV (2004) Arthropoda. In: Kristensen NP, Beutel RG (Eds) Handbook of Zoology Vol. IV, Arthropoda, Part II, Insecta. Coleoptera Vol 1, Chap 12. Walter De Gruyter, Berlin.
- Scudder SH (1891) The early stages of three Coleoptera. Psyche 6: 173-175. https:// doi.org/10.1155/1891/91932
- Schuh RT (1989) Torre-Bueno Glossary of Entomology (revised edition). New York Entomological Society, New York, 840 pp. [ISBN 0-913424-13-7]
- Sekerka L, Windsor D, Dury G (2014) *Cladispa* Baly: revision, biology and reassignment of the genus to the tribe, Spilophorini (Coleoptera: Chrysomelidae: Cassidinae). Systematic Entomology 39(3): 518–530. https://doi.org/10.1111/syen.12070
- Sharp D (1899) Hymenoptera Continued (Tubulifera and Aculeata), Coleoptera, Strepsiptera, Lepidoptera, Diptera, Aphaniptera, Thysanoptera, Hemiptera, Anoplura (Part II). MacMillan & Co., New York, 626 pp. https://lccn.loc.gov/05012145
- Shivashankar T, Kumar AR, Veeresh GK, Pearson DL (1988) Angular turret-building behavior in a larval Tiger beetle Species from South India (Coleoptera: Cicindelidae). The Coleopterists Bulletin 42(1): 63–68. http://www.jstor.org/stable/4008565
- Skuhrovec J, Stejskal R, Trnka F, di Giulio A (2017) Velcro-like system used to fix a protective faecal shield on weevil larvae. PLoS ONE 12(1): e0170800. https://doi.org/10.1371/journal.pone.0170800
- Smeathman H (1781) Some accounts of the termites which are found in Africa and other hot climates. In a letter from Mr. Henry Smeathman, of Clement's Inn, to Sir Joseph Banks, Bart. P.R.S. Philosophical Transactions of the Royal Society of London LXX(I): 139–192. https://doi.org/10.1098/rstl.1781.0033
- Smith RG (2010) Wax glands, wax production and the functional significance of wax use in three aphid species (Homoptera: Aphididae). Journal of Natural History (4): 513–530. https://doi.org/10.1080/002229399300227
- Smith RC (1922) The biology of the Chrysopidae. Cornell University Agricultural Experiment Station Memoir 58: 1286–1376.
- Smith RC (1923) The life histories and stages of some hemerobiids and allied species (Neuroptera). Annals of the Entomological Society of America 16(2): 129–151. https://doi.org/10.1093/aesa/16.2.129
- Smith A, Oechsner A, Rowley-Conwy P, Moore AMT (2022) Epipalaeolithic animal tending to Neolithic herding at Abu Hureyra, Syria (12,800–7,800 calBP): Deciphering dung spherulites. PLoS ONE 17(9): e0272947. https://doi.org/10.1371/journal.pone.0272947
- Snodgrass RE (1935) Principles of Insect Morphology. McGraw-Hill Book Co., NY, republished 1993, Cornell University Press, 667 pp.
- Song N, Yin X, Zhao X, Chen J, Yin J (2017) Reconstruction of mitogenomes by NGS and phylogenetic implications for leaf beetles. Mitochondrial DNA, Part A, DNA Mapping, Sequencing, and Analysis 29(7): 1041–1050. https://doi.org/10.1080/24701394.20 17.1404044

- Spruyt FJ (1925) Observations on the egg-laying habits of *Saxinis saucia* LeC. (Coleoptera-Chrysomelidae). Pan-Pacific Entomologist 1: 176–178. [BHL: 53385839]
- Staines CL (2002) The new world tribes and genera of hispines (Coleoptera: Chrysomelidae: Cassidinae). Proceedings of the Entomological Society of Washinghton 104: 721–784. [BHL: 16186338]
- Staines CL (2015) Catalog of the hispines of the world (Coleoptera: Chrysomelidae: Cassidinae). https://naturalhistory.si.edu/research/entomology/collections-overview/ coleoptera/catalog-hispines-world [Accessed 30 Nov 2021]
- Stammer HJ (1935) Studien an Symbiosen zwischen Käfern und Mikroorganismen I. Die Symbiose der Donaciinen (Coleopt. Chrysomel.). Zeischrift für Morphologie und Ökologie Tiere 29: 585–608. https://doi.org/10.1007/BF00407434
- Stewart PD, Macdonald DW, Newman C, Cheeseman CL (2001) Boundary faeces and matched advertisement in the European badger (*Meles meles*): A potential role in range extension. Journal of Zoology 255(2): 191–198. https://doi.org/10.1017/ S0952836901001261
- Stitz H (1931) Planipennia. In: Schulze P (Ed.) Biologie der Tiere Deutschlands. Borntraeger, Berlin, Teil 35: 68–304.
- Stromberg J (2012) This insect uses its victims' carcasses as camouflage. Smithsonian Magazine. https://www.smithsonianmag.com/science-nature/this-insect-uses-its-victims-carcasses-as-camouflage-83656246/
- Suzuki T (1985) Presence of another aggregation substance(s) in the frass of the red flour beetles, *Tribolium castaneum* (Coleoptera: Tenebrionidae). Applied Entomology and Zoology 20: 90–91. https://doi.org/10.1303/aez.20.90
- Świętojańska J (2009) The immatures of tortoise beetles with review of all described taxa (Coleoptera: Chrysomelidae: Cassidinae). Polish Taxonomical Monographs 16: 157 pp. [ISBN: 9788361764045]
- Świętojańska J, Borowiec L (2007) Comparative description of last instar larva of *Cassida uniformis* Boheman, 1862 and *Cassida varians* Herbst, 1799 (Coleoptera: Chrysomelidae: Cassidinae). Genus 18(2): 297–314. [Corpus ID: 83311149]
- Świętojańska J, Moradian H, Borowiec L (2013) Description of larvae of two closely related species *Cassida palaestina* Reiche, 1858 and *Cassida rubiginosa* Müller, 1776 (Coleoptera: Chrysomelidae: Cassidinae). Zootaxa 3741(4): 511–537. https://doi.org/10.11646/zootaxa.3741.4.4
- Takizawa H (1978) Notes on Taiwanese Chrysomelid Larvae. V. Entomological Review of Japan 31(1/2): 75–85.
- Takizawa H (1980) Immature stages of some Indian Cassidinae (Coleoptera: Chrysomelidae). Insecta Matsumurana 21: 19–48. [Corpus ID: 81764404]
- Tamayo-Castillo G, Jakupovic J, Castro V, King RM (1989) (+)-α-copaen-8-one and other constituents from *Neomirandea* species. Phytochemistry 28(3): 938–940. https:// doi.org/10.1016/0031-9422(89)80149-9
- Tauber CA, Tauber MJ, Albuquerque GS (2014) Debris-carrying in larval Chrysopidae: Unraveling its evolutionary history. Annals of the Entomological Society of America 107(2): 295–314. https://doi.org/10.1603/AN13163
- Tishechkin AK, Konstantinov AS, Bista S, Pemberton RW, Center TD (2011) Review of the continental Oriental species of *Lilioceris* Reitter (Coleoptera, Chrysomelidae, Crioce-rinae) closely related to *Lilioceris impressa* (F.). ZooKeys 103: 63–83. https://doi.org/10.3897/zookeys.103.983
- Turner JS (2000) The Extended Organism: The Physiology of Animal-Built Structures. Harvard University Press, Cambridge, 235 pp. [ISBN-13: 978-0674009851]

- Vanin SA, Bená DdeC (2020) Description of larva and pupa of the weevil Hybolabus amazonicus Voss (Attelabidae: Attelabinae), a leaf roller on Brazil nut (Bertholletia excelsa). Papeis Avulsos de Zoolgia 60(Special-issue): e202060(s.i.).03. http://doi. org/10.11606/1807-0205/2020.60.special-issue.03
- Van Leerdam MB, Smith Jr JW, Fuchs JR, Fuchs TW (1985) Feces-mediated, host finding behaviour of *Cotesia flavipes*, a parasite of *Diatrea saccharalis* (Lepidoptera: Pyralidae). Annals of the Entomological Society of America 78(5): 647–650. https://doi. org/10.1093/aesa/78.5.647
- Vencl FV, Gómez NE, Ploss K, Boland W (2009) The chlorophyll catabolite, pheophorbide a, confers predation resistance in a larval tortoise beetle shield defense. Journal of Chemical Ecology 35: 281–288. https://doi.org/10.1007/s10886-008-9577-1
- Vencl FV, Morton TC (1998a) The shield defense of the sumac flea beetle, *Blepharida rhois* (Chrysomelidae: Alticinae). Chemoecology 8(1): 25–32. https://doi.org/10.1007/PL00001800
- Vencl FV, Morton TC (1998b) Did chemical change in shield defenses promote diversification of shining leaf beetles (Chrysomelidae: Criocerinae)? Proceedings of the 4<sup>th</sup> Symposium of the Chrysomelidae. Museo Regionale di Scienze Naturali 205–218.
- Vencl FV, Morton TC (1999) Macroevolutionary aspects of larval shield defences. In: Cox ML (Ed.) Advances in Chrysomelidae Biology (Vol. 1). Backhuys Publishers, Leiden, 217–238.
- Vencl FV, Morton TC, Mumma RO, Schultz JC (1999) Shield defense of a larval tortoise beetle. Journal of Chemical Ecology 25(3): 549–566. https://doi. org/10.1023/A:1020905920952
- Vencl FV, Levy A, Geeta R, Keller G, Windsor DM (2004) Observations on the natural history, systematics and phylogeny of the Criocerinae of Costa Rica and Panama. In: Jolivet P, Santiago-Blay JA, Schmitt M (Eds) New Developments in the Biology of Chrysomelidae. SPB Academic Publishing bv, The Hague, 423–454. https://doi. org/10.1163/9789004475335\_038
- Vencl FV, Nogueira-de-Sá F, Allen BJ, Windsor DM, Futuyma DJ (2005) Dietary specialization influences the efficacy of larval tortoise beetle shield defenses. Oecologia 145(3): 404–414. https://doi.org/10.1007/s00442-005-0138-9
- Vencl FV, Srygley RB (2013) Enemy targeting, trade-offs, and the evolutionary assembly of a tortoise beetle defense arsenal. Evolutionary Ecology 27: 237–252. https://doi. org/10.1007/s10682-012-9603-1
- Vencl FV, Trillo PA, Geeta R (2011) Functional interactions among tortoise beetle larval defenses reveal trait suites and escalation. Behavioral Ecology and Sociobiology 65: 227–239. https://doi.org/10.1007/s00265-010-1031-z
- Villanueva-Jimenez JA, Fasulo TR (1996 [present]) Household casebearer, *Phereoeca uterella (dubitatrix)* Walsingham (Insecta: Lepidoptera: Tineidae). Department of Entomology and Nematology, UF/IFAS Extension. https://edis.ifas.ufl.edu/publication/ IN129 [Accessed 18 May 2022]
- von Frisch K (1974) Animal Architecture. Harcourt, Brace, Jovanovich, New York and London, 306 pp.
- Wallace JB (1970) The defensive function of a case on a chrysomelid larva. Journal of the Georgia Entomological Society 5: 19–24.
- Wallace JB (1975) The larval retreat and food of *Arctopsyche*; with phylogenetic notes on feeding and adaptations in hydropsychid larvae (Trichoptera). Annals of the Entomological Society of America 68(1): 167–173. https://doi.org/10.1093/aesa/68.1.167

- Wallace JB, Sherberger FF (1975) The larval dwelling and feeding structure of Macronema transversum (Walker) (Trichoptera: Hydropsychidae). Animal Behaviour 23: 592– 596. https://doi.org/10.1016/0003-3472(75)90135-9
- Wang B, Xia F, Engel MS, Perrichot V, Shi G, Zhang H, Chen J, Jarzembowski EA, Wappler T, Rust J (2016) Debris-carrying camouflage among diverse lineages of Cretaceous insects. Science Advances 2(6): e1501918. https://doi.org/10.1126/sciadv.1501918
- Waterhouse DF (1974) The biological control of dung. Scientific American 230(4): 100– 109. https://doi.org/10.1038/scientificamerican0474-100
- Weirauch C (2006) Anatomy of disguise: Camouflaging structures in nymphs of some Reduviidae (Heteroptera). American Museum Novitates 3542(1): 1–18. https://doi.org/10.1206/0003-0082(2006)3542[1:AODCSI]2.0.C0;2
- Weise J (1893) Naturgeschichte der Insecten Deutschlands. Erste Abtheilung Coleoptera. Sechster Band. Nicolaische Verlags-Buchhandlung R. Stricker, Berlin, [XIV +] 1161 pp. [+ 1 taf.] https://doi.org/10.5962/bhl.title.8270
- Weise J (1896) Feststellung einiger Cassiden-Gattungen. Deutsche Entomologische Zeitschrift 1896: 10–15. https://doi.org/10.1002/mmnd.48018960103
- Weiss M (2006) Defecation behaviour and ecology of insects. Annual Review of Entomology 51(1): 635–661. https://doi.org/10.1146/annurev.ento.49.061802.123212
- Wheeler WM (1928) The Social Insects: Their Origin and Evolution. London, Routledge, 446 pp. https://doi.org/10.4324/9781315624860
- Wilson ED (1971) The Insect Societies. Belknap Press of Harvard University Press, Boston, 562 pp. [ISBN 9780674454903]
- Windsor DM, Riley EG, Stockwell HP (1992) An introduction to the biology and systematics of Panamanian tortoise beetles (Coleoptera: Chrysomelidae: Cassidinae). In: Quintero D, Aiello A (Eds) Insects of Panama and Mesoamerica: Selected Studies. Oxford University Press, Oxford, 372–391.
- Wood GW (1966) Life history and control of a casebearer, *Chlamisus cribripennis* (Coleoptera, Chrysomelidae), on blueberry. Journal of Economic Entomology 59: 823–825. https://doi.org/10.1093/JEE/59.4.823
- Yamamoto I [Director] (2018) Extraordinary! Survival of Rare Insects. Wildlife TV program by KAZE Co.; NHK General (Japan Broadcasting Corporation). [Broadcasted on August 19, 2018.] https://www.nhk-ondemand.jp/#/0/
- Yamamoto I [Director] (2020) Central America Costa Rica Tropical Forest Exploratory Entomologist Approaches! Transcendence Survival Techniques. Wildlife TV program by KAZE Co.; NHK BS Premium (Japan Broadcasting Corporation). [Broadcasted on April 6, 2020.] https://www.nhk-ondemand.jp/#/0/
- Yoshida T, Leschen RAB (2020) Larval descriptions and exuvial retention of Toramini (Coleoptera: Erotylidae: Cryptophilinae). The Coleopterists Bulletin 74(1): 1–14. https://doi.org/10.1649/0010-065X-74.1.1
- Zeledón R, Valerio CE, Valerio JE (1973) The camouflage phenomenon in several species of Triatominae (Hemiptera: Reduviidae). Journal of Medical Entomology 10(2): 209–211. https://doi.org/10.1093/jmedent/10.2.209
- Zhang F, Luo W, Shi Y, Fan Z, Ji G (2012) Should we standardize the 1,700-year-old faecal microbiota transplantation. American Journal of Gastroenterology 107(11): e1755. https://doi.org/10.1038/ajg.2012.251
- Ziegenhorn MA (2017) Sea Urchin Covering Behaviour: A Comparative Review. In: Agnello M (Ed.) Sea Urchin – From Environment to Aquaculture and Biomedicine. InTechOpen Limited, London, 19–33. https://doi.org/10.5772/intechopen.6846

## Supplementary material 1

#### Film 1: Cassida sphaerula (Chrysomelidae, Cassidinae, Cassidinae)

Authors: Caroline Simmrita Chaboo, Sally Adam, Kenji Nishida, Luke Schletzbaum Data type: Video (wmv file)

Explanation note: Larva moving shield over dorsum (1.21 mins; real-time speed; Sally Adam). YouTube link: https://youtu.be/bDygjys6M-0.

Copyright notice: This dataset is made available under the Open Database License (http://opendatacommons.org/licenses/odbl/1.0/). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: https://doi.org/10.3897/zookeys.1177.102600.suppl1

## Supplementary material 2

#### Film 2: Cassida sphaerula (Chrysomelidae, Cassidinae, Cassidinae)

Authors: Caroline Simmrita Chaboo, Sally Adam, Kenji Nishida, Luke Schletzbaum Data type: Video (wmv file)

Explanation note: Telescopic anus of larva excreting wet droplet (3.08 mins; real-time speed; Sally Adam). YouTube link: https://youtu.be/3vNZN60IRM8.

Copyright notice: This dataset is made available under the Open Database License (http://opendatacommons.org/licenses/odbl/1.0/). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: https://doi.org/10.3897/zookeys.1177.102600.suppl2