

Two new records and an updated checklist of freshwater crabs (Arthropoda, Malacostraca, Decapoda, Potamidae and Gecarcinucidae) from Bangladesh

Shibly Sadique Shashi^{1,2}, Da Pan¹, Nusrath Jahan Emu³, Mishal Roy³,
Md. Abu Sadek², SM Sharifuzzaman², Hongying Sun¹

¹ College of Life Sciences, Nanjing Normal University, Nanjing 210023, China

² Institute of Marine Sciences, University of Chittagong, Chattogram 4331, Bangladesh

³ Department of Fisheries, University of Chittagong, Chattogram 4331, Bangladesh

Corresponding author: Hongying Sun (sunhongying@njnu.edu.cn)

Abstract

The species diversity of freshwater crabs of Bangladesh is poorly known. In this study, *Acanthopotamon fungosum* (Alcock, 1909) and *Maydelliathelphusa edentula* (Alcock, 1909) are reported as new records from Bangladesh. The two species were identified through morphology and molecular phylogeny based on 16S rDNA gene sequences. Herein, diagnostic characters of both species are provided respective to close congeners. There is concern over the conservation status of *A. fungosum* due to its narrow distributional range. In addition, an updated checklist and a key to the freshwater crabs of Bangladesh are provided.

Key words: *Acanthopotamon*, biodiversity, conservation, *Maydelliathelphusa*, south Asia



Academic editor: Célio Magalhães

Received: 1 March 2023

Accepted: 24 May 2023

Published: 16 June 2023

ZooBank: <https://zoobank.org/3FC6EC49-97A0-477D-AE1F-839CBB258099>

Citation: Shashi SS, Pan D, Emu NJ, Roy M, Sadek MA, Sharifuzzaman SM, Sun H (2023) Two new records and an updated checklist of freshwater crabs (Arthropoda, Malacostraca, Decapoda, Potamidae and Gecarcinucidae) from Bangladesh. ZooKeys 1167: 211–222. <https://doi.org/10.3897/zookeys.1167.102766>

Copyright: © Shibly Sadique Shashi et al. This is an open access article distributed under terms of the Creative Commons Attribution License ([Attribution 4.0 International – CC BY 4.0](https://creativecommons.org/licenses/by/4.0/)).

Introduction

Freshwater crabs are common in the tropical and subtropical regions where they inhabit a wide range of habitats such as rivers, swamps, lakes, and caves (Yeo et al. 2008). Bangladesh is a riverine country and has various bodies of freshwater suitable for crabs. For example, the hilly areas of the districts of Khagrachari, Rangamati, Bandarban, Chattogram, Cox's Bazar, Mymensingh, Netrokona, Sylhet, Moulvibazar, and Habiganj have watersheds consisting of rivers, lakes, streams, and waterfalls that harbour unique aquatic biodiversity, including crabs. The wetlands at Manikganj, Jessore, Narail and Rajshahi, and the Halda and Meghna rivers are also suitable habitats for crabs (Hasan and Rashid 2016).

More than 1,300 species of freshwater crabs have been found throughout the world (Yeo et al. 2008). Remarkably, only five species, namely *Acanthopotamon martensi* (Wood-Mason, 1875) [recorded as *Potamon martensi*], *Lotholophusa woodmasoni* (Rathbun, 1905) [recorded as *Potamon woodmasoni*], *Lamella lamellifrons* (Alcock, 1909) [recorded as *Paratelphusa lamellifrons*], *Sartoriana spinigera* (Wood-Mason, 1871) [recorded as *Paratelphusa spinigera*], and *Sartoriana trilobata* (Alcock, 1909), have been reported from Bangladesh (Shafi and Quddus 1982; Siddiqui and Zafar 2002; Ahmed et al. 2008; Rahman et al. 2008;

Hossain 2015; Mia et al. 2016; Hasan and Rashid 2016; Chakraborty et al. 2021). There are two additional reports, i.e., *Austrothelphusa transversa* (von Martens, 1868) (Ahmed et al. 2008; Rahman et al. 2008; Hossain 2015; Chakraborty et al. 2021) and *Pyxidognathus fluviatilis* (Alcock, 1900) (Ahmed et al. 2008; Rahman et al. 2008; Hossain 2015; Hasan and Rashid 2016; Chakraborty et al. 2021), but the former is native to Australia and Papua New Guinea (Esser and Cumberlidge 2008) and the latter species inhabits fresh, brackish, and marine waters and is not regarded as true freshwater crab (Yeo et al. 2008). Therefore, the record of *A. transversa* is inaccurate or questionable. Overall, the species diversity of freshwater crabs of Bangladesh and their natural history remains largely unknown. Based on morphological and molecular data, the present study newly reports *Acanthopotamon fungosum* (Alcock, 1909) and *Maydelliathelphusa edentula* (Alcock, 1909) as part of freshwater crab fauna of Bangladesh.

Material and method

During a field survey in August 2021, two specimens of freshwater crab were collected from a small hilly stream near the Chittagong University Campus, Chattogram and from the Kangsa River, Mymensingh (Fig. 1). Crabs were collected by hand and with a fishing net. Both specimens were photographed, diagnosed based on their morphometric characters, preserved in 95% ethanol, and had tissue sampled for molecular study. These specimens were transferred to the laboratory of Institute of Marine Sciences (IMS), University of Chittagong (CU) for further study. The specimens were identified following the taxonomic keys of Pati et al. (2019) and Das (2021). The two crab specimens were deposited in the IMSCU (voucher numbers IMSCU/FW-crab2108.01 and IMSCU/FW-crab2108.02) for future reference. The following abbreviations used are: G1 for the male first gonopods; G2 for the male second gonopods. The terminology of morphological characteristics used follows that of Ng (1988) and Davie et al. (2015).




Figure 1. Map showing the collection points of two crab species from Bangladesh. Station 1: *Acanthopotamon fungosum* collected from Chittagong University Campus, Chattogram. Station 2: *Maydelliathelphusa edentula* collected from the Kangsa River, Mymensingh.

Genomic DNA was extracted from the gill tissue using Trelief Animal Genomic DNA kit (Tsingke, China) according to the manufacturer's protocol and sequenced with an Illumina HiSeq X Ten platform (150 bp paired end). Full-length mitochondrial 16S sequences were assembled using MITOZ (Meng et al. 2019). Sequences were uploaded to NCBI GenBank with accession numbers [OQ788486](#) and [OQ788487](#). Together with downloaded sequences, the matrix was aligned with MAFFT (Kato and Standley 2013) using the default setting. IQ-TREE (Nguyen et al. 2015) was employed to conduct a maximum-likelihood (ML) analysis. Node supports were obtained through 1,000 ultra-fast bootstrap replicates (Minh et al. 2013). The phylogenetic tree was visualized using ITOL (Letunic and Bork 2019).

Taxonomy

Family Potamidae Ortmann, 1896

Genus *Acanthopotamon* Kemp, 1918

Type species. *Paratelphusa martensi* Wood-Mason, 1875, by original designation.

***Acanthopotamon fungosum* (Alcock, 1909)**

Figs 2, 4a, b

Potamon (*Paratelphusa*) *fungosum* Alcock, 1909: 250.

Potamon (*Acanthopotamon*) *fungosum*—Alcock 1910: 65, fig. 12.

Lobothelphusa fungosa—Bott 1970: 148, pl. 38 fig. 25, pl. 46 fig. 23.

Paratelphusa fungosum—Brandis and Sharma 2005: 14.

Acanthopotamon fungosum—Yeo and Ng 2007: 274; Ng et al. 2008: 159.

Material examined. 1 male, 28.84 × 24.28 mm (Table 1), Chittagong University Campus, Chattogram, Bangladesh, 22°28'07"N, 91°46'48"E; 15 August 2021, collected by Shibly Sadique Shashi.

Description of the male. Carapace subhexagonal, convex, covered by short spongy fur, dorsal surface rough, region distinct; ca 1.19× broader than long. H-shaped groove distinct. Epigastric cristae broad, blunt, well advance of postorbital cristae; post orbital cristae short, not confluent with first epibranchial tooth (Fig. 2a); external orbital tooth blunt, broadly triangular; anterolateral margin convex, with 4 epibranchial teeth, first epibranchial tooth distinctly triangular, others sharp. Eyes moderate in size, outer margin of eye with U- to V-shaped incision (Fig. 2c). Third maxilliped elongated, rectangular; with well-developed flagellum present distally on exopod; exopod not distally tapering and longer than merus width. Epistome lateral margins slightly sinuous, medial lobe triangular (Fig. 2c).

Chelipeds unequal in size, right larger; carpus and merus of cheliped with distinct subdistal and subterminal spine, fingers longer than palm, distinct gap with dactyl and pollex closed, both movable and immovable fingers with 3 or 4 large, rounded teeth (Fig. 2d); fingers of minor chela slightly gaping when closed (Fig. 2e). Ambulatory legs bearing short setae; second pair of ambulatory legs longest, fourth pair shortest; dactylus slender, styliiform, with spinules (Fig. 2a).




Figure 2. *Acanthopotamon fungosum* **a** dorsal view **b** ventral view **c** frontal view **d** major chela **e** minor chela. Scale bars: 1 cm.

Table 1. The morphometric features of *Acanthopotamon fungosum* (IMSCU/FW-crab2108.01) and *Maydelliathelphusa edentula* (IMSCU/FW-crab2108.02). Lengths and widths in mm; weight in g.

Morphometric data	<i>Acanthopotamon fungosum</i> (♂)	<i>Maydelliathelphusa edentula</i> (♂)
Carapace width	28.84	65.89
Carapace length	24.28	49.16
Frontal width	9.11	14.84
Pleon width	11.13	24.72
Pleon length	15.89	35.29
Telson length	4.26	10.37
Major chela length	22.23	65.52
Cheliped length	41.65	125.57
Dactyl length	13.81	45.77
Merus length	10.44	25.97
1 st ambulatory leg length	33.31	72.96
2 nd ambulatory leg length	38.99	88.87
3 rd ambulatory leg length	38.01	84.55
4 th ambulatory leg length	32.24	65.64
Weight	5.99	97.00
No. of epibranchial tooth on each side	4	1

Thoracic sternum smooth, pitted, suture between s1/s2 completely fused to form triangular structure; suture between s3/s4 indistinct, suture between s4/s5, s5/s6, s6/s7, s7/s8 distinct (Fig. 2b).

Pleon broadly triangular; all segments rectangular. Telson tongue-shaped, length and width almost equal (Fig. 2b).

G1 curved outwardly, gradually tapering towards tip; terminal segment sub-cylindrical, slender, covered by short setae; nearly 3× shorter than subterminal segment (Fig. 4a, b). G2 elongated, shorter than G1.

Remarks. *Acanthopotamon fungosum* was originally described as *Potamon* (*Paratelphusa*) *fungosum* Alcock, 1909 from Cachar, India (Bott 1970). Previously, *A. fungosum* was only known from the states of Arunachal Pradesh, Assam, Mizoram and Manipur, in India (Pati et al. 2019; Mitra and Pati 2021; Rath et al. 2022). This species is recorded here for the first time from Bangladesh. *Acanthopotamon martensi* is also distributed in Bangladesh, i.e., in Manikganj district and estuaries of the Chakaria Sundarban area (Shafi and Quddus 1982; Rahman et al. 2008).

The IUCN conservation status of *A. fungosum* was assessed as Data Deficient (DD) (Cumberlidge 2008a). The species is distributed over a small geographical area, i.e., eastern Bangladesh and southern Assam, India. Due to restricted distributional range and increasing threats to freshwater habitats of this region from various human activities, *A. fungosum* is likely more threatened than *M. edentula*. Further field surveys are needed to determine population size and threats.

Until now, four species have been described for the genus *Acanthopotamon* (Pati et al. 2019). *Acanthopotamon fungosum* can be easily differentiated by having four epibranchial teeth, compared to two epibranchial teeth in *A. panningi* (Bott 1970: fig. 19; Pati et al. 2019), three in *A. horai* (Pati et al. 2019), and three in *A. martensi* (Bott 1970: fig. 20; Rahman et al. 2008; Pati et al. 2019).

Family Gecarcinucidae Rathbun, 1904

Genus *Maydelliathelphusa* Bott, 1969

Type species. *Telphusa masoniana* Henderson, 1893, by original designation.

Maydelliathelphusa edentula Alcock, 1909

Figs 3, 4c, d

Potamon lugubre edentulum Alcock, 1909: 247.

Paratelphusa (*Barytelphusa*) *edentula*—Alcock 1909: 376; Alcock 1910: 84, fig. 19.

Barytelphusa (*Maydelliathelphusa*) *edentula*—Bott 1970: 34.

Maydelliathelphusa edentula—Ng et al. 2008: 68.

Material examined. 1 male, 65.89 × 49.16 mm (Table 1), Kangsa River, Netrokona, Mymensingh, Bangladesh, 25°00'45"N, 90°38'54"E, 10 August 2021, collected by Shibly Sadique Shashi and Nusrath Jahan Emu.

Description of the male. Carapace slightly depressed, ca 1.34× broader than long; epigastric cristae distinct, epigastric and postorbital cristae on either side united (Fig. 3a); postorbital cristae distinct, sharp, subparallel to frontal margin; frontal region deflexed, relatively wide; external orbital tooth prominent, epibranchial tooth present, prominent; frontal margin bilobed, frontal median triangle not complete; cervical groove well developed; mesogastric furrow very distinct, deep, slightly bifurcated posteriorly. Anterolateral and posterolateral regions rugose. Eyes smaller than orbital floor; eyestalk short. Third maxilliped




Figure 3. *Maydelliathelphusa edentula* **a** dorsal view **b** ventral view **c** frontal view **d** major chela and **e** minor chela. Scale bars: 2 cm.




Figure 4. G1 **a, b** *Acanthopotamon fungosum* **c, d** *Maydelliathelphusa edentula*. Scale bars: 2 mm.

with ischium subrectangular, longer than broad, with distinct narrow medial groove; merus pentagonal, broader than long; exopod slender, longer than ischium, reaching base of merus, with long flagellum (Fig. 3c). Epistome lateral margins slightly sinuous, medial lobe triangular (Fig. 3c).

Chelipeds surface smooth, unequal, right cheliped larger; carpus with distinct spine on inner angle; fingers longer than palm, movable finger strongly curved downward, immovable finger smoothly curved upward, wide gap between dactyl and pollex when closed, movable finger comparatively larger than immovable finger, inner margin of fingers lined with numerous round and blunt teeth (Fig. 3d); ambulatory legs stout; second pair of ambulatories longest while the fourth pair shortest (Fig. 3b); dactylus slender, longer than propodus, with 4 rows of spines on the margin (Fig. 3b).

Male thoracic sternum smooth, pitted. Sternites s1/s2 completely fused forming triangular structure; suture between s3/s4 shallow; suture between s4/s5, s5/s6, s6/s7, s7/s8 distinct (Fig. 3b).

Male pleon T-shaped, somites 5 and 6 constricted medially. Telson tongue-shaped, length and width almost equal (Fig. 3b).

G1 stout, straight; terminal segment tapering gradually, almost 2× shorter than subterminal segment (Fig. 4c, d). G2 elongated, shorter than G1.

Remarks. *Maydelliathelphusa edentula* was originally described as *Potamon lugubre* var. *edentulum* Alcock 1909 from Assam, India and subsequently transferred to *Barytelphusa* (*Maydelliathelphusa*) Bott, 1969 (Bott 1970). The present record is the first report of the genus *Maydelliathelphusa* from Bangladesh. Previously, *M. edentula* was documented in India (Assam, Nagaland, Mizoram) and Bhutan (Samchi) (Cumberlidge 2008b; Valarmathi 2017; Ray et al. 2018; Das 2021). Suitable habitats include freshwater rivers and streams (Cumberlidge 2008b). Therefore, it is not surprising that *M. edentula* occurs in eastern Bangladesh, near its known distribution range.

The IUCN conservation status of *M. edentula* is Near Threatened (NT) because of its limited distribution range and vulnerable habitat (Cumberlidge 2008b). Although the current study has expanded its known geographic distribution, the conservation status of this species is still not optimistic. In western Bangladesh, some local people, especially fishermen, eat this crab on a limited scale. In addition, various types of human activities, like pollution, urbanization and sand mining are impacting freshwater habitats in that area and, consequently, are threats to the population of *M. edentula*.

Until now, five species belonging to the genus *Maydelliathelphusa* have been recorded from India, Bhutan, and Nepal (Cumberlidge 2008c; Valarmathi 2017; Das 2021). The morphology of the carapace of *M. edentula* is superficially similar to that of other four species. However, *M. edentula* can be easily distinguished externally by its large, distinctively asymmetric chelipeds and its united epigastric and postorbital cristae (Bott 1970; Das 2021).

Check list

Herein, two freshwater crab species, *A. fungosum* and *M. edentula*, are documented for the first time from Bangladesh. A molecular phylogeny based on 16S rDNA sequences confirmed their identification (Fig. 5). Specifically, *A. fungosum* and *M. edentula* are clustered separately with *A. panningi* and *M. lugubris*. With these two species included, there are now seven species of true freshwater crab known to inhabit Bangladesh, namely *Acanthopotamon fungosum*, *A. martensi*, *Lobothelphusa woodmasoni*, *Lamella lamellifrons*, *Maydelliathelphusa edentula*, *Sartoriana spinigera*, and *S. trilobata* (Table 2).




Figure 5. Reconstructed maximum-likelihood tree based on 16S rDNA sequences. Newly obtained sequences are shown in red. Numbers on branches represent bootstrap values.

Table 2. Checklist of the freshwater crabs from Bangladesh, with the inclusion of species identified in this study.

Family/genus/species	Locality	IUCN Status	References
Family Gecarcinucidae Rathbun, 1904			
Genus <i>Lamella</i> Bahir & Yeo, 2007			
1) <i>Lamella lamellifrons</i> (Alcock, 1909) ^a	5, 7, 8, 10, 11, 13	—	D–F
Genus <i>Maydellia</i> thelphusa Bott, 1969			
2) <i>Maydellia thelphusa edentula</i> (Alcock, 1909)*	5	—	This study
Genus <i>Sartoriana</i> Bott, 1969			
3) <i>Sartoriana spinigera</i> (Wood-Mason, 1871) ^b	1, 4, 7, 8, 9, 10, 12	Least Concern	A–C, E, G, H
4) <i>Sartoriana trilobata</i> (Alcock, 1909)	1, 2, 3	—	A
Family Potamidae Ortmann, 1896			
Genus <i>Acanthopotamon</i> Kemp, 1918			
5) <i>Acanthopotamon fungosum</i> (Alcock, 1909)*	6	—	This study
6) <i>Acanthopotamon martensi</i> (Wood-Mason, 1875) ^c	4, 8, 11, 12	Least Concern	B–E, G, H
Genus <i>Lobothelphusa</i> Bouvier, 1917			
7) <i>Lobothelphusa woodmasoni</i> (Rathbun, 1905) ^d	1, 2, 3, 4, 11, 12	Least Concern	A–E, G, H

*Indicates a new record.

Originally reported as:

^a*Paratylphusa lamellifrons* (see Shafi and Quddus 1982; Siddiqui and Zafar 2002; Mia et al. 2016);

^b*Paratylphusa spinigera* (see Shafi and Quddus 1982);

^c*Potamon martensi* (see Shafi and Quddus 1982; Siddiqui and Zafar 2002);

^d*Potamon woodmasoni* (see Shafi and Quddus 1982; Siddiqui and Zafar 2002).

Locality: 1: Cox's Bazar; 2: Bandarban; 3: Maulovi Bazar; 4: Manikganj; 5: Mymensingh; 6: Chittagong University Campus, Chattogram; 7: Rajshahi; 8: Jessore; 9: Narail; 10: Sylhet; 11: Chakaria Sundarban; 12: Mogra River, Netrokona; 13: Dhaka.

References: A: Hasan and Rashid (2016); B: Rahman et al. (2008); C: Hossain (2015); D: Siddiqui and Zafar (2002); E: Shafi and Quddus (1982); F: Mia et al. (2016); G: Ahmed et al. (2008); H: Chakraborty et al. (2021).

Key to freshwater crabs from Bangladesh

- 1 Mandibular palp with single terminal lobe; male pleon T-shaped2
- Mandibular palp with bilobed terminal segment; male pleon triangular.....5

- 2 Epibranchial tooth situated well posterior to level of postorbital cristae, at widest part of carapace; G1 subterminal segment relatively broad**3**
- Epibranchial tooth situated in line with or slightly posterior to level of postorbital cristae, anterior to widest part of carapace; G1 subterminal segment relatively narrow.....**4**
- 3 Anterolateral margin without protrusion between external orbital angle and epibranchial tooth; cervical groove relatively narrow; G1 terminal segment straight***Sartoriana spinigera***
- Anterolateral margin with one lobe-like protrusion between external orbital angle and epibranchial tooth; cervical groove relatively wide; G1 terminal segment bend outward ***Sartoriana trilobata***
- 4 Epigastric cristae sharp; outer margin of external orbital angle comparatively long; G1 with inner margin characteristically curved or angled just below juncture between terminal and subterminal segment
.....***Lamella lamellifrons***
- Epigastric cristae rugose; outer margin of external orbital angle comparatively short; G1 straight***Maydelliathelphusa edentula***
- 5 Carapace dorsal surface uneven, setose; male sternopleonal cavity reaches up to the level of the imaginary line joining the anterior part of the cheliped coxae.....**6**
- Carapace dorsal surface smooth, glabrous; male sternopleonal cavity reaches up to the level of the imaginary line joining the median part of the cheliped coxae.....***Lobothelphusa woodmasoni***
- 6 Carapace with four epibranchial teeth on each anterolateral margin.....
.....***Acanthopotamon fungosum***
- Carapace with three epibranchial teeth on each anterolateral margin
.....***Acanthopotamon martensi***

References: Bahir and Yeo (2007); Yeo and Ng (2012); Pati et al. (2019); Chetia et al. (2021).

Acknowledgements

The authors express their sincere gratitude to the local people and fishermen for their support. Special thanks to the staffs of Institute of Marine Sciences, University of Chittagong for their sincere support. We thank the editor and reviewer for their careful reading of our manuscript and thoughtful comments. Financial support from the National Natural Science Foundation of China (no. 31772427 to SHY and no. 32200356 to PD) and the Natural Science Foundation of the Jiangsu Higher Education Institutions (no. 22KJB180018 to PD) are acknowledged.

Additional information

Conflict of interest

No conflict of interest was declared.

Ethical statement

No ethical statement was reported.

Funding

The National Natural Science Foundation of China (No. 31772427 to SHY and No. 32200356 to PD), the Natural Science Foundation of the Jiangsu Higher Education Institutions (No. 22KJB180018 to PD).

Author contributions

Conceptualization: DP, HS. Data curation: SSS, MAS, MR, SS, NJE. Funding acquisition: HS, DP. Investigation: SSS, SS, NJE, MR. Methodology: MAS, DP. Supervision: SS, HS. Writing – original draft: SSS. Writing – review and editing: HS, DP, SS, MAS.

Author ORCIDs

Da Pan  <https://orcid.org/0000-0001-5445-6423>

Mishal Roy  <https://orcid.org/0000-0001-7560-4392>

SM Sharifuzzaman  <https://orcid.org/0000-0003-1280-7947>

Hongying Sun  <https://orcid.org/0000-0003-2311-1814>

Data availability

All of the data that support the findings of this study are available in the main text or Supplementary Information.

References

- Ahmed ATA, Kabir SMH, Ahmed M, Rahman AKA, Haque EU, Begum ZNT, Hasan MA, Khondker M [Eds] (2008) Encyclopedia of Flora and Fauna of Bangladesh (Vol. 18): Arthropoda: Crustacea. Asiatic Society of Bangladesh, Dhaka, 480 pp.
- Alcock A (1900) Material for a carcinological fauna of India. No. 6: The Brachyura Catometopa, or Grapsoidea. Journal of the Asiatic Society of Bengal 69(2): 279–456. <https://doi.org/10.5962/bhl.title.15344>
- Alcock A (1909) Diagnoses of new species and varieties of freshwater crabs. Nos 1–4. Records of the Indian Museum 3: 243–252. [375–381.] <https://doi.org/10.26515/rzsi/v3/i3/1909/163275>
- Alcock A (1910) Catalogue of the Indian Decapoda Crustacea in the Collection of the Indian Museum. Part I. Brachyura. Fasciculus II. The Indian Freshwater Crabs – Potamonidae. Trustees of the Indian Museum, Calcutta, 134 pp.
- Bahir MM, Yeo DCJ (2007) The gecarcinucid freshwater crabs of southern India (Crustacea: Decapoda: Brachyura). The Raffles Bulletin of Zoology 16: 309–354.
- Bott R (1969) Flußkrabben aus Asien und ihre Klassifikation (Crustacea, Decapoda). Senckenbergiana Biologica 50: 359–366.
- Bott R (1970) Die Süßwasserkrabben von Europa, Asien, Australien und ihre Stammgeschichte. Abhandlungen der Senckenbergischen Naturforschenden Gesellschaft 526: 1–338.
- Bouvier EL (1917) Sur la classification des Eupotamonea, crabes d'eau douce de la famille des potamonidés. Comptes Rendus de l'Académie de Sciences 165: 613–621. <https://doi.org/10.5962/bhl.part.1215>
- Brandis D, Sharma S (2005) Taxonomic revision of the freshwater crab fauna of Nepal with description of a new species (Crustacea, Decapoda, Brachyura, Potamoidae and Gecarcinucidae). Senckenbergiana Biologica 82: 1–30.

- Chakraborty BK, Verma AK, Muniya S (2021) Present status of aquatic resource and its catch of Mogra River in Bangladesh. *Sustainable Marine Structures* 3: 26–38. <https://doi.org/10.36956/sms.v3i2.436>
- Chetia A, Mitra S, Das DN (2021) Redescription of *Sartoriana trilobata* (Alcock, 1909) (Crustacea: Decapoda: Brachyura: Gecarcinucidae) from Assam, Northeast India, with notes on the morphology of male gonopods. *Zootaxa* 5026(1): 136–144. <https://doi.org/10.11646/zootaxa.5026.1.6>
- Cumberlidge N (2008a) *Acanthopotamon fungosum*. The IUCN Red List of Threatened Species 2008: e.T134781A4012582.
- Cumberlidge N (2008b) *Maydelliathelphusa edentula*. The IUCN Red List of Threatened Species 2008: e.T134064A3905366.
- Cumberlidge N (2008c) *Maydelliathelphusa lugubris*. The IUCN Red List of Threatened Species 2008: e.T134282A3931648.
- Das RK (2021) First record of a freshwater crab, *Maydelliathelphusa masoniana* (Henderson, 1893) (Decapoda: Brachyura: Gecarcinucidae) from West Bengal, India. *Journal of Threatened Taxa* 13(13): 20084–20089. <https://doi.org/10.11609/jott.7465.13.13.20084-20089>
- Davie PJF, Guinot D, Ng PKL (2015) Anatomy and functional morphology of Brachyura. In: Castro P, Davie PJF, Guinot D, Schram FR, von Vaupel Klein JC (Eds) *Treatise on Zoology—Anatomy, Taxonomy, Biology. The Crustacea*, 9(C-I), Decapoda: Brachyura (Part 1). Brill, Leiden, 11–163. https://doi.org/10.1163/9789004190832_004
- Esser L, Cumberlidge N (2008) *Austrothelphusa transversa*. The IUCN Red List of Threatened Species 2008: e.T134922A4036588.
- Hasan R, Rashid HO (2016) A study and availability assessment of freshwater crabs in the hill streams of Bangladesh. *International Journal of Aquaculture and Fishery Sciences* 2: 018–022. <https://doi.org/10.17352/2455-8400.000014>
- Henderson JR (1893) A contribution to Indian carcinology. *Transactions of the Linnean Society of London* 5(10): 325–458. <https://doi.org/10.1111/j.1096-3642.1893.tb00653.x>
- Hossain MAR (2015) Red List of Bangladesh (Vol. 6): Crustaceans. IUCN, International Union for Conservation of Nature, Bangladesh Country Office, Dhaka, 256 pp.
- Katoh K, Standley DM (2013) MAFFT multiple sequence alignment software version 7: Improvements in performance and usability. *Molecular Biology and Evolution* 30(4): 772–780. <https://doi.org/10.1093/molbev/mst010>
- Kemp S (1918) Crustacea Decapoda of the Inle Lake Basin. *Records of the Indian Museum* 14: 81–102. <https://doi.org/10.5962/bhl.part.18606>
- Letunic I, Bork P (2019) Interactive Tree Of Life (iTOL) v4: Recent updates and new developments. *Nucleic Acids Research* 47(W1): W256–W259. <https://doi.org/10.1093/nar/gkz239>
- Meng G, Li Y, Yang C, Liu S (2019) MitoZ: A toolkit for animal mitochondrial genome assembly, annotation and visualization. *Nucleic Acids Research* 47(11): e63–e63. <https://doi.org/10.1093/nar/gkz173>
- Mia MB, Islam MB, Sarker MM, Rahman MR, Jalil MA, Rahim MA, Roy DK (2016) Morphology and biochemical composition of crab (*Paratethusa lamellifrons*) in Bangladesh. *Academic Research Journal of Biological Sciences and Medicinal Plants* 1: 1–13.
- Minh BQ, Nguyen MAT, Von Haeseler A (2013) Ultrafast approximation for phylogenetic bootstrap. *Molecular Biology and Evolution* 30(5): 1188–1195. <https://doi.org/10.1093/molbev/mst024>

- Mitra S, Pati SK (2021) A new species of freshwater crab, *Teretamon kapota* sp. nov. (Decapoda: Brachyura: Potamidae) and a new record from Arunachal Pradesh, North-Eastern India. Records of the Zoological Survey of India 121: 1–10.
- Ng PKL (1988) The Freshwater Crabs of Peninsular Malaysia and Singapore. Shinglee Press, Singapore, 156 pp.
- Ng PKL, Guinot D, Davie PJF (2008) Systema Brachyurorum: Part I. An annotated checklist of extant brachyuran crabs of the world. The Raffles Bulletin of Zoology 17: 1–286.
- Nguyen LT, Schmidt HA, Von Haeseler A, Minh BQ (2015) IQ-TREE: A fast and effective stochastic algorithm for estimating maximum-likelihood phylogenies. Molecular Biology and Evolution 32(1): 268–274. <https://doi.org/10.1093/molbev/msu300>
- Ortmann AE (1896) Das System der Decapoden-Krebse. Zoologische Jahrbücher. Abtheilung für Systematik. Geographie und Biologie der Thiere 9: 409–453.
- Pati SK, Mitra S, Yeo DC (2019) A new species of *Acanthopotamon* Kemp, 1918 (Decapoda: Brachyura: Potamidae: Potaminae) from northeastern India, with a key to the species of the genus and notes on their distribution in relation to freshwater ecoregions. Journal of Crustacean Biology 39(4): 450–458. <https://doi.org/10.1093/jcbiol/ruz040>
- Rahman MA, Rahman MM, Ahmed ATA, Mollah AR, Hossain MA (2008) A survey on the diversity of freshwater crabs in some wetland ecosystems of Bangladesh. International Journal of Sustainable Crop Production 3: 10–17.
- Rath S, Bhaumik R, Mitra S, Kosygin L (2022) First report of *Acanthopotamon fungosum* (Alcock, 1909) (Decapoda: Potamidae) from Zeilad Wild Life Sanctuary, Manipur. Biological Forum : An International Journal 14: 807–810.
- Rathbun MJ (1905) Les crabes d’eau douce (Potamonidae) (seconde partie). Nouvelles Archives du Muséum National d’Histoire Naturelle 4(7): 159–322.
- Ray D, Mitra S, Shaw S, Roy D, Bhandari SK, Jana T (2018) Freshwater crab of North East India, its importance and conservation. International Journal of Engineering. Science and Mathematics 4: 96–103.
- Shafi M, Quddus MMA (1982) Bangladesher Matsho Sampad (Fisheries Resources of Bangladesh). Bangla Academy, Dhaka, 444 pp.
- Siddiqui MZH, Zafar M (2002) Crabs in the Chakaria Sundarban area of Bangladesh. Journal of Noami 19: 61–77.
- Valarmathi K (2017) Crustacea: Decapoda. In: Chandra K, Gopi KC, Rao DV, Valarmathi K, Alfred JRB (Eds) Current Status of Freshwater Faunal Diversity in India. Zoological Survey of India, Kolkata, India, 313–331.
- Von Martens E (1868) Über einige neue Crustaceen. Monatsberichte der Königlischen Preussischen Akademie der Wissenschaften zu Berlin 868: 608–615.
- Wood-Mason J (1871) Contributions to Indian carcinology. Journal of the Asiatic Society of Bengal 40: 189–200. [201–207, 449–454.]
- Wood-Mason J (1875) On new or little-known crustaceans. Proceedings of the Asiatic Society of Bengal 1875: 230–232.
- Yeo DCJ, Ng PKL (2007) On the genus “*Potamon*” and allies in Indochina (Crustacea: Decapoda: Brachyura: Potamidae). The Raffles Bulletin of Zoology 16: 273–308.
- Yeo DCJ, Ng PKL (2012) *Sodhiana*, a new genus of freshwater crab from south Asia (Decapoda: Brachyura: Gecarcinucidae). The Raffles Bulletin of Zoology 25: 279–284.
- Yeo DCJ, Ng PKL, Cumberlidge N, Magalhães C, Daniel SR, Campos MR (2008) Global diversity of crabs (Crustacea: Decapoda: Brachyura) in freshwater. Hydrobiologia 595(1): 275–286. <https://doi.org/10.1007/s10750-007-9023-3>