

Peruvian nudibranchs (Mollusca, Gastropoda, Heterobranchia): an updated literature review-based list of species

Alessandra Grández¹ , André Ampuero¹ , Sergio P. Barahona¹ 

1 Carrera de Biología Marina, Universidad Científica del Sur, Lima, Perú

Corresponding author: Sergio P. Barahona (sbarahona@cientifica.edu.pe)

Abstract

Nudibranchs, as a group, have received limited attention in terms of scientific study along the coastline of Peru. Here, an updated and comprehensive list of nudibranch species found in the Peruvian sea is presented, compiled through an extensive review of relevant literature. This compilation encompasses a total of 31 species, classified into two suborders, 10 superfamilies, 20 families, and 28 genera. With respect to the biogeographic provinces along the Peruvian coast, 23 species inhabit the Warm Temperate Southeastern Pacific province, 18 species occur in the Tropical Eastern Pacific province, and 10 species are found in both provinces, crossing the transitional zone between them. In terms of distribution patterns, two species exhibit a cosmopolitan distribution (*Glaucus atlanticus* and *Fiona pinnata*), while two species display a circumtropical distribution (*Cephalopyge trematooides* and *Phylliroe bucephala*). One species exhibits a bipolar distribution in the Eastern Pacific and possesses an amphi-South American distribution (*Rostanga pulchra*). Additionally, six species exhibit an amphi-South American distribution (*Rostanga pulchra*, *Diadumene punctuolata*, *Doto uva*, *Tyrrinna evelinae*, *Tyrrinna delicata*, and *Doris fontainii*), and two species are endemic to Peru (*Corambe mancorensis* and *Felimare sechurana*). This study provides comprehensive information on biogeographical aspects, geographical distributions, and taxonomic updates within the nudibranch species documented in Peru. Furthermore, we discuss the status of species listed in previous literature that have not been confirmed by collections, referring to them as potentially occurring species.



Academic editor: Nathalie Yonow

Received: 8 March 2023

Accepted: 29 June 2023

Published: 23 August 2023

ZooBank: <https://zoobank.org/>

DE7EC71C-BDDE-4A9A-B958-

F1B99633D11D

Citation: Grández A, Ampuero A, Barahona SP (2023) Peruvian nudibranchs (Mollusca, Gastropoda, Heterobranchia): an updated literature review-based list of species. ZooKeys 1176: 117–163. <https://doi.org/10.3897/zookeys.1176.103167>

Copyright: © Alessandra Grández et al.
This is an open access article distributed under terms of the Creative Commons Attribution License ([Attribution 4.0 International – CC BY 4.0](https://creativecommons.org/licenses/by/4.0/)).

Key words: biogeography, bibliographic compilation, geographic distribution, Nudibranchia, Peru, sea slug, taxonomy

Introduction

Nudibranchia Cuvier, 1817 (Subclass Heterobranchia, Infraclass Euthyneura, Superorder Nudipleura) represents an order of exclusively marine gastropod mollusks, distinguished by the absence of shells in the adult stage (Behrens et al. 2005; Wägele and Klussmann-Kolb 2005). These remarkable organisms exhibit striking aposematic body colorations, making them frequent targets of underwater photography. Approximately 3000 species of nudibranchs have been described worldwide inhabiting both cold and tropical regions (Shields 2009; Almada et al. 2016), primarily in shallow waters ranging from 0 to 30 m in depth (Wägele and Klussmann-Kolb 2005). However, novel species discov-

eries have expanded our knowledge of nudibranchs in deeper habitats (Valdés 2001a, 2001b; Gosliner et al. 2008). The ecological importance of nudibranchs stems from their role in controlling populations of cnidarians and poriferans, from which they acquire and incorporate toxins for their own defense (Greenwood 2009; Faulkner and Ghiselin 1983). Furthermore, recent research has elucidated symbiotic associations between nudibranchs and bacteria, wherein the bacteria provide essential nutrients to the hosts (Zhukova et al. 2022). Certain species have been found to synthesize secondary metabolites with high biotechnological potential (Pereira et al. 2012; Dean and Prinsep 2017). Additionally, nudibranch species possess the potential to serve as environmental indicators, as they are sensitive to ocean stressors such as coastal pollution (Caballer et al. 2008).

The most recent inventory of aquatic mollusks in Peru, as documented by Ramírez et al. (2003), recorded a total of 1018 marine species, primarily comprising gastropods and bivalves, with only a limited number of nudibranch species reported. In fact, the Peruvian sea is recognized as one of the impoverished regions worldwide in terms of nudibranch diversity (Schrödl 1997, 2002, 2003; Schrödl and Hooker 2014). In comparison, other regions such as the Caribbean Sea, the Tropical Eastern Pacific, the Indian Ocean, the Mediterranean Sea (Sachidhanandam et al. 2000; Valdés et al. 2006; Chavanich et al. 2013; Ah Shee Tee et al. 2019; Furfaro et al. 2020; Londoño-Cruz 2021), as well as other South American countries including Brazil, Chile, and Venezuela (Fischer and Cervera 2005b; Ardila et al. 2007; Aldea et al. 2011; Padula et al. 2011; Alvim and Pimenta 2013; Gutiérrez et al. 2015; Araya and Valdés 2016; Londoño-Cruz 2021) exhibit considerably higher richness of nudibranch species.

The earliest records of nudibranchs in Peru can be attributed to d'Orbigny (1835–1846) and later to Dall (1909). Following a significant period without the discovery of new species, Millen et al. (1994) reported the presence of *Okenia luna* in Peruvian and Chilean waters. Subsequently, the first list of Peruvian aquatic mollusks was published, which included some nudibranch species (Álamo and Valdivieso 1997), and two years later, another list was published (Paredes et al. 1999), in which Sandra Millen was acknowledged for the preliminary list of species belonging to the infraclass Opisthobranchia (a taxonomic category that has since been abandoned and deprecated; see Jörger et al. 2010; Schrödl et al. 2011; Wägele et al. 2014). In 2003, an updated inventory of Peruvian aquatic mollusks was published (Ramírez et al. 2003), which included the nudibranch species reported up to that time. Several years later, four new species were reported on the northern coast of Peru, *Felimida baumanni*, *Doriopsilla janaina*, *Kynaria cynara*, and *Cuthona* sp. (Nakamura 2006), with the species *Corambe mancorensis* identified as endemic (Martynov et al. 2011), and the species *Spurilla neapolitana* (later corrected as *Spurilla brasiliiana*) (Uribe and Pacheco 2012). Subsequently, four additional nudibranch species were documented for the Peruvian coast (Uribe et al. 2013), and another study focused on the species shared with Chile (Schrödl and Hooker 2014). *Felimare sechurana* was identified as an endemic species in the transition zone of the northern coast (Hoover et al. 2017) and, additionally, two new species of planktonic nudibranchs were described (Quesquen 2017).

We must emphasize that certain species have been listed in previous publications as occurring in Peru without sufficient evidence, such as assumptions of geographic continuity (e.g., *Cadlina sparsa*; Álamo and Valdivieso 1997), reliance on personal communications only (e.g., *Polycera cf. alabe*; Paredes et al. 1999; Uribe et al. 2013), and misinterpretations (e.g., *Gargamella immaculata* and *Thecacera darwini*; Nakamura 2006), which has created problems as these listings have persisted in the literature. To distinguish them from the confirmed species, the term “potentially occurring” is used hereafter. However, such statuses can be revised in the future, as exemplified by the species *Rostanga pulchra*, which was initially predicted for Peruvian waters for many years until its confirmation (Schrödl and Hooker 2014).

El Niño-Southern Oscillation (ENSO) warm events have been observed to induce southward displacement of tropical species (Velez and Zeballos 1985; Paredes et al. 1998) while cold events tend to enhance the intensity of the Humboldt Current, resulting in the northward transport of larvae. Specifically, the northward transport of larvae by the Humboldt Current or the southward transport facilitated by warm ENSO events may introduce Magellanic or tropical species, respectively, into Peruvian waters, thereby influencing distribution ranges. In addition, the susceptibility of nudibranchs to temperature fluctuations, particularly during their larval stages (Leatherman 2019) due to the aragonite-based internal structure found in several species (Ehrlich 2010), their small body size, limited populations (Nybakken 1978), and the sensitivity of the Humboldt Current Ecosystem to oceanic stressors (Echevin et al. 2012) such as warming and acidification (Barnosky et al. 2011; Ceballos et al. 2015; Pievani 2014), collectively suggest that the diversity and distribution of nudibranchs could be impacted (Nimbs and Smith 2018).

The available information on Peruvian nudibranchs remains limited primarily due to a lack of research effort (Uribe and Pacheco 2012). Explorations specifically targeting nudibranchs have been extremely scarce, and most sightings and reports are sporadic (Nakamura 2006; Schrödl and Hooker 2014; Uribe et al. 2013). Given that the species richness of nudibranchs is likely underestimated in Peru (Hooker pers. comm.), this taxonomic order warrants further attention. Our aim was to update and revise the list of nudibranchs in the Peruvian sea, based on a comprehensive review of the scientific literature.

Materials and methods

A comprehensive review was conducted to compile all available literature pertaining to the order Nudibranchia in Peru. The literature search encompassed diverse sources of information, including peer-reviewed journal articles, books, book chapters, “grey literature” (such as scientific reports and theses), and the Sea Slug Forum (<http://www.seaslugforum.net/>). Key terms such as ‘Opisthobranchia,’ ‘Heterobranchia,’ ‘Nudibranch,’ ‘Nudibranchia,’ ‘sea slug,’ ‘phylogeny,’ ‘checklist,’ ‘Peru,’ ‘Humboldt,’ and ‘taxonomy’ were employed. Pertinent data, such as type material, geographic distribution, sampling/reporting sites, bathymetric distribution, and biogeographical provinces, were meticulously included. The most up-to-date scientific names were validated through the World Register of Marine Species (WoRMS, <https://www.marinespecies.org/>), and

reports (occurrences) were cross-referenced using the Global Biodiversity Information Facility (GBIF, <https://www.gbif.org/>) and the iNaturalist database (<https://www.inaturalist.org/>). Any modifications, revalidations, or refutations pertaining to taxonomy are concisely presented as “Remarks”, accompanied by justifications as needed. Endemic species of Peru are also duly indicated. The distribution map was made using QGIS 3.22.8 software (QGIS Development Team 2022), while VENNY 2.1 online software (Oliveros 2016) was employed to visualize the number of species shared with some neighboring countries. Potentially occurring species were clearly distinguished from the confirmed ones.

The acronyms corresponding to the collections where the type material for certain species is deposited have been included, as follows:

CASIZ	California Academy of Sciences Invertebrate Zoology, San Francisco
CZA	Colección de Zoología Acuática, Universidad Peruana Cayetano Heredia, Lima
NHMUK	Natural History Museum, London
MHNURP	Museo Historia Natural Vera Alleman Haeghebaert, Universidad Ricardo Palma, Santiago de Surco
RMNH	Naturalis Biodiversity Center, Leiden
SMNH	Swedish Museum of Natural History, Stockholm
USNM	Smithsonian National Museum of Natural History, Washington DC
ZMB	The Berlin Zoological Museum, Berlin
ZSM	The Bavarian State Collection of Zoology, Munich

Results

Overview

A total of 31 species, encompassing two suborders, ten superfamilies, 20 families, and 28 genera (Table 1), has been confirmed within Peruvian waters. The suborder Cladobranchia comprises 16 species, spanning five superfamilies and 13 families. The suborder Doridina consists of 15 species, distributed among five superfamilies and seven families (Table 1). Notably, the families Chromodorididae and Discodorididae, both belonging to the suborder Doridina, exhibit remarkable species richness with five and four species, respectively (Fig. 1). Additionally, potentially occurring species ($n = 9$) are distributed across five superfamilies, eight families, and nine genera (Table 2). A chronological overview of articles documenting nudibranch species in the Peruvian sea can be found in Table 3, revealing the progressive increase in reported species and the latest updates in scientific nomenclature. Regarding the distribution of species along the Peruvian coast, according to the coastal marine biogeographical classification proposed by Spalding et al. (2007), 23 species inhabit within the Warm Temperate Southeastern Pacific province, 18 species are found within the Tropical Eastern Pacific province, and ten species are common to both provinces (Table 4, Fig. 2).

Peru's inventory of nudibranch species is comparatively modest in comparison to other South American countries, such as Chile, Colombia, and Brazil (Fig. 3A). Concerning species shared with these countries, of the 31 species that have been conclusively documented in Peruvian waters, 19 species are found

Table 1. Nudibranch species confirmed for Peruvian waters according to the bibliographic compilation of this study.

Suborders (n = 2)	Superfamilies (n = 10)	Families (n = 20)	Species (n = 31)
Cladobranchia	Aeolidioidea	Aeolidiidae	<i>Spurilla brasiliiana</i> MacFarland, 1909
		Facelinidae	<i>Phidiana lottini</i> (Lesson, 1831)
		Glaucidae	<i>Bajaeolis bertschi</i> Gosliner & Behrens, 1986
	Arminoidea	Arminidae	<i>Armina californica</i> (J.G. Cooper, 1863)
		Dendronotidae	<i>Dendronotus cf. venustus</i> MacFarland, 1966
		Dotidae	<i>Doto uva</i> Er. Marcus, 1955
		Hancockiidae	<i>Hancockia schoeferti</i> Schrödl, 1999
		Phylliroiidae	<i>Cephalopyge trematoides</i> (Chun, 1889) <i>Phylliroe bucephala</i> Lamarck, 1816
		Cuthoniidae	<i>Cuthona</i> sp.
		Fionidae	<i>Fiona pinnata</i> (Eschscholtz, 1831)
		Flabellinidae	<i>Kynaria cynara</i> (Ev. Marcus & Er. Marcus, 1967) <i>Coryphellina cerverai</i> (M. A. Fischer, van der Velde & Roubos, 2007)
	Proctonotoidea	Janolidae	<i>Janolus rebeccaiae</i> Schrödl, 1996
	Tritonioidea	Tritoniidae	<i>Tritonia</i> sp.
Doridina	Chromodoridoidea	Chromodorididae	<i>Tyrinna delicata</i> (Abraham, 1877) <i>Tyrinna evelinae</i> (Er. Marcus, 1958) <i>Felimare agassizii</i> (Bergh, 1894) <i>Felimare sechurana</i> Hoover, Padula, Schrödl, Hooker & Valdés, 2017 <i>Felimida baumanni</i> (Bertsch, 1970)
			<i>Baptodoris peruviana</i> (d'Orbigny, 1837)
			<i>Diaulula variolata</i> (d'Orbigny, 1837)
			<i>Diaulula punctulata</i> (d'Orbigny, 1837)
			<i>Rostanga pulchra</i> MacFarland, 1905
	Doridoidea	Dorididae	<i>Doris fontainii</i> d'Orbigny, 1837
		Corambidae	<i>Corambe lucea</i> Er. Marcus, 1959
			<i>Corambe mancorensis</i> Martynov et al., 2011
	Onchidoridoidea	Goniodorididae	<i>Okenia luna</i> Millen, Schrödl, Vargas & Indacochea, 1994
	Phyllidioidea	Dendrodorididae	<i>Doriopsilla janaina</i> Er. Marcus & Ev. Marcus, 1967
	Polyceroidea	Polyceridae	<i>Polycera priva</i> Er. Marcus, 1959

in Chilean waters, four species in Colombian waters, and four species in Brazilian waters (Fig. 3B). The recorded collection/reporting sites of nudibranch species found in Peruvian waters, limited to South America, within the framework of the coastal-marine biogeographical classification proposed by Spalding et al. (2007), are illustrated in Fig. 4.

Table 2. Nudibranch species that could potentially occur in Peruvian waters based on the bibliographic compilation of this study.

Suborder (n = 2)	Superfamilies (n = 5)	Families (n = 8)	Species (n = 9)
Cladobranchia	Aeolidioidea	Aeolidiidae	<i>Aeolidia campbellii</i> (Cunningham, 1871)
		Glaucidae	<i>Glaucus</i> sp.
		Phylliroiidae	<i>Phylliroe lichensteinii</i> Eschscholtz, 1825
Doridina	Fionoidea	Coryphellidae	<i>Itaxia falklandica</i> (Eliot, 1907)
	Flabellinoidea	Flabellinidae	<i>Coryphellina marcusorum</i> (Gosliner & Kuzirian, 1990)
Dordina	Chromodoridoidea	Cadlinidae	<i>Cadlina sparsa</i> (Odhner, 1922)
		Discodorididae	<i>Gargamella immaculata</i> Bergh, 1894
	Polyceroidea	Polyceridae	<i>Polycera cf. alabe</i> Collier & Farmer, 1964
			<i>Thecacera darwini</i> Pruvot-Fol, 1950

Table 3. Chronologically ordered publications listing nudibranch species in the Peruvian sea. Legend: First reports for Peruvian waters: ^ad'Orbigny (1835–1846), ^bDall (1909), ^cMillen et al. (1994), ^dNakamura (2006), ^eSchrödl (2003), ^fSchrödl (1999), ^gSchrödl (2000), ^hMartynov et al. (2011), ⁱUribe et al. (2013), ^jUribe and Pacheco (2012), ^kSchrödl and Hooker (2014), ^lHoover et al. (2017), ^mQuesquen (2017), sp = The total count of nudibranch species mentioned in each checklist. Single asterisk (*) shows potentially occurring species and double asterisk (**) shows recent confirmation of previously predicted species in Peruvian waters. [§] shows that the scientific name has undergone changes.

d'Orbigny (1835–1846) (sp = 5)	Dall (1909) (sp = 6)	Álamo and Valdivieso (1997) (sp = 7)	Paredes et al. (1999) / Ramírez et al. (2003) (sp = 18)	(Nakamura 2006) (sp = 23)	Uribe et al. (2013) (sp = 25)	Schrödl and Hooker (2014) (sp = 14)	This study (sp = 31) * nine potentially occurring species
<i>Doriopsis peruviana</i> ^a	<i>Doris peruviana</i>	<i>Dendrodoris peruviana</i>	<i>Doris peruviana</i>	<i>Baptodoris peruviana</i>	<i>Baptodoris peruviana</i>	<i>Baptodoris?</i> <i>peruviana</i>	<i>Baptodoris peruviana</i> [§]
<i>Diphyllidia cuvieri</i>	<i>Pleurophyllidia cuvieri</i>			<i>Armina cuvieri</i>	<i>Armina californica</i>		<i>Armina californica</i> [§]
<i>Phidiana natans</i> ^a	<i>P. natans/ Fiona pinnata</i>	<i>Phidiana natans</i>	<i>P. natans / Fiona pinnata</i>	<i>Fiona pinnata</i>	<i>Fiona pinnata</i>		<i>Fiona pinnata</i> [§]
<i>Phidiana inca</i> ^a	<i>Phidiana inca</i>	<i>Phidiana inca</i>	<i>Phidiana lottini</i>	<i>Phidiana lottini</i>	<i>Phidiana lottini</i>	<i>Phidiana lottini</i>	<i>Phidiana lottini</i> [§]
<i>Glaucus distichoicus</i>	<i>Glaucus distichoicus</i>		<i>Glaucus atlanticus</i>		<i>Glaucus atlanticus</i>		<i>Glaucus atlanticus</i> [§]
	<i>Doris punctuolata</i> ^b	<i>Doris punctuolata</i>	<i>Anisodoris punctuolata</i>	<i>Diadulula punctuolata</i>			<i>Diadulula punctuolata</i> [§]
		<i>Okenia luna</i> ^c	<i>Okenia luna</i>	<i>Okenia luna</i>	<i>Okenia luna</i>	<i>Okenia luna</i>	<i>Okenia luna</i>
		<i>Cadlina?</i> <i>sparsa*</i>	<i>Cadlina sparsa*</i>	<i>Cadlina sparsa*</i>			<i>Cadlina sparsa*</i>
		<i>Rostanga pulchra*</i>	<i>Rostanga pulchra*</i>	<i>Rostanga pulchra*</i>		<i>Rostanga pulchra**</i>	<i>Rostanga pulchra</i>
			<i>Aeolidia serotina*</i>	<i>Aeolidia serotina*</i>			<i>Aeolidia campbellii</i> [§]
			<i>Hypselodoris cf. agassizii</i>	<i>Hypselodoris agassizii</i>	<i>Felimare agassizii</i>		<i>Felimare agassizii</i> [§]
			<i>Flabellina cf. falklandica*</i>	<i>Flabellina falklandica</i>			<i>Itaxia falklandica</i> [§]
			<i>Dendronotus frondosus</i>	<i>Dendronotus frondosus</i>	<i>Dendronotus cf. venustus</i>		<i>Dendronotus cf. venustus</i> [§]

d'Orbigny (1835–1846) (sp = 5)	Dall (1909) (sp = 6)	Álamo and Valdivieso (1997) (sp = 7)	Paredes et al. (1999) / Ramírez et al. (2003) (sp = 18)	(Nakamura 2006) (sp = 23)	Uribe et al. (2013) (sp = 25)	Schrödl and Hooker (2014) (sp = 14)	This study (sp = 31) * nine potentially occurring species
			<i>Doto cf. uva</i>	<i>Doto uva</i>	<i>Doto uva</i>	<i>Doto uva</i>	<i>Doto uva</i>
			<i>Polycera cf. alabe</i>	<i>Polycera alabe</i>	<i>Polycera alabe</i>		<i>Polycera cf. alabe</i>
			<i>Tyrinna evelinae</i>	<i>Tyrinna evelinae</i>	<i>Tyrinna evelinae</i>		<i>Tyrinna evelinae</i>
			<i>Bajaeolus bertschi</i>	<i>Bajaeolis bertschi</i>	<i>Bajaeolis bertschi</i>		<i>Bajaeolis bertschi</i>
			<i>Phylliroe lichensteini*</i>				<i>Phylliroe lichensteini**</i>
				<i>Flabellina cynara^d</i>	<i>Flabellina cynara</i>		<i>Kynaria cynara[§]</i>
				<i>Glossodoris baumanni^d</i>	<i>Glossodoris baumanni</i>		<i>Felimida baumanni[§]</i>
				<i>Cuthona sp.^d</i>	<i>Cuthona sp.</i>		<i>Cuthona sp.</i>
				<i>Doriopsilla janaina^d</i>	<i>Doriopsilla janaina</i>		<i>Doriopsilla janaina</i>
				<i>Flabellina sp. 2^e</i>	<i>Flabellina cerverai</i>	<i>Flabellina cf. cerverai</i>	<i>Coryphellina cerverai[§]</i>
				<i>Gargamella immaculata^{*f}</i>			<i>Gargamella immaculata*</i>
				<i>Doris fontainei^g</i>	<i>Doris fontainei</i>	<i>Doris fontainei</i>	<i>Doris fontainii[§]</i>
					<i>Corambe mancorensis^h</i>		<i>Corambe mancorensis</i>
					<i>Diaulula variolataⁱ</i>	<i>Diaulula variolata</i>	<i>Diaulula variolata</i>
					<i>Tyrinna nobilis^j</i>		<i>Tyrinna delicata[§]</i>
					<i>Tritonia sp.^j</i>		<i>Tritonia sp.</i>
					<i>Spurilla cf. neapolitanaⁱ</i>	<i>Spurilla braziliiana</i>	<i>Spurilla braziliiana</i>
				<i>Thecacera darwini*</i>	<i>Thecacera darwini*</i>	<i>Thecacera darwini*</i>	<i>Thecacera darwini*</i>
						<i>Polycera priva^k</i>	<i>Polycera priva</i>
						<i>Corambe lucea^k</i>	<i>Corambe lucea</i>
						<i>Janolus rebecca^k</i>	<i>Janolus rebecca</i>
						<i>Hancockia schoeferti^k</i>	<i>Hancockia schoeferti</i>
							<i>Felimare sechurana^l</i>
							<i>Cephalopyge trematoides^m</i>
							<i>Glaucus sp.*^m</i>
							<i>Phylliroe bucephala^m</i>
							<i>Coryphellina marcusorum*</i>

Table 4. Presence of nudibranch species inhabiting Peruvian waters along several marine coastal biogeographic provinces according the reporting sites. Legend: the asterisk (*) indicates potentially occurring species in Peruvian waters.

Families	Species	Magellanic	Pacific				Atlantic				Lusitanian	Mediterranean Sea	Circumtropical	Cosmopolitan
			Warm Temperate Southeastern Pacific	Tropical Eastern Pacific	Warm Temperate Northeast Pacific	Cold Temperate Northeast Pacific	Galapagos	North Brazil Shelf	Tropical Northwestern Atlantic	Tropical Southwestern Atlantic				
Aeolidiidae	<i>Spurilla brasiliiana</i>		X	X	X			X	X					
	<i>Aeolidia campbellii</i> (*)	X	X											
Facelinidae	<i>Phidiana lottini</i>	X	X											
	<i>Bajaeolis bertschi</i>			X	X									
Glaucidae	<i>Glaucus atlanticus</i>		X	X	X		X		X	X			X	X
	<i>Glaucus</i> sp. (*)		X											
Arminidae	<i>Armina californica</i>			X	X	X								
Dendronotidae	<i>Dendronotus cf. venustus</i>		X		X	X								
Dotidae	<i>Doto uva</i>	X	X								X			
Hancockiidae	<i>Hancockia schoeferti</i>	X	X											
Phyllioidae	<i>Cephalopyge trematoides</i>		X	X	X					X		X		X
	<i>Phylliroe bucephala</i>		X	X					X			X	X	X
	<i>Phylliroe lichensteinii</i> (*)		X							X				X
Coryphellidae	<i>Itaxia falklandica</i> (*)	X	X								X			
Cuthonidae	<i>Cuthona</i> sp.	X		X										
Fionidae	<i>Fiona pinnata</i>		X										X	
Flabellinidae	<i>Kynaria cynara</i>		X	X	X									
	<i>Coryphellina cerverai</i>		X	X							X			
	<i>Coryphellina marcusorum</i> (*)			X										
	<i>Janolidae</i>	<i>Janolus rebecca</i>	X	X										
Tritoniidae	<i>Tritonia</i> sp.		X	X										
Cadlinidae	<i>Cadlina sparsa</i> (*)	X	X		X									
Chromodorididae	<i>Tyrinna delicata</i>	X	X											
	<i>Tyrinna evelinae</i>			X	X									
	<i>Felimare agassizii</i>		X	X	X		X							
	<i>Felimare sechurana</i>			X										
	<i>Felimida baumanni</i>			X	X									

Families	Species	Magellanic	Pacific			Atlantic				Mediterranean Sea	Circumtropical	Cosmopolitan
			Warm Temperate Southeastern Pacific	Tropical Eastern Pacific	Warm Temperate Northeast Pacific	Cold Temperate Northeast Pacific	Galapagos	North Brazil Shelf	Tropical Northwestern Atlantic	Tropical Southwestern Atlantic	West African Transition	Warm Temperate Southwestern Atlantic
Discodorididae	<i>Baptodoris peruviana</i>	X										
	<i>Diaulula variolata</i>	X	X									
	<i>Diaulula punctuolata</i>		X									
	<i>Rostanga pulchra</i>	X	X		X	X						
	<i>Gargamella immaculata</i> (*)	X	X									
Dorididae	<i>Doris fontainii</i>	X	X									
Corambidae	<i>Corambe lucea</i>	X	X	X								
	<i>Corambe mancorensis</i>			X								
Goniodorididae	<i>Okenia luna</i>		X									
Dendrodorididae	<i>Doriopsilla janaina</i>			X	X		X					
Polyceridae	<i>Polycera priva</i>	X	X									
	<i>Polycera cf. alabe</i> (*)			X	X							
	<i>Thecacera darwini</i> (*)	X	X									




Figure 1. Number of species by family and suborder of Nudibranchia in Peruvian waters.




Figure 2. Venn diagram showing the number of species shared between the two coastal marine biogeographic provinces present in Peru.




Figure 3. **A** Nudibranch species richness by country **B** Venn diagram illustrating the shared species count among countries. Only the confirmed species from Peru were considered. The counts of nudibranch species for neighboring countries were derived from a comprehensive literature review (data not shown).




Figure 4. Reporting sites for nudibranch species found in Peruvian waters whose geographic ranges are limited to South America. The gray band highlights the locations where these species have been reported along the Peruvian coastline. The coastal marine biogeographic classification introduced by Spalding et al. (2007) is presented to provide context.

Confirmed species in Peruvian waters

Suborder Cladobranchia

Superfamily Aeolidioidea Gray, 1827

Family Aeolidiidae Gray, 1827

***Spurilla brasiliiana* MacFarland, 1909**

Peruvian specimen photographs: Uribe and Pacheco (2012), Uribe et al. (2013)
Common name: Brazilian Aeolid

Habitat. Benthic.

Depth. 0–10 m (Gosliner 1979; Uribe and Pacheco 2012).

Type material. Holotype CASIZ 019731—Alagoas, Brazil (Carmona et al. 2014).

Distribution. This species exhibits a distribution range spanning the western Atlantic, extending from Florida to Brazil (Behrens and Hermosillo 2005; Marcus 1959), as well as the Pacific Ocean.

Sampling/reporting sites. In Peru, it was reported in Ferrol Bay (Chimbote, 09°06'S) (Uribe and Pacheco 2012) and Pucusana (Lima, 12°25'S) (Uribe et al. 2013) under the name *Spurilla neapolitana*. In the western Pacific, this species has been reported in Japan (Hamatani 2000), China (Lin 1992), and Australia (Willan 2006). Within the Eastern Pacific, it was reported in Mexico, Costa Rica, and Colombia (Carmona et al. 2013). Additionally, a specimen was reported from Hawaii (Gosliner 1979).

Remarks. Carmona et al. (2013), based on mitochondrial and nuclear sequences, revealed that *Spurilla neapolitana* MacFarland, 1909 comprises a complex of five cryptic species. In the light of this discovery, the name *Spurilla brasiliiana* has been assigned to populations found in the western Atlantic and Pacific regions. Carmona et al. (2013) further speculated that the occurrence of this species in the Pacific Ocean might be attributed to human-mediated introductions.

Family Facelinidae Bergh, 1889

Phidiana lottini (Lesson, 1831)

Peruvian specimen photographs: Fig. 5, iNaturalist

Common name: Flameback Sea Slug

Habitat. Benthic.

Depth. 0–15 m (Schrödl and Hooker 2014).

Type material. Not available.

Distribution. From Puerto Malabriga (La Libertad, Peru, 07°42'S) (Flores 2014) to Melinka (Guaitecas islands, Chilean fjord region, 43°52'S) (Schrödl 2003).

Sampling/reporting sites. In Peru, it was initially reported in Callao as *Phidiana inca* by d'Orbigny, (1835–1846) and Dall (1909). It was also reported in Isla Santa (Ancash, 09°01'S), Ancon (Lima, 11°47'S), San Bartolo (12°22'S), Pucusana (Lima, 12°25'S) and San Juan de Marcona (Ica, 15°21'S) (Uribe et al. 2013). Schrödl and Hooker (2014) also collected individuals in Pucusana (12°25'S), Paracas (13°48'S) and La Punta (Callao). The report by Flores (2014) in Puerto Malabriga (La Libertad, 07°42'S) revealed the northernmost locality of this species distribution. Valdivia-Chavez et al. (2020) presented a recent report of this species in Arequipa (15–17°S). Other reports (occurrences) for



Figure 5. *Phidiana lottini*, courtesy of Yuri Hooker.

this species in the Peruvian coast are Playa Tartacay (El Paraíso, Huaura, Lima, 11°13'44.9"S) (Zanabria 2020e), Isla Chuncho (Pucusana, Lima, 12°28'S) (Guzman 2018a; Cuba 2019) and Puerto General San Martín (Paracas Bay, Pisco, Ica, 13°48'37.3"S) (Zanabria 2020d). In Chile, it was reported in Playa Brava (Caldera, Atacama, 27°03'S), Calderilla (Atacama, 27°05'S) (Araya and Valdés 2016), Punta Blanca (Arica, 18°29'S), Comau Fjord (42°15'S), and Melinka (Guaitecas islands, Chilean fjord region, 43°52'S) (Schrödl 2003). Other reports (occurrences) for this species in the Chilean coast include Chascos Bay (Coquimbo, Atacama, 27°40'S), Reserva Nacional Pingüino de Humboldt (Huasco, Atacama, 29°01'S), Coquimbo (29°58'S), Valparaíso (33°21'S), Concepción (36°45'S), Valdivia (39°57'S), and Chiloé (41°53'S) (iNaturalist 2023a).

Remarks. Initially designated as *Phidiana inca* (d'Orbigny, 1837) until research by Schrödl (1997) who considered it conspecific with *Eolidia lottini* Lesson, 1831, proposing *Phidiana lottini* as a valid name.

***Bajaeolis bertschi* Gosliner & Behrens, 1986**

Peruvian specimen photographs: Uribe et al. (2013), Nakamura (2007)

Habitat. Benthic.

Depth. 3–8 m (Nakamura 2006).

Type material. Holotype CASIZ 059589—Punta la Gringa, Baja California (Gosliner and Behrens 1986).

Distribution. Eastern Pacific, from Baja California (Mexico, 28°N) to the northern coast of Peru (04°S) (Nakamura 2006).

Sampling/reporting sites. In Peru, it was reported in Playa Las Pocitas (Mancora, Piura, 04°06'S) (Nakamura 2006) and Playa El Rubio (Tumbes) (Uribe et al. 2013, based on a personal communication with Sandra Millen). It was also reported in Panama (Camacho-García et al. 2005) and Mexico (Baja California) (Gosliner and Behrens 1986).

Family Glaucidae Gray, 1827

***Glaucus atlanticus* Forster, 1777**

Peruvian specimen photographs: Uribe et al. (2013), iNaturalist

Common name: Sea Swallow

Habitat. Pelagic.

Depth. Neustonic (Churchill et al. 2014b).

Type material. Not available.

Distribution. Cosmopolitan and circumtropical (Churchill et al. 2014b; Thompson and McFarlane 1967).

Sampling/reporting sites. Off the northern coast of Chile (Schrödl 2003). On the coast of El Salvador (13°N) (Segovia and López 2015). In Peru, it was mentioned by Paredes et al. (1999) and Ramírez et al. (2003) based on the records of d'Orbigny (1854) in Callao (10°15'S). Recently reported in Isla Santa, Ancash (09°01'S) by Uribe et al. (2013).

Remarks. Included in Paredes et al. (1999), probably based on a personal communication with Sandra Millen. The records of *Glaucus distichoicus* d'Or-

bigny, 1837 (d'Orbigny 1854; Dall 1909; Paredes et al. 1999; Ramírez et al. 2003) do not have enough evidence to formalize the species within the genus *Glaucus* and could refer to *G. atlanticus*.

Superfamily Arminoidea Iredale & O'Donoghue, 1923 (1841)

Family Arminidae Iredale & O'Donoghue, 1923 (1841)

***Armina californica* (J.G. Cooper, 1863)**

Peruvian specimen photographs: not available

Common name: Striped Nudibranch

Habitat. Benthic.

Depth. 11–268 m (Báez et al. 2011).

Type material. Not available.

Distribution. Eastern Pacific, from the Gulf of Alaska (Báez et al. 2011) to Piura (northern coast of Peru) (Dall 1909; Baez et al. 2011).

Sampling/reporting sites. In Peru, it was reported in Paita (Piura) (Dall 1909). Baez reported a Peruvian specimen (USNM 805043 South Pacific Ocean, Peru) but lacks geographic reference. It was also reported in Alaska (Central Aleutian Islands), Canada (Scott Islands), United States (California), Mexico (Isla Tortuga, Baja California, Socorro Island), and Panama (Islas Ladrones) (see Báez et al. 2011).

Remarks. In Peru, it was initially reported in Paita (Piura) (Dall 1909) under the name *Pleurophyllidia* (*Diphyllidia*) *cuvieri* d'Orbigny, 1837 and later considered as *Armina cuvieri* (d'Orbigny, 1837). However, *D. cuvieri* currently corresponds to *Armina tigrina* Rafinesque, 1814, a species from the Mediterranean Sea (Thompson et al. 1990). Nakamura (2006), referring to Dall (1909), listed *A. californica* as *A. cuvieri*. Báez et al. (2011) examined specimens of *A. cuvieri* and discovered an identification error, as it was actually *A. californica*. This correction is adopted in Uribe et al. (2013).

Superfamily Dendronotoidea Allman, 1845

Family Dendronotidae Allman, 1845

***Dendronotus* cf. *venustus* MacFarland, 1966**

Peruvian specimen photograph: Uribe et al. (2013)

Common name: Branched Dendronotid

Habitat. Benthic.

Depth. 5–20 m (Korshunova et al. 2020).

Type material. Not available.

Distribution. From Alaska (Stout et al. 2010) to Coliumo Bay (Chile, 36°32'S) (Schrödl 2003).

Sampling/reporting sites. in Peru, it was reported in Pucusana (12°25'S) as *Dendronotus* cf. *venustus* (Uribe et al. 2013). It was also reported in Alaska (Stout et al. 2010), United States (Morro Bay and Crescent City, California) (MacFarland 1966), and Chile (Coliumo Bay, 36°32'S) (Schrödl 2003).

Remarks. Paredes et al. (1999) listed this species as *Dendronotus frondosus* (Ascanius, 1774), a North Atlantic species (Ekimova et al. 2015), based

on personal communication with Sandra Millen. *Dendronotus frondosus* was confirmed to have morphological and molecular differences with *Dendronotus venustus* (Stout et al. 2010; Ekimova et al. 2015) that is exclusive to the Pacific.

Family Dotidae Gray, 1853

Doto uva Er. Marcus, 1955

Peruvian specimen photographs: Uribe et al. (2013), Schrödl and Hooker (2014)
Common name: Grape-cluster Nudibranch

Habitat. Benthic.

Depth. 0–15 m (Schrödl and Hooker 2014).

Type material. Not available.

Distribution. Amphi-South American. On the Pacific side of South America, it extends from Callao (Peru, 12°S) to Comau Fjord (Chile, 42°S) (Schrödl et al. 2005). On the Atlantic side it ranges from San Matías gulf (Argentina, 41°26'S) (Cetra and Roche 2023) to São Paulo (Brazil, 21°S) (Marcus 1959).

Sampling/reporting sites. In Peru, it was reported in Callao (12°S), San Juan de Marcona (15°21'S), Islas Ballestas (13°44'S) (Schrödl and Hooker 2014) and Independencia Bay (14°14'S) (Uribe et al. 2013). In Chile, it was collected in Tocopilla (22°05'S), La Herradura (Coquimbo, 29°59'S), Tongoy (Coquimbo, 30°15'S) (Fischer and Cervera 2005b), Coliumo Bay (36°32'S) (Schrödl 2003), Canal de Calbuco (41°45'S) (Marcus 1959), and in Comau Fjord (42°22'S) (Schrödl et al. 2005). In Argentina, it was collected in San Matías Gulf (Patagonia, 41°30'S) (Cetra and Roche 2023). In Brazil, it was reported in São Paulo (21°21'S) (Marcus 1959).

Remarks. Molecular studies are needed to clarify the genetic identities of the populations on both sides of South America (Schrödl 2003; Uribe et al. 2013).

Family Hancockiidae MacFarland, 1923

Hancockia schoeferti Schrödl, 1999

Peruvian specimen photograph: Schrödl and Hooker (2014)

Habitat. Benthic.

Depth. 0–3 m (Schrödl and Hooker 2014).

Type material. Holotype ZSM Moll 19983471—Coliumo Bay (36°32'S), Chile (Schrödl 1999a).

Distribution. San Juan de Marcona, Peru (15°21'S) (Schrödl and Hooker 2014) and southern Chile (37–43°S) (Schrödl 2009).

Sampling/reporting sites. In Peru, it was reported for the first time in San Juan de Marcona (Ica, 15°21'S) (Schrödl and Hooker 2014). In Chile, it was reported in Coliumo Bay (36°32'S) and Queule (39°23'S) (Schrödl 1999a).

Family Phyllioidae Menke, 1830

Cephalopyge trematoides (Chun, 1889)

Peruvian specimen photograph: Quesquen (2017)

Habitat. Pelagic.

Depth. 40 m (Fernández-Alamo 1997).

Type material. Not available.

Distribution. Circumtropical (van der Spoel et al. 1997).

Sampling/reporting sites. In Peru, it was reported in Piura (Quesquen 2017).

It was also reported in the Canary Islands and Cape Verde (Hernández et al. 2001, 2017), New South Wales (Steinberg 1956), Gulf of California (Fernández-Alamo 1997), and central and northern Chilean coast (Tokioka 1963).

Remarks. Originally described as *Phylliroe trematoides* Chun, 1889. The samples described in Quesquen (2017) were reported in grey literature (Quesquen 2008) and had been previously reported by Quesquen and Guzmán (1999).

***Phylliroe bucephala* Lamarck, 1816**

Peruvian specimen photograph: Quesquen (2017)

Habitat. Pelagic.

Depth. 40–60 m (Fernández-Alamo 1997).

Type material. Not available.

Distribution. Circumtropical (van der Spoel et al. 1997).

Sampling/reporting sites. In Peru, it was reported in Tumbes and Piura (Quesquen 2017). It was also reported in the Canary Islands (Hernández and Jiménez 1996), off the coasts of Florida and Bermuda (Abbott 1974), in northeastern Atlantic waters near the African coast (van der Spoel 1970), and in the western Atlantic Ocean (Spencer et al. 2009). In the Mediterranean Sea there are reports from France and Syria (Durgham et al. 2016; Durgham and Ikhtiyar 2020; Pruvot-Fol 1954). Recorded south of the Pacific Ocean in Australia and New Zealand (Powell 1979; Spencer and Willan 1995). In the Indo-Pacific it has been reported from Vietnam (Sachidhanandam et al. 2000).

Family Cuthoniidae Odhner, 1934

***Cuthona* sp.**

Peruvian specimen photograph: Nakamura (2006)

Habitat. Benthic.

Depth. 5–7 m (Nakamura 2006).

Distribution. Northern coast of Peru.

Sampling/reporting sites. Cancas (Tumbes, 03°56'S) (Nakamura 2006).

Remarks. Description provides a length of 5 mm, body completely white, including rhinophores and oral tentacles with a translucent base. In addition, the specimen had dark, reddish-brown cerata without the white tip, which would differentiate it from other species of the genus (Nakamura 2006).

Family Fionidae Gray, 1857

***Fiona pinnata* (Eschscholtz, 1831)**

Peruvian specimen photographs: not available

Common name: Feather Aeolid

Habitat. Pelagic.

Depth. Neustonic (Willan 1979).

Type material. Not available.

Distribution. Cosmopolitan (Gosliner 1987; Schmekel and Portmann 1982).

Sampling/reporting sites. In Peru, it was reported in Lima (d'Orbigny 1835–1846; Álamo and Valdivieso 1997; Dall 1909). It was also reported in Chile (Meihuín, Valdivia, Juan Fernández Islands and Talcahuano) (Fischer and Cervera 2005b; Bergh 1898).

Remarks. Originally named *Eolidia pinnata* Eschscholtz 1831 (type specimen from Alaska), until its current designation as *F. pinnata*; it is a species with a wide latitudinal range (Trickey 2013).

Family Flabellinidae Bergh, 1889

Kynaria cynara (Ev. Marcus & Er. Marcus, 1967)

Peruvian specimen photographs: Fig. 6, Uribe et al. (2013), Nakamura (2006)

Common name: Swimming Cynara

Habitat. Benthic.

Depth. 5–8 m (Nakamura 2006).

Type material. Holotype USNM 678417—La Choya Bay ($31^{\circ}20'30''N$, $113^{\circ}38'06''W$), Puerto Peñasco, Sonora, Mexico.

Distribution. Eastern Pacific, from Gulf of California (Mexico, $28^{\circ}N$) (Millen and Hermosillo 2007) to Isla Tortuga (Peru, $09^{\circ}S$) (Uribe et al. 2013).

Sampling/reporting sites. In Peru, it was reported in Punta Sal ($03^{\circ}56'S$), Cancas ($03^{\circ}56'S$), Mancora ($04^{\circ}6'S$), Chimbote ($09^{\circ}4'S$), and Ancash ($09^{\circ}S$) (Nakamura 2006; Uribe et al. 2013). It was recently reported in Isla La Viuda ($09^{\circ}20'57''S$) and Isla Tortuga ($09^{\circ}21'48''S$) in Ancash (Uribe et al. 2019). It was also reported in Mexico (Gulf of California) (Millen and Hermosillo 2007) and in



Figure 6. *Kynaria cynara*, courtesy of Yuri Hooker.

other several points (occurrences) such as Mexico (Guerrero), Costa Rica (Tamarindo Bay, Parque Nacional Santa Rosa), Ecuador (Salinas) (iNaturalist 2023b).

Remarks. The species was originally described as *Coryphella cynara* Ev. Marcus & Er. Marcus, 1967 and reported along the Peruvian coast as *Flabellina cynara* (Nakamura 2006; Uribe et al. 2013, 2019). These designations are currently invalid and have been replaced by *Kynaria cynara* (Korshunova et al. 2017).

***Coryphellina cerverai* (M. A. Fischer, van der Velde & Roubos, 2007)**

Peruvian specimen photograph: Schrödl and Hooker (2014)

Habitat. Benthic.

Depth. 0–10 m (Schrödl and Hooker 2014).

Type material. Holotype RMNH Moll. 98130—La Herradura, Coquimbo (29°58'S, 071°22'W), Chile.

Distribution. From Sechura Bay (Peru, 05°49'S) to Coliumo Bay (Chile, 36°32'S).

Sampling/reporting sites. In Peru, it was reported in Sechura Bay (05°49'S) (Schrödl and Hooker 2014), Ancon (Lima, 11°47'S) and Pucusana (Lima, Peru, 12°25'S) (Schrödl 1996a). In Chile, it was reported in La Herradura (Coquimbo, 29°59'S) (Fischer et al. 2007), Peñón de Vida Marina (Montemar, Viña del Mar, 32°57'26.8"S) (Molina 2021), Coliumo Bay (36°32'S) (Schrödl 1996a).

Remarks. The species was first reported as *Flabellina* sp. 2 (Schrödl 1996a) and then as *Flabellina cerverai* van der Velde & Roubos, 2007 (Fischer et al. 2007; Schrödl and Hooker 2014).

Superfamily Proctonotoidea Gray, 1853

Family Janolidae Pruvot-Fol, 1933

***Janolus rebeccaiae* Schrödl, 1996**

Peruvian specimen photographs: Fig. 7, Schrödl and Hooker (2014)

Habitat. Benthic.



Figure 7. *Janolus rebeccaiae*, courtesy of Yuri Hooker.

Depth. 2–12 m (Schrödl 1996a, b; Schrödl and Hooker 2014).

Type material. Holotype ZSM 19960557—English Bay (27°07'S, 070°53'W), Chile.

Distribution. From Sechura Bay (Peru, 05°49'S) (Schrödl and Hooker 2014) to English Bay (Chile, 27°07'51"S) (Schrödl 1996b, 2003).

Sampling/reporting sites. In Peru, it was reported in Sechura Bay (05°49'S) and Paracas (13°43'S) (Schrödl and Hooker 2014). In Chile, it was reported in Juan Lopez (23°30'S) and English Bay (27°07'S) (Schrödl 1996a, 1996b).

Superfamily Tritonioidea Lamarck, 1809

Family Tritoniidae Lamarck, 1809

Tritonia sp.

Peruvian specimen photographs: Fig. 8, Uribe et al. (2013), Uribe et al. (2019), [iNaturalist](#)

Habitat. Benthic.

Depth. 5–15 m (Uribe et al. 2013).

Distribution. From Foca Island (Piura, Peru, 05°12'S) to Punta Picata (Tacna, Peru) (Uribe et al. 2013).

Sampling/reporting sites. In Peru, it was reported in Foca Island (Piura, 05°12'S), Santa Island (09°01'S), Ferrol Bay (Chimbote, 09°06'S), Punta El Huaro (Casma, Ancash, 09°37'S), La Gramita (Casma, Ancash, 09°43'S), Punta Patillos (Huarmey, 09°53'S), Punta Colorado (Huarmey, Ancash, 10°29'S), Pucusana (Lima, 12°25'S), Isla Asia (Lima, 12°47'S), Isla La Vieja (Independencia Bay, Pisco, Ica, 14°16'S) and Punta Picata (Tacna, 17°52'S) (Uribe et al. 2013).

Remarks. It bears resemblance to *Tritonia odhneri* (common in Chile) in terms of its external morphology, while displaying similarities to *Tritonia festiva* (found in Alaska, Baja California, and Japan; Uribe et al. 2013). Anatomical and molecular analyses are necessary to describe this species (Uribe et al. 2013).



Figure 8. *Tritonia* sp., courtesy of Jaime Calvo-Pérez.

Suborder Doridina

Superfamily Chromodoridoidea Bergh, 1891

Family Chromodorididae Bergh, 1891

***Tyrinna delicata* (Abraham, 1877)**

Peruvian specimen photographs: Fig. 9, Uribe et al. (2013)

Habitat. Benthic.

Depth. 0–22 m (Uribe et al. 2013).

Type material. Holotype NHMUK 1995038—Chiloé Island, Chile.

Distribution. Amphi-South American. From Pucusana (Lima, Peru, 12°25'S) (Fabián Avilés pers. comm.) to Strait of Magellan (Chile, 53°S) and Peninsula Valdés (Argentina, 42°S) (Schrödl and Millen 2001; Uribe et al. 2013; Araya and Valdés 2016).

Sampling/reporting sites. In Peru, two specimens were collected (MHNURP, specimens currently lost) in Playa Las Ninfas (Pucusana, Lima, 12°28'49"S) on 23 October 2019, at 1.5–2.0 m depth, 55–60 mm length (Fig. 9) (Fabián Avilés pers. comm.). In Peru, this species was also reported in San Juan de Marcona (Ica, 15°21'S) and Isla Blanca (Arequipa, 17°00'S) (Uribe et al. 2013). In Chile, it was reported in Caleta Obispito (Caldera, Atacama, 26°45'51"S) (Araya and Valdés 2016) and in several points (occurrences) such as Reserva Nacional Pingüino de Humboldt (29°15'S), Playa El Francés (Coquimbo, 30°05'S), Pichicuy (32°20'S), Caleta Cocholgüe-Tomé (36°35'S), Caleta Chome (Península de Hualpén, 36°48'S), Caleta Chaihuín (39°56'S), Faro San Isidro (Strait of Magellan, 53°28'S) (iNaturalist 2023a). It was also reported in San Juan Fernández Islands (33°38'S) (Araya and Valdés 2016). In Argentina it was observed (occurrences) in Baliza Davison (Tierra del Fuego, 54°56'S), Grand Jason (Ja-



Figure 9. *Tyrinna delicata*, courtesy of Fabián Avilés.

son Islands, 51°04'S), La Tranquera (46°02'S), Golfo San Jorge (46°00'S), Punta del Marqués (Rada Tilly, 45°57'S), Parque Interjurisdiccional Marino Costero Patagonia Austral (PIMCPA, 45°02'S), Camarones Bay (44°46'S), Golfo Nuevo (42°47'S), and Golfo San José (42°24'S) (iNaturalist 2023c).

Remarks. This species was originally identified as *Tyrinna nobilis* Bergh, 1898, a name that is currently not accepted.

***Tyrinna evelinae* (Er. Marcus, 1958)**

Peruvian specimen photographs: not available

Habitat. Benthic.

Depth. 0–5 m (Welch 2010).

Type material. Not available.

Distribution. Amphi-American and West Africa.

Sampling/reporting sites. In Peru, it was reported in El Rubio (Tumbes, 03°52'S) (Schrödl and Millen 2001). It was also reported in Mexico (Gulf of California, 28°N), Jamaica (Schrödl and Millen 2001). In the Atlantic, it has been sampled in the Gulf of Mexico (de la Cruz-Francisco et al. 2017), Panama (Goodheart et al. 2016), Brazil (Marcus 1958), Ghana, and Cape Verde (Camarcho-García et al. 2005).

Remarks. It was initially listed by Paredes et al. (1999). According to Uribe et al. (2013) molecular studies are necessary to confirm the consistency of reports of *T. evelinae* in both the Atlantic and Pacific populations that are morphologically difficult to distinguish (Valdés et al. 2006).

***Felimare agassizii* (Bergh, 1894)**

Peruvian specimen photographs: Uribe et al. (2013), Nakamura (2007)

Common name: Agassiz's Chromodorid

Habitat. Benthic.

Depth. 7–8 m (Nakamura 2006).

Type material. Not available.

Distribution. From the Gulf of California to the coast of northern Peru.

Sampling/reporting sites. In Peru, it was reported in Cancas (Tumbes, 03°56'S) (Nakamura 2006) and Lobos de Tierra Island (Lambayeque, 06°25'S) (Uribe et al. 2013). It was also reported in Mexico (Gulf of California, 28°N), Ecuador (Galapagos Islands, 0°S), Colombia (Malpelo Island, 04°N) (Behrens and Hermosillo 2005).

Remarks. Originally described as *Chromodoris agassizii* Bergh, 1894 and reported in Peruvian waters as *Hypselodoris agassizii* by Nakamura (2006).

***Felimare sechurana* Hoover, Padula, Schrödl, Hooker & Valdés, 2017**

Peruvian specimen photographs: Fig. 10, iNaturalist, Hoover et al. (2017)

Habitat. Benthic.

Depth. 6–15 m (Hoover et al. 2017; Bravo et al. 2020)



Figure 10. *Felimare sechurana*, courtesy of Yuri Hooker.

Type material. Holotype CZA 402—Foca Island ($05^{\circ}12'13.8''S$, $81^{\circ}12'38.0''W$), Piura, Peru.

Distribution. Northern coast of Peru's transition zone.

Sampling/reporting sites. In Peru, it was reported in Punta Veleros (Los Organos, Piura, $04^{\circ}10'28.7''S$) (Zavala, 2022), Quebrada Verde (Piura, $04^{\circ}13'34.8''S$), Foca Island (Piura, $05^{\circ}12'13.8''S$) (Hoover et al. 2017), and Lobos de Afuera Islands (Lambayeque, $06^{\circ}56'S$) (Bravo et al. 2020).

Remarks. This species is endemic to the northern coast of Peru. It was initially reported as *Felimare ghiselini* (Bertsch 1978) by Thompson (2006) but later corrected by Hoover et al. (2017) and described as a new species.

***Felimida baumanni* (Bertsch, 1970)**

Peruvian specimen photograph: Fig. 11

Common name: Baumann's Chromodorid

Habitat. Benthic.

Depth. 5–8 m (Nakamura 2006).

Type material. Not available.

Distribution. Eastern Pacific, from Gulf of California ($28^{\circ}N$) to Cancas (Tumbes, Peru) (Nakamura 2006).

Sampling/reporting sites. In Peru, it was reported in Cancas (Tumbes, $03^{\circ}56'S$) (Nakamura 2006). It was also reported in Mexico (Gulf of California, $28^{\circ}N$), Panama ($08^{\circ}N$), Ecuador (Galapagos Islands, $00^{\circ}S$) and Colombia (Malpelo Islands, $04^{\circ}N$) (Behrens and Hermosillo 2005).

Remarks. The species was originally reported as *Chromodoris baumanni* Bertsch, 1970 in the Eastern Pacific (Rudman 1983). Years later, Gosliner et al. (2004) discussed the anatomical characteristics of this species to be like those attributed to the genus *Glossodoris*, suggesting a reclassification. Finally, the phylogenetic study by Johnson and Gosliner (2012) defined its new classification as part of the genus *Felimida*.



Figure 11. *Felimida baumanni*, courtesy of Fabián Encinas.

Superfamily Doridoidea Rafinesque, 1815

Family Discodorididae Bergh, 1891

***Baptodoris peruviana* (d'Orbigny, 1837)**

Peruvian specimen photographs: Uribe et al. (2013), Schrödl and Hooker (2014)

Habitat. Benthic.

Depth. 4–15 m (Schrödl and Hooker 2014).

Type material. Holotype ZMB 50748—Isla de Pajargo (Pájaros), Chile (as *Platydoris punctatella* Bergh, 1898), poorly preserved. Neotype SSUC 6977 (*Doris peruviana*), Iquique (Chile, 20°12'S), February 1965, undissected specimen (Fischer and Cervera 2005a).

Distribution. From San Lorenzo Island (Callao, Peru, 12°S) (d'Orbigny 1835–1846) to Los Molles (Valparaíso, 32°15'S) (Fischer and Cervera 2005).

Sampling/reporting sites. In Peru it was reported for first time as *Doriopsis peruviana* in San Lorenzo Island (12°05') by d'Orbigny (1835–1846). It was also reported in Callao (12°S, as *Doris peruviana*, Dall 1909), Pucusana (Lima, 12°28'S, as *Platydoris punctatella*, Schrödl 1996a), Tortugas Bay (Ancash, 09°21'S), Independencia Bay (Ica, 14°14'S), and San Juan de Marcona (Ica, 15°21'S) (Uribe et al. 2013). In Chile, it was collected in Iquique (20°12'S), Mejillones (23°20'S), La Portada (Antofagasta, 23°39'S), and Los Molles (Valparaíso, 32°15'S) (Fischer and Cervera 2005a).

Remarks. Initially reported as *Doris peruviana* d'Orbigny 1836, transferred to *Platydoris* Bergh, 1877 by Schrödl (2003), who also included Bergh's (1898) description of *P. punctatella* as a junior synonym. Dorgan et al. (2002) ruled out that these reports were about a species belonging to *Platydoris*, based on a photograph of a live specimen (Schrödl 1996a). Based on the description of the radular teeth, this species was transferred from *Platydoris* to the genus *Baptodoris* (Fischer and Cervera 2005a). Regarding its northernmost distribution, Fischer and Cervera (2005a) considered the reports from Tagus Cove (Albermarle, Galapagos Islands) (Pilsbry and Vanatta 1902) as doubtful.

***Diaulula variolata* (d'Orbigny, 1837)**

Peruvian specimen photographs: Fig. 12, Schrödl and Hooker (2014), iNaturalist

Habitat. Benthic.

Depth. 2–15 m (Schrödl and Hooker 2014).

Type material. Not available.

Distribution. From Pucusana (Lima, Peru, 12°28'S) (Guzman 2018b) to Punta Hualpén (Concepción, Chile, 36°44'S) (Marcus 1959).

Sampling/reporting sites. In Peru, it was reported in Pucusana (Lima, 12°28'S) (Guzman 2018b), El Chaco (Ica, 13°49'S), Caleta Atenas (Ica, 13°49'S), Independencia Bay (Ica, 14°14'S), San Juan de Marcona (Ica, 15°21'S) (Uribe et al. 2013), Dos Playas (Arequipa, 17°00'S) and Playa Calera (Arequipa, 17°15'S) (Tejada-Pérez et al. 2018; Valdivia-Chavez et al. 2020), Caleta La Huata (Camaná, Arequipa, 16°50'S) (Zanabria 2020a), Terminal Portuario Matarani (Islay, Arequipa, 16°59'S) (Zanabria 2020b), and Playa Jaboncillo (Ilo, Moquegua, 17°59'S) (Zanabria 2020c).

In Chile, it was reported in Arica (18°26'S) (Schrödl 2003), Coquimbo Bay (29°57'S) (Bergh 1898; Valdés and Muniaín 2002), English Bay-Guanaqueiros-Los Hornos (30°10'S), El Tabo (Valparaíso, 33°27'S) (Fischer and Cervera 2005b), Coliumo Bay (36°32'S) (Schrödl 1996a, 1997), San Vicente Bay (36°44'S), and Punta Hualpén (Concepción, 36°44'S) (Marcus 1959). Within its Chilean distribution, several other recent observations (occurrences) have been reported (iNaturalist 2023d).

Remarks. This species had not been recorded outside Chile (Fischer and Cervera 2005b) until the first reports from Peru (Uribe et al. 2013). The specimen collected in Bernardo O'Higgins National Park (Chilean fjord region, 51°S) (Aldea et al. 2011) needs confirmation (Uribe et al. 2013).



Figure 12. *Diaulula variolata*, courtesy of Jaime Calvo-Pérez.

***Diaulula punctulata* (d'Orbigny, 1837)**

Peruvian specimen photographs: not available

Common name: Dotted Bristle Sea Slug

Habitat. Benthic.

Depth. 0–7 m.

Type material. ZSM Moll 20040984—Ipún Island (44°33'S, 74°48'W), Aysén, Chile.

Distribution. Amphi-South American. It is frequently found on the Magellanic coasts of Chile and Argentina.

Sampling/reporting sites. In Peru, it was collected in Callao (12°S) (Dall 1909; Schrödl 2003). In Chile, it was collected in Lota (37°05'S), Lacuy Peninsula (Greater Island of Chiloé, 41°49'S) (Valdés and Muníaín 2002), Ipún Island (Chonos Archipelago, 44°33'S) (Schrödl and Grau 2006) and Strait of Magellan (53°35'S) (Roche et al. 2023). In Argentina, it was collected in San Matías Gulf (41°30'S) (Roche et al. 2023; Cetra 2019), Gulf Nuevo (42°42'S) (Valdés and Muníaín 2002), Peninsula Valdés (42°30'S), Punta Pardelas (42°36'S), Puerto Madryn (42°46'S) (Roche et al. 2023; Cetra 2019), Comodoro Rivadavia (45°51'S) (Valdés and Muníaín 2002), and Tierra del Fuego (54°21'S) (Roche et al. 2023).

Remarks. This species was listed as *Anisodoris punctulata* (d'Orbigny, 1836) and *Doris punctulata* d'Orbigny, 1837 in previous Peruvian articles listing nudibranch species. Both names are currently not accepted.

***Rostanga pulchra* MacFarland, 1905**

Peruvian specimen photographs: Schrödl and Hooker (2014)

Common name: Red Sponge Dorid

Habitat. Benthic.

Depth. 6–12 m (Schrödl and Hooker 2014).

Type material. Holotype USNM 181292—Monterey Bay, California, United States.

Distribution. This species presents a bipolar distribution in the Eastern Pacific and an amphi-South American distribution (Schrödl 2003; Schrödl and Grau 2006).

Sampling/reporting sites. In Peru it was considered a predicted species (Uribe et al. 2013) until the first specimen was finally reported in Punta San Juan (Marcona, Ica, 15°22'S) (Schrödl and Hooker 2014). It has been also reported in Alaska (Point Craven) (Lee and Foster 1985), Mexico Los Angeles Bay (Lance 1966), Chile (Coliumo Bay, Playa Brava and Chonos archipelago) (Marcus 1959; Schrödl 2003; Schrödl and Grau 2006), and Argentina (Camarones Bay) (Marcus and Marcus 1969).

Remarks. Its distribution in Peruvian Waters was not certain; however, it was listed by Álamo and Valdivieso (1997) and Paredes et al. (1999). *Rostanga pulchra* is the only species of the genus in Peruvian and Chilean waters. Schrödl and Hooker (2014) mentioned that populations in the northern and southern hemispheres are likely to be distinct species based on preliminary unpublished molecular data.

Family Dorididae Rafinesque, 1815

Doris fontainii d'Orbigny, 1837

Peruvian specimen photographs: Fig. 13, iNaturalist

Common name: Large Warted Doris

Habitat. Benthic.

Depth. 8–17 m (Schrödl and Hooker 2014).

Type material. Holotype ZSM 19983417. Coliumo Bay, Chile (36°32'S, 72°57'W).

Distribution. Amphi-South American.

Sampling/reporting sites. In Peru it was reported in Independencia Bay (14°14'S) (Schrödl 1996a Schrödl and Grau 2006; Rudman 2002a, 2002b; Uribe et al. 2013), San Fernando National Reserve (14°58'S) (Ampuero 2010), and recently in Arequipa (15–17°S) (Valdivia-Chavez et al. 2020). Other reports (occurrences) for this species in the Peruvian coast are Pucusana, Ilo, Marcona, Pisco, Islay and Camaná, (iNaturalist 2023e). In Chile it was reported in Arica (18°28'S), Coquimbo (29°57'S), Dichato (36°32'S), Puerto Montt (41°27'S) (Valdés and Muniaín 2002; Schrödl 2003), and the Chonos Archipelago (45°08'S) (Schrödl and Grau 2006). In Argentina, it was collected from 37°50'S to Rada Tilly (45°55'S) (Valdés and Muniaín 2002).

Remarks. Erroneously named as *Doris fontainei* in previous articles.

Superfamily Onchidoridoidea Gray, 1827

Family Corambidae Bergh, 1871

Corambe lucea Er. Marcus, 1959

Peruvian specimen photograph: Schrödl and Hooker (2014)

Habitat. Benthic.




Figure 13. *Doris fontainii*, courtesy of Jaime Calvo-Pérez.

Type material. Neotype ZSM 1912—Caleta Buena (Chile, 22°25'S, 70°15'W).

Depth. 0–27 m (Schrödl and Hooker 2014)

Distribution. From Bayóvar (Sechura Bay, Peru, 05°49'S) (Schrödl and Hooker 2014) to Gulf of Corcovado (Chile, 42°46'50"S) (Schrödl 1996a).

Sampling/reporting sites. In Peru, it was reported in Bayóvar (Sechura Bay, 05°49'S), Callao (pier of IMARPE, 12°03'59"S), Ballestas Islands (Paracas, 13°43'54"S) and San Juan de Marcona (Ica, 15°21'S) (Schrödl and Hooker 2014). In Chile, it was reported in Caleta Buena (22°25'S), Coliumo Bay (36°32'S) (Schrödl 1997), Faro Punta Corona (Chiloé, 41°50'S) (Marcus, 1959), Ancud Bay (41°52'S) (Schrödl 1996a), Comau Fjord (42°22'S) (Schrödl et al. 2005) and Gulf of Corcovado (42°46'S) (Schrödl 1996a).

Remarks. It was first described as *Neocorambe lucea* (Schrödl 1996a). A morphological analysis confirmed its status as a valid species (Schrödl and Wägele 2001).

***Corambe mancorensis* Martynov, Brenzinger, Hooker & Schrödl, 2011**

Peruvian specimen photograph: Martynov et al. (2011)

Habitat. Benthic.

Depth. 0–3 m (Martynov et al. 2011).

Type material. ZSM 20080543—Mancora (04°06'36"S, 81°04'02"W), Piura, Peru.

Distribution. Species only reported off the coast of Mancora (Piura, Peru) (Martynov et al. 2011).

Remarks. Endemic species of northern coast of Peru.

Family Goniodorididae H. Adams & A. Adams, 1854

***Okenia luna* Millen, Schrödl, Vargas & Indacochea, 1994**

Peruvian specimen photographs: Fig. 14, Schrödl and Hooker (2014)



Figure 14. *Okenia luna*, courtesy of Yuri Hooker.

Habitat. Benthic.

Depth. 4–20 m (Schrödl 1996a).

Type material. Holotype CASIZ 089293—Coliumo Bay ($36^{\circ}32'S$, $73^{\circ}57'W$), north of Concepción, Chile.

Distribution. From Ancon Bay (Lima, Peru) to Coliumo Bay (Chile).

Sampling/reporting sites. In Peru, it was collected in Ancon Bay (Lima, $11^{\circ}47'S$) (Millen et al. 1994). In Chile, it was collected in Iquique ($22^{\circ}13'S$) (Fischer 2006) and Coliumo Bay ($36^{\circ}32'S$) (Schrödl 2003).

Remarks. First record of the genus *Okenia* reported in the Southeast Pacific (Millen et al. 1994). It is distributed in Peruvian and Chilean waters.

Superfamily Phyllidoidea Rafinesque, 1814

Family Dendrodorididae O'Donoghue, 1924 (1864)

***Doriopsilla janaina* Er. Marcus & Ev. Marcus, 1967**

Peruvian specimen photograph: Nakamura (2007)

Habitat. Benthic.

Depth. 0–3 m (Nakamura 2006).

Type material. Holotype USNM 576269—Panama Canal ($09^{\circ}05'N$, $79^{\circ}41'W$), Panama.

Distribution. From the Gulf of California ($28^{\circ}N$) to Cancas (Peru, $03^{\circ}56'S$).

Sampling/reporting sites. In Peru, it was reported in Cancas (Tumbes, $03^{\circ}56'S$) (Nakamura, 2006). It was also reported in Mexico (Punta Lobos, Sonora), Panama ($08^{\circ}N$), and Ecuador (Galapagos Islands, $00^{\circ}S$) (Gosliner 1991).

Superfamily Polyceroidea Alder & Hancock, 1845

Family Polyceridae Alder & Hancock, 1845

***Polycera priva* Er. Marcus, 1959**

Peruvian specimen photographs: Schrödl and Hooker (2014)

Habitat. Benthic.

Depth. 10 m (Schrödl 1996a).

Type material. Holotype ZSM Moll. 20041005—Punta Llonco, Comau Fjord ($42^{\circ}22'10"S$, $072^{\circ}27'18"W$), Chile.

Distribution. From Paracas (Ica, Peru, $13^{\circ}43'S$) (Schrödl and Hooker 2014) to Chilean Patagonia ($54^{\circ}S$) (Schrödl 2009; Schrödl et al. 2005).

Sampling/reporting sites. In Peru, it was reported in Ballestas Islands (Lima, $13^{\circ}43'S$) (Schrödl and Hooker 2014). In Chile, it was reported in Gulf of Ancud ($42^{\circ}06'S$) (Marcus 1959) and Comau Fjord ($42^{\circ}22'S$) (Schrödl et al. 2005).

Remarks. Peruvian presence of this species was surprising for the discoverers (Schrödl and Hooker 2014) since it was considered a Magellanic species, being very abundant in the fjords of southern Chile (41 – $44^{\circ}S$) (Schrödl 1996a, 1999b, 2003, 2009).

Potentially occurring species (unconfirmed)

Thecacera darwini Pruvot-Fol, 1950

Common name: Dalmatian Sea Slug

Habitat. Benthic.

Type locality. Orange Bay (55°31'S), Nassau Bay, Chile (Valdés and Héros 1998).

Distribution. From Juan López, northern coast of Chile, to Strait of Magellan (Schrödl 2003).

Remarks. This species was listed by Nakamura (2006) based on Schrödl (1999b), who listed this species as present in the Peruvian province, referring to Chilean waters. Fischer and Cervera (2005b) argued against its occurrence off the Chilean coast, although they acknowledge a high likelihood of its presence in Peruvian waters. Uribe et al. (2013) and Fischer (2006) referred to Zagal and Hermosilla's (2001) presumed finding of this species in Peru.

Glaucus sp.

Habitat. Pelagic.

Type material. Not available.

Distribution. Ica (Peru).

Sampling/reporting sites. In Peru, Quesquen (2017) reported specimens twice off the coast of Ica in 1995 and 1998. Currently, the only valid species of the genus *Glaucus* is *G. atlanticus*, reported in Peru by Uribe et al. (2013) from Isla Santa, Ancash.

Remarks. Quesquen (2017) offered a description of the specimens collected in Ica. He described a slender body that was ventrally flattened, with a small head and two cephalic tentacles. The dorsum was navy blue and green, and the ventral area was white. Additionally, the specimens had three or four branches on both sides of the body, and their body length could reach up to 43 mm. According to Valdés and Campillo's (2004) description, *G. atlanticus* is characterized by its slim and elongated body, along with a small head and sleek oral tentacles and rhinophores. The coloration of its dorsum can vary from deep blue to brown hues. It possesses up to three groups of cerata, and its ventral region exhibits a silver shade.

A confirmation of the taxonomic status of these specimens is necessary, using morphological and molecular analyses.

Aeolidia campbellii (Cunningham, 1871)

Habitat. Benthic.

Type material. ZSM 20020700 (Chile), designated as the neotype due to absence of the holotype (Kienberger et al. 2016).

Distribution. Falkland Islands (50°S), Argentine and Chilean Patagonia (41°S) to Valparaíso (32°S) (Schrödl 2003). In Peru, this species was listed by Paredes et al. (1999) as *Aeolidia serotina* Bergh, 1873 and replicated by Ramírez et al. (2003).

Remarks. It was reported off the coast of Chile by Schrödl (2003) as *Aeolidia papillosa* (Linnaeus, 1761), later reassigned to the species *A. campbellii* according to molecular studies by Kienberger et al. (2016).

***Gargamella immaculata* Bergh, 1894**

Common name: Unicolored Sea Lemon

Habitat. Benthic.

Type material. SMNH 1015—Tierra del Fuego, Chile.

Distribution. A common species on the southern coast of Chile and Argentina.

Sampling/reporting sites. Cabo Metalqui, Chiloé (Fischer and Cervera 2005b; Odhner 1926), in Última Esperanza, Tierra del Fuego (Marcus 1959), in Cabo San Antonio; Cabo Delgado; Gulf of Ancud, between Isla Quenu and Calbuco; Seno Otway, Queule and Coliumo Bay (Schrödl 1996a). In Argentina, in the north (Bergh 1894; Odhner 1926), also in Argentine Patagonia and on the Burdwood Bank (Odhner 1926; Schrödl 2003).

Remarks. According to Schrödl (2003), the Peruvian records of this species by Zagal and Hermosilla (2001) are doubtful. However, in that compilation and in the most up-to-date publication (Zagal and Hermosilla 2007), *Gargamella immaculata* is not mentioned from Peruvian waters, but as an inhabitant of the Peruvian zoogeographic province from Juan López (Atacama) to the south. The species was mistakenly included in Kentrodorididae by Schrödl (1996a), until Valdés (2002) transferred it back to the Discodorididae.

***Cadlina sparsa* (Odhner, 1922)**

Common name: Dark-spot Cadlina

Habitat. Benthic.

Type locality. Juan Fernández Islands and Desventuradas Islands, Chile (Odhner 1922).

Distribution. It presents disjunct populations with a bipolar distribution in the eastern Pacific and an amphi-South American pattern.

Sampling/reporting sites. In the Pacific, the northernmost location is Baja California (Behrens 1991; Jaeckle 1983) and the southernmost location is the Comau Fjord in southern Chile. In the Atlantic, it was recorded in Camarones Bay in the central region of Argentina (Schrödl 2000, 2003). In Chile, it has also been sampled in the Juan Fernández Islands (Odhner 1922), Chiloé Islands (Marcus 1959), and Coliumo Bay (Schrödl 1996a, 2003).

Remarks. *Cadlina sparsa* was initially proposed as probable species in Peruvian water by Álamo and Valdivieso (1997). Subsequently, its presence was consistently mentioned in the lists compiled by Paredes et al. (1999), Ramírez et al. (2003), and Nakamura (2006). However, no actual specimens have been collected from intermediate Pacific locations, including Peru. Despite the absence of direct observations, a hypothetical distribution for Peru has been predicted through extrapolation, assuming a continuous geographic range (Uribe et al. 2013).

Studies have demonstrated that *C. sparsa* does not fall within the family Chromodorididae, as initially suggested (Johnson 2011), but rather belongs to the family Cadlinidae. However, there have been no other revisions or updates regarding its scientific name. The taxonomy of the genera involved has been thoroughly elucidated in previous literature (Schrödl 2000).

***Polycera cf. alabe* Collier & Farmer, 1964**

Habitat. Benthic.

Type material. CASIZ 18190—Cedros Island, Baja California ($28^{\circ}12'13''N$, $115^{\circ}15'28''W$), Mexico.

Distribution. From Baja California (Behrens 2004; Behrens and Hermosillo 2005; Camacho-García et al. 2005), Puerto Vallarta ($20^{\circ}40'N$) in Mexico to Costa Rica (Behrens 2004); and northern Chile with a single isolated record (Schrödl 2003).

Remarks. In Peru, Paredes et al. (1999) listed this species as *Polycera cf. alabe*, likely based on information obtained through personal communication with Sandra Millen. Subsequently, Uribe et al. (2013) included this species in their listing, also citing personal communication with Sandra Millen, who observed this species at El Rubio (Tumbes).

***Phylliroe lichtensteinii* Eschscholtz, 1825**

Habitat. Pelagic.

Type material. Not available.

Distribution. Cosmopolitan (Padula 2015).

Sampling/reporting sites. Espírito Santo, southeastern Brazil (Ralph 1959).

Remarks. For Peru, the species was listed in Ramírez et al. (2003). There are no reports of collections of this species in Peruvian waters.

***Itaxia falklandica* (Eliot, 1907)**

Habitat. Benthic.

Depth. 1–15 m (Schrödl 1996a).

Type material. Not available.

Distribution. Abundant in the Magellanic Province, in the Southeast Pacific (Aldea et al. 2011; Velasco-Charpentier et al. 2021) and with records in the South Atlantic (Eliot 1907; Odhner 1926, 1944), Pacific Ocean (Marcus 1959; Meyers-Muñoz et al. 2009; Schrödl 1996a, 2003), and Indian Ocean (Odhner 1944).

Remarks. It was included in the list of mollusks of Peru by Paredes et al. (1999) as *Flabellina falklandica* (Eliot 1907); this information was repeated by Ramírez et al. (2003).

According to Uribe et al. (2013), its presence in Peru requires confirmation since the inclusion by Paredes et al. (1999) was based on unfounded records.

Coryphellina marcusorum (Gosliner & Kuzirian, 1990)

Common name: Marcus's Aeolid

Habitat. Benthic.

Depth. 3–22 m (Welch 2010).

Type material. CASIZ 066151—San Diego Reef (25°12'N, 110°42'W), Gulf of California, Mexico.

Distribution. From Brazil to Gulf of California (Mexico) (Gosliner and Kuzirian 1990; Fischer et al. 2007).

Sampling/reporting sites. In Peru, this species was recently photographed in Los Organos (Piura, 04°10'S) on 11 March 2022 (Torrejón 2023).

Remarks. Originally named *Flabellina marcusorum* Gosliner & Kuzirian, 1990. Its presence in Peru needs be confirmed by future surveys.

Discussion

Overview

This article presents an updated compilation of nudibranchs found in Peru, derived from an extensive literature review. The revised and updated scientific names are presented, while species not verified or erroneously listed in previous articles, referred to here as 'potentially occurring', are separated from those confirmed.

Despite the presence of two distinct coastal marine biogeographic provinces and a transitional zone between them (Schrödl and Hooker 2014; Ibanez-Erquiaga et al. 2018; Barahona et al. 2019), the species richness of nudibranchs remains relatively low ($n = 31$) compared to other marine regions, such as the Gulf of Mexico, the Caribbean, the South Atlantic, the Indian Ocean, and the Mediterranean Sea (Behrens 2004; Chavanich et al. 2013). In neighboring South American countries, such as Colombia ($n > 40$) (Ardila et al. 2007; Londoño-Cruz 2021), Chile ($n > 50$) (Schrödl 1996a, 1999b, 2003; Fischer and Cervera 2005b; Schrödl et al. 2005; Schrödl and Grau 2006; Aldea et al. 2011), and Brazil ($n > 80$) (Marcus and Marcus 1969; Padula et al. 2011; Pereira et al. 2012; Padula 2015), a greater diversity of nudibranch species is found (Fig. 3A, B).

It is worth noting that there are areas along the Peruvian coast that remain unexplored. In the Tropical Eastern Pacific, only a limited number of locations have been sampled, including Pocitas, Punta Sal, Mancora, and Cancas. Within the transition zone, Sechura Bay and Foca Island are the common reporting sites, while within the Warm Temperate Southeastern Pacific, Santa, Casma, Huarmey (Ancash), Ancon, Callao, San Lorenzo Island, Pucusana (Lima), Pisco, Independencia Bay, San Juan de Marcona (Ica), Matarani, and Isla Blanca (Arequipa) are frequently mentioned. Factors such as limited exploration efforts, challenging diving conditions, a scarcity of nudibranch taxonomists, and a general lack of interest in this group in Peru should be highlighted. In addition, it is highly likely that several species remain unreported and undescribed, particularly in deeper waters. Therefore, the confirmed number of nudibranch species in Peruvian waters ($n = 31$) is presumed to represent only a fraction of the actual diversity present.

Potential influence of the Humboldt Current and El Niño

Several confirmed species exhibit a biogeographical affinity for the Warm Temperate Southeastern Pacific ($n = 23$) (Fig. 2). The Humboldt Current plays a significant role as an oceanographic factor, facilitating the influx of various species from the Chilean sea into Peruvian waters, with 19 of them occurring in both. For instance, *Phidiana lottini*, found in Callao and Puerto Malabriga, may extend into tropical waters due to the influence of the Humboldt Current (Uribe et al. 2013). The occurrence of *Polycera priva* in Peru came as a surprise to researchers, considering that this species is typically Magellanic and endemic to the Patagonian fjords (Schrödl 1996a, 1999b, 2003, 2009). Therefore, its presence in Peru is attributed to the influence of the Humboldt Current (Schrödl 1996a, 1999b, 2003, 2009; Schrödl and Hooker 2014). Species such as *Corambe lucea*, *Coryphellina cerverai*, *Janolus rebecca*, and *Tritonia* sp., collected in the Warm Temperate Southeastern Pacific province, have also been reported in Sechura Bay and Foca Island, situated in the biogeographical transition zone. This suggests their adaptation to slightly warmer waters (Fig. 4).

El Niño events can induce shifts in the distribution ranges of sea slugs (Goddard et al. 2018) and even result in species turnover (Valqui et al. 2021). During strong El Niño events, it is noteworthy that mollusk species endemic to tropical areas have been observed in Peruvian waters due to the displacement of warm-water masses (Velez and Zeballos 1985; Paredes et al. 1998; Ramírez et al. 2003). Species such as *Armina californica*, *Felimida baumanni*, *Felimare agassizii*, *Tyrinna evelinae*, and *Bajaeolis bertschi*, which are abundant in the Tropical Eastern Pacific province, Warm Temperate Northeast Pacific, or Cold Temperate Northeast Pacific, have only been recorded in Tumbes or Piura (the northernmost coastal areas of Peru) (Table 4). This suggests that their presence may be temporary, resulting from the displacement of warm-water masses, or they may have permanently adapted to similar conditions following multiple El Niño events (Ashton et al. 2008).

The distribution patterns of cosmopolitan and circumglobal species can be attributed to various biological factors, including their remarkable dispersal capabilities. Take, for instance, *Glaucus atlanticus*, which possesses intriguing adaptations for dispersal such as larval gas bubbles and specialized anatomy enabling it to exploit water surface tension (Thompson and McFarlane 1967; Valdés and Campillo 2004; Churchill et al. 2014a). Additionally, abiotic factors like ocean currents (Miller 1993) and indirect human influences such as buoy rafting (Astudillo et al. 2009), ballast water, or shipping activities can facilitate the dispersal of these species, expanding their geographic ranges (Borg et al. 2009).

Potentially occurring species

This group of species poses a challenge as they have been consistently listed and referenced in several previous articles (indicated by asterisks in Table 3) despite lack of substantiated records. One notable example is *Cadlina sparsa*. Initially mentioned as a probable species by Alamo and Valdivieso (1997), it persisted in the subsequent publications of Paredes et al. (1999), Ramírez et al.

(2003), and Nakamura (2006) without acknowledging its speculative status. In the work of Uribe et al. (2013), the authors discussed the predicted occurrence of *Cadlina sparsa* in Peruvian waters based on extrapolation, assuming a continuous distribution, due to its presence in Baja California and Chile.

Paredes et al. (1999), based on personal communication with Sandra Millen, included *Aeolidia campbellii* (referred to as *Aeolidia serotina*) and *Itaxia falklandica* (referred to as *Flabellina falklandica*), which persisted in the subsequent lists of Nakamura (2006) and Ramírez et al. (2003), respectively. However, no additional published reports have surfaced to substantiate their presence. Ramírez et al. (2003) also included the species *Phylliroe lichensteinii* without providing any justification for its inclusion. On the other hand, Nakamura (2006) mistakenly listed *Gargamella immaculata* and *Thecacera darwini* as present in Peruvian waters, referencing Schrödl (1999b). Nevertheless, Schrödl (1999b) only listed these species in a table as occurring in the ‘zoogeographic Peruvian province’, indicating their presence in the Chilean waters corresponding to this biogeographical province, but not in Peruvian waters.

A solitary specimen of *Coryphellina marcusorum* was recently documented through photography on the northern coast of Peru (Los Organos, Piura) (Torrejón 2023). Given the species’ tropical distribution and a previous record near countries such as Ecuador (GBIF 2022), its occurrence in Peru is plausible. However, these two records do not provide evidence of an established population in this region, suggesting the possibility that it may have arrived independently through shipping or been displaced by ocean currents. Further collections and taxonomic verifications are necessary to confirm its presence definitively.

Potentially occurring species should not be included in the official list of Peruvian nudibranch species. However, considering their disjointed or patchy geographic distributions or unique observations, they may be reported in Peruvian waters in forthcoming papers. *Rostanga pulchra*, for instance, was a long-standing predicted species for Peruvian waters until its initial sighting in San Juan de Marcona (Ica) by Schrödl and Hooker (2014). As a cosmopolitan species, the presence of *Phylliroe lichensteinii* in Peru is plausible. Fischer and Cervera (2005b) have deemed the presence of *Thecacera darwini* in Peruvian waters highly probable. The term “potentially occurring” signifies a provisional status, indicating that these species have not yet been officially confirmed.

Identification uncertainties

The genus *Polydora* displays remarkable color variability, seemingly correlated with its geographic range (Behrens and Hermosillo 2005). A genetic investigation identified clades with overlapping distributions in the Northeastern Pacific, strongly suggesting the existence of a species complex (Santander and Valdés 2013). One year later, a morphology-based study by Pola et al. (2014) revealed that specimens previously collected by Camacho-García et al. (2005) in Costa Rica, Panama, and Mexico actually belong to a new species, *Polydora anae*. Considering that *P. alabe* was solely “observed” by Millen in Tumbes, it is possible that it represents another *Polydora* species, such as *P. anae*, given its inclusion in a species complex (Santander and Valdés 2013).

The report of *Glaucus* sp., documented by Quesquen (2017), is based on a single poorly preserved specimen sampled in 1995, warranting verification

through molecular methods. Currently, within the genus *Glaucus*, only the species *G. atlanticus* has been confirmed, as many previous records turned out to be synonyms (e.g., *G. distichoicus*) or were later reassigned to the genus *Glaucilla*. Similar circumstances apply to the reports of *Tritonia* sp. and *Cuthona* sp., as their species-level recognition is still pending.

Concluding remarks

This research contributes to the dissemination and diffusion of this understudied group of organisms in Peru. It is imperative to intensify monitoring efforts to verify the presence of doubtful species, evaluate anthropogenic impacts, and El Niño-driven displacements. Furthermore, considering the intricate nature of external morphological identification, frequent variability in coloration, and the probable existence of cryptic species, it is possible that a considerable number of species remain undiscovered. Consequently, there is an urgent need for comprehensive research involving detailed internal anatomy and the application of molecular tools, such as DNA barcoding and phylogenetic analyses. These methodologies will play a vital role in shedding light on the taxonomy and evolutionary relationships within this group.

Acknowledgements

We express our profound gratitude to Yuri Hooker, Fabián Avilés, Fabián Encinas, and Jaime Calvo-Pérez for their kindness in providing us with their wonderful photographs of these beautiful marine creatures. We are also grateful for the partial support received from Universidad Científica del Sur (UCSUR). Likewise, we are immensely grateful to the reviewer and editor who have made valuable contributions to the improvement of this manuscript.

Additional information

Conflict of interest

The authors have declared that no competing interests exist.

Ethical statement

No ethical statement was reported.

Funding

No funding was reported.

Author contributions

Conceptualization: AG. Data curation: AA. Investigation: AG. Methodology: SPB, AG. Software: SPB. Supervision: SPB. Validation: AA. Writing - original draft: AG. Writing - review and editing: SPB.

Author ORCIDs

Alessandra Grández  <https://orcid.org/0000-0002-0142-9357>

André Ampuero  <https://orcid.org/0000-0001-6929-5423>

Sergio P. Barahona  <https://orcid.org/0000-0002-0136-7205>

Data availability

All of the data that support the findings of this study are available in the main text or Supplementary Information.

References

- Abbott RT (1974) American seashells: The marine Mollusca of the Atlantic and Pacific coasts of North America. Van Nostrand Reinhold, New York, 663 pp.
- Ah Shee Tee L, Puchooa D, Bhoyroo V, Appadoo C (2019) A review of nudibranch (Mollusca: Euthyneura) diversity from the Republic of Mauritius: Status and future work. *Western Indian Ocean Journal of Marine Science* 18(1): 83–93. <https://doi.org/10.4314/wiojms.v18i1.8>
- Álamo V, Valdivieso V (1997) Lista sistemática de moluscos marinos del Perú (Segunda edición, revisada y actualizada). Publicación Especial. Instituto del Mar del Perú, 183 pp. <https://repositorio.imarpe.gob.pe/handle/20.500.12958/1436>
- Aldea C, Cesped T, Rosenfeld S (2011) Opisthobranchs from Bernardo O'Higgins National Park (S. Chile). *Thalassas* 27(2): 37–48.
- Almada F, Levy A, Robalo JI (2016) Not so sluggish: The success of the *Felimare picta* complex (Gastropoda, Nudibranchia) crossing Atlantic biogeographic barriers. *PeerJ* 4: e1561. <https://doi.org/10.7717/peerj.1561>
- Alvim J, Pimenta AD (2013) Taxonomic review of the family Discodorididae (Mollusca: Gastropoda: Nudibranchia) from Brazil, with descriptions of two new species. *Zootaxa* 3745(2): 152–198. <https://doi.org/10.11646/zootaxa.3745.2.2>
- Ampuero A (2010) Nudibranch from Ica, Peru. Sea Slug Forum. <http://www.seaslugforum.net/find/23633>
- Araya JF, Valdés Á (2016) Shallow water heterobranch sea slugs (Gastropoda: Heterobranchia) from the Region de Atacama, northern Chile. *PeerJ* 4: e1963. <https://doi.org/10.7717/peerj.1963>
- Ardila NE, Báez D, Valdés Á (2007) Babosas y Liebres de mar (Mollusca: Gastropoda: Opisthobranchia) de Colombia. *Biota Colombiana* 8(2): 185–197. <http://revistas.humboldt.org.co/index.php/biota/article/view/190>
- Ashton T, Riascos JM, Pacheco A (2008) First record of *Cymatium keenae* Beu, 1970 (Mollusca: Ranellidae) from Antofagasta Bay, northern Chile, in connection with El Niño events. *Helgoland Marine Research* 62(S1): 107–110. <https://doi.org/10.1007/s10152-007-0084-5>
- Astudillo J, Bravo M, Dumont C, Thiel M (2009) Detached aquaculture buoys in the SE Pacific: Potential dispersal vehicles for associated organisms. *Aquatic Biology* 5: 219–231. <https://doi.org/10.3354/ab00151>
- Báez DP, Ardila N, Valdés Á, Acero PA (2011) Taxonomy and phylogeny of *Armina* (Gastropoda: Nudibranchia: Arminidae) from the Atlantic and eastern Pacific. *Journal of the Marine Biological Association of the United Kingdom* 91(5): 1107–1121. <https://doi.org/10.1017/S0025315410002109>
- Barahona SP, Velez-Zuazo X, Santa-Maria M, Pacheco AS (2019) Phylogeography of the rocky intertidal periwinkle *Echinolittorina paytensis* through a biogeographic transition zone in the Southeastern Pacific. *Marine Ecology (Berlin)* 40(4). <https://doi.org/10.1111/maec.12556>
- Barnosky AD, Matzke N, Tomaia S, Wogan G, Swartz B, Quental TB, Marshall C, McGuire JL, Lindsey EL, Maguire KC, Mersey B, Ferrer EA (2011) Has the Earth's sixth mass extinction already arrived? *Nature* 471(7336): 51–57. <https://doi.org/10.1038/nature09678>

- Behrens DW (1991) Pacific coast nudibranchs: A guide to the opisthobranchs, Alaska to Baja California (2nd edn.). Sea Challengers, Monterey, CA, 107 pp.
- Behrens DW (2004) Pacific Coast Nudibranchs. Supplement II New Species to the Pacific Coast and New Information on the Oldies 55(2): 11–54.
- Behrens DW, Hermosillo A (2005) Eastern Pacific nudibranchs: A guide to the opisthobranchs from Alaska to Central America. Sea Challengers, Monterrey, California, 137 pp.
- Behrens DW, Petrinos C, Schrurs C (2005) Nudibranch behavior. New World Publications Incorporated, Jacksonville, Florida, 176 pp.
- Bergh R (1894) Die Opisthobranchien, Reports on the dredging operations off the west coast of Central America to the Galápagos, to the west coast of Mexico, and in the Gulf of California, in charge of Alexander Agassiz, carried on by the U.S. Fish Commission steamer "Albatross", during 1891, Lieut. Commander Z. L. Tanner, U.S. N. commanding (Vol. 25). Museum of Comparative Zoölogy. <https://www.biodiversitylibrary.org/item/48546>
- Bergh R (1898) Die Opisthobranchier der Sammlung Plate. Zoologische Jahrbücher 4: 481–582.
- Borg JA, Schembri PJ, Evans J (2009) Occurrence of the alien nudibranch *Melibe viridis* (Kelaart, 1858) (Opisthobranchia, Tethydidae), in the Maltese Islands. Mediterranean Marine Science 10(1): 131–136. <https://doi.org/10.12681/mms.127>
- Bravo N, Carbalal W, Chanamé-Céspedes J, Fupuy J, Alfaro J (2020) Biodiversidad marina en las islas Lobos de Afuera, Lambayeque. SINGAN 1: 337–352.
- Caballer M, Canteras JC, Ortea J (2008) Babosas marinas (Sacoglosos y Opistobranquios) de la bahía de Santander. En Locustella: Anuario de la Naturaleza de Cantabria, 44–57.
- Camacho-García Y, Gosliner TM, Valdés Á (2005) Guía de Campo de las Babosas Marinas del Pacífico Este Tropical. California Academy of Sciences, San Francisco, California, 130 pp.
- Carmona L, Pola M, Gosliner TM, Cervera JL (2013) A Tale That Morphology Fails to Tell: A Molecular Phylogeny of Aeolidiidae (Aeolidida, Nudibranchia, Gastropoda). PLoS ONE 8(5): e63000. <https://doi.org/10.1371/journal.pone.0063000>
- Carmona L, Lei BR, Pola M, Gosliner TM, Valdés Á, Cervera JL (2014) Untangling the *Spurilla neapolitana* (Delle Chiaje, 1841) species complex: A review of the genus *Spurilla* Bergh, 1864 (Mollusca: Nudibranchia: Aeolidiidae). Zoological Journal of the Linnean Society 170(1): 132–154. <https://doi.org/10.1111/zoj.12098>
- Ceballos G, Ehrlich PR, Barnosky AD, García A, Pringle RM, Palmer TM (2015) Accelerated modern human-induced species losses: Entering the sixth mass extinction. Science Advances 1(5): e1400253. <https://doi.org/10.1126/sciadv.1400253>
- Cetra N (2019) Abundancia, diversidad y variabilidad temporal de un ensamble de nudibranchios en un arrecife costero del golfo San Matías. Honor Thesis. Universidad Nacional del Comahue. <http://rdi.uncoma.edu.ar/handle/uncomaid/16855>
- Cetra N, Roche A (2023) Nudibranchia (Mollusca: Gastropoda) from San Matías Gulf, Northern Argentine Patagonia. Zootaxa 5244(5): 455–473. <https://doi.org/10.11646/zootaxa.5244.5.3>
- Chavanich S, Viyakarn V, Sanpanich K, Harris LG (2013) Diversity and occurrence of nudibranchs in Thailand. Marine Biodiversity 43(1): 31–36. <https://doi.org/10.1007/s12526-012-0141-4>
- Churchill CKC, Valdés Á, Ó Foighil D (2014a) Molecular and morphological systematics of neustonic nudibranchs (Mollusca: Gastropoda: Glaucidae: Glaucus), with descrip-

- tions of three new cryptic species. *Invertebrate Systematics* 28(2): 174–195. <https://doi.org/10.1071/IS13038>
- Churchill CKC, Valdés Á, Ó Foighil D (2014b) Afro-Eurasia and the Americas present barriers to gene flow for the cosmopolitan neustonic nudibranch *Glaucus atlanticus*. *Marine Biology* 161(4): 899–910. <https://doi.org/10.1007/s00227-014-2389-7>
- Cuba D (2019) Flameback (*Phidiana lottini*). iNaturalist. <https://www.inaturalist.org/observations/20735738>
- d'Orbigny A (1854) List of the shells of South America in the collection of the British Museum. Printed by order of the trustees. <https://doi.org/10.5962/bhl.title.15821>
- d'Orbigny AD, Bibron G, Blanchard E, Brullé GA, Gervais P, Guizot F, Lucas PH, von Martius KFP, Milne-Edwards H, Montagne JFC, Valenciennes M (1835) Voyage dans l'Amérique méridionale (le Brésil, la république orientale de l'Uruguay, la République argentine, la Patagonie, la république du Chili, la république de Bolívia, la république du Pérou), exécuté pendant les années 1826, 1827, 1828, 1829, 1830, 1831, 1832, et 1833: Vol. t.1. Chez Pitois-Levrault et ce., libraires-éditeurs, rue de la Harpe, no. 81, 1835–1847. <https://doi.org/10.5962/bhl.title.8353>
- Dall WH (1909) Report on a collection of shells from Peru, with a summary of the littoral marine Mollusca of the Peruvian zoological province. *Proceedings of the United States National Museum* 37(1704): 147–294. <https://doi.org/10.5479/si.00963801.37-1704.147>
- De la Cruz-Francisco V, Ortigosa D, González-González M (2017) First records of marine sea slugs (Gastropoda: Heterobranchia) of the Tuxpan Reef System, Mexico, with extended distribution records. *Biodiversity and Natural History* 3(1): 15–23. <https://www.biodiversnathist.com/index.php/bnh/article/view/37>
- Dean L, Prinsep MR (2017) The chemistry and chemical ecology of nudibranchs. *Natural Product Reports* 34(12): 1359–1390. <https://doi.org/10.1039/C7NP00041C>
- Dorgan KM, Valdés Á, Gosliner TM (2002) Phylogenetic systematics of the genus *Platydoris* (Mollusca, Nudibranchia, Doridoidea) with descriptions of six new species. *Zoologica Scripta* 31(3): 271–319. <https://doi.org/10.1046/j.1463-6409.2002.00105.x>
- Durgham H, Ikhtiyar S (2020) First record of *Phylliroe bucephala* Péron & Lesueur, 1810 in the Ras-Ibn-Hani (Lattakia-Syria). *International Journal of Agriculture & Environmental Science* 7(01): 39–42. <https://doi.org/10.14445/23942568/IJAES-V7I1P108>
- Durgham H, Ikhtiyar S, Ibraheem R (2016) First record of *Pelagia noctiluca* (Forsskål, 1775) on the coast of Syria. *Marine Biodiversity Records* 9(1): 39. <https://doi.org/10.1186/s41200-016-0060-3>
- Echevin V, Goubanova K, Belmadani A, Dewitte B (2012) Sensitivity of the Humboldt Current system to global warming: A downscaling experiment of the IPSL-CM4 model. *Climate Dynamics* 38(3): 761–774. <https://doi.org/10.1007/s00382-011-1085-2>
- Ehrlich H (2010) Molluscs Spicules. In: Ehrlich H (Ed.) *Biological Materials of Marine*. Springer Netherlands, 211–242. <https://doi.org/10.1007/978-94-007-5730-1>
- Ekimova I, Korshunova T, Schepetov D, Neretina T, Sanamyan N, Martynov A (2015) Integrative systematics of northern and Arctic nudibranchs of the genus *Dendronotus* (Mollusca, Gastropoda), with descriptions of three new species. *Zoological Journal of the Linnean Society* 173(4): 841–886. <https://doi.org/10.1111/zoj.12214>
- Eliot C (1907) Nudibranchs from New Zealand and the Falkland Islands. *The Journal of Molluscan Studies* 7(6): 327–361. <https://doi.org/10.1093/oxfordjournals.mollus.a066204>
- Eschscholtz JF (1831) Zoologischer Atlas enthaltend Abbildungen und Beschreibungen neuer Thierarten während des Flottcapitains von Kotzebue zweiter Reise um die Welt

- auf der Russisch-Kaiserlichen Kriegesschlupp Predpriaetië in den Jahren 1823–1826, G. Reimer. <https://doi.org/10.5962/bhl.title.38058>
- Faulkner DJ, Ghiselin MT (1983) Chemical defense and evolutionary ecology of dorid nudibranchs and some other opisthobranch gastropods. *Marine Ecology Progress Series* 13: 295–301. <https://doi.org/10.3354/meps013295>
- Fernández-Alamo A (1997) Distribución horizontal y vertical de *Phylliroe bucephala* y *Cephalopyge trematoides* (Opistobranchia-Nudibranchia) en el Golfo de California. *Hidrobiológica* 7: 75–80.
- Fischer MA (2006) Opisthobranchs from the Chilean Coast. A taxonomic, morphological and histological study of nudibranch species. *Cellular Animal Physiology*. Radboud University Nijmegen.
- Fischer MA, Cervera JL (2005a) *Baptodoris peruviana* (D'Orbigny, 1837) comb. nov., an alternative taxonomic placement for *Doris peruviana* (Gastropoda: Nudibranchia: Doridoidea). *Journal of Conchology* 38(5): 513–528.
- Fischer MA, Cervera JL (2005b) Checklist of the opisthobranchs (Mollusca: Gastropoda) from the Chilean coast deposited at the "Colección de Flora y Fauna Profesor Patricio Sanchez Reyes, Pontificia Universidad Católica de Chile". *Iberus : Revista de la Sociedad Espanola de Malacologia* 23(2): 165–181.
- Fischer MA, Roubos E, van der Velde G (2007) Morphology, anatomy, and histology of a new species of *Flabellina* Voigt, 1834 (Opisthobranchia: Aeolidoidea) from the Chilean coast. *Animal Biology* (Leiden, Netherlands) 57(4): 363–391. <https://doi.org/10.1163/157075607782232116>
- Flores D (2014) Diversidad de macrozoobentos en Puerto Malabriga, La Libertad, abril a setiembre 2014. Tesis para optar el título de Biólogo pesquero. Universidad Nacional de Trujillo, Trujillo, Perú. <http://dspace.unitru.edu.pe/handle/UNITRU/12345>
- Furfaro G, Vitale F, Licchelli C, Mariottini P (2020) Two Seas for One Great Diversity: Checklist of the Marine Heterobranchia (Mollusca; Gastropoda) from the Salento Peninsula (South-East Italy). *Diversity* (Basel) 12(5): 171. <https://doi.org/10.3390/d12050171>
- GBIF (2022) *Coryphellina marcusorum* (Gosliner & Kuzirian, 1990). GBIF Backbone Taxonomy. Checklist dataset. <https://www.gbif.org/species/9259595> [Accessed on 2023-05-27]
- Goddard JH, Treneman N, Prestholdt T, Hoover C, Green B, Pence WE, Mason DE, Dobry P, Sones JL, Sanford E, Agarwal R, McDonald G, Johnson R, Gosliner TM (2018) Heterobranch sea slug range shifts in the northeast Pacific Ocean associated with the 2015–16 El Niño. *Proceedings of the California Academy of Sciences* 65(4): 107–131.
- Goodheart JA, Ellingson RA, Vital XG, Galvão H, McCarthy J, Medrano S, Bhave V, García-Méndez K, Jiménez L, López G, Hoover C, Awbrey J, De Jesús J, Gowacki W, Krug P, Valdés Á (2016) Identification guide to the heterobranch sea slugs (Mollusca: Gastropoda) from Bocas del Toro, Panama. *Marine Biodiversity Records* 9(56): 56. <https://doi.org/10.1186/s41200-016-0048-z>
- Gosliner TM (1979) The systematics of the Aeolidacea (Nudibranchia: Mollusca) of the Hawaiian Islands, with descriptions of two new species. University of Hawaii Press 33(1): 37–77.
- Gosliner TM (1987) Nudibranchs of southern Africa: A guide to opisthobranch molluscs of southern Africa. Sea Challengers Inc; J. Hamann in association with the California Academy of Sciences, 136 pp. <https://doi.org/10.1163/9789004631496>
- Gosliner TM (1991) The opisthobranch gastropod fauna of the Galápagos Islands. In: James MJ (Ed.) Galápagos marine invertebrates: Taxonomy, biogeography, and evo-

- lution in Darwin's islands. Plenum Press, New York, and London, 281–305. https://doi.org/10.1007/978-1-4899-0646-5_14
- Gosliner TM, Behrens DW (1986) Two new species and genera of Aeolid nudibranchs from the Tropical Eastern Pacific. *The Veliger* 29(1): 101–113.
- Gosliner TM, Kuzirian AM (1990) Two new species of Flabellinidae (Opisthobranchia: Aeolidacea) from Baja California. *Proceedings of the California Academy of Sciences*. 4, 47(1): 1–15. <https://biodiversitylibrary.org/page/15776021>
- Gosliner TM, Ortea J, Valdés Á (2004) New data on tropical eastern Pacific Chromodorididae (Nudibranchia: Doridina) with description of a new species of Mexichromis Bertsch, 1977. *Proceedings-California academy of sciences* 55(26/35): 588.
- Gosliner TM, Behrens DW, Valdés Á (2008) Indo-Pacific nudibranchs and sea slugs: A field guide to the world's most diverse fauna. California Academy of Sciences/Sea Challengers Natural History Books, San Francisco, 425 pp.
- Greenwood PG (2009) Acquisition and use of nematocysts by cnidarian predators. *Toxicon* 54(8): 1065–1070. <https://doi.org/10.1016/j.toxicon.2009.02.029>
- Gutiérrez MC, Ortea J, Rivero N, Tucker GC, Malaquias MA, Narciso S (2015) The opisthobranch gastropods (Mollusca: Heterobranchia) from Venezuela: An annotated and illustrated inventory of species. *Zootaxa* 4034(2): 201–256. <https://doi.org/10.11646/zootaxa.4034.2.1>
- Guzman R (2018a) Flameback (*Phidiana lottini*). iNaturalist. <https://www.inaturalist.org/observations/19010160>
- Guzmán R (2018b) *Diaulula variolata*. iNaturalist. <https://www.inaturalist.org/observations/19010219>
- Hamatani I (2000) A New Recorded Species of the Genus *Spurilla* Bergh, 1864 from Osaka Bay, Middle Japan (Opisthobranchia, Aeolidacea). *Venus, the Japanese Journal of Malacology*, 59: 263–265.
- Hernández F, Jiménez S (1996) Nota sobre la presencia de *Phylliroe bucephala* (Mollusca, Opistobranchia, Nudibranchia) en aguas de la isla de El Hierro (Canarias). *Revista de la Academia Canaria de Ciencias* 8: 173–181.
- Hernández F, Navarro SJ, Fernández-Alamo MA, Tejera E, Arbelo E (2001) Sobre la presencia de moluscos nudibranquios planctónicos en el archipiélago de Cabo Verde. *Revista de la Academia Canaria de Ciencias* 12: 49–54.
- Hernández F, de Vera A, Soldevilla FL, García-Talavera F, Collazo N, Fraile-Nuez E (2017) Scientific note on the presence of *Cephalopyge trematoides* (Chun, 1889) from Canary Islands waters. *Spixiana* 40: 6.
- Hoover CA, Padula V, Schrödl M, Hooker Y, Valdés Á (2017) Integrative taxonomy of the *Felimare californiensis* and *F. ghiselini* species complex (Nudibranchia: Chromodorididae), with description of a new species from Peru. *The Journal of Molluscan Studies* 83(4): 461–475. <https://doi.org/10.1093/mollus/eyx035>
- Ibanez-Erquiaga B, Pacheco AS, Rivadeneira MM, Tejada CL (2018) Biogeographic zonation of rocky intertidal communities along the coast of Peru (3.5–13.5°S Southeast Pacific). *PLoS ONE* 13(11): e0208244. <https://doi.org/10.1371/journal.pone.0208244>
- iNaturalist (2023a) Flameback (*Phidiana lottini*). https://www.inaturalist.org/observations?place_id=any&subview=map&taxon_id=487858
- iNaturalist (2023b) Swimming Cynara (*Kynaria cynara*). https://www.inaturalist.org/observations?place_id=any&subview=map&taxon_id=949423
- iNaturalist (2023c) Tyrinna delicata. https://www.inaturalist.org/observations?place_id=any&subview=map&taxon_id=50230

- iNaturalist (2023d) *Diaulula variolata*. https://www.inaturalist.org/observations?place_id=any&subview=map&taxon_id=491332
- iNaturalist (2023e). Large Warted Doris (*Doris fontainii*). https://www.inaturalist.org/observations?place_id=any&subview=map&taxon_id=543510
- Jaeckle WB (1983) Additions to the opisthobranch mollusk fauna of Marin County, California, with notes on the biology of certain species. *The Veliger* 26(2): 93–95.
- Johnson RF (2011) Breaking family ties: Taxon sampling and molecular phylogeny of chromodorid nudibranchs (Mollusca, Gastropoda): *Cadlina* is not a chromodorid. *Zoologica Scripta* 40(2): 137–157. <https://doi.org/10.1111/j.1463-6409.2010.00457.x>
- Johnson RF, Gosliner TM (2012) Traditional taxonomic groupings mask evolutionary history: A molecular phylogeny and new classification of the chromodorid nudibranchs. *PLoS ONE* 7(4): e33479. <https://doi.org/10.1371/journal.pone.0033479>
- Jörger KM, Stöger I, Kano Y, Fukuda H, Knebelsberger T, Schrödl M (2010) On the origin of Acochlidia and other enigmatic euthyneuran gastropods, with implications for the systematics of Heterobranchia. *BMC Evolutionary Biology* 10(1): 323. <https://doi.org/10.1186/1471-2148-10-323>
- Kienberger K, Carmona L, Pola M, Padula V, Gosliner TM, Cervera JL (2016) *Aeolidia pillosa* (Linnaeus, 1761) (Mollusca: Heterobranchia: Nudibranchia), single species or a cryptic species complex? A morphological and molecular study. *Zoological Journal of the Linnean Society* 177(3): 481–506. <https://doi.org/10.1111/zoj.12379>
- Korshunova T, Martynov A, Bakken T, Evertsen J, Fletcher K, Mudianta IW, Saito H, Lundin K, Schrödl M, Picton B (2017) Polyphyly of the traditional family Flabellinidae affects a major group of Nudibranchia: Aeolidacean taxonomic reassessment with descriptions of several new families, genera, and species (Mollusca, Gastropoda). *ZooKeys* 717: 1–139. <https://doi.org/10.3897/zookeys.717.21885>
- Korshunova T, Bakken T, Grøtan VV, Johnson KB, Lundin K, Martynov A (2020) A synoptic review of the family Dendronotidae (Mollusca: Nudibranchia): a multilevel organismal diversity approach. *Contributions to Zoology* (Amsterdam, Netherlands) 90(1): 93–153. <https://doi.org/10.1163/18759866-BJA10014>
- Lance JR (1966) New distributional records of some northeastern Pacific Opisthobranchiata (Mollusca: Gastropoda) with descriptions of two new species. *The Veliger* 9(1): 69–81.
- Leatherman M (2019) Developmental differences in Monterey Sea Lemon (*Doris montereyensis*) veliger larvae with temperature variation. Friday Harbor Laboratories, University of Washington. <https://digital.lib.washington.edu/researchworks/handle/1773/45646>
- Lee RS, Foster NR (1985) A distributional list with range extensions of the opisthobranch gastropods of Alaska. *The Veliger* 27: 440–448.
- Lin G (1992) Three new species of Opisthobranchia from Hong Kong. In: Morton B (Ed.) *The Marine Flora and Fauna of Hong Kong and southern China III. Proceedings of the Fourth International Marine Biological Workshop: The Marine Flora and Fauna of Hong Kong and Southern China*, Hong Kong, 11–29 April 1989, vol. III. Hong Kong University Press, 181–186.
- Londoño-Cruz E (2021) The contribution of Heterobranchia (Mollusca: Gastropoda) to the biodiversity of the Colombian Tropical Eastern Pacific. *Marine Biodiversity* 51(6): 93. <https://doi.org/10.1007/s12526-021-01230-8>
- MacFarland FM (1966) Studies of opisthobranchiate mollusks of the Pacific coast of North America. *Memoirs of the California Academy of Sciences* 6: 1–72.

- Marcus E (1958) On Western Atlantic opisthobranchiate gastropods. American Museum Novitates 1906: 1–82. <https://digilibRARY.amnh.org/handle/2246/5128>
- Marcus E (1959) Lamellariacea und Opisthobranchia. Reports from the Lund University Chile expedition 1948–1949, No. 36. Vol. 55(9): 1–133. Lunds Universitets Arsskrift (Ny Foljd).
- Marcus E, Marcus E (1969) Opisthobranchian and lamellarian gastropods collected by "Vema". American Museum Novitates 2368: 1–31.
- Martynov A, Brenzinger B, Hooker Y, Schrödl M (2011) 3D-anatomy of a new tropical Peruvian nudibranch gastropod species, *Corambe mancorensis*, and novel hypotheses on dorid gill ontogeny and evolution. The Journal of Molluscan Studies 77(2): 129–141. <https://doi.org/10.1093/mollus/eyq047>
- Meyers-Muñoz MA, Roubos EW, van der Velde G (2009) Morphology and distribution of *Flabellina falklandica* (Eliot, 1907) (Nudibranchia, Aeolidina) from the Chilean coast. Basteria 73(4–6): 123–132.
- Millen S, Hermosillo A (2007) The genus *Flabellina* Voight, 1834 (Mollusca: Opisthobranchia) from Bahía de Banderas (Pacific Coast of Mexico with ecological observations, the description of a new species and the redescription of *Flabellina cynara*. Proceedings of the California Academy of Sciences 58: 543–559.
- Millen S, Schrödl M, Vargas N, Indacochea A (1994) A new species of Okenia (Nudibranchia: Doridacea) from the Peruvian Faunal Province. The Veliger 37(3): 312–318.
- Miller SE (1993) Larval period and its influence on post-larval life history: Comparison of lecithotrophy and facultative planktotrophy in the aeolid nudibranch *Phestilla sibogae*. Marine Biology 117(4): 635–645. <https://doi.org/10.1007/BF00349776>
- Molina V (2021) *Coryphellina cerverai*. iNaturalist. <https://www.inaturalist.org/observations/92505003>
- Nakamura K (2006) New records of opisthobranch mollusks from the Guayaquil marine ecoregion: Northern Peru. The Festivus 38(7): 75–83. <https://www.biodiversitylibrary.org/page/50520051#page/231/mode/1up>
- Nakamura K (2007) Especies bentónicas de Opisthobranchia (Mollusca: Gastropoda) presentes en el litoral del norte peruano. Revista Peruana de Biología 13(3): 255–257. <https://doi.org/10.15381/rpb.v13i3.2352>
- Nimbs MJ, Smith SDA (2018) Beyond Capricornia: Tropical Sea Slugs (Gastropoda, Heterobranchia) Extend Their Distributions into the Tasman Sea. Diversity (Basel) 10(3): 99. <https://doi.org/10.3390/d10030099>
- Nybakk J (1978) Abundance, diversity and temporal variability in a California intertidal nudibranch assemblage. Marine Biology 45(2): 129–146. <https://doi.org/10.1007/BF00390549>
- Odhner NH (1922) Mollusca from Juan Fernandez and Easter Island. In: Skottsberg C (Ed.) The Natural History of Juan Fernández and Easter Island. Almqvist y Wiksell Boktryckeri, Uppsala, 219–254. <https://www.biodiversitylibrary.org/page/28680849>
- Odhner NH (1926) Die Opisthobranchien. Further Zoological Results of the Swedish Antarctic Expedition 1901–1903 under the direction of Dr. Otto Nordenskjöld. <https://www.marinespecies.org/aphia.php?p=sourcedetails&id=285344>
- Odhner NH (1944) Mollusca: Nudibranchia and Scaphopoda with zoogeographical remarks and explanations. Det Norske Videnskaps Akademi i Oslo. Scientific Results of the Norwegian Antarctic Expeditions 1927–1928(2): 1–48.
- Oliveros JC (2016) Venny 2.1. An Interactive Tool for Comparing Lists with Venn's Diagrams. (2007–2015). <https://bioinfogp.cnb.csic.es/tools/venny/>
- Padula V (2015) Testing traditional concepts: Biodiversity and integrative taxonomy of Brazilian opisthobranchs (Mollusca, Heterobranchia). Dissertation, LMU München.

- Padula V, Bahia J, Vargas C, Lindner A (2011) Mollusca, Nudibranchia: New records and southward range extensions in Santa Catarina, southern Brazil. Check List 7(6): 806–808. <https://doi.org/10.15560/11037>
- Paredes C, Tarazona J, Canahuire E, Romero L, Cornejo O, Cardoso F (1998) Presencia de moluscos tropicales de la provincia panameña en la costa central del Perú y su relación con los eventos «El Niño». Revista Peruana de Biología 5(2): 123–128. <https://doi.org/10.15381/rpb.v5i2.8330>
- Paredes C, Huamán P, Cardoso F, Vivar R, Vera V (1999) Estado actual del conocimiento de los moluscos acuáticos en el Perú. Revista Peruana de Biología 6(1): 005–047. <https://doi.org/10.15381/rpb.v6i1.8298>
- Pereira FR, Berlinck RG, Rodrigues Filho E, Veloso K, Ferreira AG, Padula V (2012) Metabólitos secundários dos nudibrânquios *Tambja stegosauriformis*, *Hypselodoris lajensis* e *Okenia zoobotryon* e dos briozoários *Zoobotryon verticillatum* e *Bugula dentata* da costa do Brasil. Quimica Nova 35(11): 2194–2201. <https://doi.org/10.1590/S0100-40422012001100018>
- Pievani T (2014) The sixth mass extinction: Anthropocene and the human impact on biodiversity. Rendiconti Lincei. Scienze Fisiche e Naturali 25(1): 85–93. <https://doi.org/10.1007/s12210-013-0258-9>
- Pilsbry HA, Vanatta EG (1902) Papers from the Hopkins Stanford Galapagos Expedition, 1898–1899. XIII. Marine Mollusca. Proceedings of the Washington Academy of Sciences 4: 549–560. <https://doi.org/10.5962/bhl.part.18574>
- Pola M, Sánchez-Benítez M, Ramiro B (2014) The genus *Polycera* Cuvier, 1817 (Nudibranchia: Polyceridae) in the eastern Pacific Ocean, with redescription of *Polycera alabe* Collier & Farmer, 1964 and description of a new species. The Journal of Molluscan Studies 80(5): 551–561. <https://doi.org/10.1093/mollus/eyu049>
- Powell AWB (1979) New Zealand Mollusca: Marine, land, and freshwater shells. Collins, Auckland, 500 pp.
- Pruvot-Fol A (1954) Mollusques – Opisthobranches. Faune de France 58: 1–460. [pl. 1.] <http://faunedefrance.org/bibliotheque/docs/A.PRUVOT-FOL%28FdeFr58%29Mollusques.pdf>
- QGIS Development Team (2022) QGIS Geographic Information System. QGIS Association. <https://www.qgis.org>
- Quesquen R (2008) Nudibranchios holoplancónicos (Familia Phylliroidae Ferussac, 1821) en el mar peruano. Libro de Resúmenes, 280.
- Quesquen R (2017) Moluscos gasterópodos holoplancónicos en el mar peruano recolectados entre 1994 y 2005. Boletín Instituto del Mar del Perú 32(2): 260–308. <https://repositorio.imarpe.gob.pe/handle/20.500.12958/3204>
- Quesquen R, Guzmán S (1999) Nuevo registro de *Phylliroe bucephala* (Mollusca, Gastropoda: Nudibranchia) para el mar peruano In: Tresierra A, Culquichicon Z (Eds) VIII Congreso Latinoamericano sobre Ciencias del Mar, Trujillo, 374–375.
- Ralph P (1959) Notes on an abnormality in the liver caeca of the nudibranch *Phylliroe lichtensteini* Eschscholtz, 1825. Proceedings of the Malacological Society of London 33(5): 186–192. <https://doi.org/10.1093/oxfordjournals.mollus.a064821>
- Ramírez R, Paredes C, Arenas J (2003) Moluscos del Perú. Revista de Biología Tropical 51(Suppl. 3): 225–284.
- Roche A, Cetra N, Núñez V, Darrigran G (2023) Catálogo de las especies de Nudibranchia (Gastropoda: Heterobranchia) de la colección de Moluscos del Museo de La Plata, Argentina. Acta Zoológica Lilloana 67(1): 159–175. <https://doi.org/10.30550/j.azl/2023.67.1/2023-04-13>

- Rudman WB (1983) The Chromodorididae (Opisthobranchia: Mollusca) of the Indo-West Pacific: *Chromodoris splendida*, *C. aspersa* and *Hypselodoris placida* colour groups. *Zoological Journal of the Linnean Society* 78(2): 105–173. <https://doi.org/10.1111/j.1096-3642.1983.tb00864.x>
- Rudman WB (2002a) Doris fontainei d'Orbigny, 1837. Sea Slug Forum. <http://www.seaslugforum.net/find/dorifont>
- Rudman WB (2002b) Concentration of dorids in Peru by Jorge Luque. Sea Slug Forum. <http://www.seaslugforum.net/find/8071>
- Sachidhanandam U, Willan RC, Chou LM (2000) Checklist of the nudibranchs (Opisthobranchia: Nudibranchia) of the South China sea. *The Raffles Bulletin of Zoology* 7: 513–537.
- Santander M, Valdés Á (2013) Genetic structure of *Polycera alabe* and *P. atra* (Mollusca: Opisthobranchia: Nudibranchia) in the Pacific coast of North America. *Bulletin of the Southern California Academy of Sciences* 112(3): 176–184. <https://doi.org/10.3160/0038-3872-112.3.176>
- Schmekel L, Portmann A (1982) Opisthobranchia des Mittelmeeres: Nudibranchia u. sacoglossa. Springer-Verlag, Berlin, 410 pp. <https://doi.org/10.1007/978-3-642-61817-8>
- Schrödl M (1996a) Nudibranchia y Sacoglossa de Chile: Morfología externa y distribución. *Gayana. Zoología* 60(1): 17–62.
- Schrödl M (1996b) *Janolus rebeccaee*, a new species of arminacean nudibranchs from northern Chile. *Spixiana* 19: 293–300.
- Schrödl M (1997) Range extensions of Magellanic Nudibranchs (Opisthobranchia) into the Peruvian faunal province. *The Veliger* 40(1): 38–42.
- Schrödl M (1999a) *Hancockia schoeferti* spec. nov., a new dendronotoidean nudibranch species from central Chile (Gastropoda, Nudibranchia, Hancockiidae). *Spixiana* 22: 247–254.
- Schrödl M (1999b) Zoogeographic relationships of Magellan Nudibranchia (Mollusca: Opisthobranchia) with particular reference to species from adjacent regions. *Scientia Marina* 63(S1): 409–416. <https://doi.org/10.3989/scimar.1999.63s1409>
- Schrödl M (2000) Revision of the nudibranch genus *Cadlina* (Gastropoda: Opisthobranchia) from the Southern Ocean. *Journal of the Marine Biological Association of the United Kingdom* 80(2): 299–309. <https://doi.org/10.1017/S0025315499001873>
- Schrödl M (2002) Peruvian dorid. <http://www.seaslugforum.net/message/8283>
- Schrödl M (2003) Sea Slugs of Southern South America. ConchBooks, Hackenheim, 165 pp.
- Schrödl M (2009) Opisthobranchia – Babosas de Mar. In: V Häussermann, G Förster-ra (Eds) Fauna Marina Bentónica de la Patagonia Chilena. Nature in Focus, Chile, 505–542.
- Schrödl M, Grau JH (2006) Nudibranchia from the remote southern Chilean Guamblin and Ipún islands (Chonos Archipelago, 44–45°S), with re-description of *Rostanga pulchra* MacFarland, 1905. *Revista Chilena de Historia Natural* 79(1): 3–12. <https://doi.org/10.4067/S0716-078X2006000100001>
- Schrödl M, Hooker Y (2014) Sea slugs of Peru: Peruvian-Chilean faunal elements. *Spixiana* 37(1): 45–59.
- Schrödl M, Millen SV (2001) Revision of the nudibranch gastropod genus *Tyrinna* Bergh, 1898 (Doridoidea: Chromodorididae). *Journal of Natural History* 35(8): 1143–1171. <https://doi.org/10.1080/00222930152434472>
- Schrödl M, Wägele M (2001) Anatomy and histology of *Corambe lucea* Marcus, 1959 (Gastropoda, Nudibranchia, Doridoidea), with a discussion of the systematic po-

- sition of Corambidae. *Organisms, Diversity & Evolution* 1(1): 3–16. <https://doi.org/10.1078/1439-6092-00002>
- Schrödl M, Alarcón MA, Bedriñana LR, Bravo FJ, Bustamante CM, Carvalho R, Försterra G, Gallardo C, Häussermann V, Salmen A (2005) Nudipleura (Gastropoda: Opisthobranchia) from the southern Chilean Comau Fjord, with redescription of *Polycera priva* Marcus, 1959. *Vita Malacologica* 3: 23–33.
- Schrödl M, Jörger KM, Klussmann-Kolb A, Wilson NG (2011) Bye bye 'Opisthobranchia'? A review on the contribution of Mesopsammic sea slugs to euhyneuran systematics. *Thalassas* 27(2): 101–112.
- Segovia J, López G (2015) Registro de *Glaucus atlanticus* en la costa de El Salvador, Pacífico de Centroamérica. *Revista Mexicana de Biodiversidad* 86(4): 1089–1090. <https://doi.org/10.1016/j.rmb.2015.05.015>
- Shields C (2009) Nudibranchs of the Ross Sea, Antarctica: Phylogeny, diversity, and divergence. MSc Thesis, Clemson University, SC, USA. https://tigerprints.clemson.edu/all_theses/637/
- Spalding MD, Fox HE, Allen GR, Davidson N, Ferdaña ZA, Finlayson M, Halpern BS, Jorge MA, Lombana A, Lourie SA, Martin KD, McManus E, Molnar J, Recchia CA, Robertson J (2007) Marine Ecoregions of the World: A Bioregionalization of Coastal and Shelf Areas. *Bioscience* 57(7): 573–583. <https://doi.org/10.1641/B570707>
- Spencer HG, Willan RC (1995) The marine fauna of New Zealand: Index to the fauna. 3: Mollusca. New Zealand Oceanographic Institute. New Zealand Oceanographic Institute, Wellington, 125 pp.
- Spencer HG, Marshall BA, Willan RC (2009) Checklist of New Zealand living Mollusca. Pp 196–219. In: Gordon DP (Ed.) New Zealand inventory of biodiversity, vol. 1. Kingdom Animalia: Radiata, Lophotrochozoa, Deuterostomia. Canterbury University Press, Christchurch, 568 pp. <https://www.marbef.org/data/aphia.php?p=sourcedetails&id=132606>
- Steinberg JE (1956) The pelagic nudibranch, *Cephalopyge trematoides* (Chun, 1889), in New South Wales with a note on other species in this genus. *Proceedings of the Linnean Society of New South Wales* 81: 184–192.
- Stout CC, Pola M, Valdés Á (2010) Phylogenetic analysis of *Dendronotus* nudibranchs with emphasis on northeastern Pacific species. *The Journal of Molluscan Studies* 76(4): 367–375. <https://doi.org/10.1093/mollus/eyq022>
- Tejada-Pérez CA, Villasante F, Luque-Fernández C, Tejada-Begazo CL (2018) Riqueza y distribución vertical de los moluscos del litoral rocoso de la provincia de Islay, Arequipa, al sur del Perú. *Revista de Ciencias Marinas y Costeras* 10(1): 47. <https://doi.org/10.15359/revmar10-1.4>
- Thompson TE, McFarlane ID (1967) Observations on a collection of *Glaucus* from the Gulf of Aden with a critical review of published records of Glaucidae (Gastropoda, Opisthobranchia). *Proceedings of the Linnean Society of London* 178(2): 107–123. <https://doi.org/10.1111/j.1095-8312.1967.tb00967.x>
- Thompson TE, Cattaneo R, Wong YM (1990) Eastern Mediterranean Opisthobranchia: Dotidae (Dendronotacea), Arminidae and Madrellidae (Arminacea). *The Journal of Molluscan Studies* 56(3): 393–413. <https://doi.org/10.1093/mollus/56.3.393>
- Thompson D (2006) *Hypselodoris ghiselini* – new record from Peru. <http://www.seaslug-forum.net/message/15852>
- Tokioka T (1963) Notes on *Cephalopyge trematoides* (Chun) collected off the central and northern Chilean coast (Nudibranchia: Phyllirhoiidae). *Publications of the Seto Marine Biological Laboratory* 11(2): 373–387. <https://doi.org/10.5134/175337>

- Torrejón R (2023) Marcus's Aeolid (*Coryphellina marcusorum*). iNaturalist. <https://www.inaturalist.org/observations/150823752>
- Trickey JS (2013) Phylogeography and molecular systematics of the rafting aeolid nudibranch *Fiona pinnata* (Eschscholtz, 1831). Thesis, Master of Science. University of Otago. <http://hdl.handle.net/10523/3822>
- Uribe RA, Pacheco AS (2012) First record of *Spurilla neapolitana* (Mollusca: Nudibranchia: Aeolidiidae) on the central coast of Peru (Humboldt Current Upwelling Ecosystem). Marine Biodiversity Records 5: e14. <https://doi.org/10.1017/S1755267211001138>
- Uribe RA, Nakamura K, Indacochea A, Pacheco AS, Hooker Y, Schrödl M (2013) A review on the diversity and distribution of opisthobranch gastropods from Peru, with the addition of three new records. *Spixiana* 36(1): 43–60.
- Uribe RA, Perea Á, García V, Huerto M (2019) Biodiversidad marina en el norcentro de la costa de Perú: Un enfoque para la evaluación de planes de manejo. Boletín Instituto del Mar del Perú 34(2): 20. <https://repositorio.imarpe.gob.pe/handle/20.500.12958/3401>
- Valdés Á (2001a) Deep-sea cryptobranch dorid nudibranchs (Mollusca, Opisthobranchia) from the tropical West Pacific, with descriptions of two new genera and eighteen new species. *Malacologia* 43: 237–311.
- Valdés Á (2001b) Depth-related adaptations, speciation processes and evolution of color in the genus *Phyllidiopsis* (Mollusca: Nudibranchia). *Marine Biology* 139(3): 485–496. <https://doi.org/10.1007/s002270100596>
- Valdés Á (2002) A phylogenetic analysis and systematic revision of the cryptobranch dorids (Mollusca, Nudibranchia, Anthobranchia). *Zoological Journal of the Linnean Society* 136(4): 535–636. <https://doi.org/10.1046/j.1096-3642.2002.00039.x>
- Valdés Á, Campillo OA (2004) Systematics of pelagic aeolid nudibranchs of the family Glaucidae (Mollusca, Gastropoda). *Bulletin of Marine Science* 75(3): 381–389. <http://www.ingentaconnect.com/content/umrsmas/bullmar/2004/00000075/00000003/art00003>
- Valdés Á, Héros V (1998) The types of recent and certain fossil opisthobranch molluscs in the Muséum national d'Histoire naturelle, Paris. *Zoosystema* 1998(4): 695–742.
- Valdés Á, Muniaín C (2002) Revision and taxonomic reassessment of Magellanic species assigned to *Anisodoris* Bergh, 1898 (Nudibranchia: Doridoidea). *The Journal of Molluscan Studies* 68(4): 345–351. <https://doi.org/10.1093/mollus/68.4.345>
- Valdés Á, Hamann J, Behrens D, DuPont A (2006) Caribbean sea slugs: A field guide to the opisthobranch mollusks from the tropical northwestern Atlantic. Sea Challengers Natural History Books, Gig Harbor, 289 pp. <https://www.nhbs.com/caribbean-sea-slugs-book>
- Valdivia-Chavez D, Santamaría J, Campos-León S, Carbajal P, Barriga E, Clemente S, Gálecio Z, Kameya A (2020) Lista taxonómica de invertebrados bentónicos marinos de Arequipa, Perú. Boletín Instituto del Mar del Perú 35(2): 378–395. <https://revistas.imarpe.gob.pe/index.php/boletin/article/view/311>
- Valqui J, Ibañez-Erquiaga B, Pacheco AS, Wilbur L, Ochoa D, Cardich J, Pérez-Huaranga M, Salas-Gismondi R, Pérez A, Indacochea A, Avila-Peltroche J, Rivera M, Carré M (2021) Changes in rocky intertidal communities after the 2015 and 2017 El Niño events along the Peruvian coast. *Estuarine, Coastal and Shelf Science* 250: 107142. <https://doi.org/10.1016/j.ecss.2020.107142>
- van der Spoel S (1970) The pelagic Mollusca from the "Atlantide" and "Galathea" expeditions collected in the East Atlantic. *Atlantide Report II*: 99–139.
- van der Spoel S, Newman LJ, Estep KW (1997) Pelagic Molluscs of the World. Expert-Center for Taxonomic Identification, Amsterdam. World Biodiversity Database. CD-ROM Series.

- Velasco-Charpentier C, Pizarro-Mora F, Navarro NP, Valdivia N (2021) Disentangling the links between habitat complexity and biodiversity in a kelp-dominated subantarctic community. *Ecology and Evolution* 11(3): 1214–1224. <https://doi.org/10.1002/ece3.7100>
- Velez J, Zeballos J (1985) Ampliación de la distribución de algunos peces e invertebrados durante el fenómeno «El Niño» 1982–1983. In: Arntz W, Landa A, Tarazona J (Eds) “El Niño”: Su impacto en la fauna marina. Boletín Extraordinario IMARPE, Volumen Extraordinario, 173–180. <https://repositorio.imarpe.gob.pe/handle/20.500.12958/1206>
- Wägele H, Klussmann-Kolb A (2005) Opisthobranchia (Mollusca, Gastropoda) – More than just slimy slugs. Shell reduction and its implications on defence and foraging. *Frontiers in Zoology* 2(1): 3. <https://doi.org/10.1186/1742-9994-2-3>
- Wägele H, Klussmann-Kolb A, Verbeek E, Schrödl M (2014) Flashback and foreshadowing – A review of the taxon Opisthobranchia. *Organisms, Diversity & Evolution* 14(1): 133–149. <https://doi.org/10.1007/s13127-013-0151-5>
- Welch JJ (2010) The “Island Rule” and Deep-Sea Gastropods: Re-Examining the Evidence. *PLoS ONE* 5(1): e8776. <https://doi.org/10.1371/journal.pone.0008776>
- Willan R (1979) New Zealand locality records for the aeolid nudibranch *Fiona pinnata* (Eschscholtz). *Tane* 25: 141–147.
- Willan R (2006) Globe-trotting nudibranch arrives in Australia. *Australasian shell news*, 130(1): 1–3.
- Zagal C, Hermosilla C (2001) Guía de invertebrados marinos del litoral Valdiviano. Querbecor World Chile, Santiago de Chile, 218 pp.
- Zagal C, Hermosilla C (2007) Guía de invertebrados marinos del Sur de Chile. Fantástico Sur, Punta Arenas, Chile, 264 pp.
- Zanabria U (2020a) *Diaulula variolata*. iNaturalist. <https://www.inaturalist.org/observations/60520768>
- Zanabria U (2020b) *Diaulula variolata*. iNaturalist. <https://www.inaturalist.org/observations/60527433>
- Zanabria U (2020c) *Diaulula variolata*. iNaturalist. <https://www.inaturalist.org/observations/60528808>
- Zanabria U (2020d) Flameback (*Phidiana lottini*). iNaturalist. <https://www.inaturalist.org/observations/60649340>
- Zanabria U (2020e) Flameback (*Phidiana lottini*). iNaturalist. <https://www.inaturalist.org/observations/60555449>
- Zavala A (2022) *Felimare sechurana*. iNaturalist. <https://www.inaturalist.org/observations/139130309>
- Zhukova NV, Eliseikina MG, Balakirev ES, Ayala FJ (2022) Multiple bacterial partners in symbiosis with the nudibranch mollusk *Rostanga alisae*. *Scientific Reports* 12(1): 169. <https://doi.org/10.1371/journal.pone.0008776>