




Notes on Dysderidae (Arachnida, Araneae) of Armenia and Iran, with new species and records

Armine Kosyan¹, Alireza Zamani², Yuri M. Marusik^{3,4}

¹ Faculty of Biology, Yerevan State University, Yerevan, Armenia

² Zoological Museum, Biodiversity Unit, FI-20014 University of Turku, Finland

³ Department of Zoology & Entomology, University of the Free State, Bloemfontein 9300, South Africa

⁴ Altai State University, Lenina Pr., 61, Barnaul, RF-656049, Russia

Corresponding author: Alireza Zamani (zamani.alireza5@gmail.com)

Abstract

New data are provided on dysderid spiders of Armenia and Iran. *Dysdera haykana* sp. nov. is described based on male specimens collected in Kotayk and Lor provinces, central and northern Armenia. The female of *Dysdera nakhchivanica* Beydizade, Shafaie & Guseinov, 2018 is described for the first time, and the species is newly recorded from Armenia. Furthermore, new distribution data are provided for *Harpactea armenica* Dunin, 1989, including the first record of the species from Iran. Photographs and a distribution map are provided for all three species.

Key words: Aranei, Caucasus, red devil spiders, woodlouse spiders



Academic editor: Shuqiang Li

Received: 28 May 2023

Accepted: 27 June 2023

Published: 25 July 2023

ZooBank: <https://zoobank.org/C8B5DE1C-0482-491D-A61C-4BFB55C93A13>

Citation: Kosyan A, Zamani A, Marusik YM (2023) Notes on Dysderidae (Arachnida, Araneae) of Armenia and Iran, with new species and records. ZooKeys 1172: 117–130. <https://doi.org/10.3897/zookeys.1172.107112>

Copyright: © Armine Kosyan et al.

This is an open access article distributed under terms of the Creative Commons Attribution License ([Attribution 4.0 International – CC BY 4.0](https://creativecommons.org/licenses/by/4.0/)).

Introduction

The spider family Dysderidae C.L. Koch, 1837 currently comprises 612 extant species in 25 genera distributed in the Western Palaearctic (WSC 2023). The dysderid fauna of the Caucasus is relatively well studied thanks to the publication of several large-scale, family-level revisions and smaller taxonomic contributions (e.g., Charitonov 1956; Dunin 1982, 1987, 1990, 1991, 1992). However, the Armenian dysderids, with only 11 currently known species (Otto 2022), remain poorly documented, especially when compared to adjacent countries of Azerbaijan (30 species; Otto 2022), Georgia (33 species; Otto 2022), Iran (15 species; Zamani et al. 2023), and Turkey (69 species; Danişman et al. 2023). This paper aims to contribute to the knowledge of dysderid diversity in this region by providing new taxonomic and faunistic data on three species occurring in Armenia and Iran.

Materials and methods

Photographs of specimens and their copulatory organs were obtained using an Olympus Camedia E-520 camera attached to an Olympus SZX16 stereomicroscope, and a JEOL JSM-5200 scanning electron microscope. Digital images of different focal planes were stacked with Helicon Focus™ v. 8.1.1. Illustrations of internal genitalia were made after digesting tissues off in a 10% KOH aqueous solution.

Body measurements exclude the chelicerae and spinnerets. Leg segments were measured on the dorsal side. Measurements and characters of the palp used in the diagnoses are based on the retrolateral view. Measurements of legs are listed as: total length (femur, patella, tibia, metatarsus, tarsus). All measurements are given in millimetres. The map was prepared using SimpleMappr (Shorthouse 2010).

Abbreviations: Eyes: AME – anterior median eye, PLE – posterior lateral eye, PME – posterior median eye. **Spination:** d – dorsal, Fe – femur, Mt – metatarsus, Pa – patella, pl – prolateral, rl – retrolateral, Ti – tibia, v – ventral.

Depositories: MHNG – Muséum d'histoire naturelle, Genève, Switzerland (P.J. Schwendinger, L. Monod); ZMMU – Zoological Museum of the Moscow State University, Russia (K.G. Mikhailov).

Taxonomy

Family Dysderidae C.L. Koch, 1837

Subfamily Dysderinae C.L. Koch, 1837

Genus *Dysdera* Latreille, 1804

***Dysdera haykana* sp. nov.**

<https://zoobank.org/F47E0340-1B68-488A-8B19-4291FADF47F7>

Figs 1A, 2A–D, 3A, B

Type materials. Holotype ♂ (ZMMU), ARMENIA: Lori Province: Mets Parni, 40°49'15.4"N, 44°04'33.2"E, 30.04.2022 (A. Kosyan). **Paratypes:** 1♂ (ZMMU), same data as for the holotype; 1♂ (ZMMU), Kotayk Province: env. of Solak Vill., 40°28'34"N, 44°42'57"E, 14.05.2021 (Y.M. Marusik).

Etymology. The specific epithet is an adjective referring to Hayk Nahapet (in Armenian: Ռայկ), the legendary patriarch and founder of the Armenian nation.

Diagnosis. The new species belongs to the *asiatica* species-group and is most similar to *D. asiatica* Nosek, 1905 (see Deeleman-Reinhold and Deeleman 1988: figs 226, 227) and *D. ukrainensis* Charitonov, 1956 (as illustrated by Charitonov 1956: fig. 16) in possessing a spine-like median process (*Mp*) (vs either lacking or of a different shape (with 2 or more teeth) in other species). *Dysdera haykana* sp. nov. can be distinguished from both species by the almost indistinct spine-like outgrowth (*So*) on the median crest (vs distinct). From *D. ukrainensis*, it also differs by having median crest (*Mc*) ca 2.5 times longer than high (vs as long as high), the median process extending the median crest (vs not extending), and the relatively larger posterior apophysis (*Pa*) (cf. Fig. 2A and Charitonov 1956: fig. 16).

Description. Male (Holotype). Habitus as in Fig. 1A. Total length 9.22. Carapace 4.16 long, 3.33 wide. Eye sizes: AME 0.22, PME 0.16, PLE 0.19. Carapace and chelicerae dark reddish; carapace slightly lighter posteriorly. Sternum, labium, and maxillae reddish. Legs yellowish orange. Abdomen light beige, without any pattern. Spinnerets uniformly beige. Measurements of legs: I: 11.00 (3.09, 2.11, 2.59, 2.49, 0.72), II: 10.91 (3.13, 1.90, 2.52, 2.69, 0.67), III: 8.23 (2.48, 1.34, 1.52, 2.26, 0.63), IV: 10.85 (3.16, 1.80, 2.23, 2.91, 0.75). Spination: I, II: no spines; III: Ti: 2pl, 1rl, 2v; Mt: 3pl, 3rl, 6v. IV: Fe: 1d; Ti: 2pl, 2rl, 2v; Mt: 2d, 3pl, 3rl, 4v.

Palp as in Figs 2A–D, 3A, B; bulb 3 times longer than its maximal width; psempolus 1.5 times longer than tegulum; tegulum 1.27 times longer than wide; median crest (*Mc*) elongate, ca 2.5 times longer than high, with almost




Figure 1. Habitus of the holotype male of *Dysdera haykana* sp. nov. (A) and the female of *D. nakhchivanica* (B), dorsal.

indistinct spine-like outgrowth (So), middle part of psembolus with spine-like process (Mp), its tip extending dorsal margin of median crest.

Female. Unknown.

Note. The record of *D. ukrainensis* in central Georgia (cf. Otto 2022) may belong to this species. Kovblyuk et al. (2008) redescribed the types of *D. ukrainensis*, although their figures of the lectotype male differs from those provided in Charitonov (1956) in the shape of the median process. Moreover, according to Kovblyuk et al. (2008), the median process can vary in shape and may be bifurcated on the tip.

Distribution. Known only from the listed localities in Kotayk and Lori provinces, central and northern Armenia (Fig. 10).




Figure 2. Male of *Dysdera haykana* sp. nov., bulb. **A** retrolateral **B** prolateral **C** dorsal **D** ventral. Abbreviation: *Mc* – median crest, *Mp* – median process, *Pa* – posterior apophysis, *So* – spine-like outgrowth. Scale bar: 0.2 mm.




Figure 3. Male of *Dysdera haykana* sp. nov., bulb. **A** prolateral **B** dorso-prolateral. Scale bar: 0.1 mm.

***Dysdera nakhchivanica* Beydizade, Shafaie & Guseinov, 2018**

Figs 1B, 4A–D, 5A, B, 6A–D, 7

Dysdera nakhchivanica Beydizade et al., 2018: 1112, figs 1–4, 7–9 (♂).

Material examined. ARMENIA: Vayots Dzor Province: 1♂ 1♀ 2 juv. (ZMMU), env. of Gnishik Vill., 39°40'18"N, 45°17'40"E, ca 2030 m, 11.05.2021 (Y.M. Marusik); 3♂ 3♀ 6 juv. (ZMMU), Shatin, 39°50'N, 45°19'E, 9.05.2021 (Y.M. Marusik).

Diagnosis. For the male, see Beydizade et al. (2018). The female of *D. nakhchivanica* is most similar to that of *D. collucata* Dunin, 1991, a species restricted




Figure 4. Male of *Dysdera nakhchivanica*, bulb. **A** retrolateral **B** prolateral **C** dorsal **D** ventral. Scale bar: 0.2 mm.




Figure 5. Male of *Dysdera nakhchivanica*, bulb. **A** dorsal **B** retrolateral. Scale bar: 0.1 mm.

to southernmost Armenia. The two species have tips of receptacular lateral edges (*Rl*) bent posteriorly (vs not bent or directed anteriorly in other species) and have membranous posterior diverticula (*Pd*) unknown in other *Dysdera* species in the Caucasus or adjacent regions. The female of *D. nakhchivanica* differs from that of *D. collucata* by the relatively longer (i.e., three times wider than long) and distinctly trapezoidal dorsal arch (*Da*).

Description. Male. See Beydizade et al. (2018).




Figure 6. Males of *Dysdera nakhchivanica*, bulbs of four specimens in retrolateral view, showing variations. Abbreviation: *Mp* – median process. Scale bar: 0.2 mm.




Figure 7. Female of *Dysdera nakhchivanica*, endogyne, dorsal. Abbreviations: *Da* – dorsal arch, *Pd* – posterior diverticulum, *Re* – receptacle, *Rl* – receptacular lateral edge, *Tb* – transverse bar. Scale bar: 0.2 mm.

Female. Habitus as in Fig. 1B. Total length 11.87. Carapace 4.12 long, 3.10 wide. Eye sizes: AME 0.16, PME 0.15, PLE 0.17. Carapace and chelicerae dark reddish; carapace slightly lighter posteriorly. Sternum, labium, and maxillae reddish. Legs yellowish orange. Abdomen light beige, without any pattern. Spinnerets uniformly beige. Measurements of legs: I: 12.10 (3.41, 2.20, 3.00, 2.90, 0.59), II: 10.90 (3.06, 1.93, 2.60, 2.65, 0.66), III: 7.87 (2.27, 1.25, 1.67, 2.05, 0.63), IV: 11.05 (3.24, 1.65, 2.43, 2.96, 0.77). Spination: I, II: no spines. III: Ti: 2pl, 2rl, 2v; Mt: 3pl, 2rl, 6v. IV: Fe: 3d; Ti: 2rl, 2v; Mt: 3pl, 3rl, 6v.

Endogyne as in Fig. 7; receptacle (*Re*) with posteriorly bent lateral edges (*Rl*), ca 8 times longer than wide and as wide as transverse bar (*Tb*); dorsal arch (*Da*) ca 2.5 times wider than long, with rounded anterior edges; transverse bar (*Tb*) long and thin, ca 15 times longer than wide; posteriorly with pair of diverticula (*Pd*).

Note. The holotype of *D. nakhchivanica* has four teeth on the median process (*Mp*), while Armenian specimens have 5–7. We have tentatively considered this as an intraspecific variation. Certain variations in the male palp, some of which depend on the photography angle, are illustrated in Figs 4A and 6A–D.

Distribution. Azerbaijan (Nakhchivan) and Armenia (new record) (Fig. 10).

Subfamily Harpacteinae Cooke, 1965

Genus *Harpactea* Bristowe, 1939

***Harpactea armenica* Dunin, 1989**

Figs 8A, B, 9A–D

Harpactea armenica Dunin, 1989: 142, figs 1–3 (♂♀).

Harpactea armenica: Dunin 1992: 68, fig. 13 (♂♀).

Materials examined. ARMENIA: Aragatsotn Province: 24♂ 34♀ 29 juv. (ZMMU), foothills of Aragates Mt., 40°23'N, 44°13'E, 2200 m, 16.05.2021 (Y.M).




Figure 8. Habitus of *Harpactea armenica*, male (A) and female (B), dorsal.




Figure 9. Copulatory organs of *Harpactea armenica* **A–C** bulb, prolateral, retrolateral and ventral **D** endogyne, dorsal. Abbreviations: Ad – anterior diverticulum, Eb – base of embolus, Re – receptacle. Scale bars: 0.2 mm.




Figure 10. Distribution records of *Dysdera haykana* sp. nov. (circles), *D. nakhchivanica* (squares), and *Harpactea armenica* (triangles). Type localities are marked with red.

Marusik); *Ararat Province*: 1♂ (ZMMU), env. of Urtsadzor Vill., 39°55'26"N, 44°48'53"E, 1040 m, 6.05.2021 (Y.M. Marusik); 1♂ 2♀ (ZMMU), env. of Urtsalanj Vill., 39°49'N, 44°59'E, 1800 m, 11.05.2021 (Y.M. Marusik); 5♂ 1♀ (ZMMU), Garni gorge, Azat river, 40°06'32"N, 44°43'57"E, 1240 m, 17.05.2021 (Y.M. Marusik); *Kotayk Province*: 3♂ 2♀ (ZMMU), env. of Geghadir, 40°09'N, 44°38'E, 15.05.2021 (Y.M. Marusik); 2♂ 4♀ (ZMMU), env. of Solak Vill., 40°28'24"N, 44°42'57"E, 14.05.2021 (Y.M. Marusik); 1♂ 1♀ (ZMMU), env. of Aghveran Vill., 40°29'54"N, 44°35'24"E, 7–8.05.2021 (Y.M. Marusik); 5♂ 7♀ 4 juv. (ZMMU), same locality, 7.05.2021 (Y.M. Marusik); *Vayots Dzor Province*: 21♂14♀ (ZMMU), Shatin, 39°50'N, 45°19'E, 9.05.2021 (Y.M. Marusik); 2♂ 3♀ (ZMMU), Shatin, 39°50'N, 45°19'E, 9.05.2021 (Y.M. Marusik); *Yerevan*: 7♂ 9♀ (ZMMU), Yerevan, botanical garden, 40°12'43"N, 44°33'21"E, 17.05.2021 (Y.M. Marusik). *IRAN*: *East Azerbaijan Province*: 1♂ 3♀ (MHNG), NE of Sofi-an, 38°21'N, 45°50'E, 5.06.1975 (A. Senglet); *Gilan Province*: 1♂ 3♀ (MHNG), Asalem, 37°45'N, 48°57'E, 11.06.1975 (A. Senglet); 1♂ 3♀ (MHNG), Lahidjan, 37°11'N, 49°54'E, 5.07.1973 (A. Senglet); *Mazandaran Province*: 1♂ 3♀ (MHNG), road to Djavaherdeh, 36°55'N, 50°33'E, 1200 m, 7.08.1974 (A. Senglet); *West Azerbaijan Province*: 1♂ 3♀ (MHNG), Qara Kelisa, 39°04'N, 44°38'E, 31.05.1975 (A. Senglet).

Comments. This species is most similar to *H. secunda* Dunin, 1989, a species restricted to northern Armenia, but can be differentiated from it by the base of embolus (*Eb*) (slightly and roundly bent, vs sharply bent to about 70°). It is possible that this small difference is only an intraspecific variation; in order to verify this, it is necessary to collect additional material from northern Armenia or examine the type specimens of *H. secunda*. Furthermore, it is likely that the female illustrated by Dunin (1989, 1992) belongs to another species. All females collected in Armenia and Iran examined here have a longer receptacle, longer than anterior diverticulum (*Ad*) *sensu* Dunin (1992), and lack bulbous thickening of the receptacle (*Re*, Fig. 9D).

Distribution. This species was previously known from only two localities in Armenia: Yerevan and the southernmost part of Kotayk Province. The new records slightly extend the known range of the species to the north, approximately 5° to the east, and around 4° to the south. Additionally, this species is newly recorded from Iran (Fig. 10).

Acknowledgements

We are grateful to Shuqiang Li (the subject editor) and Dragomir Dimitrov, Mikhail Omelko, and an anonymous reviewer for the valuable comments on the manuscript. Furthermore, we thank Peter J. Schwendinger and Lionel Monod for arranging a loan of the material deposited in MHNG. YM is thankful to Noushig Zarijian (Yerevan, Armenia) for providing assistance with arranging his expedition to Armenia. We express our gratitude to Prof. Marine Arakelyan (Yerevan, Armenia) for providing assistance in collecting material.

Additional information

Conflict of interest

The authors have declared that no competing interests exist.

Ethical statement

No ethical statement was reported.

Funding

This work was supported by the Science Committee of the Republic of Armenia within the frames of the research projects 21AG-1F033, 22AA-1F018.

Author contributions

Conceptualization: AZ, YM. Material - collection: AK, YM. Material - sorting and identification: AZ, AK. Funding acquisition: AK. Writing - original draft: AZ, YM. Writing - review and editing: AZ, YM, AK.

Author ORCIDs

Armine Kosyan  <https://orcid.org/0000-0001-5692-5369>

Alireza Zamani  <https://orcid.org/0000-0002-8084-9666>

Yuri M. Marusik  <https://orcid.org/0000-0002-4499-5148>

Data availability

All of the data that support the findings of this study are available in the main text.

References

- Beydizade N, Shafaie S, Guseinov E (2018) A new *Dysdera* species (Araneae: Dysderidae) from Azerbaijan. *Linzer Biologische Beiträge* 50(2): 1111–1118.
- Bristowe WS (1939) The Comity of Spiders. Vol. 1. Ray Society, London, 228 pp.
- Charitonov DE (1956) Obzor paukov semeistva Dysderidae fauny SSSR. *Uchenye Zapiski Molotovskogo Gosudarstvennogo Universiteta Imeni A. M. Gor'kogo* 10: 17–39. [In Russian]
- Cooke JAL (1965) Spider genus *Dysdera* (Araneae, Dysderidae). *Nature* 205(4975): 1027–1028. <https://doi.org/10.1038/2051027b0>
- Deeleman-Reinhold CL, Deeleman PR (1988) Revision des Dysderinae (Araneae, Dysderidae), les espèces méditerranéennes occidentales exceptées. *Tijdschrift voor Entomologie* 131: 141–269.
- Dunin PM (1982) New data on *Dysdera concinna* L. Koch (Aranei, Dysderidae). *Zoologicheskii Zhurnal* 61: 605–607. [In Russian]
- Dunin PM (1987) *Dysdera ghilarovi* sp. n. (Aranei, Dysderidae) from Azerbaijan. *Zoologicheskii Zhurnal* 66: 949–951. [In Russian]
- Dunin PM (1989) New spider species of the *Harpactea* genus from Armenia (Aranei, Dysderidae). *Zoologicheskii Zhurnal* 68(7): 142–145. [In Russian]
- Dunin PM (1990) Spiders of the genus *Dysdera* (Aranei, Haplogynae, Dysderidae) from Azerbaijan. *Zoologicheskii Zhurnal* 69(6): 141–147. [In Russian]
- Dunin PM (1991) New spider species of genus *Dysdera* from the Caucasus (Aranei, Haplogynae, Dysderidae). *Zoologicheskii Zhurnal* 70(8): 90–98. [In Russian]
- Dunin PM (1992) The spider family Dysderidae of the Caucasian fauna (Arachnida Aranei Haplogynae). *Arthropoda Selecta* 1(3): 35–76. [In Russian]
- Koch CL (1837) Übersicht des Arachnidensystems. Heft 1. C. H. Zeh'sche Buchhandlung, Nürnberg, 39 pp [6 pls]. <https://doi.org/10.5962/bhl.title.39561>
- Kovblyuk MM, Prokopenko EV, Nadolny AA (2008) Spider family Dysderidae of the Ukraine (Arachnida, Aranei). *Euroasian Entomological Journal* 7(4): 287–306. <https://doi.org/10.23885/1814-3326-2008-4-1-3-40>
- Latreille PA (1804) Tableau methodique des Insectes. Nouveau Dictionnaire d'Histoire Naturelle, Paris 24: 129–295.
- Otto S (2022) Caucasian Spiders. A faunistic database on the spiders of the Caucasus. Version 02.2022. <https://caucasus-spiders.info/>
- Shorthouse DP (2010) SimpleMappr, an online tool to produce publication-quality point maps. <http://www.simplemappr.net> [Accessed on 23.5.2023]
- WSC (2023) World Spider Catalog. Version 24. Natural History Museum Bern. <http://wsc.nmbe.ch> [Accessed on 23.5.2023]
- Zamani A, Marusik YM, Szűts T (2023) A survey of the spider genus *Dysdera* Latreille, 1804 (Araneae, Dysderidae) in Iran, with fourteen new species and notes on two fossil genera. *ZooKeys* 1146: 43–86. <https://doi.org/10.3897/zookeys.1146.97517>