

A conspectus of Australian *Apotropina* (Diptera, Chloropidae) with the description of two new species

Yuchen Ang¹, James Lumbers^{2,3}, Paula R. Riccardi⁴

¹ Lee Kong Chian Natural History Museum, National University of Singapore, 2 Conservatory Dr., 117377 Singapore, Singapore

² Australian National Insect Collection (ANIC), CSIRO Black Mountain, 1 Clunies Ross St, Acton Black Mountain, Canberra, ACT 2601, Australia

³ Research School of Biology, Australian National University, Canberra, ACT 2601, Australia

⁴ Center for Integrative Biodiversity Discovery, Museum für Naturkunde, Leibniz-Institute for Evolution and Biodiversity Science, Invalidenstr. 43, 10115 Berlin, Germany

Corresponding authors: Yuchen Ang (nhmay@nus.edu.sg); Paula R. Riccardi (paulariccardi@gmail.com)

Abstract

The genus *Apotropina* (Diptera, Chloropidae) has a global distribution with more than 80 valid described species, of which 22 are known to occur in Australia. The Australian *Apotropina* fauna is poorly studied, with many species known from single type specimens, more with the morphology of the other sex unknown, and there have been no new species descriptions since 1959. Here, we describe two new species from Australia, *A. maculigena* Riccardi, **sp. nov.** and *A. popeye* Ang, **sp. nov.**, and provide an updated illustrated key. We also provide a conspectus of the known Australian *Apotropina* with images of types and collate all original descriptions and subsequent taxonomic notes of relevance as supplementary information. Finally, we discuss the validity of two known syntype specimens of *A. bispinosa* due to incongruencies with the species description.



Key words: *Apotropina*, Australia, Chloropidae, Diptera, key, Sexual Dimorphism, taxonomy

Academic editor: Owen Lonsdale

Received: 26 June 2023

Accepted: 6 November 2023

Published: 21 December 2023

ZooBank: <https://zoobank.org/919C320F-AA72-4F1D-9028-1ECB12948B8D>

Citation: Ang Y, Lumbers J, Riccardi PR (2023) A conspectus of Australian *Apotropina* (Diptera, Chloropidae) with the description of two new species. ZooKeys 1187: 261–299. <https://doi.org/10.3897/zookeys.1187.108497>

Copyright: © Yuchen Ang et al.

This is an open access article distributed under terms of the Creative Commons Attribution License ([Attribution 4.0 International – CC BY 4.0](https://creativecommons.org/licenses/by/4.0/)).

Introduction

The genus *Apotropina* Hendel (Diptera, Chloropidae) has a worldwide distribution, being the most speciose member of the subfamily Siphonellopsinae Duda with more than 80 described species, largely concentrated in the Southern Hemisphere (Ismay et al. 2021). Thirty-seven species of *Apotropina* are described from the Australasian-Oceanian region, with 22 species known to occur in Australia (Sabrosky 1989).

Apotropina are small (1.5–5.0 mm), dark to yellowish flies, sometimes with distinct tomentosity on their bodies, and with hyaline or patterned wings. *Apotropina* are associated with water bodies, sandy river margins and sea-shores; some adults form congregations in rock shelters (Ismay et al. 2021). While little else is known about their biology, and a majority of chloropids are recorded to be saprophages and phytophages. Evidence of saprophagy in *Apotropina* is inferred from various life-history records: specimens are observed with the infestation of fungi Laboulbeniales (Ascomycota: Laboulbeniomycetes) (primarily on antennae, mouth parts, and female ovipositor) and mites

(likely Pyemotidae [Acari: Acariformes]) (usually found on neck, coxae and posterior margin of tergites) – two parasitic groups which are largely associated with flies that visit decaying matter (M.V. Tschirnhaus, pers. comm. 2023). A more direct observation is the unidentified species shown in Fig. 1 visiting a dead snail shell (S. Grove, pers. comm. 2023). Additionally, some chloropids specialize as predators of other invertebrate eggs or larvae [see Ismay and Ang (2019) for a review]. Indeed, two Australian species *A. proxima* (Rayment, 1959) and *A. raymenti* (Curran, 1930) are known to prey on the immature stages of hymenopterans (Rayment 1935, 1959), and it is likely that there are other predatory species in the genus.

There is also known sexual dimorphism in the genus, largely in differences on the wing venation and coloration between males and females [e.g., *A. anomala* (Malloch, 1925) and *A. maculigena* Riccardi sp. nov.] and antennal modifications in *A. australis* (Malloch). Given that most Australian species are only known from a single sex, it is likely that there are more species with undocumented sexual dimorphism: for example, Fig. 1 shows a species observation [likely of *A. ornatipennis* (Malloch, 1923) or similar, see species remarks] from iNaturalist (Grove 2019) where there are distinct differences in wing color patterning between males and females.

Australian *Apotropina* are poorly studied, with at least ten described species known solely from their type specimen, and thus the morphology for the other sex unknown. The type species for *Apotropina* is *A. viduata* (Schiner, 1868), an Australian species originally described under *Ectropa* Schiner as a member of the family Ephydriidae Zetterstedt. However, *Ectropa* was already preoccupied by the lepidopteran genus *Ectropa* Wallengren, and was renamed *Apotropina* Hendel (1907). Most Australian *Apotropina* were described (predominantly by Malloch) under *Parahippelates* Becker and *Lasiopleura* Becker, with the former synonymized to the latter by Malloch (1936); a few other species were also described under *Ephydroscinis* Malloch. Sabrosky (1980) was able to examine the type specimen for *A. viduata* and transferred *Apotropina* to Chloropidae; he further synonymized *P. fuscipes* Malloch, 1924 with *A. viduata*, effectively synonymizing *Lasiopleura* under *Apotropina*. *Ephydroscinis* was also synonymized under *Apotropina* (Sabrosky 1989). No new Australian species in this genus have been described since 1959 (Sabrosky 1989), and it is likely that many more species remain undescribed.

Here we describe two new species of *Apotropina* to Australia: *A. maculigena* Riccardi, sp. nov. and *A. popeye* Ang, sp. nov.; we also provide an updated illustrated key to the Australian *Apotropina* species, and a conspectus of the known species with images of type specimens for most of these species. Original descriptions and subsequent taxonomic notes of relevance are also collated in a supplementary file. Finally, we propose disregarding the current type specimens [specimen no. 547474 (5c8582 and 5c85bb)] for *Apotropina bispinosa* (Becker, 1911) due to their incongruence with the species descriptions.

Materials and methods

The type series for the two new species are deposited in the following collections: the Australian Museum Research Institute, Sydney, Australia (AMRI),



Figure 1. In situ photo of *Apotropina* sp. showing sexual dimorphism. iNaturalist observation (Grove 2019) of two individuals (likely of *A. ornatipennis* (Malloch) or similar where ♂ wings (bottom left) have color pattern extending to wing apices (blue arrow), while the ♀ color pattern (top right) does not (red arrow).

the Queensland Museum, Brisbane, Australia (**QM**), the Museu de Zoologia da Universidade de São Paulo, São Paulo, Brazil (**MZUSP**), the Museum für Naturkunde, Berlin, Germany (**MfN**) and the Zoological Reference Collection, Lee Kong Chian Natural History Museum, Singapore (**ZRC**). Images of type material were also obtained for 20 *Apotropina* species deposited in the South Australian Museum, Adelaide, Australia (**SAMA**), the Naturhistorisches Museum Wien, Vienna, Austria (**NHMW**), the California Academy of Sciences San Francisco, California, USA (**CAS**), the AMRI, and the MfN. The protocol for preparing wing and terminalia follows Riccardi et al. (2018). Images at different depths of field were taken with either a Leica M205 C or M205 A trinocular stereomicroscope coupled to the software LasX and focus-stacked in either Helicon Focus 6 or Zerene Stacker v. 1.04. Scanning electron micrographs (SEM) were obtained using a JEOL JCM-6000plus high vacuum equipment. Illustrations were treated with Illustrator CS6 and images were further processed and plated in Photoshop CS6. The morphological nomenclature follows primarily Ismay et al. (2021) and also Andersson (1977) (specifically, scapular setae). Three 312-bp COI DNA barcodes for the two new species were also generated: DNA extraction and sequencing procedures followed protocols as described in Mortelmans et al. (2022); the DNA barcodes were then uploaded to GenBank.

Results

Genus diagnosis for *Apotropina* Hendel

Apotropina species are predominantly dark to yellowish species, with various patterns of pruinosity. For chaetotaxy they have three to four long fronto-orbital proclinate setae but with at least (usually the most posterior) one laterocline; long, proclinate and divergent ocellar setae; two postpronotal setae with the inner one inclinate and the outer one reclinate; notopleural setae 1+1; a row of dorsocentrals rarely reduced to the posterior pair. Fore femur or tibia not enlarged; hind tibia usually with tibial organ and tibial spur present. Wing vein C extends to M_1 ; M_4 usually with basal sinuosity as a kink (as a plesiomorphy for Chloropidae). Male with postabdomen asymmetrical sclerites; epandrium usually large and with lateral extensions (but minute epandria can also occur; von Tschirnhaus, pers. comm. 2023); hypandrial complex developed.

Illustrated key to the Australian species of *Apotropina* (with parts adapted from Malloch 1924, 1940; Rayment 1959)

- 1 Dark body largely pruinose; pleuron with 2 longitudinal strips on scutum, gena, occiput and abdomen with fine white setulae, femora additionally with longer white setae; frons dark but yellowish near anterior half; 2 short proclinate interfrontal setae; arista almost bare, only the posterior pair of dorsocentrals developed; katapisternal setae distinct; hind tibial spur indistinct, pale and short; wing hyaline, with bm+dm cell wide and dm-m vein approximately perpendicular (Fig. 11) ***A. dasypleura* (Malloch, 1928)**
- Other combination of characters **2**
- 2 Gena patterned with dark macula medially below eye; male wings with an elongate dark spot across in the middle of cells r_1 and r_{2+3} but never beyond R_{2+3} (Fig. 2C) and veins light brown basally but darkening towards apex; female wings may be hyaline..... **3**
- Gena without dark macula below eye; wings not as above **4**
- 3 Gena yellow with black setae only; genal macula reaches or nears ventral margin of gena; proboscis yellowish brown; ocellar triangle reaching $\sim\frac{1}{2}$ length of head dorsally; frons yellowish except for light brown posterior region; 2 vibrissae; 2 decussate interfrontal setae; 1 postpronotal seta; femora yellow to light brown (Fig. 2) ***A. maculigena* Riccardi, sp. nov.**
- Gena whitish with white and black setae; genal macula does not end near ventral margin of gena; proboscis dark; ocellar triangle reaching at least length of head dorsally; 3 vibrissae; 3 strong decussate interfrontal setae; femora whitish yellow but slightly infuscate near apices (Fig. 10) ***A. costomaculata* (Malloch, 1924)**
- 4 At least medial region of wing with distinct black/dark brown marking(s) in radial and medial sectors that extend from costa to beyond R_{2+3} , may extend to apex of wing; alula usually whitish **5**
- Wing without well-defined dark area, at most with a linear brownish costal suffusion **8**

- 5 Proboscis long and geniculate; scutum with silvery green metallic pattern divided by three longitudinal black lines; only 2 distal segments of tarsi dark (Fig. 25E, F).....**A. proxima (Rayment, 1959)**
- Proboscis short and capitate; scutum not as above; ≥ 2 distal tarsal segments dark.....**6**
- 6 Scutal color pattern with paired white lateral vittae on postscutum and none on prescutum; arista completely brown; 1 vibrissa; 2 to 3 weak proclinate interfrontal setae; pleuron completely white (Fig. 20)**A. raymenti (Curran, 1930)**
- Scutal color pattern with white patterning predominantly on prescutum and sometimes reaching towards postscutum; arista brown on basal bulge and apical half but yellow medially; pleuron not completely white; 1 vibrissa; 2 largely proclinate interfrontal setae**7**
- 7 Wing veins lighter basally, darker brown near apex; katapisternum usually completely white; femora dark but broadly yellow at least quarter to apices (Fig. 16)**A. ornatipennis (Malloch, 1923)**
- Wing veins dark brown throughout; katapisternum usually white with dark patches; femora dark with extreme apices yellow (Fig. 13)**A. exquisita (Malloch, 1940)**
- 8 Costal margin of the wing browned from apex of R_1 to M_1 ; body and legs largely fulvous-yellow with tarsi darker near apex, thorax shiny with a very faint dusting; postpedicel largely brown; arista with short pilosity; proboscis geniculate; 1 vibrissa; 3 or 4 pairs proclinate interfrontal setae; scutellum with 1 discal setula (Fig. 8)**A. brunneicosta (Malloch, 1923)**
- Wing hyaline or with a faint yellowish marking; other combination of characters**9**
- 9 Arista white with dense whitish pilosity; postpedicel yellowish but brown at base of arista; gena yellow, approx. as high as postpedicel; 2 vibrissae; 2 strong decussate interfrontal setae; large geniculate proboscis dark; femora and tibiae largely or entirely brown; halter brown; wing hyaline; thorax dark fulvous with distinct but not dense pruinosity; scutellum with 1 discal setula (Fig. 5).....**A. albiseta (Malloch, 1924)**
- Arista and its pilosity not white; other combination of characters**10**
- 10 Arista plumose, with basal pilosity distinctly long (almost the length of the postpedicel), shorter apically; 2 vibrissae; 3 or 4 proclinate interfrontal setae; body entirely brown and pruinose; legs and gena yellow; hind tibial spur longer than width of tibia, curved (Fig. 21).....**A. rufescens (Duda, 1934)**
- Arista with short pubescence or bare; other combination of characters**11**
- 11 Arista pubescent, with at least the basal pilosity $2\times$ basal width of the arista**12**
- Arista almost bare, or with pilosity approx. as long as the basal width of the arista (Fig. 25C).....**15**
- 12 Thorax almost uniformly fulvous, entirely pruinose with microtrichia, additionally with distinct grey-dusted central stripe extending beyond dorsocentral row; scutellum yellowish on margin; abdomen much uniformly darker than thorax; 1 katapisternal seta; legs yellow with last 2 tarsal segments dark; hind tibial organ distinctly darkened; hind tibial spur curved,

- not longer than width of tibia; antenna entirely pale yellow; M_4 reaches wing margin (Fig. 14) ***A. griseovitta* (Malloch, 1936) ♂ [part]**
- Other combination of characters **13**
- 13 Body largely covered in distinct white setulae; ventral part of katepisternum, most of scutum and discal region of scutellum black, with margins of scutum and scutellum, and rest of pleuron reddish yellow; in male hypopygium glossy black; gena deeper than half of eye; short geniculate proboscis dark; postpedicel yellow but darker at base of arista; arista densely pilose and dark; 2 vibrissae; 3 inclinate interfrontal setae; scutellum with 2 discal setulae; 1 katepisternal seta; dm-m almost 2× its own length from apex of M_4 ; hind tibial spur $\leq$ width of tibia (Fig. 12) ***A. duplicata* (Malloch, 1923)**
- Body not covered in distinct white setulae; other combination of characters **14**
- 14 Head, thorax and legs almost entirely reddish yellow without distinct markings, tarsi apices slightly darker, abdomen entirely brown; postpedicel brownish; gena shallower than half of eye; short geniculate proboscis dark; 2 vibrissae; 3 or 4 inclinate interfrontal setae; 1 katepisternal seta; mid coxa with 1 row of long setae; hind tibial spur longer than width of tibia (Fig. 15) ***A. nigripila* (Duda, 1934)**
- Thorax dorsum dusted brown grey with 2 closely associated narrow grey stripes; thorax pleuron completely grey pruinose; abdomen dark with distinct whitish posterior margins; legs brownish with darkened femora and hind legs, hind tibial spur distinctly large and curved; frons dark red-brown that becomes red anteriorly; frontal triangle dull, reaches middle of head; 2 vibrissae; 1 decussate interfrontal seta (Figs 4, 25A, B) ***A. aequalis* (Becker, 1911)**
- 15 Gena very deep, subequal in height to eye; frons strongly projected anteriorly; parafacial as wide as the postpedicel; bright orangish to yellow species but with darker abdomen; vibrissa very short/missing; 3 decussate interfrontal setae; hind tibial spur shorter than tibia diameter, but strong and curved; proboscis geniculate, longer than head when extended (Fig. 9) ***A. conopsea* (Duda, 1934)**
- Parafacial $\leq \frac{1}{2}$ as wide as the postpedicel; proboscis not longer than head when extended **16**
- 16 Body largely reddish yellow, legs paler yellow; gena deep, subequal in height to eye; 1 vibrissa; 2 inclinate interfrontal setae; 1 katepisternal seta; tibial spur short and slightly curved, slightly longer than width of tibia; males with hind tibia grossly enlarged, bearing a large and oval tibial organ (Fig. 3) ***A. popeye* Ang, sp. nov.**
- Hind tibia not enlarged; body not largely reddish yellow; other combination of characters **17**
- 17 Hind tibial spur almost indistinct and not nearly $\frac{1}{2}$ as long as the tibial diameter; all presutural acrostical setulae minute **18**
- Hind tibial spur curved and stout, usually at least as long as the tibial diameter; presutural acrostical setae distinct **20**
- 18 Frons strongly projected anteriorly; postpedicel dark; males with long, light-colored geniculate arista, females with normal bare arista; no vibrissae; 5 or 6 proclinate interfrontal setae; thorax with dorsum dark and traces of blackish lines and laterals yellow pruinose; scutellum black on the

- disc and grey laterally; a pair of grey spots on the apex of the abdominal syntergite 1+2; male mid tarsi with the 3 apical segments slightly broadened (Fig. 7)..... **A. australis (Malloch, 1924)**
- Male arista and mid tarsi with no modifications; other combination of characters **19**
- 19 Legs at least with hind femur and tibia largely fuscous; thorax with dorsum glossy black; tergites dark but with silver pruinosity along posterior margins; male wings with R_{2+3} curved anteriorly and R_{4+5} curved posteriorly creating a much wider r_{2+3} cell, but parallel in female; presutural setae absent; face dark yellow with a silvery pruinosity, $<2\times$ longer than wide; gena as deep as postpedicel length; parafacial not visible in profile; 1 white vibrissa; 3 proclinate interfrontal setae (Fig. 6).....
- **A. anomala (Malloch, 1925)**
- Legs pale yellow, the 2 apical tarsomeres of mid and hind legs slightly darkened; thorax with dorsum evenly and densely covered with grey pruinosity; last abdominal segment and male terminalia of male yellow; thorax with dorsum covered with grey pruinosity; anterior half of head yellow, face $2\times$ longer than wide; gena not as deep as postpedicel length; parafacial visible in profile; proboscis capitate; 1 vibrissa; 1 proclinate interfrontal seta (Fig. 19)..... **A. pruinosa (Thomson, 1869)**
- 20 Scutum reddish to dark brown, with a broad grey-dusted stripe extending beyond dorsocentral row; 1 katepisternal seta; scutellum grey-dusted on disc; hind tibial spur approx. as long as diameter of tibia; antenna pale yellow; M_4 reaches wing margin (Fig. 14).....
- **A. griseovitta (Malloch, 1936) ♂ [part]**
- Scutum without a grey-dusted vitta covering the dorsocentral area; another combination of characters **21**
- 21 Scutum with the anterior acrostichals very short and fine, biseriate but not decussate; most of thorax and head except ventral half of katepisternum and frons heavily pruinose with white microtrichia; 3 vibrissae; 2 or 3 proclinate interfrontal setae; thorax uniformly dark, abdomen dark basally and lighter towards apex; legs uniformly yellow except for brown terminal 2 segments of tarsi; M_4 ends well before wing margin (Fig. 22).....
- **A. taylori (Malloch, 1940)**
- Scutum with at least some anterior acrostichals $\sim\frac{1}{2}$ as long as the anterior dorsocentrals, decussate; hind tibial spur longer than the tibial width, other combination **22**
- 22 Gena at most as deep as $\frac{1}{2}$ of eye height; hind tibial spur similar to or shorter than width of tibia..... **23**
- Gena almost as high as the eye; arista with short pubescence thorax entirely shiny black with grey or brownish pruinosity; hind tibial spur much longer than width of tibia (Fig. 25D)..... **25**
- 23 Thorax brownish yellow, shiny; 1 katepisternal seta; scutum with 2 grey-dusted lines along the dorsocentrals; postpedicel largely fuscous; M_4 reaches wing margin (Fig. 14)..... **A. griseovitta (Malloch, 1936) ♀ [part]**
- Thorax shiny black or dark brown; stripes on the scutum, if present, almost indistinct; other combination; 2 vibrissae; 3 proclinate interfrontal setae; scutellum without discal setulae; katepisternal setae missing/indistinct **24**

- 24 Legs entirely yellow; postpedicel yellow to light brown; ocellar triangle and gena with distinctly pruinose with white microtrichia; M_4 ends well before wing margin (Fig. 17) ***A. pallipes* (Malloch, 1940)**
- Legs dark yellow to brown, all femora darkened medially; postpedicel dark brown dark; ocellar triangle and gena pruinose but not distinctly white; M_4 ends at or close to wing margin (Fig. 18) ***A. parva* (Malloch, 1928)**
- 25 Legs entirely orangish yellow; scutum pruinose grey with a central faint dark stripe that becomes more distinct towards the posterior; acrostichal row with ≥ 4 strong setae; scutellum not very convex (Fig. 25C, D) ***A. nudiseta* (Becker, 1911)**
- Legs dark yellow but femora largely dark; scutum with brownish pruinosity and 3 brownish stripes; scutellum convex; 3 vibrissae, 3 proclinate interfrontal setae, katepisternal seta white, indistinct (Fig. 23)..... ***A. viduata* (Schiner, 1868)**

***Apotropina maculigena* Riccardi, sp. nov.**

<https://zoobank.org/16906BC7-91B7-48D9-8BEC-6874EA51D48F>

Fig. 2A–H

Type locality and distribution. AUSTRALIA: New South Wales (Taree).

Material examined. **Holotype** ♂ AUSTRALIA: New South Wales, Taree, Lorien Wildlife Refuge, 3km N Landsdowne; sclerophyll forest, Dec.14–31.2011, Malaise trap; 31°45'04"S, 152°32'03"E; E.G. & B. Williams leg; [AMRI TYPE CODE]. Deposited in the AMRI.

Paratypes ♂♂♀♀ 34 same data as holotype; Deposited in the AMRI (20♂, 5♀), MfN (3♂, 1♀), MZUSP (3♂), ZRC (1♂, 1♀, same as specimens with submitted Genbank barcodes).

GenBank barcodes. Specimen no. ZRCENT0021054: [OR136427](#); specimen no. ZRCENT0021055: [OR136428](#).

Diagnosis. Gena yellow with a dark median macula that reaches ventral margin; frons dark yellow; ocellar triangle with silvery pruinosity; male terminalia with anal area bearing a pair of conical membranous extensions.

Description. **Male** (Fig. 2A, C–G). Body length, 2.4–2.7 mm. Wing length, 2.28–2.45 mm. **Head** (Fig. 2A, B). Broader than long dorsally and deeper than long in profile, dark yellow except for black ocellar tubercle and a dark occiput. Head and thoracic setae black. Ocellar seta strongly developed, as long as inner and outer vertical setae. Postocellar cruciate, $\sim 2/3$ of ocellars length. Three fronto-orbital setae developed, $\sim 1/3$ of the ocellars; the two anterior proclinate and the posterior laterocline. Inner vertical seta inclinate and outer vertical seta laterocline. Two pairs of interfrontal setae distinct; as long as fronto-orbitals; proclinate and slightly convergent. Frons as long as broad, lateral margins slightly convergent, front margin straight. Ocellar triangle with a silvery pruinosity, extending to half of frons length, posterior margin two thirds width of frons, lateral margins straight. Eye oval, long axis slightly oblique with short, very sparse pubescence. Face deeper than broad; carina knife-like, restricted to the upper half of frons; antennae dark yellow, postpedicel reniform, as deep as long, mostly yellow; arista blackish, with short sparse pubescence, $\sim 3\times$ as long as postpedicel; gena as wide as the length of postpedicel, with




Figure 2. *Apotropina maculigena* Riccardi, sp. nov. ♂♀ **A** ♂ habitus, lateral view **B** ♀ habitus, dorsal view **C** ♂ wing **D** ♂ terminalia, terminal view **E** ♂ terminalia, ventral view **F** hypandrium and phallic complex, lateral view **G** ♂ postabdomen, ventral view **H** egg. Abbreviations: bas, basiphallus; cer, cercus; dis, distiphallus; ep, epandrium; hyp, hypandrium; phal, phallapodeme; phal s, phallapodemic sclerite; pog, postgonite; prg, pregonite; S, syntergosternite; sp, spiracle; spm, sperm pump; st, sternite; sur, surstylus.

~ 2 unordered rows of setulae and two vibrissae; occiput blackish; proboscis short yellow; palpus yellow, small, equal in length to postpedicel, with brown setulae; mouth edge not protruding; clypeus dark brown. **Thorax** (Fig. 2A, B). Scutum approx. as long as broad, dark brown, entirely pruinose; one row of decussate acrostichal setulae and one pair of prescutellar acrostichal seta; four dorsocentral setae developed, the posterior one longer than the remaining three and as long as the ocellar seta; postpronotal lobe concolorous with scutum; one long basal postpronotal seta, equal to notopleurals; anterior post-

pronotal seta short and reclinate; one presutural intra-alar seta developed; notopleuron with 1+1 setae; one long presutural and three short postsutural supra-alar setae; postalar seta as long as ocellar seta. Pleuron dark brown, pruinose, katapisternum with a dorsal long seta. Scutellum concolorous with scutum, pruinose, broader than long, rounded apically with one pair of setulae on the disc; apical scutellar setae with separation greater to that of posterior ocelli and as long as half the scutal length; one pair of lateral scutellar setae inserted on the disc, as long as the prescutellar dorsocentrals; post-scutellum blackish. Halter pale yellow. **Wing** (Fig. 2A, C). Translucent with a large dark spot more than half of second sector, light brown veins covered in sparse brown microtrichia; costal ratios measured from h: R_1 : R_{2+3} : R_{4+5} is 6: 7.5: 7: 2; veins R_{4+5} and M_1 subparallel; distance between r-m and dm-m six times length of r-m. **Legs** (Fig. 2A). Brownish yellow, dark pilosity organized in rows; posterior tibial organ well developed, occupying one third of tibia, narrow, yellow; hind tibial spur subapical, as long as the width of the tibia apex. **Abdomen** (Fig. 2A). Tergites brown. **Terminalia** (Fig. 2D–G). Postabdomen sclerites asymmetric as in generic diagnosis. Epandrium well developed, with a laterobasal projection; surstylus conical, directed inwards; cerci not fused to each other, oval; anal lobe membranous, with a long bifid projection. Hypandrium with arms open; basiphallus oval; distiphallus cylindrical and membranous; pregonite with four setulae and sensorial pores on the apex; postgonite minute, elongated; phallopodeme short, not bifid basally; sperm pump present. **Female** (Fig. 2B). Same as male, except wing completely hyaline; abdominal segments 6–8 narrow; epiproct with one pair of setae; hypoproct pilose; cerci dark yellow, long, and narrow, with short setae at base.

Egg (Fig. 2H). Length 0.43 mm, width 0.14 mm, matt, milky white, elongated, slightly rounded at both ends; one apical pole with four to five spine-like structures. Chorionic surface with poorly visible pattern of small rounded microsculptures.

Etymology. The specific epithet *maculigena* is feminine derived from Latin, meaning gena with macula.

Remarks. The wings of *Apotropina maculigena* sp. nov. males are similar to *A. costomaculata*. However, a yellow gena with a distinct mesal dark spot and postgena yellow are considered distinctive features of *A. maculigena* sp. nov., while *A. costomaculata* has a whitish gena with an indistinct dark marking and dark postgena. Furthermore, the yellow mouthparts and male epandrium of *Apotropina maculigena* sp. nov. differ from the brown coloration of the same structures in *A. costomaculata*.

***Apotropina popeye* Ang, sp. nov.**

<https://zoobank.org/24E61305-A1A4-4C38-A2B3-F547213D2931>

Fig. 3A–H

Type locality and distribution. AUSTRALIA: Queensland (Dinden National Park).

Type material. **Holotype** ♂ Label transcription: “QLD Dinden NP, 20k EbS Ma-reeba, 17.034°S, 145.6064°E, 9 Nov 2017, Kahlpahlim Rock trail, J A & J G Lumbers”; 710 m a.s.l.; a single male specimen was collected via sweep-netting at the edge of a forest clearing. ZRC issued specimen code ZRCENT0021052. Deposited in the QM.




Figure 3. *Apotropina popeye* Ang, sp. nov. Holotype ♂ **A** habitus, left lateral view (wings truncated) **B** habitus, right lateral view **C** head & thorax, dorsal view **D** collection label **E** epandrium, terminal view **F** epandrium, ventral view **G** syntergosternite complex, hypandrium and phallic complex, ventral view **H** hypandrium and phallic complex, lateral view. Abbreviations: bas, basiphallus; cer, cercus; dis, distiphallus; ep, epandrium; hyp, hypandrium; phal, phallapodeme; phal s, phallapodemic sclerite; pog, postgonite; prg, pregonite; S, syntergosternite; sp, spiracle; st, sternite; sur, surstylus.

GenBank barcode. Holotype specimen (ZRCENT0021052): [OR136429](https://www.ncbi.nlm.nih.gov/nuclot/OR136429).

Diagnosis. Body largely yellowish orange except for blackened ocellar tubercle, light brown arista, brown dorsal regions on tergites and light yellow legs with darkened tarsal segments 4 and 5; gena deep, arista fulvous pectinate; wings hyaline with brown veins; hind tibial spur robust but short, male distinctive with extremely large, flattened oval hind tibial organ.

Description. Male. Body length, 4 mm. Wing length, 3.5 mm. **Head** (Fig. 3A–C). Broader than long dorsally, with deep gena in profile, light yellow except for black ocellar tubercle. Head setae black. Ocellar seta strongly developed, as long as inner and outer vertical setae. Postocellars cruciate, slightly shorter than ocellars. Three fronto-orbital setae developed, ~ 1/2 of ocellars; the two anterior proclinate and the posterior laterocline. Inner vertical seta laterocline and outer vertical seta inclinate. Three pairs of

interfrontal setae distinct; anterior two as long as fronto-orbitals, posterior half-length; all proclinate and slightly convergent. Frons as long as broad, lateral margins slightly convergent, front margin straight. Ocellar triangle with slight pruinosity, extending to half of frons length, posterior margin two thirds width of frons, lateral margins straight. Eye oval, long axis slightly oblique with short, very sparse pubescence. Face deeper than broad; antennae yellow except for brown base of arista, postpedicel reniform, as deep as long, arista with short pubescence, $\sim 3\times$ as long as postpedicel; gena $\sim 1/3$ as deep as eye height, with ~ 2 unordered rows of setulae and two vibrissae; genal dilation distinct. One row of postocular setae. Proboscis short yellow; palpus whitish, small, equal in length to postpedicel, with black setulae; mouth edge not protruding; clypeus light brown. **Thorax** (Fig. 3A–C). Entirely yellow, scutum slightly longer than broad, entirely pruinose; one row of laterocliniate acrostichal setae and one pair of parallel prescutellar acrostichal seta; five dorsocentral setae developed, the posterior one larger than the remaining four and as long as the ocellar seta; postpronotal lobe slightly lighter yellow than scutum; two long postpronotal seta, both similar length to notopleurals; prescutum with a pair of scapular seta (see Andersson 1977) on each side interior to postpronotal lobe at anterior margin; presutural intra-alar seta developed; notopleuron with 1+1 setae; one long presutural and two shorter postsutural supra-alar setae; postalar seta as long as ocellar seta. Pleuron dark brown, pruinose, katepisternum with one black seta on dorsal margin and populated with long white setulae on anterior half. Scutellum concolor, pruinose, broader than long, rounded apically with one pair of setulae on the disc; apical scutellar seta with separation greater to that of posterior ocelli and as long as half scutum; one pair of equally long lateral scutellar setae; postscutellum light brown. Halter pale yellow. **Wing** (Fig. 3B). Hyaline, covered in brown microtrichia; veins brown; costal ratios measured from h: R_1 : R_{2+3} : R_{4+5} : M_1 is 5: 7: 4.5: 1.5; veins R_{4+5} and M_1 subparallel; distance between r-m and dm-m five times length of r-m. **Legs** (Fig. 3A, B). Mostly light yellow except for dark brown on all tarsomeres 4 and 5, and light brown hind tibia, mid-coxal prong black. With dark pilosity organized in rows; posterior tibial organ extremely developed, occupying two-thirds of tibia basally and expanding it into a large, flattened tibial dilation; hind tibial spur subapical, as long as the width of the tibia apex. **Abdomen** (Fig. 3A, B). Tergites yellow with medial region light brown in dorsal view, with dark pilosity. Sternites weakly sclerotized, with white pilosity. **Male terminalia** (Fig. 3E–H). Postabdomen segments asymmetric as in generic diagnosis; sternites 5, 6 and syntergosternite 7+8 fused, spiracle 8 within sclerites on both sides. Epandrium well developed, with a laterobasal projection; surstylus conical, directed inwards; cerci fused, flattened; anal lobe membranous, short. Hypandrium with arms open; basiphallus oval; distiphallus short, cylindrical, and membranous; pregonite with three setulae; postgonite minute, elongate; phallapodeme short, not bifid basally.

Female. Unknown.

Etymology. The specific epithet *popeye* refers to the comically enlarged hind tibia, which in combination with the comparatively thin femur, resembles the distinctive arms and legs of the spinach-powered cartoon character “Popeye the Sailor”. It is a noun in the nominative singular standing in apposition.

***Apotropina aequalis* (Becker, 1911)**

Figs 4A–D, 25A, B

Parahippelates aequalis Becker, 1911: 111; Malloch 1924: 331.

Lasiopleura (*Lasiopleura*) *aequalis*: Malloch 1936: 23; 1940: 271.

Parahippelates variabilis Curran, 1936: 50 (synonymy: Sabrosky 1989: 651).

Type locality. PAPUA NEW-GUINEA: Stephansort, Astrolabe Bay (coll. Biró).

Distribution. AUSTRALIA: Australian Capital Territory (“Blundell’s, Molongo R.”; Canberra), New South Wales (Como; “Coramba-Dorrigo Rd”); PAPUA NEW-GUINEA: Bismarck Archipelago; SOLOMON ISLANDS: (Guadalcanar Is.; Santa Ana Is.; Matema Is.).

Examined material. *Allotype* [= *paratype*] ♀ Label transcription: “Guadalcanar Island, V-20-33; Kau Kau Plantation; Solomon Islands; M Willows Jr., Collector; Templeton Crocker Exped. 1933; *Parahippelates variabilis* Currani Allotype ♀; Collection of the California Academy Of Sciences, San Francisco, California”. Deposited in the CAS.

Taxonomic notes. This species was originally described from Papua New-Guinea based on a single specimen (sex not indicated); some Australian specimens were subsequently determined to the species (Malloch 1924, 1940). Note that Malloch (1940) erroneously stated that the type as “[o]riginally described from Sydney”. Type is indicated to be in the same collection as *A. nudiseta* Becker, apparently deposited in the Hungarian Natural History Museum, Hungary; however, the authors were not able to examine this material for this study, but were able to obtain an allotype of *Parahippelates variabilis* Curran (Fig. 4A–D), with the following chaetotaxy observed: 2 vibrissae; 1 weak decussate interfrontal setae; 2 postpronotal setae; 2 scapular setae; 1+1 notopleural setae; strong biseriate divergent acrostichal row; 1+3 dorsocentral setae; postalar and intrapostalar setae present; 1 weak acrostichal prescutellar setae; 2 setulae on scutellum; katapisternal seta weak. Becker (1911) did include a drawing of the head in lateral view (Fig. 25A) and Malloch (1940) illustrated the hind tibial spur (Fig. 25B). This species does not have any immediately distinctive diagnosable character sets based on existing descriptions. As such, study with more identified material from recorded localities would be useful for determining the limits of this apparently widespread species. Note that the images provided in Fig. 4A–D are of the *P. variabilis* allotype specimen. Original description and subsequent taxonomic notes in Suppl. material 1.

***Apotropina albiseta* (Malloch, 1924)**

Fig. 5A–E

Parahippelates albiseta Malloch, 1924: 330.

Type locality and distribution. AUSTRALIA: Queensland (Eidsvold; Draper).

Examined material. *Holotype* ♂ Label transcription: “Australian Museum, K 584429; Bancroft, Eidsvold Q., 19.8’23; K50096; *Parahippelates albiseta* Type, Det. J R Malloch”; 25°22’14”S, 151°7’21”E. Deposited in the AMRI.




Figure 4. *Parahippelates variabilis* Curran Allotype ♀, (synonym of *Apotropina aequalis*) **A** habitus, lateral view **B** habitus, dorsal view **C** head, anterior view **D** specimen labels.

Taxonomic notes. This is a testaceous species with distinctive white pubescent arista (Fig. 5C). It was described based on a syntype series in AMRI of four specimens with accession code K50096. The specimens are now separated, and we designate one male holotype (K 584429, imaged), and three paratypes (K 584430, male; K 584431, female; K 584432, female). An iNaturalist observation (Sarna 2020) further shows a female specimen in situ from Draper, Queensland (Fig. 5E). Chaetotaxy as observed: 2 vibrissae; 3 strong inclinate interfrontal setae; 2 postpronotal setae; 2 scapular setae; 1+1 notopleural setae; strong biseriate divergent acrostichal row; 1+3 dorsocentral setae; 1+2 supra-alar setae; postalar and intrapostalar setae present; katepisternal seta missing/indistinct. Original description and images of all type specimen labels in Suppl. material 1.

***Apotropina anomala* (Malloch, 1925)**

Fig. 6A–D

Parahippelates anomala Malloch, 1925: 96.

Lasiopleura (*Lasiopleura*) *anomala*: Malloch 1940: 273.




Figure 5. *Apotropina albiseta* (Malloch) holotype ♂ (K 584429) and live specimen ♀ **A** holotype ♂ habitus, lateral view **B** holotype ♂ habitus, dorsal view **C** holotype ♂ head, anterior view **D** holotype ♂ specimen labels **E** ♀ live specimen, iNaturalist observation (Sarna 2020).




Figure 6. *Apotropina anomala* (Malloch) holotype ♀ (K 359223) **A** habitus, lateral view **B** habitus, dorsal view **C** head, anterior view **D** specimen labels.

Type locality. AUSTRALIA: New South Wales (Blue Mountains).

Distribution. AUSTRALIA: New South Wales (Blue Mountains), South Australia (Mt. Eba).

Examined material. Holotype ♀ Label transcription: "Australian Museum, K 359223; Blue Mtns., 15.1.22., Health Dept.; HOLOTYPE, *Lasiopleura anomala* Malloch; *Parahippelates anomala* Type, Det. J.R.Malloch; SPHTM Coll."; 33°39'55"S, 150°17'4"E. Deposited in the AMRI.

Taxonomic notes. This is a dark colored species with whitish tomentosity. It is noted for its sexually dimorphic wing venation. Initially described based on female specimens (Malloch 1925) with normal venation (Fig. 6B), male specimens were found to have R_{2+3} curved anteriorly and R_{4+5} curved posteriorly, creating a much wider r_{2+3} and narrower r_1 cells (Malloch 1940). Chaetotaxy as observed: 1 (white) vibrissa; 3 weak proclinate interfrontal setae; 2 postpronotal setae; at least 1 scapular seta; 1+1 notopleural setae; weak biseriate acrostichal row; 1+3 dorsocentral setae; postalar and intrapostalar setae present. Currently, the holotype and five paratypes are Deposited in the AMRI. Original description and subsequent taxonomic notes in Suppl. material 1.

***Apotropina australis* (Malloch, 1924)**

Fig. 7A–E

Ephydroscinis australis Malloch, 1924: 331.

Type locality and distribution. AUSTRALIA: New South Wales (Woy Woy).

Examined material. Holotype ♂ Label transcription: "Australian Museum, K 359224; WoyWoy, 2.Sept. '23, Mackerras; HOLOTYPE, *Ephydroscinis australis* Mall.; *Ephydroscinis australis* Type, Det. J R Malloch; SPHTM Coll."; 33°29'10"S, 151°19'24"E. Deposited in the AMRI.

Taxonomic notes. This distinctive species is dark colored with whitish tomentosity. It was described from a male (holotype; Fig. 7A–C) and female specimen from the type locality and is noted for the male's sexually dimorphic geniculate arista; female has normal bare arista (Fig. 7E). Chaetotaxy as observed: no vibrissae; 5 or 6 weak proclinate interfrontal setae; 2 postpronotal setae; 1+1 notopleural setae; weak biseriate acrostichal row; 1+3 dorsocentral setae; katepisternal seta missing/indistinct. Original description in Suppl. material 1.

***Apotropina brunneicosta* (Malloch, 1923)**

Fig. 8A–C

Parahippelates brunneicosta Malloch, 1923: 620.

Lasiopleura (*Lasiopleura*) *brunneicosta*: Malloch 1940: 272.

Type locality and distribution. AUSTRALIA: Northern Territory (Darwin).

Examined material. Holotype ♀ Label transcription: "TYPE; Darwin, G. F. Hill; *Parahippelates brunneicosta* Type, Det. J.R.Malloch"; deposited in the SAMA.




Figure 7. *Apotropina australis* (Malloch) holotype ♂ (K 359224) & female. **A** holotype ♂ habitus, lateral view **B** holotype ♂ habitus, dorsal view **C** holotype ♂ head, anterior view **D** holotype ♂ specimen labels **E** ♀ antennae, illustration adapted from Malloch, 1924: 332.

Taxonomic notes. This testaceous species is distinctive for having its costal regions of the wing browned from the apex of R_1 to M_1 (Fig. 8A). Chaetotaxy as observed: 1 to 2 vibrissae; at least 3 weak proclinate interfrontal setae; 2 postpronotal setae; 2 scapular setae; 1+1 notopleural setae; weak biseriate divergent acrostichal row; 1+3 discal setae; 1+1 supra-alar setae; postalar setae and intrapostalar setae present; scutellum with 1 dorsocentral setulae; katepisternal seta present but weak. It is described based on a single female; male morphology remains unknown. Original description in Suppl. material 1.

Apotropina conopsea (Duda, 1934)

Fig. 9A–D

Parahippelates conopseus Duda, 1934: 45.

Lasiopleura (*Lasiopleura*) *conopsea*: Malloch 1936: 23; 1940: 273.

Type locality and distribution. AUSTRALIA: Queensland (Cairns).

Examined material. **Syntype** ♀ Label transcription: "Holotypus"; Typus; Cairns, N. Queensland., 1907; coll. Lichtwardt; *Parahippelates conopseus* D., ♀ d. Duda; <http://coll.mfn-berlin.de/u/5c8591>". Deposited in the MfN.




Figure 8. *Apotropina brunneicosta* (Malloch) holotype ♀ **A** habitus, lateral view **B** habitus, dorsal view **C** specimen labels.




Figure 9. *Apotropina conopsea* (Duda) syntype ♀ (5c8591) **A** habitus, right lateral view **B** habitus, left lateral view **C** habitus, dorsal view **D** specimen labels.

Taxonomic notes. This testaceous species is distinctive for its deep gena, relatively small eyes, long dark geniculate proboscis (Fig. 9A, B) and strongly projected frons (Fig. 9C). Originally described in detail by Duda (1934) based on five specimens (2♂, 3♀), with Malloch (1936) adding more details. Chaetotaxy as observed: no vibrissae; 3 strong proclinate interfrontal setae; 2 postpronotal setae; 2 scapular setae; 1+1 notopleural setae; weak biseriate divergent acrostichal row; 1+3 dorsocentral setae; 1+1 supra-alar setae; postalar and intrapostalar setae present; scutellum with 1 discal setula; katepisternal seta missing/

indistinct. Note that Malloch (1940) stated the type locality (being N. Territory: Darwin) in error, which should be Queensland instead. Original description and subsequent taxonomic notes in Suppl. material 1.

***Apotropina costomaculata* (Malloch, 1924)**

Fig. 10A–D

Parahippelates costomaculata Malloch, 1924: 329.

Lasiopleura (*Lasiopleura*) *costomaculata*: Malloch 1940: 270.

Type locality and distribution. AUSTRALIA: New South Wales (Sydney).

Examined material. *Holotype* ♂ Label transcription: “Australian Museum, K 50094; Sydney, 31.12.22, Health Dept.; (red circle label); K50094; *Parahippelates costomaculata* Type, Det J R Malloch”; 33°52'S, 151°13'E. Deposited in the AMRI.

Taxonomic notes. This brownish species is similar to *A. maculigena* sp. nov. in that it has a genal macula and wing with an elongate macula on the costal region within cells r_1 and r_{2+3} (Fig. 10A) but can be reliably differentiated from the latter by its much smaller genal macula (never close to reaching the ventral margin: Fig. 10A), bearing both dark and whitish genal setae (all black in *A. maculigena* sp. nov.), lighter anterior gena and legs, and also a less projected frons (Fig. 10B). Three male specimens from the type locality are known thus far. Female morphology currently unknown but based on the sexual dimorphism exhibited in *A. maculigena* sp. nov., it is likely that females in *A. costomaculata* will also have completely hyaline wings. Chaetotaxy as observed: 3–4 vibrissae; 2 strong decussate interfrontal setae; 2 postpronotal setae; 2 scapular setae; 1+1 notopleural setae; weak biseriate divergent acrostichal row; 1+3 dorsocentral setae; 1+1 supra-alar setae; postalar and intrapostalar setae present; 1 strong prescutellar seta; scutellum with 1 discal setula; katapisternal seta missing/indistinct. Original description in Suppl. material 1.

***Apotropina dasypleura* (Malloch, 1928)**

Fig. 11A–D

Parahippelates (*Terraeregina*) *dasypleura* Malloch, 1928: 303.

Lasiopleura (*Terraeregina*) *dasypleura*: Malloch 1940: 270.

Type locality and distribution. AUSTRALIA: Queensland (Macknade).

Examined material. *Holotype* ♀ Label transcription: “Australian Museum, K 359225; Macknade, 1918., Q.; HOLOTYPE, *Lasiopleura* (*Terraeregina*) *dasypleura* Type, Det. J.R. Malloch”; 18°35'15"S, 146°15'38"E. Deposited in the AMRI.

Taxonomic notes. This is a dark colored species with whitish tomentosity and is distinct for its completely whitish pruinose pleura (including anepisternum: Fig. 11A), and two longitudinal strips on the scutum (Fig. 11B). Chaetotaxy as observed: no vibrissae; at least 2 weak proclinate interfrontal setae; 2 postpronotal setae; 2 scapular setae; 1+1 notopleural setae; weak biseriate divergent acrostichal row; 1+3 dorsocentral setae; 1+1 supra-alar setae; postalar




Figure 10. *Apotropina costomaculata* (Malloch) holotype ♂ (K 50094) **A** habitus, lateral view **B** habitus, dorsal view **C** head, anterior view **D** specimen labels.




Figure 11. *Apotropina dasyleura* (Malloch) holotype ♀ (K 359225) **A** habitus, lateral view **B** habitus, dorsal view **C** head, anterior view **D** specimen labels.

and intrapostalar setae present; scutellum with 1 discal setula; katapisternal seta missing/indistinct. It is described from one female specimen and is only known from the type locality.

Original description in Suppl. material 1.

***Apotropina duplicata* (Malloch, 1928)**

Fig. 12A–C

Parahippelates duplicata Malloch, 1923: 621.

Lasiopleura (*Lasiopleura*) *duplicata*: Malloch 1940: 272.

Type locality and distribution. AUSTRALIA: Northern Territory (Melville Is.).

Examined material. Holotype ♂ Label transcription: “Melville Is., N.T., G. F. Hill; *Parahippelates duplicata* Type, Det. J.R.Malloch”. Deposited in the SAMA.

Taxonomic notes. This is a relatively testaceous species with darkened dorsum, glossy hypopygium and yellowish legs, largely covered in whitish tomentosity (Fig. 12A, B). It does not have any immediately distinctive diagnosable character sets based on existing descriptions but can be identified based on the provided key. Thus far only known from the male type specimen; female morphology unknown. Chaetotaxy as observed: 2 vibrissae; 3 strong inclinate/decussate interfrontal setae; 2 postpronotal setae; 2 scapular setae; 1+1 notopleural setae; weak biseriate divergent acrostichal row; 1+3 dorsocentral setae; 1+1 supra-alar setae; postalar and intrapostalar setae present; 1 strong prescutellar seta; scutellum with 1 discal setula; katapisternal seta missing/indistinct. Original description in Suppl. material 1.

***Apotropina exquisita* (Malloch, 1940)**

Fig. 13A–D

Lasiopleura (*Lasiopleura*) *exquisita* Malloch, 1940: 270.

Type locality and distribution. AUSTRALIA: Western Australia (Geraldton).

Examined material. Holotype ♂ Label transcription: “Australian Museum, K 359226; Geraldton, W.A., 5. Sept. 1926., E.W. Ferguson; HOLOTYPE, *Lasiopleura exquisita* Mall.; *Lasiopleura exquisita* Type, det. JRMALLOCH; SPHTM Coll.”; 28°46'S, 114°37'E. Deposited in the AMRI.

Taxonomic notes. *Apotropina exquisita* belongs to a group of described species (including *A. ornatipennis*, *A. proxima*, and *A. raymenti*) that have dark bodies with shiny tomentosity, wings with distinct dark patterning covering at least the medial region from costal margin to beyond R_{2+3} vein, and usually shiny-white alula. *Apotropina exquisita* can be distinguished from other species in this group with the following combination of characters: short capitate proboscis, with more than two distal tarsal segments dark, pleuron not completely whitish tomentose, wing with completely dark venation especially basally, and femora dark with extreme apices yellow (Fig. 13A–C). This species is only known from female specimens, male morphology unknown. However, there is evidence that species in this group may have sexually dimorphic color patterns (see Fig. 1)




Figure 12. *Apotropina duplicata* (Malloch) holotype ♂ **A** habitus, lateral view **B** habitus, dorsal view **C** specimen labels.

where males may have more prominent patterns than females. Chaetotaxy as observed: 1 vibrissa; 2 weak proclinate interfrontal setae; 2 postpronotal setae; 2 scapular setae; 1+1 notopleural setae; short weak biseriate acrostichal row only in medial region of scutum; 1+3 dorsocentral setae; 1+1 supra-alar setae; postalar and intrapostalar setae present; scutellum with 1 discal setula; katepisternal seta missing/indistinct. Original description in Suppl. material 1.

***Apotropina griseovitta* (Malloch, 1936)**

Fig. 14A–C

Lasiopleura griseovitta Malloch, 1936: 25, 1940: 25.

Type locality and distribution. AUSTRALIA: Queensland (Mt Molloy).

Examined material. Holotype ♂ Label transcription: "Australian Museum, K 359227; Mt. Molloy, QUEENSLAND, F.H. Taylor; HOLOTYPE, *Lasiopleura* "*griseohirta*" [note: likely a misspelling] Mall.; *Lasiopleura griseovitta* Type, det. JRMALLOCH; SPHTM Coll."; 16°40'27"S, 145°19'50"E. Deposited in the AMRI.

Taxonomic notes. Malloch (1936) described this species based on a single male specimen, which has a predominantly tawny thorax, darker abdomen, yellowish legs, and the scutum has a broad central pruinose stripe along the dorso-centrals. Unfortunately, the male holotype specimen is currently damaged: head missing, pruinosity pattern on scutum likely abraded (Fig. 14A, B). Malloch (1940)




Figure 13. *Apotropina exquisita* (Malloch) holotype ♀ (K 359226) **A** habitus, lateral view **B** habitus, dorsal view **C** head, anterior view **D** specimen labels.




Figure 14. *Apotropina griseovitta* (Malloch) holotype ♂ (missing head; K 359227) **A** habitus, lateral view **B** habitus, dorsal view **C** specimen labels.

subsequently determined a female specimen from the type locality without providing his reasoning for assigning it to this species, and noted that the female has two narrow pruinose stripes on the scutum (as opposed to a single broad stripe in the male). We were unable to examine the female specimen for this study. Based on Malloch's (1936, 1940) descriptions, there are no autapomorphies to easily diagnose this species, and the set of characters used to delimit this species contains multiple interpretations (e.g., the basal pilosity on the arista is either twice as long [♂] or just as long as the basal width of arista [♂ & ♀], arista either pale-yellow [♂] or brownish [♀] and the aforementioned sexually dimorphic difference in scutal pruinosity pattern). Malloch (1940) acknowledges this multiplicity in the diagnosis of this species and makes three possible interpretations in his key; this is reflected in ours as well. In our opinion, the species limits for *A. griseovitta* are poorly defined and need to be further investigated with more character systems (e.g., genitalia, molecular data). Chaetotaxy as observed: 2 postpronotal setae; 1+1 notopleural setae; postalar and intrapostalar setae present; scutellum with 1 dorsocentral setula; katepisternal seta missing/indistinct. Original description and subsequent taxonomic notes in Suppl. material 1.

***Apotropina nigripila* (Duda, 1934)**

Fig. 15A–D

Parahippelates nigripilus Duda, 1934: 48.

Type locality and distribution. AUSTRALIA: Northern Territory (Darwin: Palmerston).

Examined material. **Syntype** ♂ Label transcription: "Palmerston, N. Australien, XI. 1908; coll. Lichtwardt; *Parahippelates nigripilus* Duda, ♂ d Duda; Typus; <http://coll.mfn-berlin.de/u/5c8585>". Deposited in the MfN.

Taxonomic notes. This is a testaceous species comparable to *A. rufescens* but with more distinct and stronger setation (Fig. 15A–C). It was described based on male specimens and is only known from the type locality. Female morphology unknown. This species does not have any immediately distinctive diagnosable character sets based on existing descriptions but can be identified based on the provided key. Chaetotaxy as observed: 2 vibrissae; at least 3 proclinate interfrontal setae; 2 postpronotal setae; 2 scapular setae; 1+1 notopleural setae; strong biseriate divergent acrostichal row; 1+3 dorsocentral setae; 1+1 supra-alar setae; postalar and intrapostalar setae present; scutellum with 1 discal setula; katepisternal seta missing/indistinct. Original description, translation, and subsequent taxonomic notes in Suppl. material 1.

***Apotropina nudiseta* (Becker, 1911)**

Fig. 25C, D

Parahippelates nudiseta Becker, 1911: 113.

Lasiopleura (*Lasiopleura*) *nudiseta*: Malloch 1940: 274.

Type locality and distribution. AUSTRALIA: New South Wales (Botany Bay, Sydney; Wahroonga, nr. Sydney).




Figure 15. *Apotropina nigripila* (Duda) syntype ♂ (5c8585) **A** habitus, right lateral view **B** habitus, left lateral view **C** habitus, dorsal view **D** specimen labels.

Taxonomic notes. This species was described from the type locality (Sydney) and remains known only from the area. Type material is noted to be in the same collection as *A. aequalis* Becker, apparently deposited in the Hungarian Natural History Museum, Hungary; however, the authors were not able to examine the material for this study. Malloch (1940) provides illustrations of the arista and hind tibial spur (Fig. 25C, D). This species does not have any immediately distinctive diagnosable character sets based on existing descriptions but can be identified based on the provided key. The original description and subsequent taxonomic notes in Suppl. material 1, which indicate chaetotaxy as having four or more strong decussate acrostichals and “similarly arranged” dorsocentral setae.

***Apotropina ornatipennis* (Malloch, 1940)**

Fig. 16A–C

Parahippelates ornatipennis Malloch, 1923: 620.

Lasiopleura (*Lasiopleura*) *ornatipennis*: Malloch 1940: 270.

Type locality and distribution. AUSTRALIA: Victoria (Chelsea), New South Wales (Collaroy, nr. Sydney).

Examined material. **Holotype** ♀ Label transcription: “Chelsea, V., 28.9.19; *Parahippelates ornatipennis* Type, Det. J.R.Malloch”. Deposited in the SAMA.




Figure 16. *Apotropina ornatipennis* (Malloch) holotype ♀ **A** habitus, lateral view **B** habitus, dorsal view **C** specimen labels.

Taxonomic notes. *Apotropina ornatipennis* belongs to a group of described species (including *A. exquisita*, *A. proxima* and *A. raymenti*) that have dark bodies with shiny tomentosity, wings with distinct dark patterning covering at least the medial region from costal margin to beyond R_{2+3} vein, and usually shiny-white alula. *Apotropina ornatipennis* can be distinguished from other species in this group with the following combination of characters: short capitate proboscis, with more than two distal tarsal segments dark, katepisternum completely whitish tomentose, wing veins lighter basally, darker brown near apex, femora dark but broadly yellow at least a quarter to the apices (Fig. 16A, B). This species is only known from female specimens, male morphology unknown. However, there is evidence that species in this group may have sexually dimorphic color patterns (see Fig. 1) where males may have more prominent patterns than females. Chaetotaxy as observed: 1 vibrissa; 2 moderately-strong proclinate interfrontal setae; 2 postpronotal setae; 1+1 notopleural setae; short weak biseriate acrostichal row only in medial region of scutum; katepisternal seta missing/indistinct. Original description in Suppl. material 1.

***Apotropina pallipes* (Malloch, 1940)**

Fig. 17A–D

Lasiopleura (*Lasiopleura*) *parva* var. *pallipes* Malloch, 1940: 273.

Type locality and distribution. AUSTRALIA: New South Wales (Narrabeen, Sydney).

Examined material. **Holotype** ♀ Label transcription: "Australian Museum, K 359228; Sydney, Narrabeen, 21.7.23, Health Dept.; HOLOTYPE, *L. parva pallipes* Mall.; Probable type of *pallipes*; in Mall. Colln. With paratype of *parva*.,




Figure 17. *Apotropina pallipes* (Malloch) holotype ♀ (K 359228) **A** habitus, lateral view **B** habitus, dorsal view **C** head, anterior view **D** specimen labels.

Lasiopleura parva var. *pallipes* MALL., det. Sabrosky; SPHTM Coll."; 33°42'54"S, 151°17'4"E. Deposited in the AMRI.

Taxonomic notes. This dark brown species with yellowish legs was described as a variety of *A. parva*, based on a single female specimen from the type locality as the latter (Malloch, 1940), where he noted differences from *A. parva* in having legs entirely 'honey-yellow', the gena being more narrowed anteriorly (Fig. 17A, B), but "having only one specimen of each form available I do not care to go farther into details". It was subsequently upgraded to species level (Sabrosky, 1989). The authors have also noted that *A. pallipes* additionally has M_4 ending well before wing margin (Fig. 17A), as opposed to *A. parva* where the vein ends at the wing margin (Fig. 18B). This species does not have any immediately distinctive diagnosable character sets based on existing descriptions but can be identified based on the provided key. Chaetotaxy as observed: 2 vibrissae; 3 moderately strong proclinate interfrontal setae; 2 postpronotal setae; 1+1 notopleural setae; short weak biseriate acrostichal row; 1+3 dorsocentral setae; 1+1 supra-alar setae; postalar and intrapostalar setae present; 1 weak acrostichal prescutellar setae; scutellum with 1 discal setulae; katapisternal seta missing/indistinct. Original taxonomic note in Suppl. material 1.

***Apotropina parva* (Malloch, 1928)**

Fig. 18A–D

Hippelates parva Malloch, 1928: 302.

Lasiopleura (*Lasiopleura*) *parva*: Malloch 1940: 273.

Type locality and distribution. AUSTRALIA: New South Wales (Sydney).

Examined material. *Holotype* ♀ Label transcription: “Australian Museum, K 359229; Sydney, 31.12.23, Health Dept.; HOLOTYPE, *Lasiopleura parva* Mall.; *Parahippelates parva* Type, det. JRMALLOCH; SPHTM Coll.”; 33°52'S, 151°13'E. Deposited in the AMRI.

Taxonomic notes. This dark-brown species has brown-banded yellowish legs and hyaline wings (Fig. 18A, B). It resembles *A. taylori* and *A. pallipes*, but can be distinguished from the former based on the longer, stronger hind tibial spur and presutural acrostichal setae, and from the latter based on characters described in the taxonomic notes for *A. pallipes*. This species does not have any immediately distinctive diagnosable character sets based on existing descriptions but can be identified based on the provided key. Chaetotaxy as observed: 2 vibrissae; 3 strong decussate interfrontal setae; 2 postpronotal setae; 2 scapular setae; 1+1 notopleural setae; short weak reclinate biseriate acrostichal row; 1+3 dorsocentral setae; 1+1 supra-alar setae; postalar and intrapostalar setae present; missing/indistinct prescutellar setae; scutellum with 1 discal setula; katepisternal seta missing/indistinct. Original description and subsequent taxonomic note in Suppl. material 1.

***Apotropina proxima* (Rayment, 1959)**

Fig. 25E, F

Ephydroscinis proxima Rayment, 1959: 332.

Type locality and distribution. AUSTRALIA: Victoria (Mt. Richmond Reserve).

Taxonomic notes. *Apotropina proxima* likely belongs to a group of described species (including *A. exquisita*, *A. ornatipennis* and *A. raymenti*) that have dark bodies with shiny tomentosity, wings with distinct dark patterning covering at least the medial region from costal margin to beyond R_{2+3} vein. Based on the species description, *A. proxima* can be distinguished from other species in this group with its long geniculate proboscis, having only two distal tarsal segments dark and scutum with a silvery-green metallic pattern divided by three longitudinal black lines. The description did not indicate any deposited type material for examination, but did provide drawings which depict the fly with brown macula at the radial sector, a long, geniculate proboscis (Fig. 25E) and three longitudinal black strips along the scutum (Fig. 25F). However, the illustration does not reflect any “basally angulated fore tibiae” as indicated in the description, and as such the authors have opted to exclude this ambiguous character from the key. This species was described with life history information - as a likely hyperparasitoid associated with two other predatory/parasitoid species *Sericophorus chalybeus* (F. Smith, 1851) (syn. *S. victoriensis* Rayment) and *Acanthostethus portlandensis* (Rayment, 1953) (Rayment 1959). Original description in Suppl. material 1, which only reflects the only chaetotaxy as possessing four dorsocentral setae.




Figure 18. *Apotropina parva* (Malloch) holotype ♀ (K 359229) **A** habitus, lateral view **B** habitus, dorsal view **C** head, anterior view **D** specimen labels.

***Apotropina pruinosa* (Thomson, 1869)**

Fig. 19A–D

Oscinis pruinosa Thomson, 1869: 606.

Parahippelates seticauda Malloch 1928: 302 (nomenclatural changes: Sabrosky 1955: 188 [transferred to *Lasiopleura*]; Sabrosky 1989: 651 [synonymized under *A. pruinosa*]).

Type locality and distribution. AUSTRALIA: New South Wales (Sydney); Victoria (Warburton).

Examined material. **Holotype** *Parahippelates seticauda* Malloch, 1928 ♂ Label transcription: "Australian Museum, K 359230; Sydney, 25.1.25, Health Dept.; HOLOTYPE, *Lasiopleura seticauda* Mall.; *Parahippelates seticauda* Type Mall., Det J R Malloch; SPHTM Coll."; 33°53'S, 151°13'E; Deposited in the AMRI.

Taxonomic notes. This species has a blackish thorax, brownish abdomen with white bands, as well as pale yellow legs, hypopygium and front portion of head. It is distinctive in having its body (including hypopygium) and posterior portion of head completely covered in dusty white pruinosity (Fig. 19A, B). It was originally described by Thomson (1869) under *Oscinis* Loew; Malloch




Figure 19. *Parahippelates seticauda* Malloch holotype ♂, (synonym of *Apotropina pruinosa*; K 359230) **A** habitus, lateral view **B** habitus, dorsal view **C** head, anterior view **D** specimen labels.

(1928) described another species *P. seticauda*, which was subsequently transferred to *Lasiopleura* (Sabrosky 1955) and further synonymized under *A. pruinosa* (Sabrosky 1989). Note that the images provided here are of the *P. seticauda* holotype specimen (Fig. 19A–D). Chaetotaxy as observed: 1 vibrissa; 1 strong proclinate interfrontal setae; 2 postpronotal setae; 2 scapular setae; 1+1 notopleural setae; short weak reclinate biseriate acrostichal row; 1+3 dorsocentral setae; 1+1 supraalar setae; postalar and intrapostalar setae present; missing/indistinct prescutellar setae; scutellum without discal setulae; katepis-ternal seta missing/indistinct. Original descriptions and subsequent taxonomic note in Suppl. material 1.

***Apotropina raymenti* (Curran, 1930)**

Fig. 20A–D

Ephydroscinis raymenti Curran, 1930: 1.

Neoborborus speculabundus: Rayment 1931: 191; Richards 1973: 396 (synonymization).




Figure 20. *Apotropina raymenti* (Curran) 'cotype' [=paratype] ♀ (K 559468) **A** habitus, lateral view **B** habitus, dorsal view **C** head, anterior view **D** specimen labels.

Type locality and distribution. AUSTRALIA: Victoria (Sandringham, Pt Phillip).

Examined material. **Cotype** [= paratype] series for *Ephydroscinis raymenti* Curran, 1930 ♀♀♀ K66965 (separated to three specimens individually: K 559467, K 559468, K 559469). Deposited in the AMRI. Label transcription of imaged specimen: "Australian Museum, K 559468; Pt. Phillip, Vict., OCT 1934, T. Rayment; K66965; Cotype".

Taxonomic notes. *Apotropina raymenti* belongs to a group of described species (including *A. exquisita*, *A. ornatipennis* and *A. proxima*) that have dark bodies with shiny tomentosity, wings with distinct dark patterning covering at least the medial region from costal margin to beyond R_{2+3} vein, and usually shiny-white alula. *Apotropina raymenti* can be distinguished from other species in this group with the following combination of characters: short capitate proboscis, arista completely brown; scutal color pattern with paired white lateral vittae on postscutum and none on prescutum; katepisternum completely whitish tomentose; wing veins lighter basally but darker near apex (Fig. 20A, B). This species is only known from female specimens, male morphology unknown. However, there is evidence that species in this group may have sexually dimorphic color patterns (see Fig. 1) where males may have more prominent patterns than

females. This species was described with life history information, as a likely parasitoid associated with the bee species *Lasioglossum* (*Homalictus*) *niveifrons* (Cockerell), where it visits the host's ground burrows near the coastline. Rayment (1931) further erected a new genus to describe a species, *Neoborborus speculabundus* Rayment under family Borboridae Newman [=Sphaeroceridae Macquart]; it was subsequently transferred to Chloropidae by Richards (1973: 396) and synonymized under *A. raymenti* (Sabrosky 1989). In AMRI are deposited a cotype [=paratype] series (K66965) of three female specimens (K 559467, K 559468 (imaged), K 559469). Chaetotaxy as observed: 1 vibrissa; 3 weak proclinate interfrontal setae; 2 postpronotal setae; 2 scapular setae; 1+1 notopleural setae; short weak reclinate biseriate acrostichal row only on the medial region; 1+3 dorsocentral setae; 1+1 supra-alar setae; postalar and intrapostalar setae present; missing/indistinct prescutellar setae; scutellum without discal setulae; katepisternal seta weak. Original descriptions provided in Suppl. material 1.

***Apotropina rufescens* (Duda, 1934)**

Fig. 21A–D

Parahippelates rufescens Duda, 1934: 49.

Lasiopleura (*Lasiopleura*) *rufescens*: Malloch 1936: 24; 1940: 272.

Type locality and distribution. AUSTRALIA: Northern Territory (Darwin: Palmerston).

Examined material. Syntype ♀ Label transcription: “Palmerston, N. Australien, XI. 1908; coll. Lichtwardt; *P. rufescens* D., ♀ d. Duda; Typus; <http://coll.mfn-berlin.de/u/5c8578>”; Deposited in the MfN.

Taxonomic notes. This species has a dark brown body and largely yellowish head and legs, superficially similar to *A. pruinosa*, but is distinctive for its plumose arista with the pilosity at its basal half almost the length of the postpedicel (Fig. 21A–C). Only known from the type locality. Chaetotaxy as observed: 2 vibrissae; 3 weak proclinate interfrontal setae; 2 postpronotal setae; 2 scapular setae; 1+1 notopleural setae; short weak reclinate biseriate acrostichal row; 1+3 dorsocentral setae; 1+1 supra-alar setae; scutellum with 1 discal setula; katepisternal seta missing/indistinct. Original description, translation, and subsequent taxonomic notes in Suppl. material 1.

***Apotropina taylori* (Malloch, 1940)**

Fig. 22A–D

Lasiopleura (*Lasiopleura*) *taylori* Malloch, 1940: 273.

Type locality and distribution. AUSTRALIA: New South Wales (Blue Mts; Hampton).

Examined material. Holotype ♂ Label transcription: “Australian Museum, K 359231; Blue Mtns., 13.4.22, Health Dept.; Presumed HOLOTYPE, *Lasiopleura taylori* MALL., (ex. MALL. Colln 1954), det Sabrosky; HOLOTYPE *Lasiopleura taylori*; SPHTM Coll.”; Deposited in the AMRI.




Figure 21. *Apotropina rufescens* (Duda) syntype ♀ (5c8578) **A** habitus, right lateral view **B** habitus, left lateral view **C** habitus, dorsal view **D** specimen labels.

Taxonomic notes. This species has a dark brown body and largely testaceous head and legs, largely covered in pruinosity, superficially similar to *A. pruinosa*, but can be distinguished from that species based on its brownish antennae (which is fully yellow in *A. pruinosa*; see Fig. 19A) and its overall darker color (Fig. 22A–C). It was described based on male specimen and is only known from the type locality. Female morphology unknown. This species does not have any immediately distinctive diagnosable character sets but can be identified based on the provided key. Chaetotaxy as observed: 2 vibrissae; 3 strong slightly inclinate interfrontal setae; 2 postpronotal setae; 2 scapular setae; 1+1 notopleural setae; short weak reclinate biseriate acrostichal row; 1+3 dorsocentral setae; katapisternal seta missing/indistinct. Original description in Suppl. material 1.

***Apotropina viduata* (Schiner, 1868)**

Fig. 23A–C

Ectropa viduata Schiner, 1868: 243.

Parahippelates fuscipes: Malloch, 1924: 330; Sabrosky 1980: 102 (synonymization).

Type locality and distribution. AUSTRALIA: New South Wales (Blue Mts; Hampton; Sydney, Collaroy).




Figure 22. *Apotropina taylori* (Malloch) holotype ♂ (K 359231) **A** habitus, lateral view **B** habitus, dorsal view **C** head, anterior view **D** specimen labels.




Figure 23. *Apotropina viduata* (Schiner) holotype ♀ NHMW-ZOO-DIP-0000753 **A** habitus, lateral view **B** habitus, dorsal view **C** specimen labels.

Examined material. Holotype ♀ Label transcription: “Novara. R., Sydney; *viduata*, Alte Sammlung; *Ectropa viduata* Schiner.; TYPE *Ectropa viduata* Schiner; CHLOROPIDAE, *Apotropina viduata* (SCHINER), det. Sabrosky; NHMW-ZOO-DIP-0000753”; Deposited in the AMRI.

Taxonomic notes. This is a dark-bodied species with tawny legs and lighter colored head (Fig. 23A, B). Chaetotaxy as observed: 3 vibrissae; 3 proclinate interfrontal setae; 2 postpronotal setae; 2 scapular setae; 1+1 notopleural setae; biseriate acrostichal row that gets strong in medial of scutum; 1+3 dorsocentral setae; 1+1 supra-alar setae; postalar setae and intrapostalar setae present; scutellum with 2 discal setulae; katepisternal seta whitish, indistinct. The taxonomic history for this species is already detailed in the introduction section (See Sabrosky, 1980 for details). Original descriptions provided in Suppl. material 1.

***Apotropina bispinosa* (Becker, 1911)**

Fig. 24A–C

Oscinella bispinosa Becker, 1911: 152.

Oscinelloides bispinosa: Malloch 1940: 268.

Type locality and distribution. AUSTRALIA: Queensland (Weipa). PAPUA NEW GUINEA: Huon (Sattleberg), New Britain (Rabaul).

Taxonomic notes. This species has a problematic type series assignment. It was originally described from, and limited to, Papua New Guinea based on six specimens (Becker 1911). Malloch (1940) identified an additional female of *Apotropina bispinosa* from Papua New Guinea and transferred this species to his newly erected genus *Oscinelloides* Malloch, providing further morphological description of the species. *Oscinelloides* was subsequently synonymized with *Apotropina* (Sabrosky 1989). Thereafter, specimens from Australia were identified to the *A. bispinosa* (Forster, 1992). We were able to examine two specimens in MfN marked as types (see Fig. 24 for one imaged specimen; label transcription: “Sattleberg, Huon-Golf.; 547474; Typus; N.-Guinea, Biró 1899.; Sammlung Dr. Th. Becker; *Rhodesiella* sp., det J.W. Ismay 2002; <http://coll.mfn-berlin.de/u/5c8582>”) and found that both specimens have no dorsocentral setae except for the posterior one and have completely dark brown femora and tibiae. These features do not correspond to descriptions provided by Becker (1911) and Malloch (1940), where both state that *A. bispinosa* has two strong pairs of dorsocentral setae, lighter-colored yellowish legs with darkened apices on mid and hind femora as well as fore tibiae. In fact, the morphology of the examined specimens matches that of genus *Rhodesiella* Adams. Given that the two MfN specimens [547474 (5c8582 and 5c85bb)] do not correspond to the original description (see Suppl. material 1) despite belonging to the same locality, a detailed assessment on *A. bispinosa*’s type series should be done. With that, the designation of a lectotype would clarify this species identity. Furthermore, as we could not examine Forster’s (1992) Australian specimens, we refrain from adding *A. bispinosa* to the key.




Figure 24. *Apotropina bispinosa* (Malloch) syntype ♀ 57474 (5C8582) **A** habitus, lateral view **B** habitus, dorsal view **C** specimen labels.




Figure 25. illustrations for various *Apotropina* spp. **A** *A. aequalis* head, lateral view (from Becker 1911: Tafel I, Fig. 15) **B** *A. aequalis* hind tibial spur (from Malloch 1940: 271, Fig. 15) **C** *A. nudiseta* antennal arista (from Malloch 1940: 271, Fig. 17) **D** *A. nudiseta* hind tibial spur (from Malloch 1940: 271, Fig. 18) **E** *A. proxima* habitus, lateral view (from Rayment 1959: plate XXXIX, Fig. 1) **F** *A. proxima* thorax, dorsal view (from Rayment 1959: plate XXXIX, Fig. 2).

Acknowledgements

The authors would like to thank Simon Grove (Tasmanian Museum and Art Gallery) and Julie Sarna for contributing their in situ photos of live *Apotropina*, Natalie Tees, Matthew Shaw, and Alexssandro Camargo for photographing the images of the AMRI, SAMA and NHMW type specimens respectively, and Leshon Lee (National University of Singapore) for his help in sequencing the specimens.

Additional information

Conflict of interest

The authors have declared that no competing interests exist.

Ethical statement

No ethical statement was reported.

Funding

PRR benefits from the program CAPES-HUMBOLDT, grant no. 88881.512934/2020-01. AY benefits from the LKCNHM-RCC Research Fund.

Author contributions

Conceptualization: YA. Data curation: PRRR, JL, YA. Formal analysis: YA, PRRR. Funding acquisition: YA, PRRR. Investigation: PRRR, YA. Methodology: YA. Project administration: YA. Visualization: YA. Writing - original draft: PRRR. Writing - review and editing: PRRR, JL, YA.

Author ORCIDs

Yuchen Ang  <https://orcid.org/0000-0001-5889-018X>

James Lumbers  <https://orcid.org/0009-0007-4895-0936>

Paula R. Riccardi  <https://orcid.org/0000-0003-4850-7524>

Data availability

All of the data that support the findings of this study are available in the main text or Supplementary Information.

References

- Andersson H (1977) Taxonomic and phylogenetic studies on Chloropidae (Diptera) with special reference to Old World genera. *Entomologica Scandinavica* (Supplementum 8): 1–200.
- Becker T (1911) Chloropidae. Eine monographische Studie. III. Teil. Die indo-australische Region. *Annales Historico-Naturales Musei Nationalis Hungarici* 9: 35–170. [Tab I–II.] <https://doi.org/10.5962/bhl.title.9555>
- Curran CH (1930) Four new Diptera from Australia. *American Museum Novitates* 442: 1–4.
- Duda O (1934) Weitere neue und wenig bekannte orientalische und australische Chloropiden (Diptera) des Deutschen Entomologischen Instituts in Berlin-Dahlem. *Arbeiten über morphologische und taxonomische Entomologie aus Berlin-Dahlem* 1: 39–60.

- Forster PI (1992) Insects associated with the flowers of *Marsdenia cymulosa* Benth. (Asclepiadaceae) and their possible role in pollination. *Australian Entomological Magazine* 19(1): 45–47.
- GroveS (2019) iNaturalist observation. <https://www.inaturalist.org/observations/19347756>; <https://www.inaturalist.org/photos/29751169> [Accessed 5 June 2023]
- Ismay B, Ang Y (2019) First records of *Pseudogaurax* Malloch 1915 (Diptera: Chloropidae) from Singapore, with the description of two new species discovered with NGS barcodes. *The Raffles Bulletin of Zoology* 67: 412–420. <https://doi.org/10.26107/RBZ-2019-0033>
- Ismay JW, Ismay B, Deeming JC (2021) 96. Chloropidae. In: Kirk-Spriggs AH, Sinclair BJ (Eds) *Manual of Afrotropical Diptera. Volume 3. Brachycera – Cyclorrhapha, Excluding Calypttratae – Higher Diptera. Suricata* 8(3). South African National Biodiversity Institute, Pretoria, 2035–2114.
- Malloch JR (1923) Notes on Australian Diptera with descriptions, I. *Proceedings of the Linnean Society of New South Wales* 48: 601–622.
- Malloch JR (1924) Notes on Australian Diptera, III. *Proceedings of the Linnean Society of New South Wales* 49(3): 329–338.
- Malloch JR (1925) Notes on Australian Diptera, VI. *Proceedings of the Linnean Society of New South Wales* 50(2): 80–97.
- Malloch JR (1928) Notes on Australian Diptera, XIV. *Proceedings of the Linnean Society of New South Wales* 53: 295–309.
- Malloch JR (1936) Notes on Australian Diptera, XXXV. *Proceedings of the Linnean Society of New South Wales* 61: 10–26.
- Malloch JR (1940) Notes on Australian Diptera, XXXVIII. Family Chloropidae, Part II. *Proceedings of the Linnean Society of New South Wales* 65(3–4): 261–288.
- Mortelmans J, Hussin M, Lee L, Ang Y (2022) First records of snail-killing flies (Diptera: Sciomyzidae) from Singapore with notes on morphology and DNA sequencing. *Nature in Singapore* 15: e2022014. <https://doi.org/10.26107/NIS-2022-0014>
- Rayment T (1931) The furrow-bees and a fly. *The Victorian Naturalist* 47: 184–191.
- Rayment T (1935) *A Cluster of Bees: Sixty Essays on the Life-histories of Australian Bees, with specific descriptions of over 100 new species, and an introduction by Professor E. F. Phillips. D.Ph., Cornell University, U.S.A.* Endeavour Press, Sydney, 752 pp. [+ frontispiece + colour plates 40 & 66]
- Rayment T (1959) Hyperparasitism by a minute fly and the specific description of a new species. *Australian Zoologist* 12: 330–333. [pl. XXXIX]
- Riccardi PR, Baziar Z, Lamas CE (2018) New genus of the subfamily Oscinellinae from Brazil (Diptera: Chloropidae). *Zootaxa* 4438(2): 394–400. <https://doi.org/10.11646/zootaxa.4438.2.13>
- Richards OW (1973) The Sphaeroceridae (= Borboridae or Cypselidae; Diptera Cyclorrhapha) of the Australian Region. *Australian Journal of Zoology, supplementary Series* 22: 297–401. <https://doi.org/10.1071/AJZS022>
- Sabrosky CW (1955) Notes and descriptions of Australian Chloropidae (Diptera). *Proceedings of the Linnean Society of New South Wales, Sydney* 79: 182–192.
- Sabrosky CW (1980) Unexpected synonymy in Chloropidae, from the Family Ephydriidae (Diptera). *Australian Entomological Magazine* 6(6): 101–102.
- Sabrosky CW (1989) 100. Family Chloropidae. In: Evenhuis N (Ed.) *Catalog of the Diptera of the Australasian and Oceanian Regions*. Bishop Museum Press, Honolulu & EJ Brill, Leiden, 650–665.

- Sarna J (2020) iNaturalist observation. <https://www.inaturalist.org/observations/48861099>; <https://www.inaturalist.org/photos/77545425> [Accessed 5 June 2023]
- Schiner IR (1868) 1. Diptera. Reise der österreichischen Fregatte Novara um die Erde in den Jahren 1857, 1858, 1859 unter den Befehlen des Commodore B. von Wüllerstorff-Urbair. Zoologischer Theil. Zweiter Band. 1. Abtheilung, B: VI + 388 pp. [Kaiserlich-Königliche Hof- und Staatsdruckerei (In Commission bei Karl Gerolds's Sohn), Wien]
- Thomson CG (1869) 6. Diptera. Species novas descripsit C.G. Thomson. Pp. 443–614, Taf. IX in: Kongliga svenska fregatten Eugénies resa omkring jorden under befäl af C.A. Virgin. Åren 1851–1853. Vetenskapliga iakttagelser på H.M. Konung OSCAR Den Förstes befallning utgifna af K. Svenska Vetenskaps Akademien. Andra Delen [= Vol. 2]. Zoologi. 1. Insecta: V + 617 pp. [Tafl. I–IX; P.A. Norstedt & Söner, Stockholm]

Supplementary material 1

Original descriptions and relevant subsequent taxonomic notes for Australian *Apotropina* fauna

Authors: Yuchen Ang, James Lumbers, Paula R. Riccardi

Data type: pdf

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/zookeys.1187.108497.suppl1>