

Review of the leafhopper tribe Deltocephalini Dallas, 1870 (Hemiptera, Cicadellidae, Deltocephalinae) in Pakistan with description of a new species of *Paramesodes*

Hassan Naveed^{1,2}, Bismillah Shah³, Kamran Sohail⁴, Yalin Zhang⁵, Keping Chen^{1,2}

¹ School of Food and Biological Engineering, Jiangsu University, Zhenjiang 212013, China

² School of Life Sciences, Jiangsu University, Zhenjiang 212013, China

³ Department of Forestry Protection, School of Forestry and Biotechnology, Zhejiang A&F University, 666 Wusu Street, Linan, Hangzhou, Zhejiang 311300, China

⁴ Department of Entomology, The University of Agriculture, Peshawar 25100, Pakistan

⁵ Key Laboratory of Plant Protection Resources and Pest Management of the Ministry of Education, Entomological Museum, Northwest A&F University, Yangling, Shaanxi Province 712100, China

Corresponding authors: Yalin Zhang (yalinzh@nwsuaf.edu.cn); Keping Chen (kpchen@ujs.edu.cn)

Abstract

A review of the leafhopper tribe Deltocephalini in Pakistan is provided, including comprehensive illustrated keys to all genera and species, along with their diagnosis and distribution. A new species of *Paramesodes* is described, *P. dirensis* **sp. nov.**, which was discovered in the Upper Dir District of Khyber Pakhtunkhwa, Pakistan. A detailed description of this new species is provided together with photographs for visual reference. This tribe now has three genera and 13 species from Pakistan. The genus *Deltocephalus* Burmeister includes two species, and *Maiestas* Distant has nine species; *Paramesodes* Ishihara is now represented with two species.

Key words: Auchenorrhyncha, distribution, key, morphology

Academic editor: J. A. Pinedo-Escatel

Received: 30 July 2023

Accepted: 16 November 2023

Published: 13 December 2023

ZooBank: <https://zoobank.org/7B477F6B-7729-4C53-BC55-87870895D5AA>

Citation: Naveed H, Shah B, Sohail K, Zhang Y, Chen K (2023) Review of the leafhopper tribe Deltocephalini Dallas, 1870 (Hemiptera, Cicadellidae, Deltocephalinae) in Pakistan with description of a new species of *Paramesodes*. ZooKeys 1186: 207–219. <https://doi.org/10.3897/zookeys.1186.110266>

Copyright: © Hassan Naveed et al. This is an open access article distributed under terms of the Creative Commons Attribution License ([Attribution 4.0 International – CC BY 4.0](https://creativecommons.org/licenses/by/4.0/)).

Introduction

Leafhoppers of the tribe Deltocephalini can be identified by their small to moderate size, tapering or parallel-sided clypellus, narrow lorum, linear connective with closely appressed anterior arms, connective fused to the aedeagus, and dorsal sculpturing of the first valvula imbricate. This tribe includes 74 genera and over 600 species distributed worldwide (Duan et al. 2020; Zhang et al. 2023). Until now, 12 species of Deltocephalini have been known from Pakistan (Naveed et al. 2021; Shah et al. 2021).

The genus *Paramesodes* Ishihara, comprising 16 Old World species, is poorly represented in Pakistan, with only a single previous species, *P. lineaticollis* (Distant), known. Wilson (1983) revised the genus and recognized one species from Europe and the Middle East, one species from Africa, and nine species from

Asia. Five other species were subsequently included: *P. montanus* Rao, (1989) from India, *P. wilsoni* Rao & Ramakrishnan, (1990) from India, *P. iraniensis* Webb & Viraktamath, (2009) from Iran, and *P. menghaiensis* Li, Dai & Xing, (2011) and *P. cangshanae* Duan & Zhang, (2012) from China. *Paramesodes* is externally similar to *Paramesus* Fieber, *Exitianus* Ball, and *Ctenurellina* McKamey but they can be differentiated by having the forewing with the inner anteapical cell open basally (closed in *Paramesus* and *Ctenurella*), and the veins are always pale or white (usually brown in *Exitianus*). In the male genitalia the connective and aedeagus are fused (free in *Paramesus* and *Exitianus*), and the pygofer side has a dense covering of long, stout, pale macrosetae (2–6 dark or black macrosetae along apical margin in *Exitianus*) and the pygofer side has a large process (lower edge of pygofer pectinate in *Ctenurella*). This, together with the aedeagus, is the most important character for species identification (Duan and Zhang 2012).

In this paper, we provide a list of leafhoppers of the tribe Deltocephalini from Pakistan, illustrated keys to genera and species with their diagnostic characters, and a description and illustration of a new species, *P. dirensis* sp. nov.

Material and methods

All specimens were examined with a Leica ZOOM2000 stereomicroscope. Drawings were made using an Olympus drawing tube. Photos were taken by a ZEISS SteREO Discovery.V20 stereomicroscope equipped with a ZEISS AxiocamICc 5 camera that also provided measurements. Adobe Photoshop CS was used to compile photographs. Specimens from Pakistan are deposited in the various collections as indicated in the published records. Type specimens of the new species, examined and figured for this study, are deposited in the Entomological Museum, Northwest A&F University, Yangling, Shaanxi, China.

Taxonomy

Family Cicadellidae Latreille, 1802

Subfamily Deltocephalinae Dallas, 1870

Tribe Deltocephalini Dallas, 1870

Key to genera of Deltocephalini from Pakistan

- 1 Crown with broad black transverse submarginal band between eyes (Figs 1, 9); pygofer side with large process (Figs 3, 10) ***Paramesodes***
- Crown without black transverse band between eyes (Figs 12, 15, 18, 19); pygofer side without process **2**
- 2 Aedeagal shaft short, robust, strongly curved dorsally, gonopore apical (Figs 13, 16) ***Deltocephalus***
- Aedeagal shaft long, slightly curved dorsally, with gonopore indistinct (Figs 20–25, 27, 29) ***Maiestas***

Genus *Deltocephalus* Burmeister

Key to species of *Deltocephalus* from Pakistan (male)

- 1 Crown with six brown spots on anterior margin (Fig. 12); aedeagal shaft with shallow apical notch (Fig. 13) *D. vulgaris*
- Crown with single brown spot on anterior margin adjacent to eyes (Fig. 15); aedeagal shaft without apical notch (Fig. 16)..... *D. infirmus*

Deltocephalus vulgaris Dash & Viraktamath

Figs 12–14

Deltocephalus (Deltocephalus) vulgaris Dash & Viraktamath, 1998: 4, figs 1–11 (India); Zhang and Duan 2011: 3, fig. 3A–H (China); Naveed et al. 2019: 285, figs 1A, B, 3A–D (Pakistan).

Diagnosis. This species can easily be identified by the color pattern and the aedeagus with a shallow apical notch.

Distribution. China, India, Pakistan.

Deltocephalus infirmus Melichar

Figs 15–17

Deltocephalus infirmus Melichar, 1903: 203, pl. 5 fig. 11 (Sri Lanka); *Jassargus infirmus*: Ishihara 1961: 244, figs 53–58 (misidentification); *Deltocephalus infirmus*: Webb and Viraktamath 2009: 13, fig. 10; Naveed et al. 2019: 285, figs 1C, 3D–G (Pakistan).

Diagnosis. A similar species to *D. vulgaris* but differs by having the crown with only a single brown spot and the aedeagal shaft without an apical notch.

Distribution. India, Pakistan, Sri Lanka, Thailand.

Genus *Maiestas* Distant

Key to species of *Maiestas* from Pakistan (males)

- 1 Overall color dark brown; forewing with sub basal and subapical irregular white transverse band (Fig. 18)..... *M. albomaculata*
- Color not as above (Fig. 19) 2
- 2 Crown, face, and thorax with black patches (Fig. 19) *M. maculata*
- Crown, face, and thorax without black patches 3
- 3 Forewing with extra cross-veins, at least in clavus 4
- Forewing without extra cross-veins 5
- 4 Aedeagus with a large subapical ventral process (Fig. 20)..... *M. indica*
- Aedeagus with a short apical ventral process (Figs 21, 22)..... *M. pruthii*
- 5 Aedeagus with pair of short lateral processes (Fig. 23)..... *M. trispinosa*
- Aedeagus without lateral processes..... 6

- 6 Aedeagus in lateral view similar in width in distal half (Fig. 24) ... ***M. subviridis***
- Aedeagus in lateral view evenly tapered from base to apex **7**
- 7 Style apophysis broadest sub basally (Fig. 26); aedeagal shaft in lateral view not sinuate (Fig. 25)..... ***M. tareni***
- Style apophysis broadest at base (Fig. 28); aedeagal shaft in lateral view slightly sinuate (Fig. 27)..... ***M. sinuata***

***Maiestas albomaculata* (Dash & Viraktamath)**

Fig. 18

Deltocephalus (Recilia) albomaculatus Dash & Viraktamath, 1998: 12, figs 29–34 (India); *Maiestas albomaculata*: Webb and Viraktamath 2009: 21; *Maiestas albomaculata*: Naveed et al. 2019: 287, figs 1E–I, 3H, I (Pakistan); Shah et al. 2021: 403, fig. 1A–D (Pakistan).

Diagnosis. This species differs from other species of *Maiestas* in color and male genitalia, including the dorsolateral, laminate serrations of the aedeagal shaft.

Distribution. Pakistan, India.

***Maiestas indica* (Singh-Pruthi)**

Fig. 20

Allophleps indica Singh-Pruthi, 1936: 120, fig. 132; pl. 9 fig. 3 (Pakistan); *Allophleps delhiensis* Rao & Ramakrishnan, 1990: 111, figs 1–9 (India), synonymized by Dash and Viraktamath 1998: 35; *Deltocephalus (Recilia) indicus*: Dash and Viraktamath 1998: 35–36, fig. 305 (India); *Maiestas indica*: Webb and Viraktamath 2009: 21; Naveed et al. 2019: 287; Shah et al. 2021: 403, fig. 1E (Pakistan).

Diagnosis. This species can be identified by the aedeagus, which has a distinctive, large, subapical ventral process, the forewings which have accessory cross-veins, and the shorter head.

Distribution. Pakistan, India.

***Maiestas maculata* (Singh-Pruthi)**

Figs 19, 29

Cicadula maculata Singh-Pruthi, 1930: 58–59, figs 80, 81, pl. 5 fig. 2 (India); *Thamnotettix prabha* Singh-Pruthi, 1930: 62, figs 85, 86, pl. 5 figs 6, 6a (India), synonymized by Webb and Viraktamath 2009: 41; *Recilia prabha*: Ghauri 1980: 166–169, figs 1, 3–11; *Deltocephalus (Recilia) maculata*: Dash and Viraktamath 1998: 32, figs 260–269 (India); *Maiestas maculata*: Webb and Viraktamath 2009: 22; *Maiestas maculata*: Zhang and Duan 2011: 37–39, figs 33–35, pl. 4 fig. E, pl. 5 fig. P, pl. 6 fig. P (China); Shah et al. 2021: 404, fig. 2A–I (Pakistan).

Diagnosis. This species can be distinguished from other *Maiestas* species by its habitus, which has variable black patches on the head and thorax, and the shape of its aedeagus and style.

Distribution. China, India, Pakistan.

***Maiestas pruthii* (Metcalf)**

Figs 21, 22

Deltocephalus notatus Singh-Pruthi, 1936: 128–129, fig. 139, pl. 9 fig. 10 (Pakistan) (primary homonym: *Deltocephalus notatus* Melichar, 1896); *Deltocephalus pruthii* Metcalf, 1967: 1173 (nom. nov. pro *D. notatus* Singh-Pruthi, 1936); *Deltocephalus (Recilia) pruthii*: Dash and Viraktamath 1998: 22, 23, figs 150–158 (India); *Maiestas pruthii*: Webb and Viraktamath 2009: 20, new combination; Khatri and Webb 2010: 11, pl. 2a fig. 13, misidentification; Naveed et al. 2019: 286, fig. 2A–C (incorrectly cited as *M. subviridis*; H. Naveed pers. comm.), Fig. 3i, misidentification; Shah et al. 2021: 406, fig. 4F–L (Pakistan).

Diagnosis. This species has a relatively long, acute head, with a pair of inverted U-shaped markings basally, and forewings with extra cross veins. The identity of this species is based on the figures of Dash and Viraktamath (1998).

Distribution. India, Pakistan.

***Maiestas setosa* (Ahmed, Murtaza & Malik)**

Recilia setosa Ahmed et al., 1988: 412, fig. 2 (Pakistan); *Maiestas setosa*: Webb and Viraktamath 2009: 20; Naveed et al. 2019: 287; Shah et al. 2021: 406 (Pakistan).

Diagnosis. The identity of this species remains uncertain due to the limitations of the original description and the accompanying figures. Additionally, the type series from Karachi, which was indicated in the original account as deposited in the Zoological Museum of the University of Karachi (Ahmed et al. 1988), is unavailable (Khatri and Webb 2010: 11). Until the type material can be studied, pinpointing the precise classification of this species will be challenging.

Distribution. Pakistan.

***Maiestas sinuata* Shah & Duan**

Figs 27, 28

Maiestas sinuata Shah & Duan in Shah et al., 2021: 406, fig. 3A–H (Pakistan).

Diagnosis. This species differs in appearance of its habitus, presence of fine, apical setae on the subgenital plate, the style having a thicker apical process than other species, and the aedeagus lacking a ventrobasal “heel”.

Distribution. Pakistan.

***Maiestas subviridis* (Metcalf)**

Fig. 24

Stirellus subviridis Metcalf, 1946: 125; *Deltocephalus (Recilia) subviridis*: Dash and Viraktamath 1998: 24, figs 166–172 (India); *Maiestas subviridis*: Webb and Viraktamath 2009: 19, fig. 40; *Maiestas subviridis*: Khatri and Webb 2010: 11, pl. 2b, c, fig. 12 (Pakistan); Zhang and Duan 2011: 19, fig. 17, pl. 2 fig. H (China); Naveed et al. 2019: 287; Shah et al. 2021: 408, fig. 4A–E (Pakistan).

Diagnosis. This species can be distinguished by the rounded apex of the aedeagus which bears a very short apical spine.

Distribution. China, India, Pakistan, Pacific.

***Maiestas tareni* (Dash & Viraktamath)**

Figs 25, 26

Deltocephalus (Recilia) tareni Dash & Viraktamath, 1995: 74–76, figs 1–15; Dash and Viraktamath 1998: 16, figs 78–84 (India); *Maiestas tareni*: Webb and Viraktamath 2009: 22; Khatri and Webb 2010: 11, pl. 2d, fig. 11 (Pakistan); Zhang and Duan 2011: 20 (China); Naveed et al. 2019: 290, figs 2G–I, 3N, O; Shah et al. 2021: 408, fig. 5A–Z (Pakistan).

Diagnosis. This species can be identified by its relatively straight and stout style, apophysis with a serrated inner margin, and the aedeagus in lateral view evenly tapered from base to apex and relatively straight.

Distribution. China, India, Pakistan.

***Maiestas trispinosa* (Dash & Viraktamath)**

Fig. 23

Deltocephalus (Recilia) trispinosus Dash & Viraktamath, 1998: 35, figs 296–304 (India); *Maiestas trispinosa*: Webb and Viraktamath 2009: 38; *Maiestas trispinosa* Shah et al., 2021: 408, fig. 6A–I (Pakistan).

Diagnosis. This species can easily be distinguished from the others by the lateral, spine-like processes of the aedeagus.

Distribution. India, Pakistan.

Genus *Parasmesodes* Ishihara

Parasmesodes Ishihara, 1953: 45. Type species: *Athysanus albinervosus* Matsu-mura, 1902.

Distribution. Palearctic, Oriental, and Ethiopian regions.

Remarks. Previously, only 1 species was recorded from Pakistan. This study adds one more new species to the genus, bringing the total to two for the country.

Key to species of *Paramesodes* from Pakistan (males)

- 1 Pale yellowish species (Fig. 1); pygofer broadly rounded posteriorly, with process in lateral view straight distally, apex directed posteriorly (Fig. 3); aedeagal shaft recurved distally in lateral view (Fig. 7); constricted subapically in ventral view (Fig. 6).....***P. dirensis* sp. nov.**
- Dark yellowish species (Fig. 9); pygofer oval posteriorly, with process in lateral view directed dorsally distally (Fig. 10); aedeagal shaft evenly curved dorsally in lateral view (Fig. 10), not constricted preapically in ventral view (Fig. 11)***P. lineaticollis***

Paramesodes dirensis sp. nov.

<https://zoobank.org/158C20E7-3752-4E06-A4CE-AED5721D0A7D>

Figs 1–8

Description. **Length:** male 5.2–6.1 mm, female 6.1–6.4 mm. **Coloration:** pale, with brown markings (Fig. 1). Crown with broad, black, transverse submarginal band between eyes (Fig. 1). Face pale yellow, with brown, transverse striations on clypeus (Fig. 2). Pronotum with medial dark brown longitudinal marking as well as three brown longitudinal markings on each side (Fig. 1). Scutellum with median, longitudinal, dark brown markings and pale brown lateral markings (Fig. 1). Forewings with variable brown markings; veins prominent and white (Fig. 1). Legs pale, with brown markings.


Male genitalia: pygofer lobe broad basally, narrowing apically, forming a rounded-oval apex, with long, yellowish-brown spines extending beyond apical margin, a large process arising near medial dorsal margin and straight apically, surpassing pygofer lobe, without any bend (Figs 3, 4). Subgenital plates triangular; macrosetae uniseriate laterally (Fig. 5). Valve triangular (Fig. 5). Style as in Figure 5. Connective fused to aedeagus, with arms closely appressed distally (Figs 6, 7). Aedeagus tubular, tapering apically, recurved in lateral view, constricted preapically in ventral view; gonopore apical (Figs 6, 7).

Female. Same in appearance as male. Seventh sternum with lateral margins not extended, posterior margin with median projection, rounded (Fig. 8).

Materials examined. **Holotype** ♂, PAKISTAN: Khyber Pakhtunkhwa: Upper Dir, 35°9'55.89"N, 72°2'48.54"E, 1840 m, 24.07.2019, Hassan Naveed leg., sweep net. **Paratypes** 8♂, 5♀, same data as holotype.

Etymology. This species is named after type locality, the Upper Dir in Khyber Pakhtunkhwa.

Remarks. *Paramesodes dirensis* sp. nov. is similar to its congeners in general appearance, but it differs from those species in the combination of male genitalia features, i.e., the pygofer is oval posteriorly with a relatively straight process distally and the aedeagal shaft is distally recurved in lateral view and constricted preapically in ventral view. In the Wilson's (1983) key, the new species runs to couplet 7 along with *P. lineaticollis*.



Figures 1–8. *Paramesodes dirensis* sp. nov. **1** habitus, dorsal view **2** face **3** male pygofer, lateral view **4** male pygofer, dorsal view **5** subgenital plate, valve and style, dorsal view **6** connective and aedeagus, dorsal view **7** connective and aedeagus, lateral view **8** female abdominal tip and 7th sternum, dorsal view.

***Paramesodes lineaticollis* (Distant)**

Figs 9–11


Paramesodes lineaticollis (Distant, 1908: 294, *Paramesus*) (India); Wilson 1983: 21, 22, figs 23–29.

Paramesodes ishurdii Mahmood & Meher, 1973: 135 (Pakistan), synonymized by Wilson 1983: 21.


Materials examined. ♂, PAKISTAN: Khyber Pakhtunkhwa: Shinkiari, 34°28'19.1064"N, 73°16'14.3004"E, 22.07.2018, Bismillah Shah leg., sweep net.

Distribution. Bangladesh, China, India, Indonesia, Pakistan, Philippines, Taiwan, Turkey.

Diagnosis. The male pygofer processes is distinct, directed ventrally or postero-ventrally over basal half, and turned abruptly posteriorly immediately distad of its midlength. Mahmood and Meher (1973) reported this species as *P. ishurdii* for the first time from Pakistan, but that species was later synonymized by Wilson (1983).



Figures 9–11. *Paramesodes lineaticollis* 9 habitus, dorsal view 10 male pygofer, lateral view 11 connective and aedeagus, dorsal view.



Figures 12–29. 12–14 *Deltocephalus vulgaris* Dash & Viraktamath: 12 habitus, dorsal view 13 aedeagus and connective, dorsal view 14 aedeagus and connective, lateral view 15–17 *D. infirmus* (Melichar): 15 habitus, dorsal view 16 aedeagus and connective, dorsal view 17 aedeagus and connective, lateral view 18 *Maestas albomaculata* (Dash & Viraktamath) habitus, dorsal view 19 *M. maculata* (Singh-Pruthi) habitus, dorsal view 20 *M. indica* (Singh-Pruthi) aedeagus and connective, lateral view (after Dash & Viraktamath, 1998) 21, 22 *M. pruthii* (Metcalf): 21 aedeagus and connective, dorsal view (after Dash & Viraktamath, 1998) 22 apex of aedeagus, lateral view 23 *M. trispinosa* (Dash & Viraktamath) aedeagus and connective, dorsal view 24 *M. subviridis* (Metcalf) aedeagus, lateral view (after Khatri & Webb, 2010) 25, 26 *M. tareni* (Dash & Viraktamath): 25 aedeagus and connective, lateral view 26 subgenital plate, valve and styles, dorsal view 27, 28 *M. sinuata* Shah & Duan: 27 aedeagus and connective, lateral view 28 subgenital plate, valve and styles, dorsal view 29 *M. maculata* (Singh-Pruthi) aedeagus, lateral view.



Figures 30, 31. *Paramesodes lineaticollis* **30** subgenital plates, valve and styles, dorsal view **31** aedeagus and connective, lateral view.

Acknowledgements

We gratefully acknowledge Prof. C.A. Viraktamath (University of Agricultural Science, Bangalore, India), and Dr M. Webb (Natural History Museum, London, UK) for checking figures and confirming the new species. We are very grateful for John Richard Schrock (Emporia State University, USA) for revising the manuscript.

Additional information

Conflict of interest

The authors have declared that no competing interests exist.

Ethical statement

No ethical statement was reported.

Funding

This research was funded by the National Natural Science Foundation of China (31861143051, 31872425).

Author contributions

All authors have contributed equally.

Author ORCIDs

Hassan Naveed  <https://orcid.org/0000-0002-9232-6299>

Bismillah Shah  <https://orcid.org/0000-0002-8407-8627>

Kamran Sohail  <https://orcid.org/0000-0003-1625-1130>

Yalin Zhang  <https://orcid.org/0000-0002-1204-9181>

Keping Chen  <https://orcid.org/0000-0001-5254-2299>

Data availability

All of the data that support the findings of this study are available in the main text.

References

- Ahmed M, Murtaza B, Malik KF (1988) Some new aphrodine leafhoppers from grasslands of Karachi, Pakistan. *Pakistan Journal of Zoology* 20: 409–421.
- Dash PC, Viraktamath CA (1995) Two new species of grass feeding leafhopper genus *Deltocephalus* (*Recilia*) (Homoptera, Cicadellidae) from Orissa, India. *Hexapoda, Insecta Indica* 10: 1–59.
- Dash PC, Viraktamath CA (1998) A review of the Indian and Nepalese grass feeding leafhopper genus *Deltocephalus* (Homoptera, Cicadellidae) with description of new species. *Hexapoda, Insecta Indica* 10: 1–59.
- Distant WL (1908) Rhynchota-Homoptera. In: Bingham CT (Ed.) *The Fauna of British India, including Ceylon and Burma* 4, 501 pp.
- Duan YN, Zhang YL (2012) Review of the grass feeding leafhopper genus *Paramesodes* Ishihara (Hemiptera: Cicadellidae: Deltocephalinae: Deltocephalini) from China. *Zootaxa* 3151(1): 53–62. <https://doi.org/10.11646/zootaxa.3151.1.4>
- Duan YN, Gao Y, Dietrich CH (2020) *Pascoia*, a new Deltocephalini leafhopper genus (Hemiptera: Cicadellidae: Deltocephalinae) from Peru. *Zootaxa* 4801(1): 185–190. <https://doi.org/10.11646/zootaxa.4801.1.11>
- Ghuri MSK (1980) Illustrated redescription of two of Pruthi's species of Cicadelloidea from India. *Reichenbachia Staatliches Museum für Tierkunde in Dresden* 18: 165–171. https://www.zobodat.at/publikation_articles.php?id=314312
- Ishihara T (1953) A tentative check list of the superfamily Cicadelloidea of Japan (Homoptera). *Scientific Report of the Matsuyama Agricultural College* 11: 1–72.
- Ishihara T (1961) Cicadidae. *Insecta Japonica* 1(2): 1–36. [4 pls. Tokyo, Hokuryukan] [In Japanese, with English summary]
- Khatiri I, Webb MD (2010) The Deltocephalinae leafhoppers of Pakistan (Hemiptera, Cicadellidae). *Zootaxa* 2365(1): 1–47. <https://doi.org/10.11646/zootaxa.2365.1.1>
- Li ZZ, Dai RH, Xing JC (2011) Deltocephalinae from China (Hemiptera: Cicadellidae). *Popular Science Press, Beijing, China*, 336pp.
- Mahmood SH, Meher K (1973) New species of *Paramesodes* Ishihara from Pakistan (Hemiptera: Cicadellidae). *Transactions of the Shikoku Entomological Society* 11(4): 135–137. <https://agris.fao.org/agris-search/search.do?recordID=US201303275209>
- Matsumura S (1902) *Monographie der Jassinen Japans*. *Termesztudományi Füzetek Kiadja a Magyar Nemzeti Múzeum Budapest* 25: 353–404.
- Melichar L (1896) Cicadinen (Hemiptera-Homoptera) von Mittel-Europa. Berlin (Vol. i–xxvii), 364 pp. <https://doi.org/10.5962/bhl.title.8568>
- Melichar L (1903) *Homopteren Fauna von Ceylon*. Verlag von Felix L. Dames, Berlin, 248 pp.
- Metcalf ZP (1946) Fascicle IV, Fulgoroidea. Part 8. Dictyopharidae. *General Catalogue of the Hemiptera* 4(8): 1–246.
- Metcalf ZP (1967) *General Catalogue of the Homoptera*. Fascicle VI. Cicadelloidea. Part 10. Section I. Euscelidae. U.S. Department of Agriculture, Agriculture Research Service, 1077pp.
- Naveed H, Sohail K, Islam W, Zhang YL, Bu WJ (2019) A review of the leafhopper tribe Deltocephalini (Hemiptera: Cicadellidae: Deltocephalinae) from Pakistan. *Revista Chilena de Entomologia* 45(2): 283–292. <https://doi.org/10.35249/rche.45.2.19.16>
- Naveed H, Shah B, Khan BS, Cao C, Webb M, Zhang YL (2021) Checklist and keys to Deltocephalinae leafhoppers (Hemiptera, Cicadellidae) from Pakistan. *ZooKeys* 1078: 135–188. <https://doi.org/10.3897/zookeys.1078.47616>

- Rao KR (1989) Descriptions of some new leafhoppers (Homoptera: Cicadellidae) with notes on some synonymies and imperfectly known species from India. *Hexapoda, Insecta Indica* 1: 79–82.
- Rao RSV, Ramakrishnan U (1990) New species of *Aconurella* and *Paramesodes* (Homoptera: Cicadellidae) from India. *Journal of the Bombay Natural History Society* 87(2): 268–271.
- Shah B, Naveed H, Webb M, Duan YN (2021) Taxonomic review of the grassland leafhopper genus *Maestas* Distant (Hemiptera: Cicadellidae: Deltocephalinae: Deltocephalini) from Pakistan with description of a new species and two new records. *Zootaxa* 5060(3): 401–416. <https://doi.org/10.11646/zootaxa.5060.3.6>
- Singh-Pruthi H (1930) Studies on Indian Jassidae (Homoptera). Part I. Introductory and description of some new genera and species. *Memoirs of the Indian Museum* 11(1): 1–68. <http://faunaofindia.nic.in>
- Singh-Pruthi H (1936) Studies on Indian Jassidae (Homoptera). Part III. Descriptions of some new genera and species, with first records of some known species from India. *Memoirs of the Indian Museum* 11(3): 101–131. <http://faunaofindia.nic.in>
- Webb MD, Viraktamath CA (2009) Annotated checklist, generic key and new species of Old World Deltocephalini leafhoppers with nomenclatorial changes in the *Deltocephalus* group and other Deltocephalinae (Hemiptera, Auchenorrhyncha, Cicadellidae). *Zootaxa* 2163(1): 1–64. <https://doi.org/10.11646/zootaxa.2163.1.1>
- Wilson MR (1983) A revision of the genus *Paramesodes* Ishihara (Homoptera: Auchenorrhyncha: Cicadellidae) with description of eight new species. *Insect Systematics & Evolution* 14(1): 17–32. <https://doi.org/10.1163/187631283X00380>
- Zhang YL, Duan YN (2011) Review of the *Deltocephalus* group of leafhoppers (Hemiptera, Cicadellidae, Deltocephalinae) in China. *Zootaxa* 2870(1): 1–47. <https://doi.org/10.11646/zootaxa.2870.1.1>
- Zhang XY, Xu J, Shah B, Duan YN (2023) New taxa in the leafhopper tribe Deltocephalini (Hemiptera: Cicadellidae: Deltocephalinae) from China. *Zootaxa* 5271(3): 525–537. <https://doi.org/10.11646/zootaxa.5271.3.5>