

# Taxonomic and nomenclatural notes on *Geodiapria longiceps* Kieffer, 1911 (Hymenoptera, Diapriidae) and synonymy of the genus *Geodiapria* Kieffer, 1910

Jeremy Hübner<sup>1</sup>, Vasilisa G. Chemyreva<sup>2</sup>, David G. Notton<sup>3</sup>

<sup>1</sup> Zoologische Staatssammlung München, Muenchhausenstr. 21, 81247 Munich, Germany

<sup>2</sup> Zoological Institute, Russian Academy of Sciences, 1 Universitetskaya Emb., St. Petersburg 199034, Russia

<sup>3</sup> Department of Natural Sciences, National Museums Collection Centre, 242 West Granton Road, Granton, Edinburgh EH5 1JA, UK

Corresponding author: Jeremy Hübner ([huebner@snsb.de](mailto:huebner@snsb.de))

## Abstract

This paper reviews the status of *Geodiapria* and its nominotypical and only included species *G. longiceps*. *Geodiapria* was previously understood to be very similar to, and doubtfully separated from the genus *Basalys*. We use integrative taxonomy (morphology, DNA-barcoding, phylogenetic tree building) to show that the valid name for what was *G. longiceps* Kieffer, 1911 is now *Basalys rufocinctus* (Kieffer, 1911) and that *Geodiapria* is consequently a junior synonym of *Basalys* **syn. nov.** The following taxa are new synonyms of *B. rufocinctus*: *Loxotropa longiceps* Wasmann, 1909, **syn. nov.**, *G. longiceps* Kieffer, 1911, **syn. nov.**, *L. rufosignata* Kieffer, 1911, **syn. nov.** *Basalys rufocinctus* is newly reported from Corsica, Germany, Norway and Spain.

**Key words:** *Basalys rufocinctus*, DNA-barcoding, first record, integrative taxonomy, parasitoid wasp, species concepts



Academic editor: Norman Johnson

Received: 10 August 2023

Accepted: 4 October 2023

Published: 23 October 2023

ZooBank: <https://zoobank.org/1C5D67D6-A4DD-4D00-9767-57D55C2DB9E7>

**Citation:** Hübner J, Chemyreva VG, Notton DG (2023) Taxonomic and nomenclatural notes on *Geodiapria longiceps* Kieffer, 1911 (Hymenoptera, Diapriidae) and synonymy of the genus *Geodiapria* Kieffer, 1910. ZooKeys 1183: 1–11. <https://doi.org/10.3897/zookeys.1183.110952>

**Copyright:** © Jeremy Hübner et al.

This is an open access article distributed under terms of the Creative Commons Attribution License ([Attribution 4.0 International – CC BY 4.0](https://creativecommons.org/licenses/by/4.0/)).

## Introduction

Parasitoid wasps of the family Diapriidae are speciose and distributed worldwide, and while about 50% of its diversity is estimated to be unknown to science, there are few experts working on this family. Small size (c. 1–4 mm), wide distribution, cryptic diversity, sexual dimorphism, and previous poor taxonomy and lack of critical study of types are some of the problems researchers face when dealing with Diapriidae. The taxonomy of this group still therefore presents many interesting challenges. The status of the genus *Geodiapria* and its single included species *G. longiceps* Kieffer, 1911 has been a taxonomic problem for some time because of its close relation to *Basalys*, in particular species such as *B. rufocinctus* (Kieffer, 1911) with similar distinctive reddish flattened petiolar hairs. The question this paper seeks to resolve is whether or not *Geodiapria* is valid. *Geodiapria* was first described in a key by Kieffer (1910) who separated it from *Loxotropa* auctt. (now *Basalys* in part) and *Basalys sensu stricto* simply by the lack of a basal vein, adding later (Kieffer 1911a)

that the form of the head, longer than wide and a little wider in front than behind, was also distinctive. It was clearly similar to *Basalys* because Kieffer had previously considered the same material to be a *Loxotropa* auctt. (Wasmann, 1909). Kieffer (1911b) then described two species of *Loxotropa* auctt. with the same distinctive reddish flattened petiolar hairs: *L. rufosignata* said to have a head slightly longer than wide and reduced wings without distinct veins; and *L. rufocincta* with an almost square head and with an almost hyaline basal vein. Pschorn-Walcher (1957) examined the type of *G. longiceps* and considered *Geodiapria* to be very close to *Loxotropa* auctt., noting that the absence of the basal vein could be a consequence of wing reduction, but did not make a decision on the validity of *Geodiapria* because of lack of material. Since more material is now available, it is timely to reexamine the question of the validity of *Geodiapria* using an integrative approach combining morphotaxonomy and DNA barcoding (Ratnasingham and Hebert 2007). We examined 18 examples including types of four relevant nominal species, including *L. rufosignata* and *L. rufocincta*, and provide an up to date nomenclatural summary, presenting the first genetic results, including the DNA-barcode placing *Geodiapria* in its proper context.

## Material and methods

The specimens of *B. rufocinctus* used for the CO1 DNA barcoding were collected in July 2021 in the Dammbach Valley (Spessart Nature Park) on an orchard meadow, using a Malaise trap. The sequencing was conducted at Canadian Centre for DNA Barcoding (Guelph, Canada) using a voucher recovery protocol. Tree building was undertaken using IQ TREE (server version 1.6.12, Trifinopoulos et al. 2016) using the default settings with 1000 generations. MODELFINDER determined GTR+F+I+G4 to be the best fitting substitution model. The resulting tree was edited using FIGTREE v. 1.4.4 (Rambaut 2010) and INKSCAPE v. 1.1 (<https://inkscape.org/de/>).

### Repository acronyms:

<b>DNPC</b>	David Notton personal collection, United Kingdom
<b>MCSN</b>	Museo Civico di Storia Naturale "Giacomo Doria", Genoa, Italy
<b>MNHN</b>	Muséum national d'Histoire naturelle, Paris, France
<b>NHME</b>	Natural History Museum, Maastricht, Netherlands
<b>NHMUK</b>	Natural History Museum, London, United Kingdom
<b>SNSB-ZSM</b>	Bavarian State Collection, Munich, Germany

## Taxonomy

### *Basalys* Westwood, 1833

*Basalys* Westwood, 1833: 343. Type species *Basalys fumipennis* Westwood, 1833 by monotypy.

*Loxotropa* auctt. nec Förster, 1856.

*Geodiapria* Kieffer, 1910: 707, syn. nov. Type species *G. longiceps* Kieffer, 1911 by subsequent monotypy.

**Notes.** Other generic synonyms are omitted from the above list for simplicity. A diagnosis and detailed description of *Basalys* was given by Masner and García (2002), hence, only a brief diagnosis is given here. Further information on synonyms can be obtained from Johnson (1992).

**Diagnosis.** Small, smooth and shining wasps; head and mesosoma with long scattered hairs; antennal shelf usually distinctly prominent; female antenna 12-segmented, with strongly abrupt 3- or 4-segmented clava; male antenna 14-segmented with A4 distinctly modified; fore wing with submarginal vein slightly remote from fore margin of wing, costal vein absent, stigmal vein often moderately developed, basal vein always present in macropterous forms, straight, usually strongly pigmented, perpendicular to but never contiguous with submarginal vein.

**Remarks.** We discovered that the type species of *Geodiapria*, that is *G. longiceps*, is a *Basalys*, a synonym of *B. rufocinctus* (see below) and so *Geodiapria* becomes a junior synonym of *Basalys* syn. nov.

### ***Basalys rufocinctus* (Kieffer, 1911)**

*Loxotropa longiceps* Wasmann, 1909: 68, 172, syn. nov., preoccupied nec *B. longiceps* (Ashmead, 1893).

*Geodiapria longiceps* Kieffer, 1911a: 897, syn. nov., preoccupied nec *B. longiceps* (Ashmead, 1893).


*Loxotropa rufocincta* Kieffer, 1911b: 916, 939 takes precedence over *L. rufosignata* by first revisor action.

*Loxotropa rufosignata* Kieffer, 1911b: 914, syn. nov.

**BIN number.** BOLD\_BIN: [AEW6196](#) (Ratnasingham and Hebert 2007).

**Type material.** **Holotype** ♀ of *Loxotropa longiceps* labelled: "Allotype ♂ (!)/ Solenopsis imitatrix/ Wasmann, err. det.!: Holotype ♀/ *Geodiapria longiceps*/ Kieffer, 1911; *Loxotropa/ longiceps* n. sp./ ♀ Kieff.; 5.98. Exaet./ b. Solenopsis; Solenopsis m/ Kol. 293. sang [=colony #293 of *Formica sanguinea*]." (NHME) (Fig. 2). **Holotype** ♀ of *Geodiapria longiceps* - the same specimen as the holotype of *Loxotropa longiceps* q.v. **Holotype** ♀ of *Loxotropa rufosignata* labelled: "Is. Giglio/ IV.1902/ G. Doria; *Loxotropa/ rufosignata*; ♀" (MCSN) (Fig. 3). **Syntypes** 2♀ 3♂ of *Loxotropa rufocincta*: 2♀ labelled: "Holotype [sic – there is no original designation]; Bitche; *Loxotropa/ rufocincta*; Muséum Paris/ 1957/ coll. Kieffer. 2♂ labelled: *Loxotropa/ rufocincta*; Bitche; ♂; Allotype; Muséum Paris/ 1957/ coll. Kieffer. ♂ labelled: Paratype; Muséum Paris/ 1957/ coll. Kieffer; Bitche" (MNHN).

**Other material.** **DENMARK** • ♀; N. E. Zealand, Tisvilde Hegn; 56°02'N, 12°04'E; 4 May 1994; P.N. Buhl leg. (DNPC). **FRANCE** • ♂; Corsica, Corse du Sud, Bastelicaccia nr. Ajaccio; 41°55'N, 08°30'E; 14–21 Jun. 1996; C. Villemant leg.; Malaise trap, *Quercus suber* stand (DNPC) • ♀; Gard, Mont Ventoux, Malaucène; 44°13'N, 05°08'E; 1–8 Jul. 1997; C. Villemant leg.; maquis, *Quercus ilex* (DNPC) • ♂; same locality; 5–12 Aug. 1997; C. Villemant leg.; maquis, *Quercus ilex* (DNPC). **GERMANY** • ♀; Bavaria, Dammbach, Dammbachtal; 49°51'58"N, 09°19'30"E; 338 m a.s.l.; 16 Jul. 2021; J. Hübner leg.; nutrient poor grassland; ZSM-HYM-42434-G02 (BOLDSYSTEMS Process ID: [DTIII5299-22](#); GenBank



**Figure 1.** *Basalys rufocinctus* (Kieffer, 1911) ♀: **A** habitus, dorsal view **B** habitus, lateral view **C** wing with reduced venation (arrow) **D** close-up of petiole.



**Figure 2.** Holotype ♀ of *Loxotropa longiceps* (Wasmann, 1909), the same specimen is also the holotype ♀ of *Geodiapria longiceps* Kieffer, 1911: **A** habitus, lateral view **B** labels.

accession ID: [OR450821](#)) (SNSB-ZSM) • ♀ same locality; 16 Jul. 2021; J. Hübner leg.; nutrient poor grassland; ZSM-HYM-42433-H11 (BOLDSYSTEMS Process ID: [DTIII5225-22](#); GenBank accession ID: [OR450822](#)) (DNPC). **NORWAY** • ♀; Onsøy, Hankø Bloksberg, EIS 20, Ø; 3–29 Jun. 1995; O. Hanssen & J.I.I. Båtvik leg.; pitfall trap (DNPC). **SPAIN** • ♀; Granada, Calahonda; Jul. 1987; L. Lockett leg.; Malaise trap, (DNPC) • ♀; Granada, Sierra Nevada; 1600 m a.s.l.; 10 Apr. 1959; C. Besuchet leg. (NHMUK). **UNITED KINGDOM** • ♀; Cheshire, Abbots Moss; 53°12'27"N, 02°36'23"W; 12 Oct. 1990; D.G. Notton leg.; swept, stream (DNPC) • 3♀; Norfolk, Santon Downham; 52°27'45"N, 00°40'29"E; 15 Aug. 1984; J. Field leg.; Malaise trap, heath with *Betula* and *Pinus* (DNPC) • 1♂; same locality; 18–25 Aug. 1983; J. Field leg. (DNPC).

**Diagnosis. Female** Head elongate, rounded, about 1.2 times as long as wide; frons without angles or teeth; antenna 12-segmented with abrupt 3-segmented clava; A11 transverse in lateral view, as long as wide in dorsal view; A6–A9 transverse in lateral view (Fig. 1A); mesonotum and scutellum slightly convex in longer winged individuals, almost flat in shorter winged individuals (Fig. 1B), anterior pronotum with a ruff of whitish setae; anterior scutellar pit small and transverse, less than one third the width of the scutellum; propodeum with medial keel slightly raised anteriorly, less so in short winged individuals; fore wing variable in length, at most extending well beyond apex of gaster, at least reaching anterior margin of petiole; basal vein present in longer winged individuals although hard to see as it is fine and barely pigmented, absent in shorter winged individuals; femora of all legs broadened medially, fore femora 2.2–2.3 times as long as wide in lateral view, with sharp keel ventrally; petiole densely covered dorsally and laterally with long orange flattened setae (Fig. 1D); basal margin of large tergite with two whitish hair tufts more or less concealed under petiolar setae; disc of large tergite normally bare, although the shortest winged individuals, e.g. the type of *L. rufosignata*, may have some long setae. **Male** As for female except antenna 14-segmented with A4 expanded posteriorly subtriangular with a fine flange; A5 elongate, flagellar segments becoming shorter towards apex, A13 more or less quadrate; fore wing variable in length at least reaching apex of gaster, at most extending well beyond it; basal vein present, fine, barely pigmented; femora slightly less broadened than female. Body length 1.3–2.2 mm (♀); 1.5–2.4 mm (♂).




Figure 3. Holotype ♀ of *Loxotropa rufosignata* Kieffer, 1911: **A** habitus, dorsal view **B** labels.

**Distribution.** Czechia (Macek 1989 as *B. rufocincta* [sic]); Denmark (Buhl 1998 as *B. rufocincta* [sic]) confirmed here; France - mainland France (Kieffer 1911b as *L. rufocincta*) confirmed here; France - Corsica (new record); Germany (new record); Italy (Kieffer 1911b as *L. rufosignata*); Netherlands (Wasmann 1909 as *L. longiceps*); Norway (new record); Spain (new record); Sweden (Hedqvist 2007 as *B. rufocincta* [sic]); United Kingdom (Nixon 1980 as *B. rufocincta* [sic]) confirmed here.

**Biology.** Host unknown. *Basalys rufocinctus* has previously been considered to be a myrmecophile but the evidence is weak. Of all the specimens we have seen only one, Wasmann's, was found in an ant nest, in a mixed colony of *Solenopsis fugax* and *Formica sanguinea*, and may have entered the nest by accident. Wasmann provided no ethological observations to demonstrate myrmecophily and the species has no obvious morphological adaptation for myrmecophily when compared to other *Basalys*.

**Remarks.** From the extensive material examined we recognised only one taxon, diagnosed above, and with more variation than previously understood. Most importantly we found that the head was always elongate when seen from above, also significant variation in fore wing length, and expression of the basal vein which was present and weakly pigmented in longer winged individuals, becoming hyaline and then altogether absent in shorter winged individuals. This taxon is therefore a *Basalys* since there is no significant morphological difference: some other species of *Basalys* are known to have elongate heads, also some other *Basalys* have the basal vein absent in short-winged individuals. Based on our examination of the type specimens we consider all four nominal species above, including *Geodiapria longiceps*, belong to this taxon.

Further support for the generic placement of *B. rufocinctus* is based on genetic analyses. A representative ML tree (Appendix 1; *Idiotype maritima* (Haliday, 1833) as outgroup, 1000 generations) with 76 Diapriini specimens shows *B. rufocinctus* nested within a *Basalys* clade (Appendix 1). The obtained sequences are publicly available on the BOLDSYSTEMS platform (Ratnasingham and Hebert 2007).

Some nomenclatural notes are necessary:

1. We differ from some authors in recognising *Loxotropa longiceps* as a nominal species separate from, and not just a combination of, *Geodiapria longiceps*. *Loxotropa longiceps* is available from Wasmann's (1909) paper

where the name is first used. The name is made available by indication (ICZN 1999: Code art. 12.2.1) since Wasmann refers to his description (Wasmann, 1899) of a specimen previously misidentified as a male of *Solenopsia imitatrix* Wasmann, 1899. Although Wasmann attributes the name to Kieffer, the author of the name is actually Wasmann because he was responsible for publishing the name and writing the prior description (ICZN 1999: Code art. 50.1). The oldest available name for the taxon is thus *L. longiceps* Wasmann, 1909.

2. As *L. longiceps* is transferred to *Basalys* it becomes a secondary junior homonym of *B. longiceps* (Ashmead, 1893) so is invalid.
3. The next oldest available name is *G. longiceps* described as new by Kieffer (1911a). The date of publication is early 1911: evidence comes from the NHMUK copy which has a library stamp 25 Feb. 1911, and the page bound into the end of vol. 10 of *Species des Hyménoptères d'Europe et d'Algérie* which says 1 Mar. 1911.
4. As *G. longiceps* is transferred to *Basalys* it becomes a secondary junior homonym of *B. longiceps* (Ashmead, 1893) so is invalid.
5. The next oldest available names are *L. rufosignata* Kieffer, 1911b and *L. rufocincta* Kieffer, 1911b which were published simultaneously in mid-1911: the page bound into the end of vol. 10 of *Species des Hyménoptères d'Europe et d'Algérie* says 1 Jun. 1911.
6. Since the only two remaining potentially valid names are published simultaneously, we here make a first revisor action to determine precedence thus: *L. rufocincta* has precedence over *L. rufosignata*. We have chosen *L. rufocincta* because this is the more widely used name.
7. *L. longiceps*, *G. longiceps* and *L. rufosignata* are all new synonyms of *L. rufocincta*.
8. The valid name is thus *Basalys rufocinctus*, a combination first recognised by Nixon (1980).
9. Despite previous misspellings, when in combination with *Basalys*, the correct spelling of the species epithet is *rufocinctus*; the gender of *Basalys* is masculine (Notton (2014)).

## Acknowledgements

We are grateful to: Claire Villemant (MNHN) for the opportunity to examine the type material of *L. rufocincta*; Paul Beuk (NHME) for providing images of the type of *G. longiceps*; Maria Tavano (MCSN) for providing the opportunity to examine the type of *L. rufosignata*; Joseph Monks (NHMUK) arranged for access to the collections there; and people who donated specimens used in the study Peter N. Buhl, Claire Villemant, and Jeremy Field.

## Additional information

### Conflict of interest

The authors have declared that no competing interests exist.

### Ethical statement

No ethical statement was reported.

## Funding

This work was partly supported by Russian State Research (project 122031100272-3). Our work is part of the German Barcode of Life III: Dark Taxa project and was funded by the German Federal Ministry of Education and Research (FKZ 16LI1901B).

## Author contributions

Conceptualization and methodology, all authors; resources, all authors; writing – original draft, all authors; writing – review and editing, all authors; supervision, David Notton; funding acquisition, Jeremy Hübner and Vasilisa Chemyreva. All authors have read and agreed to the published version of the manuscript.

## Author ORCIDs

Jeremy Hübner  <https://orcid.org/0009-0007-5624-8573>

Vasilisa G. Chemyreva  <https://orcid.org/0000-0002-6547-6259>

David G. Notton  <https://orcid.org/0000-0002-8933-7915>

## Data availability


All of the data that support the findings of this study are available in the main text or Supplementary Information.

## References

- Buhl PN (1998) Proctotrupeidea (Hymenoptera) from different habitats at the Mols Laboratory area, Jutland, Denmark. *Natura Jutlandica* 23(7): 79–90.
- Hedqvist K-J (2007) A provincial catalogue of Swedish Proctotrupeidea. *Entomologisk Tidskrift* 128(3): 113–126. InkScape version 1.1. <https://inkscape.org/de/>
- International Commission on Zoological Nomenclature (1999) International Code of Zoological Nomenclature (4<sup>th</sup> edition). International Trust for Zoological Nomenclature, The Natural History Museum, London. <https://www.iczn.org/the-code/the-code-online/>
- Johnson N (1992) Catalog of world species of Proctotrupeidea, exclusive of Platygasteridae (Hymenoptera). *Memoirs of the American Entomological Institute* 51: 1–825.
- Kieffer JJ (1910) Proctotrypidæ. In: André E (Ed.) *Species des Hyménoptères d'Europe et d'Algérie*. Vol. 10. Hermann et Fils, Paris, 593–752.
- Kieffer JJ (1911a) Proctotrypidæ. In: André E (Ed.) *Species des Hyménoptères d'Europe et d'Algérie*. Vol. 10. Hermann et Fils, Paris, 753–912.
- Kieffer JJ (1911b) Proctotrypidæ. In: André E (Ed.) *Species des Hyménoptères d'Europe et d'Algérie*. Vol. 10. Hermann et Fils, Paris, 913–1015.
- Macek J (1989) Proctotrupeidea, Ceraphronoidea. In: Šedivý J (Ed.) *Enumeratio insectorum Bohemoslovakiae: Check List of Czechoslovak Insects III (Hymenoptera)*. *Acta Faunistica Entomologica Musei Nationalis Pragae*, 135–142.
- Masner L, García JL (2002) The genera of Diapriinae (Hymenoptera: Diapriidae) in the New World. *Bulletin of the American Museum of Natural History* 268: 1–138. [https://doi.org/10.1206/0003-0090\(2002\)268<0001:TGODHD>2.0.CO;2](https://doi.org/10.1206/0003-0090(2002)268<0001:TGODHD>2.0.CO;2)
- Nixon GEJ (1980) Diapriidae (Diapriinae) Hymenoptera, Proctotrupeidea. *Handbooks for the Identification of British Insects* 8(3di): 1–55.
- Notton DG (2014) A catalogue of the types of Diapriinae (Hymenoptera, Diapriidae) at the Natural History Museum, London. *European Journal of Taxonomy* 75(75): 1–123. <https://doi.org/10.5852/ejt.2014.75>

- Piel WH, Chan L, Dominus MJ, Ruan J, Vos RA, Tannen V (2009) TreeBASE v. 2: A Database of Phylogenetic Knowledge. [In: e-BioSphere 2009] <http://purl.org/phylo/treebase/phylo/study/TB2:S30685>
- Pschorn-Walcher H (1957) Zur Kenntnis der Diapriinae (Proctotrupeoidea) der Wasmann-Sammlung. Mitteilungen der Schweizerische Entomologische Gesellschaft 30(1): 73–88.
- Rambaut A (2010) FigTree v1.3.1. Institute of Evolutionary Biology, University of Edinburgh, Edinburgh. <http://tree.bio.ed.ac.uk/software/figtree/>
- Ratnasingham S, Hebert PD (2007) BOLD: The Barcode of Life Data System (<http://www.barcodinglife.org>). Molecular Ecology Notes 7(3): 355–364. <https://doi.org/10.1111/j.1471-8286.2007.01678.x>
- Trifinopoulos J, Nguyen LT, von Haeseler A, Minh BQ (2016) W-IQ-TREE: A fast online phylogenetic tool for maximum likelihood analysis. Nucleic Acids Research 44(W1): W232–W235. <https://doi.org/10.1093/nar/gkw256>
- Wasmann E (1899) Die psychischen Fähigkeiten der Ameisen. Zoologica (Stuttgart) 11(26): 1–132.
- Wasmann E (1909) Die psychischen Fähigkeiten der Ameisen. Mit einem Ausblick auf die vergleichende Tierpsychologie, Zweite bedeutend vermehrte Auflage, Zoologica 11 (26): i–xi, 1–190. [5 pls]

## Appendix 1



**Figure A1.** Maximum-likelihood tree of 76 Diapriini specimens, with *Idiotypa maritima* as outgroup. The different genera are color-coded, the numbers on the nodes represent the bootstrap values. Files are openly accessible online at TREEBASE (Piel et al. 2009; <http://purl.org/phylo/treebase/phyloWS/study/TB2:S30685>).

## Supplementary material 1

### Supplementary data

Authors: Jeremy Hübner, Vasilisa G. Chemyreva, David G. Notton

Data type: xlsx

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/zookeys.1183.110952.suppl1>