

Description of a new species of the *Petrolisthes galathinus* complex from the Caribbean Sea, and resurrection of *Petrolisthes occidentalis* from the East Pacific (Crustacea, Anomura, Porcellanidae)

Alexandra Hiller¹, Bernd Werding²

¹ Smithsonian Tropical Research Institute, Apartado 0843–03092, Panama, Panama

² Senckenberg Research Institute and Natural History Museum, Frankfurt am Main, Germany

Corresponding author: Alexandra Hiller (hillera@si.edu)

Abstract

The *Petrolisthes galathinus* complex currently consists of six American species distributed in the West Atlantic, including the amphi-American *P. galathinus*. All species in the complex are similar in their adult morphology but differ in colour, size, larval morphology, and shape of the adult sternal plate. The West Atlantic species have different geographic ranges, which overlap in the southern Caribbean. Previously published molecular data support the monophyly of the complex, and the reciprocal monophyly of each described species and further clades corresponding to different colour morphs. Here, the morph *P. caribensis* “Blue” is described as *Petrolisthes coeruleus* **sp. nov.**, and *Petrolisthes occidentalis* is formally resurrected for the Pacific individuals of *P. galathinus*. By adding these two species to the *P. galathinus* complex, this now consists of eight species. Colour illustrations of all species and colour morphs are provided and their geographic distributions and ecological ranges are discussed and updated.

Key words: Caribbean, colour morphs, ecological range, geographical range, *Petrolisthes coeruleus* sp. nov., *Petrolisthes occidentalis*

Academic editor: Célio Magalhães

Received: 4 September 2023

Accepted: 25 January 2024

Published: 16 February 2024

ZooBank: <https://zoobank.org/0E5EFE30-CAFF-4C64-A441-60B5E0E57616>

Citation: Hiller A, Werding B (2024) Description of a new species of the *Petrolisthes galathinus* complex from the Caribbean Sea, and resurrection of *Petrolisthes occidentalis* from the East Pacific (Crustacea, Anomura, Porcellanidae). ZooKeys 1191: 391–407. <https://doi.org/10.3897/zookeys.1191.111570>

Copyright: © Alexandra Hiller & Bernd Werding. This is an open access article distributed under terms of the Creative Commons Attribution License ([Attribution 4.0 International – CC BY 4.0](https://creativecommons.org/licenses/by/4.0/)).

Introduction

Petrolisthes galathinus (Bosc, 1802) is an American species, described as *Porcellana galathina* because of the transverse piliferous ridges covering carapace and extremities resembling those of the galatheid genus *Galathea* (Fig. 1). For more than two centuries the species remained subject of taxonomic conflict for three main reasons: Bosc’s ambiguous description failed to mention a type locality, which was declared as unknown (“on ignore son pays natal”; Bosc 1802: 233), the drawing depicting his holotype specimen consists of a crude sketch (see Fig. 2), and type material does not seem to be traceable. This ambiguous description prompted a labyrinthic path to recreate the morphology and origin of Bosc’s *P. galathinus*. Gibbes (1854), amongst others, criticized Bosc’s drawing because it exhibited “dots instead of stripes”, a pattern which did not correspond to observations by earlier authors (e.g., Benedict 1901). Gibbes’ remarks also alluded to different colour patterns, morphologically matching




Figure 1. Scanning Electron Microscopy (SEM) image of **a** carapace and **b** right cheliped of *Petrolisthes caribensis* Werding, a member of the *Petrolisthes galathinus* complex. Scale bars: 0.2 cm.



Figure 2. *Petrolisthes galathinus* (Bosc), original drawing by Bosc (1802: pl. 6, fig. 2).

Bosc's description. Based on specimens from Puerto Rico, Benedict (1901) described two colour varieties, one with purple stripes matching Gibbes' (1854) specimens, and another with a double white cross on the carapace. Decades later, Rickner (1975) emphasised the considerable colour variation of speci-

mens from the eastern coast of Mexico, and Williams (1984) reconstructed the history of different colour forms described by previous authors.

Regarding the possible type locality of *P. galathinus*, Latreille (1803: 76) declared the Antilles as the original locality of the species: “Elle se trouve aux Antilles”. Later mentions further referred to its provenance: Desmarest (1825) mentioned Georgia and Florida, and Gray (1831) reported specimens from the British Museum as coming from “North America”. Gibbes (1850) described *Porcellana sexspinosa*, and in a later study (Gibbes 1854), based on own collections from Key West, Florida, concluded that his species was synonymous with Bosc’s, and that it also occurred further north, up to South Carolina. In this study Gibbes (1854) reviewed the literature, acknowledging two western Atlantic species with the typical transverse piliferous rugae: *Porcella galathina* Bosc from the southern Atlantic coast of the United States, and *Porcellana boscii*? Savigny sensu Dana (1852) from Brazil. Gibbes concluded that the specimens from Brazil were different from Savigny’s *P. boscii* Audouin, 1826 from Egypt (de Savigny 1809) and described *Porcellana danae* Gibbes, 1854. *Porcellana boscii* was only listed by de Savigny (1809) but was later described by Audouin (1826). The species is currently accepted as *Petrolisthes boscii* (Audouin) from the Indo-West Pacific (see Werdning and Hiller 2007).

In the decades following Bosc’s description, West Atlantic species of *Petrolisthes* with piliferous transverse ridges on carapace and extremities were reported as *P. galathinus* from localities throughout the Caribbean and southwards to Brazil. The eastern Pacific individuals, morphologically matching *P. galathinus*, were described as *Petrolisthes occidentalis* Stimpson, 1859, based on specimens collected in Panama. However, Boone (1931) and Schmitt (1935) reported specimens from the East Pacific under the name *P. galathinus*.

In her study on the Porcellanidae of the western North Atlantic, Haig (1956) extensively reviewed the literature and synonymy of *Petrolisthes galathinus*, concluding that West Atlantic and East Pacific specimens should be considered as one species. As a consequence, all subsequent authors (e.g. Chace 1956; Haig 1962, 1966; Fausto-Filho 1968; Gore 1970, 1974; Rouse 1970; Coelho and Ramos 1972; Rickner 1975; Gore and Abele 1976; Hiller et al. 2004) treated *P. galathinus* as a single widespread species distributed throughout the western Atlantic from North Carolina (U.S.A.) to Santa Catarina (Brazil), and in the eastern Pacific from Jalisco (Mexico) to Ecuador (Hiller et al. 2004).

Werdning (1977) recognised different species in the Colombian Caribbean, later describing *P. rosariensis* Werdning, 1982 (Fig. 3a). Werdning (1983) concluded that *P. galathinus* comprised a complex of several species and described *P. columbiensis* Werdning (1983) (Fig. 3b) and *P. caribensis* Werdning (1983) (Fig. 3c, d), only distinguishable from *P. galathinus* by their different colouration and two discrete morphological characters (Table 1). Later collections in the Colombian Caribbean revealed the presence of other colour types, described as *P. sanmartini* Werdning & Hiller (2002) (Fig. 4a) and *P. bolivarensis*, Werdning & Kraus (2003) (Fig. 4b), reaching a total of six species, including *P. galathinus*. Hiller et al. (2006) reconstructed the phylogenetic history of the complex based on mitochondrial DNA sequences and concluded that all species were reciprocally monophyletic, and that further clades, taxonomically matching *P. galathinus* (Fig. 5a–e), supported the presence of new species.

Here, we describe *P. coeruleus* sp. nov. (Figs 6a–c, 7, 8a), which corresponds to the “Blue” morph revealed in the phylogeny by Hiller et al. (2006),




Figure 3. Dorsal view of **a** *Petrolisthes rosariensis*, male, Islas del Rosario, Colombian Caribbean **b** *P. columbiensis*, female, Islas del Rosario, Colombian Caribbean **c** *P. caribensis*, male, Islas del Rosario, Colombian Caribbean **d** *P. caribensis*, Roatán, Honduras, photo courtesy of M. Charteris. Scale bars: 0.6 cm (**a**); 0.8 cm (**b**); 0.3 cm (**c**).

Table 1. Morphotypes and distinguishing characters of the *Petrolisthes galathinus* complex.

Species	Number of epibranchial spines	Number of spines on inner border of dactylus of all walking legs	Maximum size of male adults (carapace width in mm)
<i>P. bolivarensis</i> Werdning & Kraus	1	3	> 15.0
<i>P. caribensis</i> Werdning	1	4	9.4
<i>P. coeruleus</i> sp. nov.	1	4	12.4
<i>P. columbiensis</i> Werdning	2	4	7.2
<i>P. galathinus</i> (Bosc) "Stripes-Spots"	1	3	> 14.0
<i>P. galathinus</i> (Bosc) "White Teeth"	1	3	> 15.0
<i>P. occidentalis</i> Stimpson	1	3	> 17.0
<i>P. rosariensis</i> Werdning	2	4	6.0
<i>P. sanmartini</i> Werdning & Hiller	1	5	6.5

and resurrect *P. occidentalis* Stimpson, 1859 (Fig. 5a) for the Eastern Pacific individuals of *P. galathinus*. We update the ecological and geographic information of all members of the complex and address the most plausible identity of Bosc's *P. galathinus*.



Figure 4. Dorsal view of **a** *Petrolisthes sanmartini*, male, Islas del Rosario, Colombian Caribbean **b** *P. bolivarensis*, male, Islas del Rosario, Colombian Caribbean. Scale bars: 0.6 cm (**a**); 1.7 cm (**b**).




Figure 5. Dorsal view of colour morphs within *Petrolisthes galathinus*, as designated by Hiller et al. (2006) **a** East Pacific morph, here resurrected as *P. occidentalis* Stimpson, male, Naos Island, Panamanian Pacific **b** *P. galathinus* "Spots", male, Islas del Rosario, Colombian Caribbean **c** *P. galathinus* "White Teeth", female (ov), Isla Cubagua, Venezuela **d** "White Teeth", Roatán, Honduras, photo courtesy of M. Charteris **e** *P. galathinus* "Stripes", male, Gulf of Morrosquillo, Colombian Caribbean. Scale bars: 1.65 cm (**a**); 0.71 cm (**b**); 1.36 cm (**c**); 1.35 cm (**e**).

Material and methods

Material of *Petrolisthes coeruleus* sp. nov. collected in Belize and the Panamanian and Colombian Caribbean, and of *P. occidentalis* collected in the Panamanian East Pacific, was used for morphological examination. Type material of the new species was deposited in the Senckenberg Naturmuseum Frankfurt (SMF), Germany, and the Museo de Peces de Agua Dulce e Invertebrados (MUPADI) de la Universidad Autónoma de Chiriquí, Panamá (UNACHI). Type specimens of *P. occidentalis* are deposited in the Museum of Comparative Zoology (MCZ), Harvard University, USA. Additional material of the new species and of *P. occidentalis* was deposited in the collections of the MUPADI in Panama. We updated the geographic and ecological ranges of all species of the *P. galathinus* complex based on new records of material collected by the authors and material found in the Florida Museum of Natural History (FMNH). Measurements of specimens are given in mm and correspond to carapace length, followed by carapace width.

Results

Systematic account

Family Porcellanidae Haworth, 1825: 184.

Genus *Petrolisthes* Stimpson, 1858: 227.

***Petrolisthes coeruleus* sp. nov.**

<https://zoobank.org/1E5A0FF9-CD71-425D-87BB-439771E73B59>

Figs 6a–c, 7, 8a

Petrolisthes galathinus Werding, 1982: 443 (part, Islas del Rosario).

Petrolisthes caribensis “Blue”: Hiller et al. 2006: 552.

Type material examined. Holotype: Male, MUPADI–Crus–14, West Atlantic, Panamá, Colón, Punta Galeta, 9°24.161'N, 79°51.634'W; in *Porites porites* (Pallas, 1766), 0.5 m, leg. A Hiller, 12 Apr. 2021; 9.53 × 8.98 mm; female (ov) 8.83 × 7.78 mm; female 7.65 × 7.22 mm. **Paratypes:** MUPADI–Crus–02–16, same data as holotype; female (ov) 8.83 × 7.78 mm; female 7.65 × 7.22 mm. SMF57499, West Atlantic, Colombia, Bolívar, Islas del Rosario, Isla San Martín de Pajarales, 10°10.637'N, 75°46.234'W; in coral gravel, 1–2 m, leg. B. Werding, Sep. 2001; female (ov) 7.0 × 6.8 mm; male 7.3 × 7.0 mm; male 7.7 × 7.6 mm; male 5.4 × 5.2 mm; female (ov) 7.7 × 7.4 mm; female (ov) 7.6 × 7.6 mm; male 6.1 × 5.8 mm; female (ov) 7.5 × 7.7 mm; female (ov) 6.4 × 6.5 mm; male 5.4 × 5.3 mm; female 6.1 × 6.2 mm.

Other material examined (personal collections by the authors). West Atlantic, Belize, Carrie Bow Cay, 16°48.188'N, 88°5.067'W; under blocks of dead elkhorn coral, 1–2 m, leg. A. Hiller, Jun. 2016; male 5.1 × 4.9 mm; female 5.3 × 5.0 mm.

Description. Carapace slightly longer than broad, evenly rounded along branchial margins, broadest on midbranchial level. Surface with transverse, piliferous plications, one epibranchial spine present. Front sinuously triangular with a longitudinal depression, its borders fringed by a row of spinules, giving a serrated aspect; orbitae moderately deep, supraocular spine strong, postorbital angle produced into a spine-tipped tooth. Eyes moderately large, dorsal extension onto




Figure 6. Dorsal view of *Petrolisthes coeruleus* sp. nov. **a** male, Islas del Rosario, Colombian Caribbean **b** male, Islas del Rosario, Colombian Caribbean **c** Bocas del Toro, Panamanian Caribbean, photograph courtesy of T. Deuss. Scale bars: 0.5 cm (**a**); 0.65 cm (**b**).

cornea narrow. Basal segment of antennulae with some transverse rugae, anterior margin with teeth. First movable segment of antennae with a serrated, spine-tipped lamellar lobe; second and third segments slightly rugose, flagellum naked.

Chelipeds sub-equal, surface with piliferous striations, merus rugose with serrated, spine-tipped lobe on anterior margin; carpus about two times as long as broad, armed on anterior margin with four, seldom five broad, serrated teeth; posterior margin slightly convex, armed with a row of strong, forwardly directed curved spines. Palm of manus broad with an inconspicuous longitudinal ridge ending at angle between fixed finger and pollex. Lateral surface of dactylus with interrupted transverse, piliferous striations; lateral surface of pollex with rough, conical protuberances extending to fixed finger; outer margin of palm convex, with row of strong, forwardly directed spines, frequently fringed with feathered setae. Gape of fingers with extended ventral pubescence covering proximal portions of pollex and dactylus. Walking legs rugose; anterior margin of merus with fringe of plumose setae, all segments covered with irregularly, wide-set, simple setae; anterior margin of merus with row of spines; large posterodistal spine on merus of walking legs 1 and 2, frequently a smaller one on leg 3; carpus of all walking legs with anterodistal spine; propodus with termi-




Figure 7. Dorsal view of *Petrolisthes coeruleus* sp. nov., male, Punta Galeta, Colón, Panamanian Caribbean. Setae on outer margin of cheliped manus omitted to depict spines. Scale bar: 0.4 cm.

nal triplet of movable spines on ventral border, with one or two additional ones; dactylus large, with four movable spinules on inner margin.

Telson seven-plated with a few short, transverse, piliferous ridges.

Colouration. The overall colouration of most specimens consists of a brownish beige background, partly overlaid with iridescent blue tones towards the posterior part of the carapace and on the chelipeds; the transverse ridges of carapace and extremities are marked by blue stripes delineated by narrow dark purple lines. The distal articulations of the walking legs are spotted with blue and purple (Fig. 6a, b). Other specimens show an entirely blue background with similar dark purple delineations of transverse ridges and granules (Fig. 6c).

The new species was first perceived as a different colour morph of *P. caribensis*, as they are not distinguishable through the two main diagnostic traits (Table 1): both bear one pair of epibranchial spines and four movable spines on the dactylus of all walking legs. However, while adult males and females of *P. caribensis* are

relatively small, with carapace lengths of up to 9.4 mm in males and 7.5 mm in females, those of *P. coeruleus* sp. nov. reach significantly larger sizes, with carapace lengths of up to 12.4 mm in males, and 12.0 mm in females. The chelipeds in the new species have a more compact aspect, and the dorsal ornamentation on the outer surface of the cheliped's palm is conspicuously more pronounced than in *P. caribensis*, which only bears scattered low granules (Fig. 8a, b).

Ecology. While *Petrolisthes caribensis* is a typical inhabitant of shallow-water coralline environments, mostly in the dead bases of the finger coral *Porites porites* (Pallas), *Petrolisthes coeruleus* sp. nov. has a wider habitat spectrum and depth range, as it occurs under boulders in protected sites of the surf zone. The authors found the species on roots of the red mangrove *Rhizophora mangle* L. in a coastal lagoon in the Colombian Gulf of Morrosquillo (9°41.684'N, 75°41.135'W), and also in the furrows of the giant barrel sponge *Xestospongia muta* (Schmidt, 1870) at 8 m depth in the same locality (Table 2). Sequences of the 16S DNA gene previously published by the authors and deposited in the GenBank database (www.ncbi.nlm.nih.gov/Genbank: sequences published as *P. galathinus* "Blue", Accession No. DQ444890–DQ444898) match a sequence from the Yucatán Peninsula collected at 20–29m, published as *P. galathinus* by Bracken-Grissom et al. (2013; accession no. KF182548). This record increases the depth range of the new species, which extends from the upper subtidal at 0.5 m to 29 m depth.

Distribution. *Petrolisthes coeruleus* sp. nov. is, so far, known from the Colombian and Panamanian Caribbean, Belize, and the east coast of Mexico. R. Lasley (FMNH; pers. comm. Nov. 2021) confirmed the species to be present in the Bahamas as well. Also, through D. Knott (PoselDon Taxonomic Services, LLC, Charleston, South Carolina, U.S.A.; pers. Comm. Oct. 2021) we became aware of material of *P. galathinus* collected in South Carolina by SERTC (Southeast Regional Taxonomic Center, South Carolina). Part of the specimens exhibit four spines on the dactylus of the walking legs, thus opening the possibility that the distribution of the new species may reach northern waters along the U.S. western coast. Collections on the coast of South Carolina await further examination to confirm if the new species indeed reaches this northern locality.




Figure 8. Dorsal view of right cheliped of **a** *Petrolisthes coeruleus* sp. nov. and **b** *P. caribensis*. Scale bars: 5 mm (**a**, **b**). Specimens preserved in ethanol.

Table 2. Geographic and ecological range of the species and colour morphs comprising the *Petrolisthes galathinus* complex and allied species. WA = West Atlantic; EP = East Pacific; FMNH = Florida Museum of Natural History.

Species	Geographic range	Ecology
<i>Parapetrolisthes tortugensis</i> (Glassell, 1945)	WA: Florida, Bahamas, Gulf of Mexico, Belize, Costa Rica, Panama, Colombia, Venezuela, Antilles (Werding et al. 2003; Poupin and Lemaitre 2014; personal records AH and BW)	Coral rubble, from sponges of the genus <i>Ircinia</i> Nardo, 1833; pers. comm. F. Sanford), 0.5–54 m (Haig 1956; Werding et al. 2003; Poupin and Lemaitre 2014; personal records AH and BW)
<i>Petrolisthes bolivarensis</i>	WA: Florida, Panama, Colombia, Venezuela (Werding and Kraus 2003; personal records AH and BW)	Dead part of <i>Porites porites</i> (Pallas, 1766), under boulders in protected sites of the surf zone, under dead blocks of <i>Acropora palmata</i> (Lamarck, 1816), 0–1.5 m (Werding and Kraus 2003; personal records AH and BW)
<i>Petrolisthes caribensis</i>	WA: Florida, Bahamas, Gulf of Mexico, Belize, Panama, Colombia, Venezuela, Antilles, (Werding 1983; Werding et al. 2003; Poupin and Lemaitre 2014; FMNH portal; personal records AH and BW)	Dead part of <i>Porites porites</i> , under dead blocks of <i>Acropora palmata</i> , 0.5–22 m, on coral heads and rubble in shallow waters at 4 m, on outer reef slope \leq 11 m (Werding et al. 2003; Poupin and Lemaitre 2014; personal records AH and BW)
<i>Petrolisthes coeruleus</i> sp. nov.	WA: Bahamas, Gulf of Mexico, Belize, Panama, Colombia (Hiller et al. 2006; FMNH portal; personal records AH and BW). Possibly in Florida up to South Carolina (personal records AH and BW; pers. comm. D. Knott)	Dead part of <i>Porites porites</i> , under boulders in protected sites of the surf zone, under dead blocks of <i>Acropora palmata</i> , on roots of <i>Rhizophora mangle</i> Linnaeus, 1753, in the furrows of <i>Xestospongia muta</i> (Schmidt, 1870), 8 m (Quiceno–Cuartas 2012); 0.5–29 m (Bracken–Grissom et al. 2013; personal records AH and BW)
<i>Petrolisthes columbiensis</i>	WA: Colombia, Cuba (Werding et al. 2003)	Dead part of <i>Porites porites</i> , 1–6 m (Werding et al. 2003; Hiller et al. 2006; personal records AH and BW)
<i>Petrolisthes galathinus</i> “Stripes–Spots”	WA: Panama, Colombia, Guyana, Brazil (Hiller et al. 2006; personal records AH and BW)	Dead part of <i>Porites porites</i> , in reef of <i>Agaricia</i> Lamarck, 1801, under boulders in protected sites of the surf zone, 0.5–3 m (Hiller et al. 2006; personal records AH and BW)
<i>Petrolisthes galathinus</i> “White Teeth”	WA: Florida, Gulf of Mexico, Belize, Panama, Colombia, Venezuela, Antilles, possibly along the east coast of Florida (Hiller et al. 2006; Rodríguez et al. 2006; Poupin and Lemaitre 2014; personal records AH and BW)	Dead part of <i>Porites porites</i> , under dead, large blocks of <i>Acropora palmata</i> , under boulders in protected sites of the surf zone, on coral heads, 0.5–6 m (Hiller et al. 2006; Poupin and Lemaitre 2014; personal records AH and BW)
<i>Petrolisthes occidentalis</i>	EP: Mexico (Cuastecomate Bay, Jalisco), El Salvador, Costa Rica, Panama, Colombia, Ecuador (Haig 1960; Moran 1984; Hiller et al. 2004, 2006; Ferreira and Anker 2021; personal records AH and BW)	Under boulders at 0–2 m; dredged from sand and sand-shell bottoms at 7.2 and 18 m; among rocks with oysters (Haig 1960; Moran 1984; Hiller et al. 2004; personal records AH and BW)
<i>Petrolisthes rosariensis</i>	WA: Bahamas, Gulf of Mexico, Belize, Panama, Colombia, Venezuela, Antilles, Brazil (Werding et al. 2003; Poupin and Lemaitre 2014; FMNH portal; personal records AH and BW)	Dead part of <i>Porites porites</i> , under dead, large blocks of <i>Acropora palmata</i> , under boulders in protected sites of the surf zone, on coral heads, in <i>Agaricia</i> coral reef framework, under boulders (Hiller et al. 2006; FMNH portal); 0.5–35 m (Poupin and Lemaitre 2014; personal records AH and BW)
<i>Petrolisthes sanmartini</i>	WA: Bahamas, French Antilles, Colombia (Werding and Hiller 2002; Werding et al. 2003; FMNH portal; personal records AH and BW)	Dead part of <i>Porites porites</i> , in coral rubble; subtidal to 18 m (Werding and Hiller 2002; Werding et al. 2003; FMNH portal; personal records AH and BW)

Etymology. The name *coeruleus* alludes to the blueish tone of carapace and extremities, which comprises a reliable diagnostic character to distinguish this species from *P. caribensis*.

***Petrolisthes occidentalis* Stimpson, 1859**

Fig. 5a

Petrolisthes occidentalis Stimpson, 1858: 227 (nomen nudum; listed); Stimpson 1859: 73 (description); Streets 1871: 240; Lockington 1878: 395; Faxon 1893: 175; Faxon 1895: 69; Ferreira and Anker 2021: 107.

Petrolisthes galathinus Ortmann, 1897: 284; Schmitt 1935: 186; Haig 1960: 36; Haig 1962: 176 (part); Gore and Abele 1976: 21; Gore 1982: 13; Moran 1984: 78; Hiller et al. 2004: 5; Hiller et al. 2006: 548.

Type material examined. Syntypes: MCZ:IZ:CRU–1401, East Pacific, Panama, 1 male, 3 females (ov).

Other material examined. MUPADI–Crus–02–17, East Pacific, Panamá, Panama City, Punta Culebra, under large boulders, low intertidal, leg. A. Hiller, 21 Feb. 2015; male, 12.8 × 13.5 mm; male, 12.9 × 13.4 mm.

Diagnosis. *Petrolisthes occidentalis* morphologically resembles the other members of the *P. galathinus* complex. An extensive description was given by Haig (1960). The species shares with most members of the group the presence of one epibranchial spine and three movable spines on the ventral side of the dactylus of the walking legs. *P. caribensis*, *P. columbiensis*, and *P. coeruleus* sp. nov. bear four such spines, and *P. sanmartini* bears five (see Table 1). The borders of the carpus of the chelipeds tend to be subparallel in *P. occidentalis* giving the carpus a straight and slender look. In the Atlantic forms, the anterior margin is more convex. *Petrolisthes occidentalis* reaches larger sizes than all Atlantic forms, with carapace lengths of more than 16 mm.

Colouration. The transverse ridges and tubercles, which are typical of the members of the *Petrolisthes galathinus* species complex, are bordered with purplish red bands, the intervening grooves are yellowish, the yellow colour prevailing on the carpus teeth. The merus of the walking legs is irregularly spotted with purplish dots, the carpus and propodus show three broad, purplish bands alternating with paler yellowish ones (Fig. 5a).

Remarks. *Petrolisthes occidentalis* was listed by Stimpson (1858), and one year later it was described by the same author (Stimpson 1859) from the Pacific coast of Panama. In the description he stated that the species is “scarcely to be distinguished from *P. sexspinosus* Stimpson, 1858”, an older synonym of the western Atlantic *P. galathinus*. In the decades after Stimpson’s studies, various authors referred to the Pacific populations as *P. occidentalis* (see Streets 1871; Faxon 1893, 1895). Ortmann (1897) emphasised the lack of differences between specimens from western India and Pacific Panama, stating that since *P. galathinus* occurred likewise on the east and west of tropical America, *P. occidentalis* should be treated as a synonym of *P. galathinus* (arguments reviewed by Haig 1960).

Hiller et al. (2006) postulated that the Eastern Pacific *P. galathinus* deserves specific status, given the relatively large genetic distances between the Pacific and the most closely related Atlantic clades, which comprised the “Spots” and “Striped” morphs. The East Pacific species differs in colour and size from all Atlantic forms. In a local catalogue of the Porcellanidae of Panama, Ferreira and Anker (2021) published a short note suggesting the need to resurrect the species, based on the recommendations by Hiller et al. (2006).

Geographic range. Haig (1960) highlighted the discontinuous distribution of the species in the East Pacific, with a concentration near Panama City, and only few findings from Isla San Lucas, Costa Rica, and off La Libertad, Ecuador. Based on new records of the species in the Panamanian and Colombian East Pacific, Hiller et al. (2004) confirmed that *P. occidentalis* seems to have a continuous distribution from Jalisco, south side of Cuastecomate Bay, Mexico, throughout Central America and Colombia, reaching Salinas and La Libertad in Ecuador.

Discussion

For more than two centuries *Petrolisthes galathinus* has been viewed as a widely dispersed species in the West Atlantic and East Pacific, with a broad ecological range, inhabiting a variety of substrates like rocks, corals, and sponges, and

from the upper subtidal down to 50 m depth. This species turned out to be a complex of morphologically similar species; so far, the *P. galathinus* complex encompasses eight species, including *P. coeruleus* sp. nov. and the resurrected *P. occidentalis*. Each species is supported by distinctive mitochondrial DNA sequences, colouration, adult size, larval morphology, and shape of the sternal plate. All species, except *P. rosariensis*, share a most recent common ancestor (MRCA), which started diverging into different Atlantic and Pacific lineages before the Central American Isthmus finished rising and interrupted gene flow between populations from each ocean (Hiller et al. 2006).

Petrolisthes galathinus still needs more revision, as it is unclear which of the molecular clades, designated in the phylogeny by Hiller et al. (2006) as “White Teeth”, “Stripes”, and “Spots”, corresponds to Bosc’s (1802) description. The three variants have different geographic distributions, overlapping in the southern Caribbean. However, while the “Stripes” and “Spots” morphs appear to have a southern distribution reaching Brazil (Hiller et al. 2006), the “White Teeth” morph extends to the Gulf of Mexico, up to the Florida Keys (pers. obs.). Since the “White Teeth” morph extends its range from the southern Caribbean to Florida, it is probable that this is the morph reported from Cape Hatteras, North Carolina by Haig (1960).

The *P. galathinus* complex poses an interesting case to study speciation within allopatric and sympatric scenarios, as it comprises closely related species on either side of the Isthmus of Panama. Such a unique assemblage allows assessing the relationship between genetic divergence and reproductive isolation, given the background of a relatively recent and well dated geological barrier that resulted in sister lineages on each side of the Americas (Hiller and Lessios 2017, 2019).

The evolutionary, ecological, and geographic processes that gave rise to the formation of different West Atlantic species with similar geographic ranges and ecologies remains to be explained in the light of a multigene phylogeographic approach of the species complex. The overlapping geographic and ecological ranges in the southern Caribbean (Table 2) are suggestive of ecological speciation driven by different microhabitats offered by coral reefs, where all species occur.

Acknowledgements

We thank T. Deuss (Bocas Mariculture, Panama) for supporting our collections in Bocas del Toro and for photographs of the new species. P. Quiceno (Universidad de Antioquia, Colombia), A. Baldinger (MCZ), A. Allspach, K. Arnold and B. Trautwein (SMF), and G. Vargas and A. Serrano (MUPADI) helped to locate and curate specimens. G. Paulay and R. Lasley (FMNH) helped identify the specimen of the new species from Bahamas; D. Knott (PoselDon TS) helped with images and examined material from the SERTC collections; M. Charteris shared professional pictures from Roatán. F. Sanford (Coe College, Iowa, U.S.A) shared ecological information on porcellanids collected in Belize and Florida. We thank J. Ceballos (STRI) for his help with the scanning electron microscope and Z. Foltz (Smithsonian National Museum of Natural History) for logistical support at the Carrie Bow Cay station in Belize. J. Poupin and an anonymous reviewer helped to improve the manuscript.

Additional information

Conflict of interest

The authors have declared that no competing interests exist.

Ethical statement

No ethical statement was reported.

Funding

No funding was reported.

Author contributions

Both authors have contributed equally.

Author ORCIDs

Alexandra Hiller  <https://orcid.org/0000-0001-9812-2126>

Bernd Werding  <https://orcid.org/0000-0002-1311-6139>

Data availability

All of the data that support the findings of this study are available in the main text.

References

- Audouin V (1826) Explication sommaire des planches de Crustacés de l'Égypte et de la Syrie, publiées par Jules-César Savigny. Description de l'Égypte, ou recueil des observations et des recherches qui ont été faites en Égypte pendant l'expédition de l'armée française. Histoire naturelle 1: 77–98. [Imprimerie impériale, Paris.]
- Benedict JE (1901) The anomuran collections made by the Fish Hawk Expedition to Porto Rico. Bulletin of the United States Fish Commission 20: 129–148. <https://research.nhm.org/pdfs/29386/29386.pdf>
- Boone L (1931) A collection of anomuran and macruran Crustacea from the Bay of Panama and the fresh waters of the Canal Zone. Bulletin of the American Museum of Natural History 63: 137–189. <https://research.nhm.org/pdfs/13727/13727.pdf>
- Bosc LAG (1802) Histoire naturelle des Crustacés, contenant leur description et leurs mœurs; avec figures dessinées d'après nature. Vol. 1, Déterville Publisher, Paris, 258 pp. <https://doi.org/10.5962/bhl.title.64025>
- Bracken-Grissom HD, Cannon ME, Cabezas P, Feldmann RM, Schweitzer CE, Ah Yong ST, Felder DL, Lemaitre R, Crandall KA (2013) A comprehensive and integrative reconstruction of evolutionary history for Anomura (Crustacea: Decapoda). BMC Evolutionary Biology 13(1): 1–28. <https://doi.org/10.1186/1471-2148-13-128>
- Chace FA (1956) Crustáceos decápodos y stomatópodos del Archipiélago de los Roques e Isla de Orchila. El Archipiélago Los Roques y la Orchila, Sociedad de Ciencias Naturales La Salle, Caracas, Venezuela, 145–168. <https://research.si.edu/publication-details/?id=13394>
- Coelho PA, Ramos MA (1972) A constituição e a distribuição de fauna de decápodos do litoral leste da América do Sul entre as Latitudes de 5° N e 39° S. Trab. Oceanogr. Univ. Fed. PE. Recife 13: 133–236, Fig. 1–4. Trabalhos Oceanográficos da Universidade Federal de Pernambuco 13: 133–236. <https://doi.org/10.5914/tropocean.v13i1.2555>

- Dana JD (1852) Crustacea. Part I. United States Exploring Expedition. During the years 1838, 1839, 1840, 1841, 1842. Under the command of Charles Wilkes, U.S.N. Vol. 13, C. Sherman, Philadelphia, 685 pp. <https://www.biodiversitylibrary.org/item/124831#page/2/mode/1up>
- de Savigny JC (1809) Crustacés, pls. 1–11. Description de l'Égypte, ou recueil des observations et des recherches qui ont été faites en Égypte pendant l'expédition de l'armée française. Histoire naturelle, Planches 2. Imprimerie impériale, Paris.
- Desmarest AG (1825) Considérations générales sur la classe des Crustacés, et description des espèces de ces animaux, qui vivent dans la mer, sur les côtes, ou dans les eaux douces de la France. F.G. Levrault, Paris and Strasbourg, 446 pp. <https://doi.org/10.5962/bhl.title.10029>
- Fausto-Filho J (1968) Terceira contribuição ao inventário dos crustáceos decápodos marinhos do nordeste brasileiro. Arquivos da Estação de Biologia Marinha da Universidade Federal do Ceará, Vol. 8: 43–45. <http://periodicos.ufc.br/arquivosdeciencia-domar/article/view/33290>
- Faxon W (1893) Reports on the dredging operations off the west coast of Central America to the Galapagos, to the west coast of Mexico, and in the Gulf of California, in charge of Alexander Agassiz, carried on by the U.S. Fish Commission steamer "Albatross," during 1891, Lieut.-Commander Z.L. Tanner, U.S.N., commanding. VI. Preliminary descriptions of new species of Crustacea. Bulletin of the Museum of Comparative Zoology at Harvard College 24: 149–220. <https://decapoda.nhm.org/pdfs/10539/10539.pdf>
- Faxon W (1895) Reports on an exploration off the west coasts of Mexico, Central and South America, and off the Galapagos Islands, in charge of Alexander Agassiz, by the U.S. Fish Commission steamer "Albatross," during 1891, Lieut.-Commander Z.L. Tanner, U.S.N. commanding. XV. The stalk-eyed Crustacea. Memoirs of the Museum of Comparative Zoology at Harvard College 18: 1–292. <https://www.biodiversitylibrary.org/page/30696747#page/81/mode/1up>
- Ferreira LA, Anker A (2021) An annotated and illustrated checklist of the porcelain crabs of Panama (Decapoda: Anomura). Zootaxa 5045: 001–154. <https://doi.org/10.11646/zootaxa.5045.1.1>
- Gibbes LR (1850) On the carcinological collections of the cabinets of natural history in the United States with an enumeration of the species contained therein, and description of new species. Proceedings of the American Association for the Advancement of Science 3rd meeting: 167–201. <https://digital.tcl.sc.edu/digital/collection/hsn/id/7625/>
- Gibbes LR (1854) Description, with figures, of six species of *Porcellana*, inhabiting Eastern Coast of North America. Proceedings of the Elliott Society of Natural History 1: 6–14. <https://www.biodiversitylibrary.org/item/135040#page/39/mode/1up>
- Glassell SA (1945) Four new species of North American crabs of the genus *Petrolisthes*. Journal of the Washington Academy of Sciences 35: 223–229. <https://decapoda.nhm.org/pdfs/12742/12742.pdf>
- Gore RH (1970) *Pachycheles cristobalensis* sp. nov. with notes on the porcellanid crabs of the southwestern Caribbean. Bulletin of Marine Science 20: 957–970. <https://www.ingentaconnect.com/contentone/umrsmas/bullmar/1970/00000020/00000004/art00009;jsessionid=gxc9ut0ikj6o.x-ic-live-03>
- Gore RH (1974) On a small collection of porcellanid crabs from the Caribbean Sea (Crustacea, Decapoda, Anomura). Bulletin of Marine Science 24: 700–721. <https://www.ingentaconnect.com/contentone/umrsmas/bullmar/1974/00000024/00000003/art00013>

- Gore RH (1982) Porcellanid crabs from the coasts of Mexico and Central America (Crustacea: Decapoda: Anomura). *Smithsonian Contributions to Zoology* 363(363): 1–34. <https://doi.org/10.5479/si.00810282.363>
- Gore RH, Abele LG (1976) Shallow water porcelain crabs from the Pacific coast of Panama and adjacent Caribbean waters (Crustacea: Anomura: Porcellanidae). *Smithsonian Contributions to Zoology* 237(237): 1–30. <https://doi.org/10.5479/si.00810282.237>
- Gray JE (1831) Description of the species of *Porcellana*, in the collection of the British Museum. *The Zoological Miscellany*, London, Part I: 14–16. <https://www.biodiversitylibrary.org/item/198027#page/1/mode/1up>
- Haig J (1956) The Galatheidea (Crustacea Anomura) of the Allan Hancock Atlantic Expedition with a review of the Porcellanidae of the western North Atlantic. *Allan Hancock Atlantic Expedition Report* 8: 1–44. <https://research.nhm.org/pdfs/29649/29649.pdf>
- Haig J (1960) The Porcellanidae (Crustacea Anomura) of the eastern Pacific. *Allan Hancock Pacific Expeditions*, University of Southern California Press, Los Angeles, California 24: 1–440.
- Haig J (1962) Papers from Dr. Th. Mortensen's Pacific expedition 1914–1916. 79. Porcellanid crabs from eastern and western America. *Videnskabelige Meddelelser fra Dansk Naturhistorisk Forening i Kjobenhavn* 124: 171–192. <https://research.nhm.org/pdfs/26886/26886.pdf>
- Haig J (1966) Campagne de la Calypso au large des côtes atlantique de L'Amérique du Sud (1961–1962) 2. Porcellanid crabs (Crustacea Anomura). *Annales de l'Institut Océanographique* 44: 351–358.
- Haworth AH (1825) A new binary arrangement of the macrurous Crustacea. *The Philosophical magazine and journal* 65: 183–184. <https://doi.org/10.1080/14786442508628417>
- Hiller A, Lessios HA (2017) Phylogeography of *Petrolisthes armatus*, an invasive species with low dispersal ability. *Scientific Reports* 7(1): 1–15. <https://doi.org/10.1038/s41598-017-03410-8>
- Hiller A, Lessios HA (2019) Marine species formation along the rise of Central America: The anomuran crab *Megalobrachium*. *Molecular Ecology* 29(2): 413–428. <https://doi.org/10.1111/mec.15323>
- Hiller A, Lazarus JF, Werding B (2004) New records and range extensions for porcellanid crabs in the eastern Pacific (Crustacea: Anomura: Porcellanidae). *Contributions to the study of East Pacific crustaceans (Vol. 3)*. Instituto de Ciencias del Mar y Limnología, Universidad de México, Ciudad de México, 127–138.
- Hiller A, Kraus H, Almon M, Werding B (2006) The *Petrolisthes galathinus* complex: Species boundaries based on color pattern, morphology and molecules, and evolutionary interrelationships between this complex and other Porcellanidae (Crustacea: Decapoda: Anomura). *Molecular Phylogenetics and Evolution* 40(2): 547–569. <https://doi.org/10.1016/j.ympev.2006.03.030>
- Lamarck JBP (1816) *Histoire naturelle des Animaux sans Vertèbres, présentant les caractères généraux et particuliers de ces animaux, leur distribution, leurs classes, leurs familles, leurs genres, et la citation des principales espèces qui s'y rapportent; précédée d'une Introduction offrant la Détermination des caractères essentiels de l'Animal, sa distinction du Végétal et des autres corps naturels, enfin, l'exposition des principes fondamentaux de la Zoologie*. Déterville & Verdrière, Paris, 568 pp. <https://doi.org/10.5962/bhl.title.40014>

- Latreille PA (1803) Histoire Naturelle, Générale et Particulière, des Crustacés et des Insectes. Dufart Publisher, Paris (Vol. 5), 406 pp. <https://www.biodiversitylibrary.org/page/15704286#page/507/mode/1up>
- Linnaeus C (1753) Species Plantarum. Laurentii Salvii, Holmiae, Stockholm (Vol. 1), 560 pp. <https://www.biodiversitylibrary.org/item/84235#page/6/mode/1up>
- Lockington WN (1878) Remarks upon the Porcellanidea of the west coast of North America. Annals & Magazine of Natural History 2(11): 394–406. <https://doi.org/10.1080/00222937808682444>
- Moran DA (1984) Additions to the known anomuran fauna of El Salvador, Central America (Crustacea: Decapoda). Journal of Crustacean Biology 4(1): 72–84. <https://doi.org/10.2307/1547898>
- Nardo GD (1833) Auszug aus einem neuen System der Spongiarien, wonach bereits die Aufstellung in der Universitäts-Sammlung zu Padua gemacht ist. Isis von Oken, Jena, Germany, 519–523.
- Ortmann AE (1897) Carcinologische Studien. Zoologische Jahrbücher, Abtheilung für Systematik, Geographie und Biologie der Thiere 10: 258–372. <https://research.nhm.org/pdfs/28108/28108.pdf>
- Pallas PS (1766) Elenchus Zoophytorum Sistens Generum Adumbrationes Generaliores et specierum Cognitarum Succintas Descriptiones cum Selectis Auctorum Synonymis. Hagae-Comitum, Apud Petrum van Cleef, The Hague, 451 pp. <https://doi.org/10.5962/bhl.title.6595>
- Poupin J, Lemaitre R (2014) Porcellanid crabs from Guadeloupe Island (Crustacea, Decapoda, Anomura), with an updated list of species from the Lesser Antilles. Zoosystema 36(1): 5–27. <https://doi.org/10.5252/z2014n1a1>
- Quiceno-Cuartas PA (2012) Historia de vida de cuatro morfotipos del complejo *Petrolisthes galathinus* (Bosc, 1902) (Crustacea, Decapoda, Anomura: Porcellanidae) en la punta norte del Golfo de Morrosquillo (San Onofre-Sucre). PhD Thesis, Universidad de Antioquia, Medellín, Colombia.
- Rickner AA (1975) Notes on members of the family Porcellanidae (Crustacea: Anomura) collected on the east coast of Mexico. Proceedings of the Biological Society of Washington 88: 159–166. <https://www.biodiversitylibrary.org/partpdf/44178>
- Rodríguez IT, Hernández G, Felder D (2006) Phylogenetic relationships among western Atlantic Porcellanidae (Decapoda: Anomura), based on partial sequences of the mitochondrial 16s rRNA gene, with comments on morphology. Crustacean Research 2006(Special Number 6): 151–166. https://doi.org/10.18353/crustacea.Special2006.6_151
- Rouse WL (1970) Littoral Crustacea from Southwest Florida. Quarterly Journal of the Florida Academy of Sciences 32: 127–152. <https://www.jstor.org/stable/24314935>
- Schmidt O (1870) Grundzüge einer Spongien-Fauna des Atlantischen Gebietes. Wilhelm Engelmann, Leipzig, 88 pp.
- Schmitt WL (1935) Crustacea Macrura and Anomura of Porto Rico and the Virgin Islands. Scientific Survey of Porto Rico and the Virgin Islands 15: 125–227. <https://doi.org/10.5962/bhl.title.10217>
- Stimpson W (1858) Prodromus descriptionis animalium evertibratorum, quae in Expeditione ad Oceanum Pacificum Septentrionalem, a Republica Federata missa, Cadwaladaro Ringgold et Johanne Rodgers Ducibus, observavit et descripsit. Pars VII. Crustacea Anomura. Proceedings of the Academy of Natural Sciences of Philadelphia 10: 225–252. <https://doi.org/10.5962/bhl.title.51447>

- Stimpson W (1859) Notes on North American Crustacea, No. I. Annals of the Lyceum of Natural History of New York 7(1): 49–93. <https://doi.org/10.1111/j.1749-6632.1862.tb00142.x>
- Streets TH (1871) Catalogue of Crustacea from the Isthmus of Panama. Collected by J. A. McNeil. Proceedings. Academy of Natural Sciences of Philadelphia 23: 238–243. <https://www.jstor.org/stable/4624183>
- Werdning B (1977) Los porcelánidos (Crustacea, Anomura, Porcellanidae) de la región de Santa Marta, Colombia. Anales del Instituto de Investigaciones Marinas Punta de Betín 9: 173–214. <https://doi.org/10.25268/bimc.invemar.1977.9.0.527>
- Werdning B (1982) Porcellanid crabs of the Islas del Rosario, Caribbean coast of Colombia, with a description of *Petrolisthes rosariensis* new species (Crustacea: Anomura). Bulletin of Marine Science 32: 439–447. <https://www.ingentaconnect.com/content/umrsmas/bullmar/1982/00000032/00000002/art00005?crawler=true>
- Werdning B (1983) Der *Petrolisthes galathinus*-Artenkomplex im karibischen Raum mit Beschreibung von *P. caribensis* n. sp. und *P. columbiensis* n. sp. (Crustacea: Anomura: Porcellanidae). Senckenbergiana biologica 63(1982) (5/6): 407–418.
- Werdning B, Hiller A (2002) A new species of the *Petrolisthes galathinus* complex from the southern Caribbean Sea, with a discussion on the identity of *P. galathinus* (Bosc, 1802) (Decapoda, Porcellanidae). Crustaceana 75(7): 849–857. <https://doi.org/10.1163/156854002321210703>
- Werdning B, Hiller A (2007) The Porcellanidae (Crustacea: Decapoda: Anomura) of the Red Sea with description of a new species of *Petrolisthes*. Zootaxa 1460(1): 1–24. <https://doi.org/10.11646/zootaxa.1460.1.1>
- Werdning B, Kraus H (2003) A new species of the *Petrolisthes galathinus* complex from the southern Caribbean Sea (Crustacea: Anomura: Porcellanidae). Crustaceana 75(9): 1141–1147. <https://doi.org/10.1163/156854002763270536>
- Werdning B, Hiller A, Lemaitre R (2003) Geographic and depth distributional patterns of western Atlantic Porcellanidae (Crustacea: Decapoda: Anomura), with an updated list of species. Memoirs of the Museum of Victoria 60(1): 79–85. <https://doi.org/10.24199/j.mmv.2003.60.11>
- Williams AB (1984) Shrimps, Lobsters, and Crabs of the Atlantic Coast of the Eastern United States, Maine to Florida. Smithsonian Institution Press, Washington D.C., 550 pp. <https://research.nhm.org/pdfs/11393/11393.pdf>