

Epitomapta aumakua sp. nov., a new species of apodous sea cucumber from Hawai`i (Echinodermata, Holothuroidea, Apodida)

Francisco Alonso Solís-Marín¹, Carlos Andrés Conejeros-Vargas^{2,3}, Andrea Alejandra Caballero-Ochoa^{3,4}, Sheila Colleen Byers⁵

1 Colección Nacional de Equinodermos “Dra. Ma. Elena Caso Muñoz”, Laboratorio de Sistemática y Ecología de Equinodermos, Instituto de Ciencias del Mar y Limnología (ICML), Universidad Nacional Autónoma de México (UNAM), Mexico City, C.P. 04510, Mexico

2 Posgrado en Ciencias del Mar y Limnología, Universidad Nacional Autónoma de México, Av. Ciudad Universitaria 3000, C.P. 04510, Coyoacán, Mexico City, Mexico

3 Facultad de Ciencias, Universidad Nacional Autónoma de México. Circuito exterior s/n, Mexico City, C.P. 04510, Mexico

4 Posgrado en Ciencias Biológicas, Instituto de Geología, Universidad Nacional Autónoma de México, C.P. 04510, Mexico City, Mexico

5 Beaty Biodiversity Museum, University of British Columbia, Vancouver, Canada

Corresponding author: Francisco Alonso Solís-Marín (fasolis@cmarl.unam.mx)

Abstract

Epitomapta aumakua sp. nov. occurs at a depth of 2.5 m in Kualoa, O`ahu, Hawai`i, living in coarse sand. It is distinctive in having 12 pinnate tentacles, each tentacle with three pairs of digits and 6–8 sensory cups. The body wall bears papillae or oval bumps, and the length of body reaches a maximum length of 18.4 mm after relaxation.

Key words: Leptosynaptinae, Synaptidae, taxonomy



Academic editor: Didier V. Spiegel

Received: 24 August 2023

Accepted: 26 September 2023

Published: 31 October 2023

ZooBank: <https://zoobank.org/934B8473-A31B-44CB-8056-EE5D6A41AB17>

Citation: Solís-Marín FA, Conejeros-Vargas CA, Caballero-Ochoa AA, Byers SC (2023) *Epitomapta aumakua* sp. nov., a new species of apodous sea cucumber from Hawai`i (Echinodermata, Holothuroidea, Apodida). ZooKeys 1183: 73–80. <https://doi.org/10.3897/zookeys.1183.111620>

Copyright: © Francisco Alonso Solís-Marín et al. This is an open access article distributed under terms of the Creative Commons Attribution License ([Attribution 4.0 International – CC BY 4.0](https://creativecommons.org/licenses/by/4.0/)).

Introduction

The apodous sea cucumbers of the subfamily Leptosynaptinae Smirnov, 1989 (Apodida Brandt, 1835; Synaptidae Burmeister, 1837) are interstitial organisms that inhabit intertidal and shallow waters. They have a vermiform, translucent integument (Hendler et al. 1995; Woo et al. 2021). Heding (1928) placed the genus *Epitomapta* Heding, 1928 in the subfamily Synaptinae and included *Epitomapta roseola* (Verrill, 1873) from Connecticut and Massachusetts, USA, and *E. tabogae* Heding, 1928 from Taboga and Taboguilla, Panama. Heding based the new genus on the presence of notched rather than perforated radial pieces of the calcareous ring. Later, Solís-Marín et al. (2019) described a new species from the Tropical East Pacific in Mexico, *E. simentalae* Solís-Marín, Conejeros-Vargas, Caballero-Ochoa & Arriaga-Ochoa, 2019, based on having 12 tentacles, with each tentacle with two or three pairs of digits and 4–6 sensory cups, and a body lacking papillae or oval bumps. *Epitomapta* is, thus, currently represented by four nominal species, including the new one described here. Smirnov (1989) placed the genus in the subfamily Leptosynaptinae.

Materials and methods

Specimens are preserved in the Marine Invertebrate Collection of the Beaty Biodiversity Museum, University of British Columbia, Canada (MI). Ossicles were extracted from the body wall (anterior, medium, and posterior region), longitudinal muscles and one whole tentacle. The tissue was dissolved in household bleach (5.0–6.5%). Bleach was washed off from the ossicles by rinsing them twice with distilled water. Hereafter the distilled water was replaced by rinsing the ossicles with 70, 80, and 95% ethanol. Finally, absolute ethanol was added to the ossicles, whereafter a small aliquot was taken and placed to dry on a scanning electron microscope (SEM) stub. The dry sample was sputter coated with 5 nm gold/palladium (80/20) using a Leica EM ACE600 and imaged with a Zeiss Crossbeam 350 SEM.

Taxonomy

Order Apodida Brandt, 1835

Family Synaptidae Burmeister, 1837

Subfamily Leptosynaptinae Smirnov, 1989

Diagnosis. Pinnate tentacles 10, 11 or 12, with 1–9 digits on each side. Digits increase in size from base to tip of tentacle. Anchor plate develops from a rod which lies at a right angle to stock of developing anchor. Anchor plates with small number of holes, usually seven (6+1) in main part of the plate: six holes form a circle around a central hole. Articular end of plate usually has a “ledge” for contact with anchor keel. Anchor arms regularly serrated, rarely smooth, and without minute knobs on the vertex (Heding 1928; Smirnov 1989).

Genus *Epitomapta* Heding, 1928

Diagnosis. Tentacles pinnate, usually 12. Digits in 2–5 pairs on each side (rarely two or none). Sense organs never in form of pigment-eyes, but as minute cups on inner face of stalk of tentacles. Calcareous ring well developed. Radial pieces not perforated for passage of nerves, but with an anterior notch. Cartilaginous ring absent. Polian vesicle usually single. Stone canal single, unbranched. Ciliated funnels of different shapes and attached to body wall, not to mesenteries. Calcareous deposits in body wall are anchors, anchor plates and miliary granules; tentacles with rods only. Stock of anchors finely toothed, but not branched; arms usually with teeth on outer edge; vertex smooth. Anchor plates oval, with large central hole, surrounded by six large holes, usually more or less dentate, and two large and several small smooth holes at narrow posterior end, but without an arched bow crossing outer surface; broad end often with additional dentate holes (Solís-Marín et al. 2019).

Type species. *Epitomapta tabogae* Heding, 1928 by original designation.

***Epitomapta aumakua* sp. nov.**

<https://zoobank.org/01CD60ED-A303-4C39-A4B7-CD0616E5FF08>

Figs 1–3

Type materials. Holotype. MI 4942, 18.4 mm total length (TL), off Kualoa, O`ahu, Hawai`i, Pacific Ocean 21°30'N, 157°50'W, 2.5 m depth, July 1975. **Paratypes.** MI 4944, 2 specimens, 1 extensively dissected, same data as the holotype.

Type locality. Off Kualoa, O`ahu, Hawai`i, Pacific Ocean 21°30'N, 157°50'W.

Diagnosis. Body wall smooth, covered with small, oval-circular bumps, especially on anterior part of body. Tentacles 12, each with three pairs of lateral digits and a terminal digit; 6–8 sensory cups on each tentacle. Polian vesicle, 1/10 of body length; stone canal single, unbranched. Anchor and anchor plates of one kind: anchors usually exceeding 170 µm in length, plates exceeding 110 µm in length. Miliary granules scarce, only present in longitudinal muscles, relatively coarse, usually in form of simple flat, often faintly undulating, stout, straight rods with enlarged ends, slightly bent but never C-shaped, usually exceeding 20 µm in length. Tentacle ossicles shaped like smooth, flattened rods, not exceeding 50 µm in length, curved, with perforated ends; some rods broad (ca 14 µm in width) with few circular peripheral holes.

Holotype description. 18.4 mm long. Specimen uniformly whitish, body wall translucent when expanded. Anchors (Fig. 1A) not projecting through body wall. Tentacles 12, each with three pairs of digits and a terminal digit; digits increase in length distally, and terminal digit longest. Inner (oral) surfaces of tentacles with double row of well-developed sensory cups; up to eight sensory cups on each tentacle. Ciliated funnels of various shapes (Fig. 2) occur on body wall, not on mesenteries. Two longitudinal rows of round-lipped, ciliated funnels present, each row attached to one side of one longitudinal muscle; a single V-shaped notch splits round lips of funnels and extends about 1/2 length of funnel. Polian vesicle single. Stone canal single, unbranched. Calcareous ring simple, stout, well developed (Fig. 3); radial pieces notched anteriorly, more conspicuous than that in interradial pieces; not pierced.

Ossicles. Body-wall deposits comprise anchors and anchor plates (Fig. 1A, B). Anchors and plates at anterior, middle, and posterior body wall essentially similar. Anchors average 170 µm in length and 55 µm in largest width (width of the arms). Arms carry up to five conspicuous teeth; vertex smooth. Stock unbranched, but equipped with numerous small, sharp projections (Fig. 1A). Anchor plates elongate, approximately oval, with numerous toothed perforations. Anchor plates average 110 µm in length and 90 µm in greatest width (Fig. 1B). Miliary granules scarce, present only in epithelium covering longitudinal muscles, variable in shape but generally flat and tending to have enlarged endings. Granules up to ca 20 µm in length (Fig. 1D). Tentacle ossicles small (40–50 µm in length), smooth, shaped like flattened, curved rods, with perforated ends (Fig. 1C). Some rods flat and broad (ca 14–16 µm in width) having 6–10 circular peripheral holes.

Paratype variations. Specimens range from 16–17 mm in length.

Etymology. The specific epithet *aumakua* refers, in Hawaiian mythology, to a person or family god that originated as a deified ancestor, who takes on physical forms as spirit vehicles. Here it is used as a non-Latin noun in apposition.




Figure 1. *Epitomapta aumakua* sp. nov. Holotype MI 4942 **A** anchors from mid-body, showing the detail of the posterior part **B** anchor plates from mid-body **C** rods from tentacles **D** military granules from the body wall.




Figure 2. *Epitomapta aumakua* sp. nov. Paratype MI 4944. Ciliated funnels showing their differing sizes and shapes.




Figure 3. *Epitomapta aumakua* sp. nov. Holotype MI 4942, calcareous ring. Abbreviations: r = radial piece, ir = interradial piece.

Ecology. *Epitomapta aumakua* sp. nov. occurs at 2.5 m depth, buried in coarse sand.

Geographical distribution. Known only from its type locality.

Discussion

Epitomapta aumakua sp. nov. is very similar to its central Eastern Pacific congener *E. simentalae* but differing in the number of sensory cups per tentacle (4–6 in *E. simentalae*, 6–8 in *E. aumakua* sp. nov.). In addition to the geographical distribution, *E. aumakua* sp. nov. is smaller (<20 mm) than *E. simentalae* (<50 mm) (Solís-Marín et al. 2019). *Epitomapta aumakua* sp. nov. clearly differs from *E. tabogae* and *E. roseola* in the number of sensory cups per tentacle (8–14 in *E. tabogae*, 2–5 in *E. roseola*), and in the number of pairs of digits present on the tentacles (5–6 in *E. tabogae*, 7 in *E. roseola*, and 2–3 in *E. aumakua* sp. nov.). *Epitomapta tabogae* was originally recorded from Taboga and Taboguilla, Panama, by Heding (1928) and is distributed throughout the Gulf of California (Solís-Marín et al. 2009), whereas *E. aumakua* sp. nov. currently is known only

from Hawai`i. In its original description (Verrill 1873), *E. roseola* was recorded from Long Island Sound, Connecticut, and Vineyard Sound, Massachusetts; it was subsequently described from Bermuda (Heding 1928) and later recorded from Connecticut and Massachusetts to Florida (USA) (Hendler et al. 1995). Most recently, *E. roseola* has been reported from the South American coast (Brazil) (Miranda et al. 2015). It has never been reported from Hawai`i.

The anchors of the body wall in *E. aumakua* sp. nov. are similar in shape to those of *E. roseola*, but they differ in size, being approximately 160–170 µm long and 76–80 µm wide in *E. aumakua* sp. nov. (Fig. 1A) versus 120–150 µm long and 70–75 µm wide in *E. roseola* (Heding 1928). The anchors of the posterior region of the body wall in both these species are similar and can reach up to 150 µm long and 70 µm wide; anchors from the anterior end of the body wall in *E. roseola* measure almost 120 µm long and 70 µm wide (Heding 1928), while in *E. aumakua* sp. nov. they are 90–150 µm long and 70 µm wide. On the other hand, the anchors of the Pacific *E. tabogae* are 200 µm in length and 100 µm in width in the posterior region of the body, and 170 µm length and 100 µm width in the anterior region of the body (Heding 1928). Today *E. tabogae* and *E. aumakua* sp. nov. possess the largest known anchors of any species in this genus.

Epitomapta aumakua sp. nov. is clearly distinguished from other species of the genus in having extremely large anchors, a character that has been used to differentiate species of the genus by various authors (see Heding 1928 and Hendler et al. 1995).

Key to species of the genus *Epitomapta*

- 1 Papillae or oval bumps present all over the body wall.....**2**
 - Papillae or oval bumps absent. With 2–3 pairs of tentacle digits, each tentacle with 4–6 sensory cups. Miliary granules in the shape of small, C- and O-shaped bodies; no papillae or oval bumps present on the body wall.....
..... ***E. simentalae***
- 2 Atlantic Ocean. With 7 pairs of tentacle digits, each tentacle with 2–5 sensory cups. Anchors of body wall exceed 120 µm in length (up to 150 µm). Miliary granules in the shape of small, oval rings and very few C-shaped bodies..... ***E. roseola***
- Pacific Ocean.....**3**
- 3 Central America (Panama). With 5–6 pairs of tentacle digits, each tentacle with 8–14 sensory cups. Anchors of body wall exceed 120 µm in length (up to 200 µm). Miliary granules in the shape of oval rings and very few C-shaped bodies..... ***E. tabogae***
- Eastern Indo Pacific (Hawai`i). With 3 pairs of tentacle digits, each tentacle with 6–8 sensory cups. Anchors of body wall exceed 150 µm in length. Scarce miliary granules in the shape of stout, flat rods.... ***E. aumakua* sp. nov.**

Acknowledgements

We thank Dr Hugh MacIntosh, collection manager of Invertebrate Zoology, Royal British Columbia for granting access to FASM to the collection under his responsibility. It is with great pleasure that we thank Dr Yves Samyn and Dr Rafael B. de Moura for critically reviewing the manuscript. We also thank Der-

rick Horne, Biolmaging Facility, University of British Columbia, for his technical support with the scanning electron microscope. Finally, we thank Matthew G. Lovegrove for his valuable comments on the manuscript's English.

Additional information

Conflict of interest

The authors have declared that no competing interests exist.

Ethical statement

No ethical statement was reported.

Funding

The first author (FASM) thanks the sabbatical fellowship from Programa de Apoyos para la Superación del Personal Académico, Dirección General de Asuntos del Personal Académicos, Universidad Nacional Autónoma de México (UNAM). Publication cost was provided by Instituto de Ciencias del Mar y Limnología, UNAM.

Author contributions

Francisco A. Solis-Marin: Investigation. Carlos A. Conejeros-Vargas: Investigation and imagen processing. Andrea A. Caballero-Ochoa: Investigation. Sheila Colleen Byres: Data analysis.

Author ORCIDs

Francisco Alonso Solís-Marín  <https://orcid.org/0000-0001-5729-3316>

Carlos Andrés Conejeros-Vargas  <https://orcid.org/0000-0003-0148-0475>

Andrea Alejandra Caballero-Ochoa  <https://orcid.org/0000-0001-5520-3823>

Sheila Colleen Byres  <https://orcid.org/0009-0003-7470-5193>

Data availability

All of the data that support the findings of this study are available in the main text.

References

- Brandt JF (1835) Echinodermata ordo Holothurina. In: Brandt JF (Ed.) *Prodromus Descriptionis Animalium ab H. Mertensio in Orbis Terrarum Circumnavigatione Observatorum*. Fasc. I. Polypos, Acalephas Discophoras et Siphonophoras, nec non Echinodermata continens. Petropoli, 42–62. <https://doi.org/10.5962/bhl.title.10196>
- Burmeister H (1837) *Handbuch der Naturgeschichte*. Zum Gebrauch bei Vorlesungen. Zweite Abtheilung: Zoologie. Theod. Chr. Friedr. Enslin, Berlin, 369–858. <https://doi.org/10.5962/bhl.title.100177>
- Heding SG (1928) Papers from Dr. Th. Mortensen's Pacific Expedition 1914–1916. Synaptidae. *Vidensk Meddelel Dansk Naturhist Foren Kobenhavn* 85: 105–323. <https://www.biodiversitylibrary.org/item/156168#page/9/mode/1up>
- Hendler G, Miller JE, Pawson DL, Kier PM (1995) *Sea Stars, Sea Urchins and Allies: Echinoderms of Florida and the Caribbean*. Smithsonian Institution Press, Washington DC, 390 pp.
- Miranda ALS, Sovierzoski HH, Correia MD (2015) Holothuroidea (Echinodermata) from reef ecosystems on the central coast of Alagoas, Brazil, with a new record to South

- Atlantic Ocean. *Cahiers de Biologie Marine* 56: 111–125. <https://doi.org/10.21411/CBM.A.5C397049>
- Smirnov AV (1989) Coordination of the system of Recent and Extinct holothurian of the family Synaptidae (in Russian with English summary). In: Kalio DL (Ed.) Fossil and Recent Echinoderm Researches. Academy of Sciences of the Estonian Socialist Soviet Republic, Tallinn, 203–217.
- Solís-Marín FA, Arriaga-Ochoa J, Laguarda-Figueras A, Frontana-Uribe S, Durán-González A (2009) Holoturoideos (Echinodermata: Holothuroidea) del Golfo de California. CONABIO, Instituto de Ciencias del Mar y Limnología, Universidad Nacional Autónoma de México, México Distrito Federal, México, 164 pp.
- Solís-Marín FA, Conejeros-Vargas CA, Caballero-Ochoa AA, Arriaga-Ochoa JA (2019) *Epitomapta simentalae* sp. n., a new species of apodous sea cucumber from the Central Eastern Pacific coast of Mexico (Echinodermata, Holothuroidea, Apodida). *ZooKeys* 817: 1–9. <https://doi.org/10.3897/zookeys.817.29406>
- Verrill AE (1873) Report upon the invertebrate animals of Vineyard Sound and the adjacent waters, with an account of the physical characters of the region. *Reports of the United States Commission of Fisheries 1871–1872*: 295–778. <https://doi.org/10.2475/ajs.s3-5.26.98>
- Woo SP, Tan SH, Nooraini I, Jaya-Ram A, Fujita T (2021) A review on knowledge and research of interstitial sea cucumber. *Paleontological Journal* 55(9): 1063–1071. <https://doi.org/10.1134/S0031030121090148>