

# A new bamboo-feeding species of the genus *Pseudosymplanella* Che, Zhang & Webb, 2009 (Hemiptera, Caliscelidae, Ommatidiotinae) from China

Nian Gong<sup>1</sup>, Xiang-Sheng Chen<sup>2</sup>, Lin Yang<sup>2</sup>

<sup>1</sup> Guizhou Provincial Engineering Research Center of Medical Resourceful Healthcare Products, Guiyang Healthcare Vocational University, Guiyang, Guizhou, 550081, China

<sup>2</sup> Institute of Entomology, Guizhou University, Guiyang, Guizhou, 550025, China

Corresponding author: Lin Yang ([yanglin6626@163.com](mailto:yanglin6626@163.com))

## Abstract

A new planthopper species, *Pseudosymplanella maxima* **sp. nov.**, belonging to the genus *Pseudosymplanella* (Hemiptera: Fulgoromorpha: Caliscelidae: Augilini), is described and illustrated, from China. In common with other Chinese Augilini, the new species feeds exclusively on bamboo. Additionally, a key to the two species of *Pseudosymplanella* is provided.

**Key words:** Augilini, bamboo, identification key, morphology, Oriental region, planthopper, taxonomy



Academic editor: Ilija Gjonov  
Received: 29 August 2023  
Accepted: 15 November 2023  
Published: 11 December 2023

ZooBank: <https://zoobank.org/7A11C606-ED1F-4064-BA2F-EBA6539D4BD5>

**Citation:** Gong N, Chen X-S, Yang L (2023) A new bamboo-feeding species of the genus *Pseudosymplanella* Che, Zhang & Webb, 2009 (Hemiptera, Caliscelidae, Ommatidiotinae) from China. ZooKeys 1186: 97–104. <https://doi.org/10.3897/zookeys.1186.111838>

**Copyright:** © Nian Gong et al.  
This is an open access article distributed under terms of the Creative Commons Attribution License ([Attribution 4.0 International – CC BY 4.0](https://creativecommons.org/licenses/by/4.0/)).

## Introduction

The family Caliscelidae Amyot & Audinet-Serville, 1843, a worldwide distributed group, is divided into two subfamilies: Caliscelinae (including tribes Caliscelini and Peltonotellini) and Ommatidiotinae (including tribes Ommatidiotini, Augilini and Adenissini) (Gnezdilov and Wilson 2006; Emeljanov 2008; Gnezdilov 2008). The two subfamilies can be separated by nymphal characters (Gnezdilov and Wilson 2006), disparities in the first metatarsomere and the degree of aedeagus reduction (Gnezdilov and Bourgoïn 2009).

Modern fauna of the tribe Augilini Baker, 1915 contains 16 genera and 41 species, including the new species described below (Zhang et al. 2020; Gong et al. 2021; Bourgoïn 2023). A fossil genus and species was documented in Dominican amber in the New World (Bourgoïn et al. 2015b). Now, seven genera and 21 species are recorded from southern China. It is worth noting that all these species have been documented to be bamboo-feeders (Chen et al. 2014; Gong et al. 2018, 2020, 2021; Zhang et al. 2020).

The planthopper genus *Pseudosymplanella* was established by Che, Zhang and Webb (2009) based on a single species, *Pseudosymplanella nigrifasciata*, from China and Thailand, and placed in the tribe Augilini of the subfamily Ommatidiotinae (Hemiptera: Fulgoroidea: Caliscelidae).

In the present paper, a new species, *Pseudosymplanella maxima* sp. nov. is described from Yunnan Province, China. Descriptions and illustrations are given, generic characteristics are redefined, and a key to species of *Pseudosymplanella* is provided.

## Material and methods

Terminology used for the external morphology and the male genitalia mainly follows the classifications proposed by Fennah (1987) and Chan and Yang (1994). The standard terminology for hind and forewing venation adheres to the principle outlined by Bourgoïn et al. (2015a). The methodology for describing the description of the female genitalia mainly follows Bourgoïn's (1993) approach. Dry specimens were used for the descriptions and illustrations. External morphology was observed under a stereoscopic microscope and characters were measured with an ocular micrometer. All dimensions were expressed in millimeters (mm); the body length was measured from the apex of the head to the apex of the forewing in repose. The genital segments of the examined specimens were subjected to maceration in a 10% NaOH and subsequently observed in glycerin jelly for illustration using a Leica MZ 12.5 stereomicroscope. Photographs were captured with a KEYENCE VHX-1000 system. Illustrations were scanned with CanoScan LiDE 200 and imported into Adobe Photoshop CS6 for labeling and plate composition. The dissected male genitalia were preserved in glycerine and then stored in small plastic tubes, which were pinned together with the specimens.

The type specimens and materials examined were deposited in the Institute of Entomology, Guizhou University, Guiyang, China (IEGU).

## Taxonomy

**Order Hemiptera Linnaeus, 1758**

**Suborder Fulgoromorpha Evans, 1946**

**Family Caliscelidae Amyot & Audinet-Serville, 1843**

**Ommatidiotinae Fieber, 1875**

**Tribe Augilini Baker, 1915**

***Pseudosymplanella* Che, Zhang & Webb, 2009**

Figs 1–21

*Pseudosymplanella* Che, Zhang & Webb, 2009: 49.

**Type species.** *Pseudosymplanella nigrifasciata* Che, Zhang & Webb, 2009, by original designation.

**Diagnosis.** Head with eyes as wide as or slightly narrower than pronotum; vertex with anterior margin a little convex or straight; second segment of antenna with a black transverse spot near apex. Mesonotum sometimes with pit along lateral margin, maximum width wider than medial length. Male with pygofer in lateral view, dorsal margin shorter than ventral margin, posterior margin with a rather slender and long process; genital style in lateral view elongate or

rather broad; aedeagus simple, tubular, slightly ventrally curved. Female genitalia with gonoplacs rounded or triangular in lateral view.

**Host plant.** Bamboo.

**Distribution.** Southern China (Yunnan Province) and Thailand (Fig. 22).

**Key to species of genus *Pseudosymplanella* Che, Zhang & Webb, 2009**

- 1 Body mainly brown, anal segment with anal pore located at mid-length, posterior margin of pygofer in profile with thick and short process near dorsal margin..... *P. nigrifasciata*
- Body grass green, anal segment with anal pore located in apical half, posterior margin of pygofer in profile with slender and long process near mid-length ..... *P. maxima* sp. nov.

***Pseudosymplanella maxima* sp. nov.**


<https://zoobank.org/A2F51374-787E-4703-8C71-694657D3A9B2>

Figs 1–21


**Description. Measurements.** Body length including forewing: male 5.1 mm ( $N = 1$ ), female 6.0–6.2 mm ( $N = 3$ ); forewing length: male 4.2 mm ( $N = 1$ ), female 5.0–5.2 mm ( $N = 3$ ).

**Coloration.** Body (Figs 1–4) grass green. Eyes reddish brown, ocelli orange red. Second segment of antenna (Fig. 6) with a black transverse spot near apex. Clypeus (Fig. 6) with basal half brown.

**Head and thorax.** Head (Fig. 5) with eyes as wide as pronotum. Vertex (Fig. 5) with length in middle line 0.8 times than width at base. Frons (Fig. 6) with length in middle line 1.1 times than maximum width. Pronotum (Fig. 5) with length in middle line shorter than vertex (0.8:1). Mesonotum (Fig. 5) 1.2 times as long as



**Figures 1–4.** *Pseudosymplanella maxima* Gong, Yang & Chen, sp. nov. **1** male habitus, dorsal view **2** male habitus, lateral view **3** female habitus, dorsal view **4** female habitus, lateral view. Scale bars: 0.5 mm (1–4).




Figures 5–15. *Pseudosymplanella maxima* Gong, Yang & Chen, sp. nov., male 5 head and thorax, dorsal view 6 face 7 head and thorax, lateral view 8 forewing 9 hindwing 10 anal segment, dorsal view 11 male genitalia, lateral view 12 pygofer, posterior view 13 genital styles, lateral view 14 pygofer and genital styles, ventral view 15 aedeagus, lateral view. Scale bars: 1 mm (8, 9); 0.5 mm (5–7); 0.3 mm (11); 0.2 mm (10, 12–15).

vertex and pronotum together in middle line. Forewing (Fig. 8) longer in middle line than broad at widest part (3.8:1); veins distinct, without nodal line, R and MP with common stem; ScP, R and CuA single, MP with three branches, Pcu uniting A1 at basal half of clavus. Hindwing (Fig. 9) with length 1.7 times as long as broad at widest part, ScP and RP single, MP and CuA with two branches. Legs relatively long, hind tibia with a single lateral tooth; spinal formula of hind leg 6-0-0.

**Male genitalia.** Anal segment (Fig. 10) in dorsal view with length 1.3 times longer in mid-line than widest part, apical margin slightly concave; anal pore located at apical half; in lateral view (Fig. 11) dorsal margin sinuated, ventral margin slightly concave near apex with a small process, broadening distally and abruptly narrowed subapically. Pygofer in lateral view (Fig. 11) with dorsal margin distinctly shorter than ventral margin, posterior margin sinuated with a rather slender and long process near mid-length; in posterior view (Fig. 12), nearly oval, with length 1.7 times longer in mid-line than widest part; in ventral view (Fig. 14), posterior margin broadly concave. Genital style in lateral view (Fig. 13) rather broad, nearly triangle, apical margin roundly convex; a strong finger-like process apically arising from dorsal margin, slightly curved. Aedeagus (Fig. 15) simple, tubular, slightly ventrally curved.

**Female genitalia.** Anal segment small, short, in dorsal view (Fig. 18) nearly quadrangle, anal pore near apex. Abdominal sternite VII in ventral view (Fig. 19) rather large and broad, behind the posterior margin with a small oval ossification flake. Gonapophysis VIII (first valvula) (Fig. 20) elongate, with five spines



Figures 16–21. *Pseudosymplanella maxima* Gong, Yang & Chen, sp. nov., female 16 genitalia, lateral view 17 genitalia, ventral view 18 anal segment, dorsal view 19 abdominal sternite VII, ventral view 20 gonapophysis VIII, lateral view 21 gonapophysis IX, lateral view. Scale bars: 0.5 mm (16, 17); 0.3 mm (19); 0.2 mm (18, 20, 21).




Figure 22. Geographic distributions of species of *Pseudosymplanella* Che, Zhang & Webb, 2009.

at apical margin. Gonapophysis IX (second valvula) (Fig. 21) with two symmetrical lobes, each lobe with many spines at dorsal margin. Gonoploc (third valvula) (Fig. 16) triangular, apical margin rounded.

**Host plant.** Bamboo.

**Distribution.** Southwestern China (Yunnan Province) (Fig. 22).

**Type material. Holotype:** ♂, CHINA: Yunnan Province, Menghai County, Mengzhe Reservoir (22°08'N, 100°26'E), 2019-X-4, Nian Gong. **Paratypes:** 1♂3♀, data same as holotype.

**Etymology.** The specific name is derived from the Latin word “maximus”, referring to the long process of the pygofer.

**Remarks.** This new species is closely related to *P. nigrifasciata* Che, Zhang & Webb, 2009, but differs in: 1) body mainly green, without stripe (body brown, with stripe in *P. nigrifasciata*); 2) anal segment with anal pore located in apical half (anal pore located at mid-length in *P. nigrifasciata*); and 3) posterior margin of pygofer in profile with slender and long process near the mid-length (posterior margin of pygofer in profile with thick and short process near dorsal margin in *P. nigrifasciata*).

## Acknowledgements

The authors are grateful to the editor and the reviewers for their valuable comments and suggestions. We wish to express our sincere thanks to Prof. Yan Zhi (Laboratory Animal Center, Guizhou Medical University, China) for her critical proofreads on the manuscript.

## Additional information

### Conflict of interest

The authors have declared that no competing interests exist.

### Ethical statement

No ethical statement was reported.

### Funding

This study was supported by the Science and Technology Support Program of Guizhou Province (No. 2020Y129), the Scientific Research Foundation of Guiyang Healthcare Vocational University (No. Guikangda K2023-9, K2022-6), and The research team specializing in the cultivation of medicinal insects of Guiyang Healthcare Vocational University.

### Author contributions

Nian Gong conceived and designed the experiments, performed the experiments, analyzed the data, prepared figures, authored or reviewed drafts of the paper, and approved the final draft. Xiang-Sheng Chen analyzed the data, prepared figures and tables, and approved the final draft. Lin Yang conceived and designed the experiments, authored or reviewed drafts of the paper, and approved the final draft.

### Author ORCIDs

Nian Gong  <https://orcid.org/0000-0002-8878-5337>

Xiang-Sheng Chen  <https://orcid.org/0000-0001-9801-0343>

Lin Yang  <https://orcid.org/0000-0002-7841-5156>

### Data availability

All of the data that support the findings of this study are available in the main text.

## References

- Amyot CJB, Audinet-Serville JG (1843) Deuxième Partie. Homoptères. Homoptera Latr. Histoire Naturelle des Insectes. Hemiptères. Librairie encyclopédique de Roret, Paris, 676 pp. <https://doi.org/10.5962/bhl.title.8471>
- Bourgoin T (1993) Female genitalia in Hemiptera Fulgoromorpha, morphological and phylogenetic data. *Annales de la Société Entomologique de France* 29(3): 225–244. <https://doi.org/10.1080/21686351.1993.12277686>
- Bourgoin T (2023) FLOW (Fulgoromorpha Lists on The Web): a world knowledge base dedicated to Fulgoromorpha. Version Updated 23 August 2023. <http://hemiptera-databases.org/flow/> [Accessed 26 August 2023]
- Bourgoin T, Wang RR, Asche M, Hoch H, Soulier-Perkins A, Stroinski A, Yap S, Szewo J (2015a) From micropterism to hyperpterism: recognition strategy and standardized homology-driven terminology of the fore wing venation patterns in planthoppers (Hemiptera: Fulgoromorpha). *Zoomorphology* 134(1): 63–77. <https://doi.org/10.1007/s00435-014-0243-6>
- Bourgoin T, Wang RR, Gnezdilov VM (2015b) First fossil record of Caliscelidae (Hemiptera: Fulgoroidea): a new Early Miocene Dominican amber genus extends the distribution of Augilini to the Neotropics. *Journal of Systematic Palaeontology* 14(3): 211–218. <https://doi.org/10.1080/14772019.2015.1032376>

- Chan ML, Yang CT (1994) Issidae of Taiwan (Homoptera: Fulgoroidea). Taichung, Taiwan, 188 pp.
- Che Y, Zhang Y, Webb MD (2009) A new genus and species of the planthopper tribe Augilini Baker (Hemiptera, Caliscelidae, Ommatidiotinae) from Thailand and China. *Zootaxa* 2311(1): 49–54. <https://doi.org/10.11646/zootaxa.2311.1.4>
- Chen XS, Zhang ZG, Chang ZM (2014) Issidae and Caliscelidae (Hemiptera: Fulgoroidea) from China. Guizhou Science and Technology Publishing House, Guiyang, 242 pp.
- Emeljanov AF (2008) New species of the genus *Peltonotellus* Puton (Homoptera, Caliscelidae) from Kazakhstan, Middle and Central Asia. *Tethys Entomological Research* 16: 5–12. [In Russian]
- Fennah RG (1987) A recharacterisation of the Ommatidiotini (Hem-Hom., Fulgoroidea, Issidae, Caliscelinae) with the description of two new genera. *Entomologist's Monthly Magazine* 123: 243–247.
- Gnezdilov VM (2008) On the taxonomy of the tribe Adenissini Dlabola (Hemiptera: Fulgoromorpha: Caliscelidae: Ommatidiotinae), with the description of a new genus and a new species from Vietnam. *Acta Entomologica Slovenica* 16(1): 11–18.
- Gnezdilov VM, Bourgoïn T (2009) First record of the family Caliscelidae (Hemiptera: Fulgoroidea) from Madagascar, with description of new taxa from the Afrotropical Region and biogeographical notes. *Zootaxa* 2020(1): 1–36. <https://doi.org/10.11646/zootaxa.2020.1.1>
- Gnezdilov VM, Wilson MR (2006) Systematic notes on tribes in the family Caliscelidae (Hemiptera: Fulgoroidea) with the description of new taxa from Palaearctic and Oriental Regions. *Zootaxa* 1359: 1–30.
- Gong N, Yang L, Chen XS (2018) *Youtuus*, a new bamboo-feeding genus of the tribe Augilini with two new species from China (Hemiptera, Fulgoromorpha, Caliscelidae). *ZooKeys* 783: 85–96. <https://doi.org/10.3897/zookeys.783.25135>
- Gong N, Yang L, Chen XS (2020) New genus and new species of the tribe Augilini (Hemiptera, Fulgoromorpha: Caliscelidae) from Yunnan Province in China. *Zootaxa* 4895(3): 411–420. <https://doi.org/10.11646/zootaxa.4895.3.6>
- Gong N, Yang L, Chen XS (2021) First record of the genus *Augilina* Melichar, 1914 (Hemiptera, Fulgoromorpha, Caliscelidae) from China, with descriptions of two new bamboo-feeding species. *European Journal of Taxonomy* 744: 38–48. <https://doi.org/10.5852/ejt.2021.744.1297>
- Zhang YL, Che YL, Meng R, Wang YL (2020) *Fauna sinica insecta* (Hemiptera: Caliscelidae and Issidae) (Vol. 70). Science Press, China, 655 pp.