

Two new species of Hypodontolaiminae (Nematoda, Chromadorida, Chromadoridae) from the Yellow Sea with a phylogenetic analysis in the subfamily

Huixin Liang¹, Wen Guo¹, Chunming Wang¹

1 College of Life Sciences, Liaocheng University, Liaocheng, 252059, China

Corresponding author: Chunming Wang (wangchunming@lcu.edu.cn)

Abstract

Two new species of Hypodontolaiminae, *Dichromadora media* sp. nov. and *Neochromadora parabilineata* sp. nov., were isolated and described from the Yellow Sea, China. *Dichromadora media* sp. nov. is characterized by four long cephalic setae, the amphidial fovea transverse oval in the male and slit-shaped in the female, the pharynx with a single posterior bulb, spicules curved and distally bifurcated, gubernaculum jointed, four (1+3) precloacal supplements papilliform, and the tail conical elongated with a short spinneret. *Neochromadora parabilineata* sp. nov. is characterized by the buccal cavity with one large hollow dorsal tooth and two small subventral teeth, the pharynx with an obvious posterior bulb, spicules L-shaped and widened medially, gubernaculum boat-shaped, seven cup-shaped and equidistant precloacal supplements, and a long and gradually tapering tail. The phylogenetic analysis of maximum likelihood and Bayesian inference based on rDNA sequences confirmed the taxonomic positions of *Neochromadora parabilineata* sp. nov. and *Dichromadora media* sp. nov. within Hypodontolaiminae. Tree topology in Hypodontolaiminae shows the genera *Neochromadora*, *Dichromadora*, *Ptycholaimellus*, and *Spilophorella* as polyphyletic groups, and the genus *Chromadorita* as a paraphyletic group.

Key words: China, *Dichromadora*, marine nematode, *Neochromadora*



Academic editor: Nic Smol

Received: 28 September 2023

Accepted: 8 January 2024

Published: 30 January 2024

ZooBank: <https://zoobank.org/>

EDBA3DBA-08BD-49AC-9ED5-

ABFBDCF40207

Citation: Liang H, Guo W, Wang

C (2024) Two new species of Hypodontolaiminae (Nematoda, Chromadorida, Chromadoridae) from the Yellow Sea with a phylogenetic analysis in the subfamily. ZooKeys 1190: 281–302. <https://doi.org/10.3897/zookeys.1190.113418>

Copyright: © Huixin Liang et al.

This is an open access article distributed under terms of the Creative Commons Attribution License ([Attribution 4.0 International – CC BY 4.0](https://creativecommons.org/licenses/by/4.0/)).

Introduction

Nematodes are the most widely distributed and diverse metazoans on the planet, and a large number of nematode species still remain unidentified (Hodda 2022). Chromadoridae Filipjev, 1917 is one of the largest families of nematodes and shown as a monophyletic group with synapomorphies of male monorchid with an anterior testis; precloacal supplements cup-shaped or absent but never tubular; females with two reflexed ovaries, anterior to the right of the intestine, posterior to the left of the intestine (Tchesunov 2014). Chromadoridae has been reviewed systematically by Wieser (1954a), Gerlach and Riemann (1973/1974), Lorenzen (1981, 1994), Tchesunov (2014), and Venekey et al. (2019), and its phylogenetic relationships have been analyzed based on rDNA sequences by Holterman et al. (2008), Leduc et al. (2017), Venekey et al. (2019), and Guo

et al. (2023). The number of rDNA sequences in Chromadoridae have rapidly increased in GenBank: up to now, 250 sequences from 20 genera of Small Subunit (SSU) and 145 sequences from 11 genera of D2–D3 fragment of Large Subunit (LSU) have been deposited in GenBank. In our study of marine nematode taxonomy from the Rizhao coast, at the Yellow China Sea, two new species from the subfamily Hypodontolaiminae are described, *Dichromadora media* sp. nov. and *Neochromadora parabilineata* sp. nov. Ribosome DNA sequences from these two new species and three other species, *Dichromadora sinica* Huang & Zhang, 2010, *Dichromadora major* Huang & Zhang, 2010, and *Dichromadora multisetosa* Huang & Zhang, 2010, are acquired for phylogenetic analysis.

Materials and methods

Sample collection

In July 2022, undisturbed samples were collected from intertidal sediments in the Rizhao coast, the Yellow China Sea. Sediment samples were vertically collected with a syringe (2.6 cm internal diameter) to a depth of 8 cm and subdivided into 0–2 and 2–8 cm depth parts. Sediments used for morphological analysis were fixed in a 10% formalin solution in seawater and for molecular analysis were preserved in 95% ethanol. Formalin-fixed samples were stained with 0.1% of Rose Bengal for more than 24 hours. Meiofauna were extracted from the sediment through Ludox centrifugation (Higgins and Thiel 1988), washed through two sieves with mesh sizes of 500 µm and 45 µm with tap water to separate meiofauna from macrofauna (larger than 500 µm), transferred to a grid-lined Petri dish, and sorted under a stereoscopic microscope. Nematodes were transferred into a mixture of ethanol (50%) and glycerin in the ratio 1:9 by volume with the ethanol slowly evaporated away (McIntyre and Warwick 1984). Nematodes were mounted in glycerin on permanent slides. Descriptions were made using an Axiscop-5 differential interference contrast microscope (Zeiss, Germany). Line drawings were made with the aid of iPad (Apple, USA), and photographs were taken with the aid of ZEN software (Zeiss). Type specimens were deposited in the Institute of Oceanology, Chinese Academy of Sciences, Qingdao.

Sediments used for molecular analysis were washed and separated as with formalin-fixed samples except without Rose Bengal dying. Seven male specimens of *D. media* sp. nov., two male specimens of *N. parabilineata* sp. nov., five male specimens of *D. sinica*, two male specimens of *D. major*, and two male specimens of *D. multisetosa* were separated and confirmed on the temporary slides.

DNA extraction, PCR amplification, and phylogenetic analysis

Genomic DNA was extracted with DNeasy Blood & Tissue kit (Qiagen, Germany) and used as amplification templates for nearly full length SSU rDNA gene, with primers of G18S4F (5' – GCT TGT CTC AAA GAT TAA GCC – 3') / 18PR (5' – TGA TCC WMC RGC AGG TTC AC – 3') (Blaxter et al. 1998), and D2–D3 fragment of LSU rDNA gene with primers of D2A (5' – ACA AGT ACC GTG AGG GAA AGT TG – 3') / D3B (5' – TCG GAA GGA ACC AGC TAC TA – 3') (Nunn 1992). PCR was conducted as described by Zhao et al. (2015). The PCR product was sequenced by Genewiz (China). The sequences were assembled in Genious v.

6.1.2. The newly obtained SSU rDNA sequences have the accession numbers as follows: *D. media* sp. nov. OR479913, *N. parabilineata* sp. nov. OR126985, *D. sinica* OR479916, *D. major* OR479911, and *D. multisetosa* OR479915; the D2–D3 fragment of LSU rDNA sequence accession numbers are *D. media* sp. nov. OR479918, *N. parabilineata* sp. nov. OR135360, *D. sinica* OR479914, *D. major* OR479912, and *D. multisetosa* OR479917. All have been deposited in GenBank.

Sequences of subfamily Hypodontolaiminae in GenBank were used for phylogenetic analysis. Forty-nine SSU rDNA sequences from seven genera (Table 1) longer than 600 bp were selected and aligned with the Muscle algorithm. Substitution models of (GTR (general time-reversible) + G (gamma distribution) + I (proportion of invariable sites)) were selected as the best-fit model and the analysis was rooted with *Latronema whataitai* Leduc & Zhao, 2015 (accession number KR048680). Sixteen D2–D3 fragment of LSU rDNA sequences from five genera (Table 2) were selected and aligned with the Muscle algorithm. Substitution models of (GTR (general time-reversible) + G (gamma distribution)) were selected as the best-fit model, and the analysis was rooted with *Latronema whataitai* (accession number KR04868).

The ML analyses were performed with Mega X with 1000 bootstrap replicates. The BI analyses were constructed with CIPRES (<http://www.phylo.org/>) and MrBayes on XSEDE v. 3.2.7a were used; the trees were run with chain length of 10,000,000, burn-in frac = 0.25. The topology of the resulting trees was visualized using FigTree v. 1.4.3 and refined with PowerPoint.

Table 1. SSU information of samples used for phylogenetic analysis.

Species	GenBank number	Reference	Locality
<i>Chromadorita aff. leuckarti</i>	MF409784.1	Schenk et al. 2018	Germany
<i>Chromadorita cf. leuckarti</i>	FJ040473.1	Holterman et al. 2008	–
<i>Chromadorita humila</i>	OQ396742.1	Sun et al. 2023	China
<i>Chromadorita leuckarti</i>	FJ969119.1	van Megen et al. 2009	–
<i>Chromadorita leuckarti</i>	KJ636254.1	Bert et al. 2014	–
<i>Chromadorita spinicauda</i>	OK317201.1	Leduc and Zhao 2023	New Zealand
<i>Dichromadora major</i>	OR479911.1	Wang et al. 2023	China
<i>Dichromadora media</i> sp. nov.	OR479913.1	Wang et al. 2023	China
<i>Dichromadora multisetosa</i>	OR479915.1	Wang et al. 2023	China
<i>Dichromadora simplex</i>	MG669747.1	Macheriotou et al. 2018	Vietnam
<i>Dichromadora sinica</i>	OR479916.1	Wang et al. 2023	China
<i>Dichromadora</i> sp.	MN250081.1	Pereira et al. 2019	Beaufort Sea (USA)
<i>Dichromadora</i> sp.	FJ040506.1	Holterman et al. 2008	–
<i>Dichromadora</i> sp.	MN250085.1	Pereira et al. 2019	Beaufort Sea (USA)
<i>Dichromadora</i> sp.	MG669748.1	Macheriotou et al. 2018	Netherlands
<i>Dichromadora</i> sp.	MN250044.1	Pereira et al. 2019	Beaufort Sea (USA)
<i>Dichromadora</i> sp.	MK626828.1	Tytgat et al. 2019	Vietnam
<i>Dichromadora</i> sp.	MG669752.1	Macheriotou et al. 2018	Vietnam
<i>Dichromadora</i> sp.	MG669751.1	Macheriotou et al. 2018	Vietnam
cf. <i>Dichromadora</i> sp.	KJ636253.1	Bert et al. 2014	–
<i>Hypodontolaimus inaequalis</i>	MG669813.1	Macheriotou 2018	Netherlands
<i>Hypodontolaimus inaequalis</i>	MG669812.1	Macheriotou et al. 2018	Netherlands

Species	GenBank number	Reference	Locality
<i>Innocuonema tentabundum</i>	AY854208.1	Meldal 2005	Southampton (United Kingdom)
<i>Innocuonema tentabundum</i>	JN968213.1	Fonseca et al. 2012	–
<i>Neochromadora bilineata</i>	OQ396744.1	Sun et al. 2023	China
<i>Neochromadora parabilineata</i> sp. nov.	OR126985.1	Wang et al. 2023	China
<i>Neochromadora poecilosomoides</i>	OQ396720.1	Chu et al. 2023	China
<i>Neochromadora</i> sp.	AY854210.1	Meldal 2005	Southampton (United Kingdom)
<i>Neochromadora</i> sp.	MG669893.1	Macheriotou et al. 2018	Netherlands
<i>Neochromadora</i> sp.	KX944147.1	Avo et al. 2017	Mira estuary (Portugal)
<i>Neochromadora</i> sp.	MN250121.1	Pereira et al. 2019	Beaufort Sea (USA)
<i>Neochromadora</i> sp.	JN968279.1	Fonseca et al. 2012	–
<i>Ptycholaimellus areniculus</i>	MG669987.1	Macheriotou et al. 2018	Vietnam
<i>Ptycholaimellus areniculus</i>	MG669988.1	Macheriotou et al. 2018	Vietnam
<i>Ptycholaimellus brevisetosus</i>	MK626833.1	Tytgat et al. 2019	Vietnam
<i>Ptycholaimellus brevisetosus</i>	MK626834.1	Tytgat et al. 2019	Vietnam
<i>Ptycholaimellus brevisetosus</i>	MK626809.1	Tytgat et al. 2019	Vietnam
<i>Ptycholaimellus brevisetosus</i>	MG669989.1	Macheriotou et al. 2018	Vietnam
<i>Ptycholaimellus ocellatus</i>	OQ538290.1	Sun et al. 2023	China
<i>Ptycholaimellus spiculuncus</i>	OK317202.1	Leduc and Zhao 2023	New Zealand
<i>Ptycholaimellus</i> sp.	FJ040472.1	Holterman et al. 2008	–
<i>Ptycholaimellus</i> sp.	KX944158.1	Avo et al. 2017	Mira estuary (Portugal)
<i>Ptycholaimellus</i> sp.	JN968285.1	Fonseca et al. 2012	–
<i>Ptycholaimellus</i> sp.	MG669992.1	Macheriotou et al. 2018	Vietnam
<i>Ptycholaimellus</i> sp.	JN968257.1	Fonseca et al. 2012	–
<i>Spilophorella aberrans</i>	MG670031.1	Macheriotou et al. 2018	Vietnam
<i>Spilophorella paradoxa</i>	AY854211.1	Meldal 2005	Southampton (United Kingdom)
<i>Spilophorella paradoxa</i>	JN968274.1	Fonseca 2012	–
<i>Spilophorella</i> sp.	MG670032.1	Macheriotou et al. 2018	Vietnam
<i>Latronema whataitai</i>	KR048680.1	Leduc and Zhao 2016	–

Table 2. D2–D3 fragment of LSU information of samples used for phylogenetic analysis.

Species	GenBank number	Reference	Locality
<i>Chromadorita humila</i>	OQ396736.1	Sun et al. 2023	China
<i>Chromadorita spinicauda</i>	OK317226.1	Leduc and Zhao 2023	New Zealand
<i>Dichromadora cucullata</i>	GU003894.1	Rodrigues et al. 2010	USA
<i>Dichromadora major</i>	OR479912.1	Wang et al. 2023	China
<i>Dichromadora media</i> sp. nov.	OR479918.1	Wang et al. 2023	China
<i>Dichromadora multiseta</i>	OR479917.1	Wang et al. 2023	China
<i>Dichromadora sinica</i>	OR479914.1	Wang et al. 2023	China
<i>Dichromadora</i> sp.	KC755220.1	Vogt et al. 2014	Wilhelmshaven (Germany)
<i>Neochromadora aff. poecilosoma</i>	KC755218.1	Vogt et al. 2014	Jadebusen (Germany)
<i>Neochromadora parabilineata</i> sp. nov.	OR135360.1	Wang et al. 2023	China
<i>Neochromadora poecilosomoides</i>	OQ417520.1	Sun et al. 2023	China
<i>Neochromadora</i> sp.	KC755219.1	Vogt et al. 2014	Wilhelmshaven (Germany)
<i>Ptycholaimellus ocellatus</i>	OQ466609.1	Sun et al. 2023	China
<i>Ptycholaimellus spiculuncus</i>	OK317227.1	Leduc and Zhao 2023	New Zealand
<i>Spilophorella</i> sp.	DQ077766.1	De Ley et al. 2009	Mexico
<i>Spilophorella</i> sp.	GU003892.1	Rodrigues et al. 2010	USA
<i>Latronema whataitai</i>	KR048681.1	Leduc and Zhao 2016	New Zealand

Results

Taxonomic account

Order Chromadorida Chitwood, 1933

Family Chromadoridae Filipjev, 1917

Subfamily Hypodontolaiminae De Coninck, 1965

Genus *Dichromadora* Kreis, 1929

Diagnosis (based on Venekey et al. 2019). Cuticle with homogeneous ornamentation and a pronounced lateral differentiation of two longitudinal rows of enlarged dots. Six outer labial papillae and four cephalic setae in separate circles. Amphideal fovea transverse slit-like and loop shaped. Buccal cavity with a triangular hollow dorsal tooth or a large dorsal tooth and two additional ventrosublateral ones; denticles can be present. Peribuccal pharyngeal tissue not swollen anteriorly or with an asymmetrical dorsal swelling; a distinct posterior pharyngeal bulb. Precloacal supplements present or absent.

Remarks. The genus *Dichromadora* was erected by Kreis in 1929 with the type species *Dichromadora microdonta* Kreis, 1929 with the genus characters the cuticle with two longitudinal rows of dots, small tooth, pharynx bulb big and round, ovaries paired symmetrical and reflexed, and males with or without precloacal supplements. Six species from the genus *Chromadora* (*C. cephalata* Steiner, 1916, *C. cricophana* Filipjev, 1922, *C. geophila* de Man, 1876, *C. parapoecilosoma* Micoletzky, 1922, *C. sabulicola* Filipjev, 1918, *C. setosa* Bütschli, 1874) were transferred to *Dichromadora* by Kreis (1929). Later, *D. hyalocheile* De Coninck & Schuurmans Stekhoven, 1933, *D. tobaensis* Schneider, 1937, *D. strandi* Allgén, 1940, *D. punctata* Schuurmans Stekhoven, 1950, *D. tenuicauda* Schuurmans Stekhoven, 1950, and *D. abnormis* Gerlach, 1953a were described. Wieser (1954a) described *D. dissipata* Wieser, 1954a, revised the genus characters based mainly on tooth shape and provided new combinations. After Wieser (1954a), *D. apapillata* Timm, 1961, *D. arcospiculum* Timm, 1961, *D. simplex* Timm, 1961, *D. islandica* Kreis, 1963, *D. scandula* Lorenzen, 1966, and *D. cucullata* Lorenzen, 1973 were described and Gerlach and Riemann (1973) presented a list of sixteen species. Later, 12 species (*D. amphidiscoides* Kito, 1981, *D. abyssalis* Bussau, 1993, *D. gathuai* Muthumbi & Vincx, 1998, *D. loiseae* Muthumbi & Vincx, 1998, *D. longicaudata* Muthumbi & Vincx, 1998, *D. quadripapillata* Muthumbi & Vincx, 1998, *D. parasimplex* Dashchenko, 2002, *D. parva* Vermeeren, Vanreusel & Vanhove, 2004, *D. polaris* Vermeeren, Vanreusel & Vanhove, 2004, *D. polarsternis* Vermeeren, Vanreusel & Vanhove, 2004, *D. southernis* Vermeeren, Vanreusel & Vanhove, 2004 and *D. weddellensis* Vermeeren, Vanreusel & Vanhove, 2004) were described. Huang and Zhang (2010) described three species from the China Sea, *D. major*, *D. multisetosa*, and *D. sinica*, and provided a short review of *Dichromadora*. *Dichromadora abyssalis* was considered as valid by Holovachov (2020) based on the high quality descriptions and illustrations despite not following the International Code of Zoological Nomenclature (Venekey et al. 2019). With the addition of *D. rigida* Thanh, Tu & Gagarin, 2016 and *D. agilis* Long, Gagarin & Tu, 2022, 35 species are currently considered as valid (based on Venekey et al. 2019):

1. *Dichromadora abnormis* Gerlach, 1953 (Italy, San Rossore and Tirrenia beaches)
2. *Dichromadora abyssalis* Bussau, 1993 (SE Pacific, Peru Basin)
3. *Dichromadora agilis* Long, Gagarin & Tu, 2022 (Vietnam, Quảng Ninh)
4. *Dichromadora amphidiscoides* Kito, 1981 (Japan, Oshoro Bay)
5. *Dichromadora antarctica* (Cobb, 1914) (Antarctica, Cape Royd; = *Spilophora antarctica* Cobb, 1914)
6. *Dichromadora apapillata* Timm, 1961 (Indian Ocean, Bay of Bengal)
7. *Dichromadora arcospiculum* Timm, 1961 (Indian Ocean, Bay of Bengal)
8. *Dichromadora cephalata* (Steiner, 1916) (Arctic Ocean, Barents Sea; = *Chromadora cephalata* Steiner, 1916, *Chromadora cricophana* Filipjev, 1922)
9. *Dichromadora cucullata* Lorenzen, 1973 (North Sea, Baltic Sea, Helgoland)
10. *Dichromadora dissipata* Wieser, 1954 (Chile, Seno de Reloncaví)
11. *Dichromadora gathuai* Muthumbi & Vincx, 1998 (Indian Ocean, Kenyan coast)
12. *Dichromadora geophila* (de Man, 1876) (North Sea, Netherlands; = *Chromadora canadensis* (Cobb, 1914), *Chromadora geophila* (de Man, 1876), *Hypodontolaimus geophilus* (de Man, 1876), *Spiliphera geophila* de Man, 1876, *Spiliphera spectabilis* Allgén, 1929)
13. *Dichromadora gracilis* (Kreis, 1929) (France, Trebeurden; = *Spilophorella gracilis* Kreis, 1929)
14. *Dichromadora hyalocheile* De Coninck & Schuurmans Stekhoven, 1933 (Belgium, Oostende)
15. *Dichromadora islandica* Kreis, 1963 (Iceland, Eyjafjörður)
16. *Dichromadora loiseae* Muthumbi & Vincx, 1998 (Indian Ocean, Kenyan coast)
17. *Dichromadora longicaudata* Muthumbi & Vincx, 1998 (Indian Ocean, Kenyan coast)
18. *Dichromadora major* Huang & Zhang, 2010 (China, Yellow Sea)
19. *Dichromadora media* sp. nov. (China, Yellow Sea)
20. *Dichromadora microdonta* Kreis, 1929 (France, English Channel)
21. *Dichromadora multisetosa* Huang & Zhang, 2010 (China, Yellow Sea)
22. *Dichromadora parasimplex* Dashchenko, 2002 (New Guinea, Astrolabe Bay)
23. *Dichromadora parva* Vermeeren, Vanreusel & Vanhove, 2004 (Antarctic Sea, Halley Bay)
24. *Dichromadora polaris* Vermeeren, Vanreusel & Vanhove, 2004 (Antarctic Sea, Halley Bay)
25. *Dichromadora polarsternis* Vermeeren, Vanreusel & Vanhove, 2004 (Antarctic Sea, Halley Bay)
26. *Dichromadora punctata* Schuurmans Stekhoven, 1950 (Mediterranean, Villefranche Bay)
27. *Dichromadora quadripapillata* Muthumbi & Vincx, 1998 (Indian Ocean, Kenyan coast)
28. *Dichromadora rigida* Thanh, Tu & Gagarin, 2016 (Vietnam, Yen River Estuary)
29. *Dichromadora scandula* Lorenzen, 1966 (North Sea, Schleswig-Holstein)
30. *Dichromadora simplex* Timm, 1961 (Indian Ocean, Bay of Bengal)
31. *Dichromadora sinica* Huang & Zhang, 2010 (China, Yellow Sea)
32. *Dichromadora southernis* Vermeeren, Vanreusel & Vanhove, 2004 (Antarctic Sea, Halley Bay)
33. *Dichromadora strandi* Allgén, 1940 (Norway, Knivskjaerodden)
34. *Dichromadora tobaensis* Schneider, 1937 (Indonesia, Sumatra)

35. *Dichromadora weddellensis* Vermeeren, Vanreusel & Vanhove, 2004 (Antarctic Sea, Halley Bay)

***Dichromadora media* sp. nov.**

<https://zoobank.org/4F7D70E0-54B3-422C-9A87-F496EA14C572>

Figs 1, 2, Table 3

Diagnosis. Medium body size, cuticle with transverse rows of dots and a lateral differentiation of two longitudinal larger dots, four long cephalic setae, buccal cavity large with one large, hollow and straight dorsal tooth and two small ventrosublateral teeth, amphidial fovea transverse oval in male and slit-shaped in female, pharynx with single posterior bulb, spicules curved and distally bifurcated, gubernaculum jointed, four (1+3) papilliform precloacal supplements, tail conical elongated with short spinneret.

Material examined. Four males and three females were measured and studied. **Holotype:** ♂ 1 on slide 22ZJT8-1-2; **Paratypes:** ♂ 2 on 22ZJT8-1-2, ♂ 3 on 22ZJT8-1-2, ♂ 4 on 22ZJT8-2-5, ♀ 1 on 22ZJT8-2-7, ♀ 2 on 22ZJT8-2-7, and ♀ 3 on 22ZJT8-2-6.

Type locality and habitat. Rizhao coast, Shandong Province, China, 35°27'N, 119°35'E, 0–2 cm sediment depth, sandy sediment.

Measurements. All measurement data are given in Table 3.

Description. Males. Body cylindrical and medium sized (881–925 µm in length). Cuticle with transverse rows of dots and a differentiation consisting of two longitudinal rows of distinct larger dots starting posterior the amphidial fovea and extending to the tail tip (2 µm in width). Transverse bars connecting the two larger dots beginning from the middle of the pharynx to the middle of tail. Somatic setae present sparsely along the lateral differentiation in two longitudinal rows, short in the head and tail (9 µm in length), long in the middle of the body (12 µm in length). Inner and outer labial sensilla papilliform. Four cephalic sensilla setiform at the level of amphidial fovea (1.20–1.36 head diameter in length). Head blunt. Amphidial fovea oval (4 µm in width and 3 µm in length), small (26.7%–28.6% corresponding body diameter) and situated 0.47–0.57 head diameter from the anterior end. Buccal cavity cuticularized with a large, hollow and straight dorsal tooth and two small ventrosublateral teeth. Cheiostoma short with longitudinal cuticularized ribs. Pharynx cylindrical, anterior region surrounding buccal cavity slightly swollen, posterior region swollen into an elongated single bulb with plasmatic interruptions resembling a double bulb (18.8–20.0% of pharynx length). Nerve ring slightly posterior to middle pharynx region (53.6–60.9% of pharynx length). Renette cell of secretory-excretory system situated posterior to pharynx bulb, excretory pore located at anterior buccal cavity (6–8 µm from anterior end). Cardia not observed.

The reproductive system monorchid, with extended testis located to the right of intestine. Spicules equal and slightly curved, 23–25 µm (1.0–1.1 cloacal body diameter) along arc, proximal end slightly widened and distal end bifurcated. Gubernaculum jointed without apophysis. Four (1 + 3) precloacal supplements papilliform, anterior three supplements closely distanced and posteriormost supplement distant from the three anterior ones, distance between the supplements 9–12 µm, 8–9 µm, 16–19 µm, respectively, and posteriormost




Figure 1. *Dichromadora media* sp. nov. **A** lateral view of male anterior region showing cuticle and amphidial fovea (holotype) **B** lateral view of male buccal cavity (holotype); **C** lateral view of male anterior region showing pharyngeal region (holotype) **D** lateral view of female anterior region showing buccal cavity, amphidial fovea and pharyngeal region (22ZJT8-2-7) **E** lateral view of female whole body (22ZJT8-2-7) **F** lateral view of male posterior body, showing precloacal supplements and tail (holotype); **G** lateral view of spicules and gubernaculum (22ZJT8-2-5) **H** lateral view of female posterior body showing tail (22ZJT8-2-7). Scale bars: 20 µm (**A–D, F, H**); 50 µm (**E**).

supplement 7–8 µm from cloaca. Tail elongated conical, gradually tapering, 4.5–5.0 cloacal body diameters. Spinneret very short, 1–2 µm in length.

Females. Similar to males in most characteristics. Amphidial fovea slit-like (50.0% corresponding body diameter). Cephalic setae short (10–11 µm in length). Reproductive system didelphic, with opposed and reflexed ovaries. Anterior ovary to right of intestine and posterior ovary to left of intestine. Eggs oval shaped, 8–10 × 10–11 µm. Vulva at the middle of the total body. Vagina short.

Etymology. Species epithet *media* refers to the medium body size.

Remarks. *Dichromadora media* sp. nov. differs from all other species of the genus *Dichromadora* by the amphidial fovea shape and jointed gubernaculum and it is similar to *D. dissipata*, *D. quadripapillata*, and *D. sinica* in body length and precloacal supplements number. However, it differs from *D. dissipata* in




Figure 2. *Dichromadora media* sp. nov. **A** lateral view of male anterior region showing tooth (arrow) (holotype) **B** lateral view of male anterior region showing cuticle, amphidial fovea (arrow) (holotype) **C** lateral view of male anterior region showing pharyngeal region (holotype) **D** lateral view of male posterior body, showing spicules (holotype) **E** lateral view of male posterior body, showing gubernaculum (22ZJT8-1-2) **F** lateral view of male posterior body, showing precloacal supplements (arrows) (holotype) **G** lateral view of female anterior region showing amphidial fovea (arrow) (22ZJT8-1-2) **H** lateral view of distal end of spicules (22ZJT8-2-5). Scale bars: 20 µm.

cephalic setae length (10–19 µm vs 9–9.5 µm), spicules length (23–25 µm vs 39 µm), gubernaculum shape (double-jointed without apophysis vs not jointed with dorsal apophysis) and precloacal supplements (1+3 vs 5); differs from *D. quadripapillata* in cephalic setae length (10–19 µm vs 4–5 µm), spicules shape (slightly curved and distally bifurcated vs curved with pointed distal end) and precloacal supplements shape (papilliform vs cup-shaped); differs from *D. sinica* in cuticle differentiation (lateral differentiation with transverse bars vs lateral differentiation without transverse bars), pharynx bulb shape (single bulb with plastic interruptions vs double bulb), spicules shape (slightly curved and distally bifurcated vs distal end with a hook), gubernaculum shape (jointed vs not jointed), and precloacal supplements arrangement (1+3 vs 3+1).

Dichromadora media sp. nov. shows a close relationship with *D. sinica* in the phylogenetic trees (Figs 5, 6) based on rDNA sequences and it differs by 2% (39 in 1656 bp, including two gaps) in SSU and 5% (38 in 770 bp, including four

Table 3. Measurements of *Dichromadora media* sp. nov. (in µm except for ratios).

Characters	Holotype	Paratypes	Paratypes
	male	males (n = 3)	females (n = 3)
Total body length	925	895±14.5(881–910)	812±44.6(761–844)
Maximum body diameter	23	22.3±0.6(22–23)	27.7±0.6(27–28)
Head diameter	15	14.3±0.6(14–15)	16±0(16–16)
Length of cephalic setae	18	18.3±0.6(18–19)	10.7±0.6(10–11)
Amphidial fovea width	4	4±0(4–4)	8±0(8–8)
Amphidial fovea from anterior end	7	7.3±0.6(7–8)	4.7±0.6(4–5)
Body diameter at amphidial fovea	15	14.3±0.6(14–15)	16±0(16–16)
Nerve ring from anterior end	64	64.3±5.1(60–70)	64±4.6(59–68)
Body diameter at nerve ring	18	18±0(18–18)	20.3±1.2(19–21)
Pharynx length	116	114.7±2.5(112–117)	119±4.4(116–124)
Pharynx bulb length	22	22.3±0.6(22–23)	25.3±1.5(24–27)
Body diameter at base of pharynx	19	19±0(19–19)	22.3±1.2(21–23)
Cloacal/anal body diameter	23	22.3±0.6(22–23)	15.7±0.6(15–16)
Spicules length along arc	23	24.3±0.6(24–25)	–
Gubernaculum length	23	22±1(21–23)	–
Vulva from anterior end	–	–	399±16.1(381–412)
Body diameter at vulva	–	–	27.3±0.6(27–28)
V%	–	–	49.2±0.8(48.6–50.1)
Precloacal supplements	1+3	1+3	–
Tail length	103	107±3.5(105–111)	106.7±6.8(99–112)
a	40.2	40.1±1.2(38.9–41.4)	29.4±1.9(27.2–30.8)
b	8.0	7.8±0.1(7.8–7.9)	6.8±0.3(6.6–7.2)
c	9.0	8.4±0.4(7.9–8.7)	7.6±0.2(7.4–7.7)
c'	4.5	4.8±0.2(4.6–5.0)	6.8±0.2(6.6–7)

gaps) in LSU D2–D3 fragment, but they can be morphologically differentiated based on pharynx bulb shape, spicule shape, gubernaculum shape and precloacal supplements.

Genus *Neochromadora* Micoletzky, 1924

Diagnosis (based on Venekey et al. 2019). Cuticle ornamentation heterogeneous and complex, with lateral differentiation visible as two or three longitudinal rows of large dots. Six small outer labial setae or papillae and four cephalic setae in separate circles. Inner labial sensilla may be conspicuous in one species (*N. munita*). Presence of somatic setae in some species. Amphidial fovea transverse slit-like and loop shaped. Buccal cavity with a dorsal tooth and two ventrosublateral teeth, in some species the dorsal one being larger than the others. Denticles can be present. Pharynx anteriorly not swollen or swollen next to the dorsal tooth. Pharynx with a single well-developed posterior bulb. Male usually with numerous precloacal supplements.

Remarks. The genus *Neochromadora* was erected by Micoletzky (1924) with the type species *Neochromadora poecilosoma* (de Man, 1893). And six species, *Neochromadora aberrans* (Cobb, 1930), *Neochromadora craspedota* (Steiner, 1916), *Neochromadora edentata* (Cobb, 1914), *Neochromadora izhorica* (Filipjev, 1929), *Neochromadora poecilosomoides* (Filipjev, 1918), *Neochromadora sabuli-*

cola (Filipjev, 1918) have been added to *Neochromadora*. Gerlach (1951) described *N. tecta* Gerlach, 1951, redescribed *N. poecilosoma*, *N. izhorica* (Filipjev, 1929), and transferred *Spiliphera trichophora* (Steiner, 1921) to *Neochromadora*. Later Gerlach (1952, 1953b) described *N. attenuata* Gerlach, 1952 and *N. complexa* Gerlach, 1953b. Wieser (1954a) divided *Neochromadora* into two subgenera *Neochromadorina* (Wieser, 1954a) and *Trichodorina* (Wieser, 1954a) based on tooth structure, cervical and somatic setal length, and pharyngeal bulb and described three new species: *N. lateralis* Wieser, 1954a, *N. calathifera* Wieser, 1954a, and *N. torquata* Wieser, 1954a. Later, Wieser (1954b) described another two species *N. amembranata* Wieser, 1954b and *N. brevisetosa* Wieser, 1954b. Afterwards, six species, *N. bonita* Gerlach, 1956, *N. coudenhoivei* Wieser, 1956, *N. notocraspedota* Allgén, 1958, *N. appiana* Wieser, 1959, *N. pugillator* Wieser, 1959 and *N. bicoronata* (Wieser, 1959) (synonym *Endeolophos spinosus* (Gerlach, 1957) were described. Wieser (1959) withdrew the subgenus *Trichodorina* with a redescription of *N. poecilosoma* found in Puget Sound. Later, 13 species (*N. alatocorpa* Hopper, 1961, *N. nitida* Timm, 1961, *N. munita* Lorenzen, 1971, *N. paratecta* Blome, 1974, *N. paramunita* Boucher, 1976, *N. angelica* Riemann, 1976, *N. bilineata* Kito, 1978, *N. oshoroana* Kito, 1981, *N. orientalis* Lemzina, 1982, *N. papillosa* Pastor de Ward, 1985, *N. lineata* Pastor de Ward, 1985, *N. nicolae* Vincx, 1986 and *N. alejandroi* Lo Russo & Pastor de Ward, 2012) were described. Hopper (1963) considered *Neochromadora trilineata* Schneider, 1943 as incertae sedis due to the unavailability of specimens. Vincx (1986) considered *N. paramunita* as a synonym of *N. munita*. Up to now, 33 species are currently considered as valid (based on Venekey et al. 2019):

1. *Neochromadora aberrans* (Cobb, 1930) (Antarctic, Commonwealth Bay; = *Spiliphera aberrans* Cobb, 1930)
2. *Neochromadora alatocorpa* Hopper, 1961 (USA, Alabama)
3. *Neochromadora alejandroi* Lo Russo & Pastor de Ward, 2012 (Argentina, San Matías gulf)
4. *Neochromadora amembranata* Wieser, 1954 (Italy, Sampieri)
5. *Neochromadora angelica* Riemann, 1976b (Germany, Helgoland)
6. *Neochromadora appiana* Wieser, 1959 (USA, Washington)
7. *Neochromadora bilineata* Kito, 1978 (Japan, Hokkaido)
8. *Neochromadora bonita* Gerlach, 1956 (Brazil, Cananeia)
9. *Neochromadora brevisetosa* Wieser, 1954 (Italy, Sampieri)
10. *Neochromadora calathifera* Wieser, 1954b (Chile, Seno Reloncavi)
11. *Neochromadora complexa* Gerlach, 1953a (Chile, Seno Ultima Esperanza)
12. *Neochromadora coudenhoivei* Wieser, 1956b (Greece, Piraeus)
13. *Neochromadora craspedota* (Steiner, 1916) (Arctic Ocean, Barents Sea; = *Chromadora craspedota* Steiner, 1916)
14. *Neochromadora edentata* (Cobb, 1914) (Antarctic, Cape Royds; = *Nygmatonchus edentata* (Cobb, 1914) Wieser, 1954, *Spiliphera edentata* Cobb, 1914)
15. *Neochromadora izhorica* (Filipjev, 1929) (Baltic Sea, Neva Bay; = *Chromadorella izhorica* Filipjev, 1929)
16. *Neochromadora lateralis* Wieser, 1954 (Chile, Seno Reloncavi)
17. *Neochromadora lineata* Pastor de Ward, 1985a (Argentina, Deseado river)
18. *Neochromadora munita* Lorenzen, 1972 (Germany, Helgoland; = *Neochromadora paramunita* Boucher, 1976)
19. *Neochromadora nicolae* Vincx, 1986 (North Sea, Southern Bight)

20. *Neochromadora nitida* Timm, 1961 (Indian Ocean, Bengal Bay)
21. *Neochromadora notocraspedota* Allgén, 1958 (Uruguay, Uruguay coast)
22. *Neochromadora orientalis* Lemzina, 1982 (Kyrgyzstan, Lake Issyk-Kul)
23. *Neochromadora oshoroana* Kito, 1981 (Japan, Oshoro Bay)
24. *Neochromadora papillosa* Pastor de Ward, 1985 (Argentina, Deseado River)
25. *Neochromadora parabilineata* sp. nov. (China, Yellow Sea)
26. *Neochromadora paratecta* Blome, 1974 (Germany, Sylt)
27. *Neochromadora poecilosoma* (de Man, 1893) (North Sea, English Channel;
= *Chromadora poecilosoma* de Man, 1893)
28. *Neochromadora poecilosomoides* (Filipjev, 1918) (Black Sea, Kruglaya Bay and
Georgievskii Monastery Bay; = *Chromadora poecilosomoides* Filipjev, 1918)
29. *Neochromadora pugilator* Wieser, 1959 (USA, Washington)
30. *Neochromadora sabulicola* (Filipjev, 1918) (Black Sea, Kruglaya Bay and
Georgievskii Monastery Bay; = *Chromadora sabulicola* Filipjev, 1918)
31. *Neochromadora tecta* Gerlach, 1951 (Germany, Amrum Island)
32. *Neochromadora torquata* Wieser, 1954 (Chile, Seno Reloncavi)
33. *Neochromadora trichophora* (Steiner, 1921a) (Spain, Canary Islands;
= *Spiliphera trichophora* Steiner, 1921, *Neochromadora longisetosa* Schuurmans-Stekhoven, 1935)

***Neochromadora parabilineata* sp. nov.**

<https://zoobank.org/27F8BC81-DA38-470D-AD2B-F4B20F302588>

Figs 3, 4, Table 4

Diagnosis. Medium body size, buccal cavity with one large hollow dorsal tooth and two small subventral teeth, spicules curved and L-shaped, gubernaculum boat-shaped, seven precloacal supplements cup-shaped, tail conical and gradually tapering.

Material examined. Four males and three females were measured and studied. **Holotype:** ♂ 1 on slide 22HSB11–2–20; **paratypes:** ♂ 2 on 22HSB11–1–21, ♂ 3 on 22HSB11–2–18, ♂ 4 on 22HSB11–2–20, ♀ 1 on 22HSB11–2–18, ♀ 2 on 22HSB11–1–18, and ♀ 3 on 22HSB11–2–18.

Type locality and habitat. Rizhao coast, Shandong Province, China, 35°5'N, 119°20'E, 0–2 cm sediment depth, sandy sediment.

Measurements. All measurement data are given in Table 4.

Description. Males. Body medium sized (864–943 µm), anterior end truncated and posterior end tapered. Cuticle heterogeneous and complex, five transverse rows of small dots present just posterior to cephalic setae, two or three longitudinal rows of larger dots posterior to the cephalic setae to middle part of body, larger dots changing to rectangular markings from middle body to posterior part of cloaca and rectangular markings changing back to larger dots until tail end. Six inner and six outer labial sensilla papilliform, four setiform cephalic sensilla (0.5–0.7 head diameter in length). Somatic setae present in pharynx and tail region (8 µm in length). Amphidial fovea situated at level of cephalic setae, transverse oval, 6–7 µm in width and 2 µm in length (50–58% corresponding body diameter). Buccal cavity shallow, 10–14 µm in depth. Cheiostoma short with cuticularized longitudinal folds. Pharyngostoma with one large hollow dorsal tooth and two small subventral teeth. Pharynx cylindrical, pos-




Figure 3. *Neochromadora parabilineata* sp. nov. **A** lateral view of male anterior region showing cuticle and amphidial fovea (holotype) **B** lateral view of male buccal cavity (holotype) **C** lateral view of male anterior region showing pharyngeal region (holotype) **D** lateral view of male cuticle at pharynx region (holotype) **E** lateral view of male cuticle at middle body (holotype) **F** lateral view of female anterior region showing buccal cavity and pharyngeal region (22HSB11-1-18) **G** lateral view of spicules and gubernaculum (22HSB11-2-20) **H** lateral view of male posterior body, showing precloacal supplements and tail (holotype) **I** lateral view of female whole body (22HSB11-2-18) **J** lateral view of female posterior body showing tail (22HSB11-2-18). Scale bars: 20 μ m (**A–H, J**); 50 μ m (**I**).




Figure 4. *Neochromadora parabilineata* sp. nov. **A** lateral view of male anterior region showing cuticle (holotype) **B** lateral view of female anterior region showing amphidial fovea (arrow) (22HSB11-2-18) **C** lateral view of cuticle at middle body (holotype) **D** lateral view of male posterior body, showing spicules (22HSB11-2-20) **E** lateral view of male posterior body, showing gubernaculum (22HSB11-2-20) **F** lateral view of male posterior body, showing precloacal supplements (arrows) (holotype) **G** lateral view of male posterior body, showing tail and cuticle (22HSB11-2-20). Scale bars: 20 µm.

terior region swollen into an oval bulb (17.5–17.7% of pharynx length). Nerve ring slightly posterior to middle pharynx region (64.2–66.9% of pharynx length). Secretory-excretory system present; renette cell situated posterior to pharynx bulb, excretory pore at level with cephalic setae. Cardia not observed.

Reproductive system with a single, outstretched testis. Spicules curved and L-shaped, widened at the middle part, 28–31 µm (0.81–0.86 cloacal body diameters) along arc. Gubernaculum short and boat-shaped, distal end tapered. Seven precloacal supplements cup-shaped, distance between the anteriormost and cloaca, the posteriormost and cloaca, 100 µm and 18 µm respectively, distance between supplements almost equal-distaned. Tail conical and gradually tapering, 4.4–4.8 cloacal body diameter in length. Spinneret short, 5 µm in length.

Females. Similar to males in most characteristics. Tail slightly longer than in males (5.8–6.8 anal body diameters in length). Reproductive system didelphic, with opposed and reflexed ovaries. Anterior ovary to left of intestine and posterior ovary to right of intestine. Spermatheca present. Vulva situated anterior to middle of body. Vagina short and muscularized.

Etymology. Species epithet *parabilineata* refers to the new species being similar to *Neochromadora bilineata*.

Differential diagnosis. *Neochromadora parabilineata* sp. nov. is similar to *N. bilineata*, *N. izhorica*, *N. complexa*, and *N. poecilosoma* in precloacal supplements number (7–9). But it differs from *N. bilineata* in body length (864–943 µm

Table 4. Measurements of *Neochromadora parabilineata* sp. nov. (in μm except for ratios).

Characters	Holotype	Paratypes	Paratypes
	male	males ($n = 3$)	females ($n = 3$)
Total body length	878	905.3±39.6(864–943)	913±27.8(881–931)
Maximum body diameter	25	25.7±1.5(24–27)	33.7±5.8(27–37)
Head diameter	12	11.7±1.5(10–13)	11.7±0.6(11–12)
Length of cephalic setae	8	7±0(7–7)	7.3±0.6(7–8)
Buccal cavity depth	10	10.7±3.1(8–14)	7±1(6–8)
Amphidial fovea width	6	6.7±0.6(6–7)	6±0(6–6)
Amphidial fovea from anterior end	3	3±0(3–3)	3±0(3–3)
Body diameter at amphidial fovea	12	11.7±0.6(11–12)	12±0(12–12)
Nerve ring from anterior end	86	80.7±3.8(78–85)	84±6.1(80–91)
Body diameter at nerve ring	23	21.3±0.6(21–22)	23.3±0.6(23–24)
Pharynx length	131	124.7±2.1(123–127)	128±2.6(125–130)
Pharynx bulb length	23	22.7±0.6(22–23)	26±3.6(23–30)
Body diameter at the base of pharynx	24	22.3±1.2(21–23)	26.3±2.9(23–28)
Cloacal/anal body diameter	25	24±0(24–24)	19.3±0.6(19–20)
Spicules length along arc	31	29±1(28–30)	–
Gubernaculum length	21	20.3±1.5(19–22)	–
Vulva from anterior end	–	–	380.3±34.1(341–401)
Body diameter at vulva	–	–	32±4.6(27–36)
Precloacal supplements	7	7	–
V%	–	–	41.6±2.5(38.7–43.1)
Tail length	113	111.3±5.5(105–115)	121.7±12.6(110–135)
a	35.1	35.4±3(32–37.9)	27.6±4.3(25.1–32.6)
b	6.7	7.3±0.3(7–7.6)	7.1±0.1(7–7.2)
c	7.8	8.2±0.7(7.6–9)	7.6±0.8(6.9–8.5)
c'	4.5	4.6±0.2(4.4–4.8)	6.3±0.5(5.8–6.8)

vs 567–852 μm), cephalic setae length (7–8 μm vs 4–6 μm), amphidial fovea width (50–58% vs 45% corresponding body diameter), spicules shape and length (L-shaped and widened in the middle portion, 28–31 μm vs arcuate and gradually narrowing, 23–26 μm), and gubernaculum length (19–22 μm vs 15–18 μm); differs from *N. izhorica* in cephalic seta length (7–8 μm vs 14 μm), pharynx shape (posterior bulb obvious vs posterior bulb weak), spicules length (28–31 μm vs 31.5–34.5 μm), gubernaculum shape (distal end tapered vs distal end with anterior-laterally curved tip) (Riemann 1966); differs from *N. complexa* in body length (864–943 μm vs 642 μm), spicules shape (curved and L-shaped with middle portion widened vs L-shaped even in width), gubernaculum shape (boat-shaped vs dorsal part slenderly extended), distance between precloacal supplements (10–15 μm vs 2–5 μm) (calculation based on Gerlach 1953b: fig. 11); differs from *N. poecilosoma* in body length (864–943 μm vs 1900–2000 μm), cephalic setae length (7–8 μm vs 10–14 μm), spicule length (28–31 μm vs 60–65 μm), gubernaculum shape (boat-shaped vs distal tip with small tooth) (de Man 1893).

Molecular phylogenetic analysis

The ML topology trees which are obtained based on the rDNA gene sequences are mostly in accordance with the BI topology trees, and only the BI trees are shown in Figs 5, 6.

Sequences of seven genera of the subfamily Hypodontolaiminae, *Chromadorita* Filipjev, 1922, *Dichromadora*, *Hypodontolaimus* de Man, 1886, *Innocuonema* Inglis, 1969, *Neochromadora*, *Ptycholaimellus* Cobb, 1920 and *Spilophorella* Filipjev, 1917 are included in the SSU and LSU rDNA analyses. At genus level, only species of *Chromadorita* cluster in one clade (posterior probability 70 in SSU and 99 in LSU, bootstrap value 99 in LSU) but *Chromadorita* is shown as paraphyletic. *Dichromadora multisetosa* (**OR479915**), *D. major* (**OR479911**), *N. poecilosomoides* (**OQ396720**), and *Innocuonema tentabunda* (**JN968213**, as *Chromadorita tentabunda*) clustered with the *Chromadorita* clade in the SSU




Figure 5. Bayesian inference tree of the subfamily Hypodontolaiminae inferred from Small Subunit (SSU) sequences under the general time-reversible (GTR) + gamma distribution (G) + proportion of invariable sites (I) model. Posterior probability (left) and bootstrap values (right) are given on corresponding clades. The sequences obtained in this study are shown in bold. The scales indicate substitutions per site.




Figure 6. Bayesian inference tree of the subfamily Hypodontolaiminae inferred from the D2-D3 fragment of Large Subunit (LSU) sequences under the general time-reversible (GTR) + gamma distribution (G) model. Posterior probability (left) and bootstrap values (right) are given on corresponding clades. The sequences obtained in this study are shown in bold. The scales indicate substitutions per site.

analysis. These species share a common character of peribuccal cavity tissue with an asymmetrical dorsal swelling and only one posterior pharynx bulb. However, *Chromadorita* can be morphologically differentiated from them by having the cuticle homogeneous without any lateral differentiation.

Six sequences of genus *Dichromadora* have been identified to species level, but they are in four different clades in both SSU and LSU analyses and therefore paraphyletic. Among the species of *Dichromadora*, *Dichromadora multiseta* is the only species with the gubernaculum not being boat-shaped but with dorsal caudal apophysis, and clustered with *Chromadorita humila* (gubernaculum possessing arched dorsal-caudal apophysis) highly supported by the LSU topology tree (posterior probability 85, bootstrap value 51 in SSU; posterior probability 100, bootstrap value 81 in LSU). *Dichromadora sinica* and *D. media* sp. nov. are highly clustered based on rDNA sequences (posterior probability 100, bootstrap value 93 in SSU; posterior probability 86, bootstrap value 71 in LSU), and can be distinguished based on the pharyngeal bulb, precloacal supplements, and gubernaculum shape.

Species of genus *Neochromadora* differ from other genera of Hypodontolaiminae mainly based on the cuticle ornamentation being heterogeneous and complex. However, the cuticle structure is sometimes seen as a variable character (e.g., Leduc et al. 2017), and sequences of *Neochromadora* present as polyphyletic clades in SSU and LSU topology trees. *Neochromadora bilineata* (OQ396744) and *N. parabilineata* sp. nov. have a close relationship in the SSU analysis (posterior probability 94, bootstrap value 68), but the LSU sequence of *N. bilineata* was missing. Relationships between these two species should be further discussed when more molecular data is available, in combination with morphological characters.

The genera *Ptycholaimellus* and *Spilophorella* are paraphyletic clades in both LSU and SSU analyses. *Ptycholaimellus*, *Hypodontolaimus*, and *Dichromadora* all show morphological similarities with each other and clades are weakly supported in the SSU analysis (posterior probability 88 in SSU). Differences between these three genera are slight, and they are clustered within one morphological group based on buccal cavity, peribuccal pharyngeal tissue, and supplements by Venekey et al. (2019).

Acknowledgements

We are grateful to Dr. Kenji Kito for his kind advice in the *Neochromadora parabilineata* sp. nov. identification, Dr. Huang Yong for sample collecting, and Dr. Wang Yanting for improving the manuscript. We are sincerely grateful to two anonymous referees and the Subject Editor Nic Smol for providing valuable improvements to the manuscript.

Additional information

Conflict of interest

The authors have declared that no competing interests exist.

Ethical statement

No ethical statement was reported.

Funding

This work was supported by A Project of Shandong Province Higher Educational Science and Technology Program (No. J18KA152) and Open Project of Liaocheng University Animal Husbandry Discipline (319312101).

Author contributions

Liang Huixin: phylogenetic analysis and description of *N. parabilineata*; Guo Wen: description of *D. media* and sample identification; Wang Chunming: Ms draft

Author ORCIDs

Huixin Liang  <https://orcid.org/0009-0006-9073-2062>

Wen Guo  <https://orcid.org/0000-0002-4452-0003>

Chunming Wang  <https://orcid.org/0000-0003-3630-0921>

Data availability

All of the data that support the findings of this study are available in the main text.

References

- Allgén CA (1940) Über einige neue freilebende Nematoden von der Nordwest- und Nordküste Norwegens. *Folia Zoologica et Hydrobiologica* 10: 443–449.
- Allgén CA (1958) Über einige freilebende marine Nematoden von der Ostküste Südamerikas (Uruguay, Nordküste Argentinas). *Zoologischer Anzeiger* Bd. 160: Heft 9/10: 205–217.
- Blaxter ML, De Ley P, Garey JR, Liu LX, Scheldeman P, Vierstraete A, Vanfleteren JR, Mackey LY, Dorris M, Frisse LM, Vida JT, Thomas WK (1998) A molecular evolutionary framework for the phylum Nematoda. *Nature* 392(6671): 71–75. <https://doi.org/10.1038/32160>
- Blome D (1974) Zur Systematik von Nematoden aus dem Sandstrand der Nordseeinsel Sylt. *Mikrofauna Meeresbodens* 33: 77–99.
- Boucher G (1976) Nématodes des sables fins infralittoraux de la Pierre Noire (Manche occidentale) II. Chromadorida. *Bulletin du Muséum National d'Histoire Naturelle*, 3e série, no 352. *Zoologie*, 25–61.
- Bussau C (1993) Taxonomische und ökologische Untersuchungen an Nematoden des Peru-Beckens. PhD Thesis, University of Kiel, German.

- Chitwood BG (1933) A revised classification of the Nematoda. In: Cort WW, Hegner R, Root FM (Eds) *The Journal of Parasitology Papers in Helminthology Ninth Annual Meeting*, 20(2). Boston, Massachusetts (USA), December 1933, 115–148.
- Cobb NA (1914) Antarctic marine free-living nematodes of the Shackleton expedition. *Contribution to a Science of Nematology* 1: 1–33.
- Cobb NA (1920) One hundred new nemas (type species of 100 new genera). *Contributions to Science of Nematology* 9: 217–343.
- Cobb NA (1930) Marine free-living nemas. *Scientific Reports of the Australasian Antarctic Expedition (1911–1914)*, Series C. Zoology & Botany 6(7): 1–28.
- Dashchenko OI (2002) Three New Free-living Marine Nematode Species of the Subfamily Hypodontolaiminae (Chromadorida, Chromadoridae). *Zoologicheskii Zhurnal* 81(7): 771–778.
- De Coninck LA, Schuurmans Stekhoven JH (1933) The free-living marine nemas of the Belgian Coast. II With general remarks on the structure and the system of nemas. *Mémoires Institut Royal Des Sciences Naturelles de Belgique* 58: 3–163.
- de Man JG (1876) Onderzoeken over vrij in de aarde levende Nematoden. *Tijdschrift der Nederlandische Dierkundige Vereeniging* 2: 78–196.
- de Man JG (1893) Cinquième Note sur les Nématodes libres de la mer du Nord et de la Manche. *Mémoires de la Société Zoologique de France* 6: 81–125.
- Filipjev IN (1917) Un Nematode libre nouveau de la mer Caspienne, *Chromadorissa* gen. nov. (Chromadoridae, Chromadorini) (Novaia Svobodnaia Nematoda iz Kaspiiskogo Moria *Chromadorissa* Gen. Nov. (Chromadoridae, Chromadorini)). *Revue Zoologique Russe* 2: 24–30. [Russkii Zoologicheskii Zhurnal]
- Filipjev IN (1918) Free-living marine nematodes of the Sevastopol area. *Transactions of the Zoological Laboratory and the Sevastopol Biological Station of the Russian Academy of Sciences* 4(2): 1–350. [In Russian]
- Filipjev IN (1922) Encore sur les Nématodes libres de la mer Noire. *Trudy Stavropol'skogo Sel'skohzoistvennogo Instituta. Zoologia* 1: 83–184.
- Filipjev IN (1929) Les Nématodes libres de l'extrême du golfe de Finlande et de la baie de la Neva (in Russian). *Études de la Neva et de son Bassin* 5: 3–22.
- Gerlach SA (1951) Nematoden aus der Familie der Chromadoridae von den deutschen Küsten. *Kieler Meeresforsch* 8(1): 106–132.
- Gerlach SA (1952) Nematoden aus dem Küstengrundwasser. *Abhandlungen Akademie der Wissenschaften und der Literatur. Mainz Mathematisch-Naturwissenschaftliche Klasse* 6: 315–372.
- Gerlach SA (1953a) Die Nematodenbesiedlung des Sandstrandes und des Küstengrundwassers an der italienischen Küste. I. Systematischer teil. *Archivio Zoologico Italiano* 37: 517–640.
- Gerlach SA (1953b) Freilebende marine Nematoden aus dem Küstengrundwasser und aus dem Brackwasser der Chilenischen Küste. Lunds Universitets Årsskrift. N. F. Avd. 2 49(10): 1–37.
- Gerlach SA (1956) Brasilianische Meeres-Nematoden I. *Boletim do Instituto Oceanográfico Tomo 5(1–2)*: 3–69.
- Gerlach SA (1957) Die Nematodenfauna des Sandstrandes na der küste von Mittelb (Brasilianische Meeres-Nematoden IV). *Mitteilungen aus dem Zoologischen Museum in Berlin* 33(2): 411–459. <https://doi.org/10.1002/mmzn.19570330206>
- Gerlach SA, Riemann F (1973/1974) The Bremerhaven Checklist of Aquatic Nematodes: A catalog of Nematoda Adenophorea excluding the Dorylaimida. *Veröffentlichungen des Instituts für Meeresforschung in Bremerhaven, Supplement 4*, 736 pp.

- Guo W, Wang MN, Li HT, Wang CM (2023) Description of a new nematode species, *Chromadorina communis* sp. nov. (Nematoda, Chromadoridae), from Changdao Island, China and phylogenetic analysis of Chromadorida based on small subunit rRNA gene sequences. ZooKeys 1159: 121–131. <https://doi.org/10.3897/zookeys.1159.100908>
- Higgins RP, Thiel H (1988) Introduction to the Study of Meiofauna. Smithsonian Institute Press, Washington DC, 488 pp.
- Hodda M (2022) Phylum Nematoda: A classification, catalogue and index of valid genera, with a census of valid species. Zootaxa 5114(1): 1–289. <https://doi.org/10.11646/zootaxa.5114.1.1>
- Holovachov O (2020) The nomenclatural status of new nematode nomina proposed in 1993 in the doctoral thesis of Christian Bussau, entitled Taxonomische und ökologische Untersuchungen an Nematoden des Peru-Beckens (Nematoda). Bionomina 19(1): 86–99. <https://doi.org/10.11646/bionomina.19.1.5>
- Hopper BE (1961) Marine nematodes from the coastline of the Gulf of Mexico. Canadian Journal of Zoology 39(2): 183–199. <https://doi.org/10.1139/z61-023>
- Hopper BE (1963) *Punctodora exochopora* n. sp. (Chromadoridae: Nematoda) from the Canadian shore of Lake Ontario. Canadian Journal of Zoology 41(6): 1121–1126. <https://doi.org/10.1139/z63-088>
- Holterman M, Holovachov O, van den Elsen S, van Megen H, Bongers T, Bakker J, Helder J (2008) Small subunit ribosomal DNA-based phylogeny of basal Chromadaria (Nematoda) suggests that transitions from marine to terrestrial habitats (and vice versa) require relatively simple adaptations. Molecular Phylogenetics and Evolution 48(2): 758–763. <https://doi.org/10.1016/j.ympev.2008.04.033>
- Huang Y, Zhang ZN (2010) Three new species of *Dichromadora* (Nematoda: Chromadorida: Chromadoridae) from the Yellow Sea, China. Journal of Natural History 44(9–10): 545–558. <https://doi.org/10.1080/00222930903471456>
- Inglis WG (1969) Convergence in the structure of the head and cuticle of *Euchromadora* species and apparently similar nematodes. Bulletin of the Natural History Museum Zoology Series 17(5): 149–204. <https://doi.org/10.5962/p.10339>
- Kreis HA (1929) Freilebende marine Nematoden von der Nordwest-Küste Frankreichs (Trébeurden Côtes du Nord). Capita Zoologica II 7: 1–98.
- Kreis HA (1963) Marine Nematoda. The Zoology of Iceland 2(14): 1–68.
- Kito K (1978) Studies on the free-living marine nematodes from Hokkaido, III. Journal of the Faculty of Science Hokkaido University Series VI. Zoology : Analysis of Complex Systems, ZACS 21(2): 248–261.
- Kito K (1981) Studies on the free-living marine nematodes from Hokkaido, IV. Journal of the Faculty of Science Hokkaido University Series VI. Zoology : Analysis of Complex Systems, ZACS 22(3): 250–278.
- Leduc D, Zhao Z (2015) *Latronema whataitai* sp. n. (Nematoda: Selachinematidae) from intertidal sediments of New Zealand, with notes on relationships within the family based on preliminary 18S and D2-D3 phylogenetic analyses. Nematology 17(8): 941–952. <https://doi.org/10.1163/15685411-00002915>
- Leduc D, Verdon V, Zhao ZQ (2017) Phylogenetic position of the Paramicrolaimidae, description of a new *Paramicrolaimus* species and erection of a new order to accommodate the Microlaimoidea (Nematoda: Chromadorea). Zoological Journal of the Linnean Society 183(1): 52–69. <https://doi.org/10.1093/zoolinnean/zlx072>
- Lemzina LV (1982) New Species of Free-living Nematodes of the Order Chromadorida from Lake Issyk-Kul (Novye Vidy Svobodnozhivushchikh Nematod Otriada Chromadorida iz Ozera Issyk-Kul'). Zoologicheskii Zhurnal 61(5): 780–783.

- Lo Russo V, Pastor de Ward CT (2012) *Neochromadora alejandroi* sp. n. (Chromadorida: Chromadoridae) and *Cobia macrodentata* sp. n. (Monhysterida: Xyalidae), two new species of free-living marine nematodes from the Patagonian coast. *Nematology* 14(7): 805–815. <https://doi.org/10.1163/156854112X627327>
- Long PK, Gagarin VG, Tu ND (2022) Two New Species of Chromadorids (Nematoda, Chromadorida) from the Water Bodies of Vietnam. *Inland Water Biology* 15(3): 238–248. <https://doi.org/10.1134/S1995082922030038>
- Lorenzen S (1966) Diagnosen einiger freilebender Nematoden von der schleswig-holsteinischen Westküste. *Veröffentlichungen des Instituts für Meeresforschung in Bremerhaven* X: 31–48.
- Lorenzen S (1971) Die Nematodenfauna im Verklappungsgebiet für Industrieabwasser nordwestlich von Helgoland: II. Desmodorida und Chromadorida. *Zoologischer Anzeiger* 187(5/6): 283–302.
- Lorenzen S (1973) Freilebende Meerenematoden aus dem Sublitoral der Nordsee und der Kieler Bucht. *Veröffentlichungen des Instituts für Meeresforschung in Bremerhaven* 14: 103–130.
- Lorenzen S (1981) Entwurf eines phylogenetischen Systems der freilebenden Nematoden. *Veröffentlichungen des Instituts für Meeresforschung in Bremerhaven* 7: 1–472.
- Lorenzen S (1994) The phylogenetic systematics of free-living nematodes. The Ray Society Institute, London, 383 pp.
- McIntyre AD, Warwick RM (1984) Meiofauna techniques. In: Holme NA, McIntyre AD (Eds) *Methods for the Study of Marine Benthos*. Blackwell Scientific Publications, Oxford, 217–244.
- Micoletzky H (1922) Die frielebenden Erdnematoden. *Archiv für Naturgeschichte* 87: 1–650.
- Micoletzky H (1924) Letzter Bericht über freilebende Nematoden aus Suez. *Sber. Akad. Wiss. Wien Mathem-naturw. Klasse. Abteilung I*, Band 133, Heft 4/6: 137–179.
- Muthumbi A, Vincx M (1998) Chromadoridae (Chromadorida: Nematoda) from the Indian Ocean: Description of new and known species. *Hydrobiologia* 364(2/3): 119–153. <https://doi.org/10.1023/A:1003109929194>
- Nunn GB (1992) Nematode molecular evolution. PhD Thesis, University of Nottingham, Nottingham.
- Pastor de Ward CT (1985) Free-living marine nematodes of the Deseado River estuary (Chromadoroidea: Chromadoridae, Ethmolaimidae, Cyatholaimidae and Choniolaimidae) Santa Cruz, Argentina. 5. Centro Nacional Patagónico Publicaciones Especiales 6: 1–83.
- Riemann F (1966) Die interstitielle Fauna im Elbe-Aestuar. *Verbreitung und Systematik. Archiv Für Hydrobiologie (Suppl.)* 31(1/2): 1–279.
- Riemann F (1976) Meeresnematoden (Chromadorida) mit lateralen Flossensäumen (Alae) und dorsoventraler Abplattung. *Zoologische Jahrbücher. Abteilung für Systematik* 103: 290–308.
- Schneider W (1937) Freilebende Nematoden der Deutschen Limnologischen Sundaexpedition nach Sumatra, Java und Bali. *Archiv für Hydrobiologie, Suppl.-Bd. 15 (Tropische Binnengewässer)*, Band VII: 30–108.
- Schneider W (1943) Freilebende Nematoden aus dem Ohridsee. *Posebna izdanja. Srpska Kralevska Akademija Nauka i Umetnosti. Nauke Prirodne i Matematicke (Spisi)* 136: 135–184.
- Schuurmans Stekhoven JH (1935) Additional notes to my monographs on the free-living marine nemas of the Belgian coast I and II, written in collaboration with W. Adam and

- L.A. De Coninck, with some remarks on the ecology of Belgian nemas. Mémoires du Musée Royal d'Histoire Naturelle de Belgique 72: 1–36.
- Schuurmans Stekhoven JH (1950) The Free-living Marine Nemas of the Mediterranean I. The Bay of Villefranche. Mémoires Institut Royal Des Sciences Naturelles de Belgique 2(37): 1–220.
- Steiner G (1916) Freilebende Nematoden aus der Barentssee. Zoologische Jahrbücher 39: 511–664.
- Steiner G (1921) Beiträge zur Kenntnis mariner Nematoden. Zoologische Jahrbücher (Systematik) 44(1–2): 1–68.
- Tchesunov AV (2014) Order Chromadorida Chitwood, 1933. In: Schmidt-Rhaesa A (Ed.) Gastrotricha, Cycloneuralia, Gnathifera (Vol. 2), Nematoda. Handbook of Zoology. De Gruyter, Berlin, 373–398. <https://doi.org/10.1515/9783110274257.373>
- Thanh NV, Tu ND, Gagarin VG (2016) *Dichromadora rigida* sp. n. (Nematoda, Chromadorida) from mangroves in the Yen River Estuary (Vietnam). Zoologicheskii Zhurnal 95(7): 779–787. <https://doi.org/10.7868/S0044513416050081>
- Timm RW (1961) The Marine Nematodes of the Bay of Bengal. Proceedings of the Pakistan Academy of Science 1(1): 2–88.
- Timm RW (1978) Redescription of the marine nematodes of Shackleton's British Antarctic Expedition of 1907–1909. American Geophysical Union. Antarctic Research Series 26(6): 237–255. <https://doi.org/10.1029/AR026p0237>
- Venekey V, Gheller PF, Kandratavicius N, Cunha BP, Vilas-Boas AC, Fonseca G, Maria TF (2019) The state of the art of Chromadoridae (Nematoda, Chromadorida): A historical review, diagnoses and comments about valid and dubious genera and a list of valid species. Zootaxa 4578(1): 1–67. <https://doi.org/10.11646/zootaxa.4578.1.1>
- Vermeeren H, Vanreusel A, Vanhove S (2004) Species distribution within the free-living marine nematode genus *Dichromadora* in the Weddell Sea and adjacent areas. Deep-sea Research. Part II, Topical Studies in Oceanography 51(14–16): 1643–1664. <https://doi.org/10.1016/j.dsr2.2004.06.028>
- Vincx M (1986) Free-living marine nematodes from the Southern Bight of the North Sea. I. Notes on species of the genera *Gonionchus* Cobb, 1920, *Neochromadora* Micoletzky, 1924 and *Sabatieria* Rouville, 1903. Hydrobiologia 140(3): 255–286. <https://doi.org/10.1007/BF00007440>
- Wieser W (1954a) Free-living marine nematodes II. Chromadoroidea. Report from the Lund University Chile Expedition Acta University Lund (N.F.2) 50(16): 1–148.
- Wieser W (1954b) Untersuchungen über die algenbewohnende Mikrofauna mariner Hartböden III. Zur Systematik der freilebenden Nematoden des Mittelmeeres. Mit einer ökologischen Untersuchung über die Beziehung zwischen Nematodenbesiedlung und Sedimentreichtum des Habitats. Hydrobiologia 6(1–2): 144–217. <https://doi.org/10.1007/BF00039417>
- Wieser W (1956) Eine Sammlung mariner Nematoden aus Piraeus (Griechenland). Österreichisch-Zoologisch Z 6(3/5): 597–630.
- Wieser W (1959) Free-living nematodes and other small invertebrates of Puget Sound beaches. University of Washington Publications in Biology (University of Washington Press, Seattle) 19: 1–179.
- Zhao ZQ, Li DM, Davies KA, Ye WM (2015) *Schistonchus zealandicus* n. sp. (Nematoda: Aphelenchoididae) associated with *Ficus macrophylla* in New Zealand. Nematology 17(1): 53–66. <https://doi.org/10.1163/15685411-00002851>