

Pisachini planthoppers of Vietnam: new records of *Pisacha* and a new *Goniopsarites* species from Central Vietnam (Hemiptera, Fulgoromorpha, Nogodinidae)

Jérôme Constant¹, Thai-Hong Pham^{2,3}, Cuong Viet Canh Le², Trung Thanh Vu⁴,
Hoai Thu Thi Nguyen⁴, Hai Nam Tran^{3,5}

1 Royal Belgian Institute of Natural Sciences, O.D. Taxonomy & Phylogeny – Entomology, Vautier street 29, B-1000 Brussels, Belgium

2 Mien Trung Institute for Scientific Research, Vietnam National Museum of Nature, VAST, 321 Huynh Thuc Khang, Hue, Vietnam

3 Graduate School of Science and Technology, Vietnam Academy of Science and Technology, 18 Hoang Quoc Viet, Hanoi, Vietnam

4 Vietnam National Museum of Nature, Vietnam Academy of Science and Technology (VAST), 18 Hoang Quoc Viet, Hanoi, Vietnam

5 Department of Biology, Hanoi National University of Education, 136 Xuan Thuy, Cau Giay, Hanoi, Vietnam

Corresponding authors: Thai-Hong Pham (phamthai@vnmn.vast.vn); Jérôme Constant (jerome.constant@naturalsciences.be)

Abstract

Two planthopper species of the family Nogodinidae are added to the fauna of Vietnam, both from two localities in Thua Thien-Hue Province: Bach Ma National Park and Phong Dien District. The first species belongs to *Goniopsarites* Meng, Wang & Wang, 2014, *G. mienrunganus* Constant & Pham, **sp. nov.**, and the second belongs to *Pisacha* Distant, 1906, *P. yinggensis* Meng, Wang & Wang, 2014. *Pisacha yinggensis* was previously recorded from Hainan Island, China. These new records greatly extend the distribution of both genera, which were known from southern China, Hainan and North Vietnam, to the south, reaching the mid area of Central Vietnam. Sexual dimorphism is reported in *P. yinggensis* for the first time. Illustrations of habitus and male terminalia of the new species are given as well as distribution maps and photographs of live specimens and their habitat. The family Nogodinidae now comprises nine species in Vietnam, with three of them present in Bach Ma National Park.

Key words: Bach Ma National Park, biodiversity, Fulgoroidea, Indochina, Phong Dien District



Academic editor: Yalin Zhang

Received: 31 October 2023

Accepted: 18 February 2024

Published: 6 March 2024

ZooBank: <https://zoobank.org/311FA5D6-CBA5-4C73-8811-861D5A27F8FF>

Citation: Constant J, Pham T-H, Le CVC, Vu TT, Nguyen HTT, Tran HN (2024) Pisachini planthoppers of Vietnam: new records of *Pisacha* and a new *Goniopsarites* species from Central Vietnam (Hemiptera, Fulgoromorpha, Nogodinidae). ZooKeys 1193: 181–194. <https://doi.org/10.3897/zookeys.1193.114957>

Copyright: © Jérôme Constant et al.

This is an open access article distributed under terms of the Creative Commons Attribution License ([Attribution 4.0 International – CC BY 4.0](https://creativecommons.org/licenses/by/4.0/)).

Introduction

The family Nogodinidae Melichar, 1898 is distributed worldwide in the tropics and subtropics and contains 378 species in 99 genera, including 11 species in Vietnam (Bourgoin 2023). Of these 11 species, five were described in the last 10 years (Constant and Pham 2014; Gnezdilov and Constant 2014; Meng et al. 2014; Constant and Pham 2016). Within the subfamily Nogodininae Melichar, 1898, the tribe Pisachini Fennah, 1978 was recently reviewed by Meng et al. (2014) and currently counts 10 species in three genera distributed in Southeast Asia (Constant and Pham 2016; Bourgoin 2023).

Study of the recent material of Nogodinidae in the collections of the Vietnam National Museum of Nature and Royal Belgian Institute of Natural Sciences revealed two species of Pisachini from Central Vietnam, which are new to the fauna of the country, including a species of *Goniopsarites* Meng, Wang & Wang, 2014 new to science.

The present paper describes this new species and provides the first Vietnamese records of *Pisacha yinggensis* Meng, Wang & Wang, 2014 as a new contribution to the Vietnamese nogodinid fauna.

Material and methods

The terminalia were extracted after soaking the abdomen overnight in a 10% solution of potassium hydroxide (KOH) at room temperature. The pygofer was separated from the abdomen, thoroughly rinsed in 70% ethanol, and the aedeagus dissected with a needle blade for examination. The whole was then placed in glycerin for preservation in a tube attached to the pin of the corresponding specimen. The hind wing was mounted, glued on a white cardboard rectangle with white glue, and the cardboard attached to the pin of the specimen. Photographs of collection specimens were taken with a Leica EZ4W stereomicroscope, stacked with CombineZ software, and optimized with Adobe Photoshop software; photographs from the field were taken with an Olympus Tough 6 camera. The maps were produced with SimpleMappr (Shorthouse 2010) and include records available from Meng et al. (2014) and Constant and Pham (2016). The external morphological terminology follows O'Brien and Wilson (1985), the wing venation terminology follows Bourgoin et al. (2015) and, for the male terminalia, Bourgoin and Huang (1990). The metatibiotarsal formula gives the number of spines on (side of metatibia) apex of metatibia / apex of first metatarsomere / apex of second metatarsomere.

The measurements were taken as by Constant (2004) and the following acronyms are used:

BB	maximum breadth of the body;
BF	maximum breadth of the frons;
BTg	maximum breadth of the tegmen;
BV	maximum breadth of the vertex;
LF	length of the frons at median line;
LT	total length (apex of head to apex of tegmina);
LTg	length of the tegmen;
LV	length of the vertex at median line.

Acronyms used for the collections:

RBINS	Royal Belgian Institute of Natural Sciences, Brussels, Belgium;
VNMMN	Vietnam National Museum of Nature, Hanoi, Vietnam.

Other abbreviations:

CCRR	Centre for Conservation of Vietnam Natural Resources and Rescue of Animals and Plants.
-------------	--

Taxonomy

Family Nogodinidae Melichar, 1898

Subfamily Nogodininae Melichar, 1898

Tribe Pisachini Fennah, 1978

Genus *Goniopsarites* Meng, Wang & Wang, 2014

Goniopsarites Meng et al., 2014: 80, figs 1–27.

Type species. *Goniopsarites fronticonvexus* Meng, Wang & Wang, 2014, by original designation.

Distribution. Southern China (Guangdong, Hainan); North and Central Vietnam.

Species included.

G. fronticonvexus Meng, Wang & Wang, 2014 – China: Hainan and Guangdong.

G. mientrunghanus Constant & Pham, sp. nov. – Vietnam: Bach Ma National Park and Phong Dien CCRR.

G. tonkinensis Constant & Pham, 2016 – Vietnam: Ba Be and Cuc Phuong national parks, and Me Linh Biodiversity Station.

Key to the species of *Goniopsarites*

- 1 Posterior processes of the periandrium (*ppp* – Fig. 2D) large and laminate, curved dorsocephalad in lateral view; posterodorsal process of the gonostyli bulging apically (*G* – Fig. 2A); basiventral lobe of the proximal half of the anal tube strongly developed in lateral view (*An* – Fig. 2A)
.....***Goniopsarites mientrunghanus* Constant & Pham, sp. nov.**
- Posterior processes of the periandrium not large and laminate, but instead indistinct (Meng et al. 2014: fig. 19) or slender and curved posterodorsad (Constant and Pham 2016: fig. 4C); posterodorsal process of the gonostyli not bulging apically (Meng et al. 2014: fig. 18; Constant and Pham 2014: fig. 4A); basiventral lobe of the proximal half of the anal tube moderately developed in lateral view (Meng et al. 2014: fig. 18; Constant and Pham 2014: fig. 4A)..... **2**
- 2 Anal tube strongly curved and very wide apically in lateral view (Constant and Pham 2016: fig. 4A) and with lateral margins strongly sinuate and narrowing towards base in dorsal view (Constant and Pham 2016: fig. 4B); apex of aedeagus more elongate and curved ventrally at apex (Constant and Pham 2016: fig. 4C); lateral processes of aedeagus strongly sinuate (Constant and Pham 2016, fig. 4C); posterior processes of the periandrium slender and curved posterodorsad (Constant and Pham 2016: fig. 4C).....***Goniopsarites tonkinensis* Constant & Pham, 2016**
- Anal tube moderately curved, and roundly truncate apically in lateral view (Meng et al. 2014: fig. 18) and with lateral margins weakly sinuate and wider towards base in dorsal view (Meng et al. 2014: fig. 14); apex of aedeagus roundly truncate and not curved ventrally at apex (Meng et al. 2014: fig. 19); lateral processes of aedeagus regularly curved in lateral view (Meng et al. 2014: fig. 19); posterior processes of the periandrium indistinct (Meng et al. 2014: figs 19, 20)
.....***Goniopsarites fronticonvexus* Meng, Wang & Wang, 2014**

***Goniopsarites mientrunganus* Constant & Pham, sp. nov.**

<https://zoobank.org/689317C8-D1E4-4526-81CC-E89B9C11E42B>

Figs 1–4A

Type material. *Holotype* ♂, VIETNAM • Thừa Thiên-Huế Province, Bach Ma National Park, near ranger station; 16°08'37"N, 107°49'36"E; 18 May 2023; alt. 300–600 m; J. Constant & L. Semeraro leg.; VNMN_E000.017.000.

Paratypes, VIETNAM • 1 ♀; Thừa Thiên-Huế Province, Bach Ma National Park; 16°13'14"N, 107°53'10"E; 9 Mar. 2023; by net; V.T. Trung leg.; VNMN_E000.017.001 • 1 ♂; Thừa Thiên-Huế Province, Bach Ma National Park, Pheasant trail; 16°13'38"N, 107°51'20"E; 10–20 May 2023; alt. 500–600 m; J. Constant & L. Semeraro leg.; I.G.: 34.640; RBINS • 2 ♂♂, 1 ♀; Thừa Thiên-Huế Province, Bach Ma National Park; low altitude; 16°13'05"N, 107°42'27"E; 17 May 2023; alt. 100–200 m; J. Constant & L. Semeraro leg.; I.G.: 34.640; RBINS • 1 ♀; Thừa Thiên-Huế Province; Phong Dien District, CCRR; 16°30'27"N, 107°16'05"E; 23 May 2023; alt. 350–400 m; J. Constant & L. Semeraro leg.; I.G.: 34.640; RBINS • 1 ♀; same collection data as for preceding; VNMN_E000.017.002.

Diagnosis. The species is very close externally to *G. fronticonvexus* Meng, Wang & Wang, 2014 and *G. tonkinensis* Constant & Pham, 2016 but has slightly more elongate tegmina, 2.2 times as long as wide (1.9 times in *G. fronticonvexus*; 2.0 times in *G. tonkinensis*). These species are better separated by the male genitalia characters as follows.

Goniopsarites mientrunganus Constant & Pham, sp. nov. can be separated from the other two species by (1) its large laminate posterior processes of the periandrium (*ppp* – Fig. 2D), which are absent in *G. fronticonvexus* and much more slender, curved the other way round, in *G. tonkinensis*; (2) the posterodorsal process of the gonostyli bulging apically (*G* – Fig. 2A), not bulging in both other species; and (3) the strongly developed basiventral lobe of the proximal half of the anal tube (*An* – Fig. 2A), clearly less developed in both other species.

From *G. tonkinensis*, *G. mientrunganus* Constant & Pham, sp. nov. also differs in (1) having the maximum width of the anal tube in the basal third (*An* – Fig. 2C), which is in the distal half of the anal tube in *G. tonkinensis* (the state of this character is not known for *G. fronticonvexus*); (2) in having a strongly developed dorsal laminate process along basal half of periandrium (*dlp* – Fig. 2D, F, G).

Description. Measurements and ratios: LT: ♂ (*n* = 4): 11.3 mm (11.1–11.7); ♀ (*n* = 4): 13.5 (13.0–14.5). LTg/BTg = 2.2; LW/BW = 1.4; LV/BV = 8.5; LF/BF = 1.4.

Head: (Fig. 1A–E) vertex dark brown, with yellowish median and lateral carinae; concave with lateral margins carinate, and with anterior and posterior margins slightly carinate and rather strongly concave in dorsal view, resulting in a very narrow central portion. Frons varying from variegated brown and yellowish, darker under V-shaped carina, carina yellowish to reddish, to nearly completely black-brown, sometimes with yellowish markings around mid-height of lateral margin; narrow, well-defined, yellowish line above frontoclypeal suture; frons elongate, broader dorsally, concave with median carina on middle of disc and with a strong projection on ventral half marked by strong, V-shaped carina extending ventrad on clypeus; projection rounded in lateral view; lateral margins of frons carinate; frontoclypeal suture grooved and rounded ventrally. Genae largely variegated yellow-brown, dark brown under antennae, with pale yellowish spot along anterior margin. Clypeus yellowish, variegated with black-brown




Figure 1. *Goniopsarites mientrunganus* Constant & Pham, sp. nov. **A–F** holotype ♂ **A** habitus, dorsal **B** habitus, ventral **C** habitus, lateral **D** habitus, perpendicular view of frons **E** habitus, anterolateral **F** right hind wing **G** paratypes ♀♀, posterior margin of sternite VII.

on sides towards apex, and with brown oblique lines on each side; clypeus elongate, narrower and shorter than frons, and showing a strong median carina; roundly convex in lateral view. Labium elongate and narrow, yellow-brown, slightly surpassing posterior coxae. Antennae black with scape ring-shaped; pedicel subglobose.

Thorax: (Fig. 1A, C–E) pronotum variegated yellow-brown, darker in middle portion, rather densely pitted, and with median carina weakly marked; lateral fields of prothorax coloured as pronotum, darker in ventral portion, widening ventrally and with ventral margin rounded. Mesonotum dark brown, with yellowish markings on sides and yellowish spot on scutellum; lateral fields moderately pitted; median carina obsolete; lateral carinae marked with yellowish, fusing anteriorly in a rounded carina parallel to posterior margin of pronotum. Tegulae brown.

Tegmina: (Fig. 1A–C) elongate, with costal margin broadly rounded on proximal half and rather strongly sinuate on posterior half; brown with large, irregular, black-brown marking on basal half, not extending on clavus; large, pale yellowish marking along costal margin in distal half; clavus with irregular, slightly darker markings on posterior half; hypocostal plate narrow, visible in proximal third; posterior margin obliquely rounded; clavus closed, extending to posterior angle of tegmen. Veins reddish brown, sometimes black in black areas.

Hind wings: (Fig. 1F) brown, darker towards posterocostal angle and with large basicostal yellow-brown area; well developed, with posterior margin trilobed; costal margin sinuate, with coupling apparatus at 2/3 of length.

Legs: (Fig. 1A–C) profemora black-brown, with narrow, pale yellow rings; mesofemora pale yellow, with brown rings; pro- and mesotibiae elongate and slender, with 6 rings alternatively pale yellow and brown, basal one brown, apical one yellow; metafemora yellow, with brown markings; metatibiae rather short, broadening towards apex, pale yellow, infuscate basally, with 2 strong lateral spines near apex. Tarsi brown, darker apically. Metatibiotarsal formula: (2) 10 / 2 (+7 on underside) / 2.

Abdomen: (Fig. 1B) pale yellowish brown.

Terminalia ♂: (Fig. 2) pygofer (*Py*) higher than long in lateral view, with anterior and posterior margins moderately sinuate; posterodorsal angles with dorsally developed, subtriangular laminate process, with posterodorsal angle right and rounded. Anal tube (*An*) massive, elongate, with lateral margins in dorsal view, widely rounded in proximal half, then sinuately tapering towards posterior, leaving widely rounded ventral margin visible on sides in distal half; 1.78 times as long in midline, as wide with maximum width at proximal 1/3 in dorsal view; basal half in lateral view with large ventral lobe; strongly curved ventrally at midlength with lateral margins strongly produced ventrally into broad lobe in distal half. Gonostyli (*G*) (Fig. 4A) elongate in lateral view, rounded apically and with strong process projecting dorsomesad at posterodorsal angle; ventral margin nearly straight and posterior margin weakly sinuate in lateral view; dorsal process slightly twisted internally, bulging in distal portion and with apex with blunt point directed anteromesad. Aedeagus (Fig. 2D–I) strongly curved in lateral view, with a pair of lateral, elongate, strongly sinuate processes (*lpa*) directed anteriorly, attached posteriorly at 2/3 of length; processes well visible and sinuate in ventral view. Periandrium posteriorly with a pair of large laminate processes (*ppp*), lobe-shaped and developed dorsally into sinuate, elongate, narrowly pointed processes directed mesodorsad; well-developed




Figure 2. *Goniopsarites mientrunganus* Constant & Pham, sp. nov., holotype ♂, terminalia **A–C** pygofer, gonostyli and anal tube **A** lateral **B** caudal **C** dorsal **D–I** aedeagus, phallobase and connective **D** left lateral **E** dorsal **F** lateroventral **G** laterodorsal **H** caudal **I** ventral. *cv* = connective; *dlp* = dorsal laminate process of periandrium; *lpa* = lateral process of aedeagus; *ppp* = posterior process of periandrium; *spp* = sclerotized process of phallus; *td* = tectiductus; *vlp* = ventral lobe of periandrium.



Figure 3. *Goniopsarites mientrunganus* Constant & Pham, sp. nov., in nature and habitat **A–C** Bach Ma National Park, low altitude, 15 May 2023 **A** adult on stem **B** habitat **C** nymph **D–F** Bach Ma National Park, Pheasant Trail, 20 May 2023 **D** habitat **E** adult on tree trunk **F** idem, general view **G, H** Phong Dien District, CCRR, 23 May 2023 **G** adult on stem **H** habitat.




Figure 4. Distribution maps **A** *Goniopsarites* species in Vietnam **B** *Pisacha yinggensis* Meng, Wang & Wang, 2014.

lamine process (*dip*) running dorsally in basal half, abruptly terminated at right angle at midlength. Ventral lobe of perianthrium (*vlp*) elongate, lanceolate in distal portion in ventral view, with sides rounded. Phallus membranous, large, with pair of curved, sclerotized processes (*spp*) directed cephalodorsad before apex; strongly, angularly projecting posterodorsad; apex narrowing, directed anteriorly, and curved ventrally. Connective (*cv*) with well-developed, elongate tectiductus (*td*) showing complete, broadly rounded dorsal crista.

Note. The female genital structures were checked and found to be similar to those of *G. fronticonvexus*, as well as to these of *G. tonkinensis*, and, as it is the

case in most nogodinids that we have observed so far, female genitalia do not help with species identification. The indentation in middle of the hind margin of sternum VII is always present but was found to vary in depth between the specimens of *G. mientrunganus* sp. nov. that we have in hand (Fig. 1G).

Etymology. The species epithet *mientrunganus* refers to the region where the new species was discovered: Central Vietnam, “Miền Trung” in Vietnamese.

Biology. The specimens were found sitting on stems of bushes (Fig. 3A, C, G) or on tree trunks (Fig. 3E, F), in subtropical evergreen forest (Fig. 3B, D, H) at the junction of the Northern Vietnam lowland rain forests, Southern Vietnam lowland rain forests, and Southern Annamites montane rain forests ecoregions, at rather low altitude (150–600 m). Some nymphs (Fig. 3C) were observed in May 2023 together with adult specimens.

Distribution. Vietnam, Thua Tinh-Hue Province, Bach Ma National Park, and Phong Dien district, CCRR (Fig. 4A).

Genus *Pisacha* Distant, 1906

Pisacha Distant, 1906: 391. Type species: *Pisacha naga* Distant, 1906, by original designation.

Soaemis Jacobi, 1915. Nomen nudum.

Soaemis Jacobi, 1916: 311. Type species: *Soaemis encaustica* Jacobi, 1916, by original designation. Synonymized by Ishihara (1965: 207).

Distribution. Southern China (Chongqing, Guangdong, Hainan, Sichuan, Zhejiang); India (Assam); Taiwan; North and Central Vietnam.

Species included.

P. baculiformis Meng, Wang & Wang, 2014 – China: Zhejiang.

P. balteiformis Meng, Wang & Wang, 2014 – Vietnam: Ninh Binh Province.

P. encaustica (Jacobi, 1916) – Taiwan.

P. falcata Meng, Wang & Wang, 2014 – China: Chongqing, Sichuan.

P. kwangsiensis Chou & Lu, 1977 – China: Guangxi.

P. naga Distant, 1906 – India: Assam.

P. yinggensis Meng, Wang & Wang, 2014 – China: Hainan; Vietnam: Thua Thien-Hue Province.

Pisacha yinggensis Meng, Wang & Wang, 2014

Figs 4B, 5

Pisacha yinggensis Meng, Wang & Wang, 2014: 93, figs 38–41, 56–69, 115, 120.

Material examined. VIETNAM • 1 ♂; Thừa Thiên-Huế Province; Phong Dien District, CCRR; 16°30'27"N, 107°16'05"E; 23 May 2023; alt. 350–400m; J. Constant & L. Semeraro leg.; I.G.: 34.640; RBINS • 1 ♂, 2 ♀♀; Thừa Thiên-Huế Province, Bach Ma National Park, near ranger station; 16°08'37"N, 107°49'36"E; 18 May 2023; alt. 300–600 m; J. Constant & L. Semeraro leg.; VNMN_ E000.017.003 • Thừa Thiên-Huế Province, Bach Ma National Park; 16°12'N, 107°52'E; 12–17 Jul. 2011; J. Constant & J. Bresseel leg.; I.G.: 31.933; RBINS • 1 ♀; Thừa Thiên-Huế Province,




Figure 5. *Pisacha yinggensis* Meng, Wang & Wang, 2014, from Bach Ma National Park, Pheasant Trail, 10–20 May 2023 **A–C** male **A** lateral **B** head and thorax, dorsal **C** frons, perpendicular **D–F** female **D** lateral **E** head and thorax, dorsal **F** frons, perpendicular **G** adult female in nature **H** habitat.

Bach Ma National Park; summit; 16°11'18"N, 107°50'56"E; 11–21 May 2023; alt. 1300–1400 m; J. Constant & L. Semeraro leg.; I.G.: 34.640; RBINS • 1 ♂, 1 ♀; Thừa Thiên-Huế Province, Bach Ma National Park, Pheasant trail; 16°13'38"N,

107°51'20"E; 10–20 May 2023; alt. 500–600 m; J. Constant & L. Semeraro leg.; I.G.: 34.640; RBINS • 2 ♀♀; same collection data as for preceding; VNMN_ E000.017.004 • 1 ♂; same collection data as for preceding; 29 Mar. 2021; Malaise trap; V.T. Trung leg.; VNMN_ E000.017.005 • 1 ♀; Thừa Thiên-Huế Province; Phong Dien District, CCRR; 16°30'27"N, 107°16'05"E; 23 May 2023; 350–400m; Trung T. Vu leg.; VNMN_ E000.017.006.

Notes. The specimens from Central Vietnam show obvious sexual dimorphism, with the tegmina of the females marked with a contrasting, incomplete, dark brown to black band at the distal 2/3 of its length, and a broad, oblique area of the same colour, or slightly paler, along the apical margin (Fig. 5D, G). These dark markings on the tegmina are weakly visible or absent in males (Fig. 5A). The markings of the head and thorax, however, are similar in both sexes (Fig. 5B, C, E, F).

Biology and distribution. The specimens were found sitting on leaves or stems on the lower vegetation (Fig. 5G), in subtropical evergreen forest (Fig. 5H) in at the junction of Northern Vietnam lowland rain forests, Southern Vietnam lowland rain forests, and Southern Annamites montane rain forests ecoregions, at altitude from 300 to 1,400 m. In China, the species is recorded from the Hainan Island monsoon rain forests ecoregion.

The species was previously only recorded from Hainan Island, China (Meng et al. 2014). It is here recorded for the first time from the mainland, in Central Vietnam (Fig. 4B).

Discussion

The present work adds two species of Nogodinidae to the fauna of Vietnam, including one species described as new. This leads to a total of 13 species for the country. As a comparison, 10 species are known from China (Bourgoin 2023). The new records also greatly extend the Vietnamese distribution of the genera *Goniopsarites* and *Pisacha* to the south but leave a gap of over 500 km without any records of these genera in the northern half of Central Vietnam.

In Phong Dien District, VNMN is conducting an ambitious project of forest restoration at the Centre for Conservation of Vietnam Natural Resources and Rescue of Animals and Plants, not far from the forest where *G. mientrunganus* Constant & Pham, sp. nov. and *P. yinggensis* were found. As already recently mentioned for a new species of Tropiduchidae, *Connelicita phongdienensis* Constant & Pham, 2023, the return of such planthopper species in this area in the future would be a great indicator of a successful project (Constant et al. 2023).

Acknowledgements

We thank Dr Linda Semeraro and Mr Joachim Bresseel (scientific collaborators, RBINS) and our guides in Bach Ma National Park, Lâm Bá Vũ Nguyễn, and Quang Thanh Nguyễn for all their help and friendship during the collecting trips, the authorities of Bach Ma National Park and Centre for Conservation of Vietnam Natural Resources and Rescue of Animals and plants, Dr Frederik Hendrickx (RBINS), and Dr Thierry Backeljau (RBINS) for their support to our projects.

Additional information

Conflict of interest

The authors have declared that no competing interests exist.

Ethical statement

No ethical statement was reported.

Funding

The present study was funded by the Vietnam Academy of Science and Technology (VAST) under the grants number NCXS02.04/22-23, CT0000.04/22-24. The 2011 record of *P. yinggensis* was obtained from fieldwork of the Global Taxonomy Initiative project “A step further in the Entomodiversity of Vietnam”, supported through a grant issued by the capacity building Programme of the Belgian Global Taxonomy Initiative National Focal Point that runs under the CEBioS programme with financial support from the Belgian Directorate-General for Development Cooperation (DGD).

Author contributions

Conceptualization, J.C. and T.-H.P.; Funding acquisition, T.-H.P. and J.C.; Investigation, J.C.; Methodology, J.C. and T.-H.P.; Project administration, T.-H.P., C.V.C.L. and J.C.; Field work, J.C., T.-H.P., C.V.C.L., T.T.V., H.T.T.N.; Field logistics, T.-H.P., C.V.C.L., T.T.V., H.T.T.N.; Supervision, J.C. and T.-H.P.; Validation, J.C. and T.-H.P.; Writing – original draft, J.C.; Writing – review and editing, J.C., T.-H.P., C.V.C.L., T.T.V., H.T.T.N. All authors have read and agreed to the published version of the manuscript.

Author ORCIDs

Jérôme Constant  <https://orcid.org/0000-0003-0254-0863>

Thai-Hong Pham  <https://orcid.org/0000-0002-4763-3679>

Cuong Viet Canh Le  <https://orcid.org/0000-0002-1430-6305>

Trung Thanh Vu  <https://orcid.org/0009-0002-8987-1672>

Hoai Thu Thi Nguyen  <https://orcid.org/0009-0004-9130-2940>

Hai Nam Tran  <https://orcid.org/0009-0003-1557-1666>

Data availability

All of the data that support the findings of this study are available in the main text.

References

- Bourgoin T (2023) FLOW (Fulgoromorpha Lists on the Web): a world knowledge base dedicated to Fulgoromorpha. V.8, updated [i.2023] <http://hemiptera-databases.org/flow/> [Accessed 26 October 2023]
- Bourgoin T, Huang J (1990) Morphologie comparée des genitalia mâles des Trypetimorphini et remarques phylogénétiques (Hemiptera: Fulgoromorpha: Tropiduchidae). *Annales de la Société entomologique de France. Nouvelle Série* 26(4): 555–564. <https://doi.org/10.1080/21686351.1990.12277614>
- Bourgoin T, Wang RR, Asche M, Hoch H, Soulier-Perkins A, Stroinski A, Yap S, Szwe-do J (2015) From micropterism to hyperpterism: recognition strategy and standardized homology-driven terminology of the fore wing venation patterns in

- planthoppers (Hemiptera: Fulgoromorpha). *Zoomorphology* 134(1): 63–77. <https://doi.org/10.1007/s00435-014-0243-6>
- Constant J (2004) Révision des Eurybrachidae (I). Le genre *Amychodes* Karsch, 1895 (Homoptera: Fulgoromorpha: Eurybrachidae). *Bulletin de l'Institut royal des Sciences naturelles de Belgique* 74: 11–28.
- Constant J, Pham HT (2014) A new species of *Orthophana* Melichar, 1923 from Vietnam (Hemiptera: Fulgoromorpha: Nogodinidae). *Belgian Journal of Entomology* 23: 1–8.
- Constant J, Pham HT (2016) Two new Nogodinidae from Vietnam in the genera *Orthophana* Melichar, 1923 and *Goniopsarites* Meng, Wang & Wang, 2014 (Hemiptera: Fulgoromorpha: Nogodinidae). *Belgian Journal of Entomology* 40: 1–16.
- Constant J, Pham T-H, Le CVC, Vu TT, Nguyen HTT (2023) The tropiduchid planthopper genus *Connelicita* Wang & Bourgoin, 2015: Two new species from Central Vietnam and new records (Hemiptera, Fulgoromorpha, Tropiduchidae). *ZooKeys* 1186: 105–122. <https://doi.org/10.3897/zookeys.1186.113701>
- Gnezdilov VM, Constant J (2014) Notes on the tribe Tongini, with description of a new species of the genus *Orthophana* from northern Vietnam (Hemiptera: Fulgoroidea: Nogodinidae). *Acta Entomologica Musei Nationalis Pragae* 54(1): 47–55.
- Ishihara T (1965) Some species of Formosan Homoptera. *Special Bulletin of the Lepidoptera Society of Japan* 1: 207–208.
- Meng R, Wang M-L, Wang Y-L (2014) On the taxonomy of the tribe Pisachini (Hemiptera: Fulgoromorpha: Nogodinidae) with the description of new taxa from China and Vietnam. *Zootaxa* 3866(1): 79–104. <https://doi.org/10.11646/zootaxa.3866.1.4>
- O'Brien LB, Wilson SW (1985) Planthoppers systematics and external morphology. In: Nault LR, Rodriguez JG (Eds) *The Leafhoppers and Planthoppers*. John Wiley & Sons, New York, 61–102.
- Shorthouse DP (2010) SimpleMappr, an online tool to produce publication-quality point maps. <http://www.simplemappr.net> [Accessed 28 September 2023]