

A new species of the subgenus *Scymnus* from Pakistan (Coleoptera, Coccinellidae)

Azad Rashid¹, Xiaosheng Chen², Baoli Qiu¹, Xingmin Wang¹

1 Key Laboratory of Bio-Pesticide Innovation and Application, Engineering Technology Research Center of Agricultural Pest Biocontrol, Guangdong Province; Department of Entomology, South China Agricultural University, Guangzhou 510640, China **2** Department of Forestry Protection, College of Forestry and Landscape Architecture, South China Agricultural University, Guangzhou, 510640, China

Corresponding author: Baoli Qiu (baileyqiu@scau.edu.cn); Xingmin Wang (wangxmcn@scau.edu.cn)

Academic editor: M. Thomas | Received 23 March 2017 | Accepted 21 July 2017 | Published 29 August 2017

<http://zoobank.org/7C6E63B1-A91F-44C8-9CE4-4456C2EBFA3B>

Citation: Rashid A, Chen X, Qiu B, Wang X (2017) A new species of the subgenus *Scymnus* from Pakistan (Coleoptera, Coccinellidae). ZooKeys 694: 31–39. <https://doi.org/10.3897/zookeys.694.12863>

Abstract

A new species, *Scymnus* (*Scymnus*) *contortubus* Rashid, Chen & Wang, **sp. n.**, is described and illustrated from Pakistan. A diagnosis, remarks, illustrations, and a distribution map are provided of the new species and its most similar congener, *S. (S.) nubilus* Mulsant.

Keywords

Coccinelloidea, Coccidulini, entomology, new species, Pakistan, taxonomy

Introduction

The subgenus *Scymnus* was established by Mulsant, 1850 with *Coccinella rufipes* Fabricius, 1798 as the type species, based on the presence of an 11-segmented antenna and the incomplete abdominal postcoxal line.

The genus *Scymnus* Kugelann, 1794 belonged to the tribe Scymnini Mulsant, 1846 in the subfamily Scymninae (Sasaji 1968; Kovář 1996). However, Ślipiński (2007) presented a two-subfamily system, moving the genus *Scymnus* to the tribe Coccidulini in the subfamily Coccinellinae. Further studies based upon molecular

and morphological characters supported Ślipiński's (2007) division of Coccinellidae in two subfamilies, Coccinellinae and Microweiseinae (Giorgi et al. 2009; Seago et al. 2011). Recent study on the Cucujoidea (Robertson et al. 2015) has recovered Coccinellidae as belonging to the superfamily Coccinelloidea together with eight other families of the former Cerylonid Series.

Members of the genus *Scymnus* Kugelann, 1794 are predatory and mostly feed on aphids, adelgids, and scale insects, playing an important role in regulating pest populations (Chen et al. 2016). Prior to the present study, 58 species were reported in the subgenus *Scymnus* (*Scymnus*) from the Oriental Region and only one species, *S. (S.) nubilus* Mulsant, 1850, has been known to exist in Pakistan (Poorani, 2002; Chen et al. 2013). In this paper, another species is described, and a diagnosis of the subgenus with distribution pattern of both species is presented.

Materials and methods

Specimens were collected from different localities in Pakistan during 2015–2016 and were preserved in 85% ethanol. A Zeiss Stemi 305 microscope was used for observing external morphology followed by the dissection of male and female genital structures. After dissection, genitalia were cleared in 10% solution of NaOH and placed in a drop of neutral balsam onto glass slides for further studies.

SteREO Discovery V20 (Zeiss) microscope with an ocular micrometer was used for all measurements, which are presented in millimetres. The following abbreviations are used:

TL	total length from clypeus to apex of elytra,
TW	total width across both elytra at widest part,
TH	total height in highest part of elytra,
HW	head width in widest part including eyes,
PL	pronotal length across the central area from anterior to basal margin of pronotum,
PW	pronotal width across widest part,
EL	elytral length along suture including scutellum,
EW	elytral width, equivalent to TW.

An AxioCam HRc digital camera attached to the stereoscope, (SteREO Discovery V20) was used for photographs of the whole bodies of beetles. Composite images were generated with AXIO VISION REL. 4.8 software and edited using ADOBE PHOTOSHOP CC. 2017.

A compound microscope, Olympus BX51 attached to a Coolsnap-Procf & CRI Micro*Color camera was used for the preparation of illustrations of the morphological characters of male genitalia. Morphological terminology of Ślipiński (2007) was followed. Type specimens are deposited in the Department of Entomology, South China Agricultural University, Guangzhou, China.

Taxonomy

Genus *Scymnus* Kugelann, 1794

Subgenus *Scymnus* Kugelann, 1794

Scymnus Kugelann, 1794: 545. Type species: designated by Westwood 1838: 43, *Scymnus nigrinus* Kugelann, 1794.

Anisoscymnus Crotch, 1874: 273. Type species: *Coccinella rufipes* Fabricius, 1798, by original designation.

Diagnosis. Members of the subgenus *Scymnus* can be easily distinguished by the following combination of characters: body small, oval or elongate oval; antennae 11-segmented (Fig. 1f); prosternum having distinct carinae, nearly reaching the anterior margin (Fig. 1b); abdominal postcoxal line incomplete, recurved forward, never reaching lateral margin of ventrite; area surrounding by postcoxal line sparsely punctate; abdomen with six ventrites, 5th and 6th abdominal ventrites in male truncate or emarginate apically in male (Fig. 1i); female genitalia with distinct infundibulum.

Distribution. Worldwide (Chen et al. 2013).

Scymnus (Scymnus) contortubus Rashid, Chen & Wang, sp. n.

<http://zoobank.org/C6D02733-EB0F-4EC9-964D-4F6FABBCC6C2>

Figs 1, 2

Etymology. The species name is derived from Latin (*contortum* = twisted and *tubus* = tube) referring to a curved, tube-like apex of penis.

Diagnosis. This species is separated by the presence of a stout penis with a curved apex (Fig. 2d) and parameres with long, dense setae on both the apex and inner side (Fig. 2f–g). In *S. (S.) nubilus*, the most similar congener, penis is slender (Fig. 3e) and parameres have sparse long setae only at apices (Fig. 3g–h).

Description. TL: 1.8 mm, TW: 1.3 mm, TH: 0.7 mm, HW: 0.6 mm, TL/TW: 1.38, PL/PW: 0.33, EL/ EW: 1.07.

Body oval, distinctly convex, dorsum with dense white pubescence. Head brown with black vertex. Antennae and mouthparts brown. Pronotum black with lateral margins reddish brown. Elytra reddish brown with broad black U-shaped sutural stripe, extending 5/6 length of elytral suture (Fig. 2a–c). Lateral margins of elytra coarsely punctate. Prothoracic hypomeron reddish brown. Prosternum, mesoventrite, and metaventrite black. Elytral epipleuron yellowish brown with both margins black. Legs reddish brown.

Head small, width 0.66 times of pronotal width (HW/PW= 0.6/0.9). Head with fine punctures, same size as eye facets, separated by 0.5–1.0 diameters (Fig. 2b). Eyes densely faceted, with dense hairs, interocular distance 0.47 times head width. Pronotum width 0.69 times of elytral width (PW/EW=0.9/1.3), pronotal punctures sparse,




Figure 1. *Scymnus (Scymnus) contortubus* Rashid, Chen & Wang, sp. n.: **a** head, frontal **b** prothorax, antero-ventral **c** maxilla **d** labium **e** mandible **f** antenna **g** fore leg **h** hind leg **i** abdomen, male.

slightly coarser than those on head, 1.0–2.0 diameters apart. Elytral punctures coarse, separated by 2.0–4.0 diameters. Prosternal carinae distinct extending to anterior margin and slightly converging anteriorly. Prosternal process T-shaped, twice as long as its width at base. Abdominal postcoxal lines incomplete, extending to 6/7 length of ventrite and slightly recurved toward base of ventrite; area surrounding postcoxal line with dense granular punctures, distributed unevenly (Fig. 2e).

Male genitalia: Penis stout (Fig. 2d). Penis capsule with long inner process and short dilated outer one. Apex of penis strongly recurved towards inner side with thread-like appendage. Tegmen stout, with penis guide parallel-sided from base to nearly half length, tapering gradually to a blunt apex in ventral view. In lateral view, penis guide with sides parallel at basal 2/3, then abruptly constricted to pointed apex. Parameres stout; slightly longer than penis guide, with dense, long setae on apical and inner margins (Fig. 2f–g).




Figure 2. *Scymnus (Scymnus) contortubus* Rashid, Chen & Wang, sp. n.: **a** dorsal view **b** anterior view **c** lateral view **d** penis **e** abdomen **f** tegmen, lateral view **g** tegmen, ventral view **h** ovipositor **i** spermatheca.

Female genitalia: Coxites elongate triangular, 2.5 times as long as wide, tapering to blunt apices, each with long terminal setae (Fig. 2h); infundibulum present; spermatheca C-shaped (Fig. 2i).

Types. Holotype, Male, **PAKISTAN:** (Kashmir). Arja Mountains, No. SCAU (E) 16579, [N33°57' 27.80, E073°39' 54.04"], ca. 940 m, 28.X.2015, Huo LZ leg; Paratypes: (86) 2♂1♀, same data as holotype; 8♂1♀ Rawalakot, [N33°52' 07.93", E073°43' 46.49"] ca. 1668 m; 28.X.2015, Huo LZ leg; 12♂ 10♀ Mirpur, [N33°28' 23.07", E073°52' 57.94"] ca. 500–610 m, 26.X.2015; Huo LZ leg; (Khyber Pakhtunkhwa) 5♂ 5♀, Balakot, [N34°33' 27.62", E073°21' 25.08"] ca. 1093 m; 15.X.2015; Wang XM leg; 1♂ Birote, [N34°03' 27.54", E073°30' 00.32"] ca. 789 m; 13.X.2015; Wang XM leg; 4♂ 2♀ Arab khan, [N34°25' 20.73", E073°18' 44.21"] ca. 104 m; 16.X.2015; Wang XM leg; 5♂ 6♀ Shung, [N34°52' 31.52", E072°54' 12.46"] ca. 656 m; 17.X.2015; Wang XM leg; 9♂ 7♀ Parihna, [N34°21' 32.17", E073°04' 53.75"] ca. 775 m; 19.X.2015; Wang XM leg; 1♂ 1♀ Paras, [N34°39' 23.53", E073°30' 13.91"] ca. 1364 m; 15.X.2015; Wang XM leg; (Punjab) 1♂ 2♀ Salgran, [N33°49', 33.96", E073°17' 09.48"] ca. 857 m; 13.X.2015; Wang XM leg; 2♂ 1♀ Gokina [N33°46', 03.26", E073°04' 37.78"] ca. 1090 m; 11.X.2015; Wang XM leg.

Distribution. Pakistan (Kashmir, Khyber Pakhtunkhwa, Punjab).




Figure 3. *Scymnus (Scymnus) nubilus* Mulsant: **a** dorsal view **b** anterior view **c** lateral view **d** abdomen **e** penis **f** apex of penis **g** tegmen, ventral view **h** tegmen, lateral view.

Scymnus (Scymnus) nubilus Mulsant, 1850

Fig. 3

Scymnus nubilus Mulsant, 1850: 972; Bielawski 1972: 293; Booth and Pope 1989: 359; Canepari 2001: 207; Chen et al. 2013: 438.

Scymnus (Scymnus) nubilus: Korschefsky 1931: 143; Kapur 1972: 311; Miyatake 1985: 9; Yu and Pang 1992: 44; Poorani 2002: 358; Kovář 2007: 590; Yu 2011: 131; Chen et al. 2013: 438.

Remarks. *Scymnus (S.) nubilus* Mulsant can be easily confused with *S. (S.) contortubus* sp. n. and other *Scymnus* species due to their similar colouration. However it can be distinguished from other species by the long and slender penis with nearly S-shaped tip (Fig. 3e–f). Diagnostic are also: the tegmen stout (Fig. 3g–h), the penis guide nearly parallel along 2/3 of basal length, converging to pointed apex in ventral view (Fig. 3g), in lateral view abruptly narrowing from apical 1/3 length to apex; parameres with long apical setae (Fig. 3h).

Material examined. PAKISTAN (Kashmir) 1♀ 2♂, Mirpur, [N33°28 23.07", E073°52 57.94"] ca. 500–610 m, 26.X.2015, Huo LZ leg. (Khyber Pakhtunkhwa) 1♀ 1♂, Besham [N34°59 51.61", E072°54 21.34"] ca. 750 m, 18.10.2015, Wang XM leg; 1♂ Arab khan, [N34°25 20.73", E073°18 44.21"] ca. 1043 m, 16.X.2015,




Figure 4. Distribution map: *Scymnus (Scymnus) contortubus* Rashid, Chen & Wang, sp. n. (▲) and *Scymnus (Scymnus) nubilus* Mulsant (●).

Wang XM leg. (Punjab) 1♀, Chakwal, [N32°55 33.23", E072°44 56.91"] ca. 515 m, 20.X.2015, Wang XM leg; 3♂, Kallar kahar, [N32°46 20.29", E072°42 56.28"] ca. 648 m, 21.X.2015, Wang XM leg; 3♀ 1♂, Margalla [N33°44 00.16", E073°02 12.16"] ca. 609 m, 10.X.2015, Wang XM leg; 1♂, Pai khel [N32°45 57", E071°34 21"] ca. 226m, 17.IX.2016, Rashid A leg; 2♂, Khurianwala, [N31°31 41.54", E073°14 50.86"], ca. 192 m, 22.X.2015, Wang XM leg.

Distribution. Pakistan (Kashmir, Khyber Pakhtunkhwa, Punjab), China, Japan, India, Sri Lanka, Burma, Nepal, Micronesia, Portugal, La Réunion, African Region.

Acknowledgements:

The authors sincerely thank Dr. Shaukat Ali (South China Agricultural University, China) for making arrangements for survey in Pakistan. We are deeply indebted to three anonymous reviewers for their constructive and valuable comments and suggestions on our manuscript. We are grateful to Mr. Huo Lizhi and Li Wenjing (South

China Agricultural University, China) for their efforts in collecting coccinellids. The research was supported by National Natural Science Foundation of China (31501884, 31601878), Guangdong Natural Science Foundation (2017 for XSC), Biodiversity survey and evaluation pilot Programme, Science and Technology Planning Project of Guangdong Province (2015A030302069), Science and Technology Partnership Program, Ministry of Science and Technology of China (KY201402014), Science and Technology Program of Guangzhou, China (201509010023).

References

- Bielawski R (1972) Die marienkafer (Coleoptera: Coccinellidae) aus Nepal. *Fragmenta Faunistica* 18: 283–312. <https://doi.org/10.3161/00159301FF1972.18.16.283>
- Booth RG, Pope RD (1989) A review of the type material of Coccinellidae (Coleoptera) described by F.W. Hope, and by E. Mulsant in the Hope Entomological Collections, Oxford. *Entomologica Scandinavica* 20: 343–370. <https://doi.org/10.1163/187631289X00366>
- Canepari C (2001) The identity of *Nephus levaillanti* (Coleoptera: Coccinellidae). *Bollettino della Societa Entomologica Italiana* 133(3): 207–211.
- Chen X, Canepari C, Wang X, Ren S (2016) Revision of the subgenus *Orthoscymnus* Canepari of *Scymnus* Kugelann (Coleoptera, Coccinellidae), with descriptions of four new species. *ZooKeys* 552: 91–107. <https://doi.org/10.3897/zookeys.552.6167>
- Chen X, Wang X, Ren S (2013) A review of the subgenus *Scymnus* of *Scymnus* from China (Coleoptera, Coccinellidae). *Annales Zoologici* 63(3): 417–499. <https://doi.org/10.3161/000-345413X672483>
- Fabricius JC (1798) *Supplementum entomologicae systematicae*. Hafniae.
- Giorgi JA, Vandenberg NJ, McHugh JV, Forrester JA, Šlipiřínski SA, Miller KB, Shapiro LR, Whiting MF (2009) The evolution of food preferences in Coccinellidae. *Biological Control* 51: 215–231. <https://doi.org/10.1016/j.biocontrol.2009.05.019>
- Kapur AP (1972) The Coccinellidae (Coleoptera) of Goa. *Records of the Zoological Survey of India* 66(1–4): 309–320.
- Korschefsky R (1931) *Coleopterorum Catalogus*. Pars 118. Coccinellidae. I. Berlin, 224 pp.
- Kovář I (1996) Phylogeny. In: Hodek I, Honěk A (Eds) *Ecology of Coccinellidae*. Kluwer Academic Publishers, Dordrecht, 19–31. https://doi.org/10.1007/978-94-017-1349-8_2
- Kovář I (2007) New nomenclatorial and taxonomic acts and comments. Coccinellidae. In: Lobl I, Smetana A (Eds) *Catalogue of Palaearctic Coleoptera*. Apollo books, Stenstrup, 568–631.
- Kugelann JG (1794) *Verzeichniss der in eigigen Gegenden Preussens bis jetzt entdeckten Käfer-Arten, nebst kurzen Nachrichten von denselben*. *Neuestes Magazin für die Liebhaber der Entomologie* 1(5): 513–582.
- Miyatake M (1985) Coccinellidae collected by the Hokkaido University expedition to Nepal Himalaya, 1968 (Coleoptera). *Insecta Matsumurana (New Series)* 30: 1–33.
- Mulsant E (1846) *Securipalpes*. *Histoire Naturelle des Coleopteres de France*, 1 pl. Paris 4: 380.
- Mulsant E (1850) *Species de Coleopteres trimeres securipalpes*. *Annales des Sciences Physiques et Naturelles d'Agriculture et d'Industrie Lyon* 2: 1–1104. <https://doi.org/10.5962/bhl.title.8953>

- Poorani J (2002) An annotated checklist of the Coccinellidae (Coleoptera) (excluding Epilachninae) of the Indian sub-region. *Oriental Insects* 36: 307–383. <https://doi.org/10.1080/00305316.2002.10417335>
- Robertson JA, Ślipiński A, Moulton M, Shockley FW, Giorgi A, Lord NP, McKenna DD, Tomaszewska W, Forrester J, Miller KB, Whiting MF, McHugh JV (2015) Phylogeny and classification of Cucujoidea and the recognition of a new superfamily Coccinelloidea (Coleoptera: Cucujiformia). *Systematic Entomology* 40: 745–778. <https://doi.org/10.1111/syen.12138>
- Sasaji H (1968) Phylogeny of the family Coccinellidae (Coleoptera). *Etizenia* 35: 1–37.
- Seago AE, Giorgi JA, Li JH, Ślipiński A (2011) Phylogeny, classification and evolution of ladybird beetles (Coleoptera: Coccinellidae) based on simultaneous analysis of molecular and morphological data. *Molecular Phylogenetics and Evolution* 60: 137–151. <https://doi.org/10.1016/j.ympev.2011.03.015>
- Ślipiński A (2007) Australian ladybird beetles (Coleoptera: Coccinellidae): their biology and classification. ABRS, Canberra, 286 pp.
- Westwood JO (1838) Synopsis of the genera of British insects [paginated separately], published with 'An introduction to the modern classification of insects, founded on the natural habits and corresponding organisations of the different families (2 vols)'. Longman, Orme, Brown, Green, and Longmans, London, 1–48.
- Yu GY, Pang XF (1992) A review of Chinese *Scymnus* Kugelann (Coleoptera: Coccinellidae). *Journal of South China Agricultural University* 13(4): 39–47. [In Chinese with English summary]
- Yu GY (2011) The Coccinellidae of Taiwan. Chemical Industry Press, Beijing, 198 pp. [In Chinese with English summary]