

New records of the soldier flies of Morocco with a bibliographical inventory of the North African fauna (Diptera, Stratiomyidae)

Driss Yimlahi¹, Turgay Üstüner², Sanae Zinebi¹, Boutaïna Belqat¹

1 Department of Biology, Faculty of Sciences, University Abdelmalek Essaâdi, Tétouan, 93030, Morocco

2 Selçuk University, Faculty of Science, Department of Biology, Campus Alaaddin Keykubat, Selçuklu, 42075 Konya, Turkey

Corresponding author: Boutaïna Belqat (belqat@gmail.com)

Academic editor: T. Dikow | Received 22 April 2017 | Accepted 14 September 2017 | Published 18 October 2017

<http://zoobank.org/FBA8EC64-060F-42B4-BA54-6EDD0C26477B>

Citation: Yimlahi D, Üstüner T, Zinebi S, Belqat B (2017) New records of the soldier flies of Morocco with a bibliographical inventory of the North African fauna (Diptera, Stratiomyidae). ZooKeys 709: 87–125. <https://doi.org/10.3897/zookeys.709.13364>

Abstract

A checklist of soldier flies species recorded from the North African countries of Morocco, Algeria, Tunisia, Libya, and Egypt is based on both literature records and material newly collected in Morocco. Four subfamilies (Stratiomyinae, Sarginae, Nemotelinae, and Pachygasterinae), and twelve species from five genera have been collected and are recognized in Morocco. *Pachygaster atra* (Panzer, 1798), *Oxycera pardalina* (Meigen, 1822), *Nemotelus danielsoni* (Mason, 1989), and *Oxycera terminata* (Meigen, 1822) are newly recorded to the North African fauna. *Nemotelus atriceps* (Loew, 1856) and *Nemotelus maculiventris* (Bigot, 1861) are reported for the first time in Morocco. The present number of soldier flies known from Morocco is 33.

Keywords

Algeria, Checklist, Egypt, Morocco, North Africa, Rif, Soldier Flies, Stratiomyidae, Tunisia

Introduction

Stratiomyids (soldier flies) constitute one of the moderately large families of Diptera that exhibit an extreme array of morphological diversity, as well as a moderate range of life histories, with about 400 genera and about 2,700 species currently recognised worldwide (<http://www.catalogueoflife.org/col/browse/tree/id/2bda68991ddbf2b55b5c7e66c8a125ad>). The soldier flies are found all over the world, but are particularly diverse in tropical regions (Woodley 2001, 2011).

The family comprises varied members, ranging from 2.0 to 20.0 mm in length. While some species are entirely slender, others are stout or evidently flattened, with coloration ranging from strikingly patterned dark with a yellow, sometimes white or greenish pattern and frequent metallic reflections (Rozkošný 1997), to rather dull and concolourous dark brownish black (Figs 22, 23).

Important studies have been done over the world, like those of Rozkošný (1977, 1982, 1983, 1998, 2004) who specially studied the European Stratiomyidae, Rozkošný and Nartshuk (1988) who provided a catalogue of the soldier flies and Woodley (2001, 2011) who provided the world catalogue of the soldier flies.

To date, the soldier fly fauna of Morocco has been poorly investigated. The first records of Moroccan Stratiomyidae were given by Becker and Stein (1912) in their study of the Diptera of Morocco. For the next two decades, they received sporadic study devoted mainly, to the records of Moroccan species amongst the Diptera of Morocco or among the Stratiomyidae of the Palaearctic Region (Séguy 1930; Lindner 1936). Till now, no comprehensive and specific study has been published on Moroccan stratiomyids.


In this present study, 12 species of soldier flies have been recorded from 23 sampling sites in Morocco (Table 1), including a total of 83 specimens (47 males and 36 females). We report the first occurrences of the species *Nemotelus atriceps* (Loew, 1856) and *Nemotelus maculiventris* (Bigot, 1861) in Morocco, in addition to *Pachygaster atra* (Panzer, 1798), *Oxycera pardalina* (Meigen, 1822), *Oxycera terminata* (Meigen, 1822) and *Nemotelus danielsoni* (Mason, 1989) for the first time not only in Morocco but for the North African continent.

Materials and methods

Three techniques were used to collect Stratiomyidae: rearing larvae and pupae in the laboratory from collected substrates in the field following the technique used by Afzan and Belqat (2016), and collecting adults with both sweep net and malaise traps. Samples were collected by two authors (DY and BB) and the specimens were either micro-pinned or preserved in ethyl alcohol. Because specimens of the subfamily Nemotelinae are small, usually less than 5 mm in length, study and illustration of the male and/or female terminalia required use of a microscope. Observation of detailed structures of the male and/or female terminalia was sometimes needed to confirm species identification. Dissections of terminalia were prepared using the methods and

Table I. Sampling sites (in alphabetical order) harbouring the species collected in Morocco with localities, geographical coordinates and altitudes.


	Site	Locality	Province	Altitude (m)	Geographical coordinates
Rif Mountains					
1. Affluent Tarmast	Parc National d'Al Hoceima	Al Hoceima	168	35°10.666'N/004°03.088'W	
2. Aïn El Mâaab	Parc National Talassemtane	Chefchaouen	1278	35°05.509'N/005°09.443'W	
3. Aïn Sidi Brahim Ben Arrif	Route Moulay Abdessalam	Larache	897	35°20.398'N/005°32.712'W	
4. Barrage Moulay Bouchta	Larbaa Beni Hassan	Tétouan	364	35°15.864'N/005°21.221'W	
5. Cascade Chrafate	Chrafate	Chefchaouen	820	35°03.997'N/005°06.434'W	
6. Daya Afrate	Tanaqoub	Chefchaouen	600	35°05.634'N/005°26.028'W	
7. Daya Aïn Jdioui	Aïn Jdioui	Tanger-Assilah	5	35°34.074'N/005°55.499'W	
8. Daya Mezine	Mezine	Chefchaouen	778	35°06.104'N/005°21.177'W	
9. Daya Rmali	El Malâab, Parc National Talassemtane	Chefchaouen	1276	35°05.563'N/005°09.488'W	
10. Daya Tazia	Route Moulay Abdessalam	Larache	721	35°20.814'N/005°33.139'W	
11. Douar Kitane	Kitane	Tétouan	52	35°32.412'N/005°20.393'W	
12. Lac Amertrasse	Chrafate	Chefchaouen	828	35°05.014'N/005°5.130'W	
13. Oued Abou Brar	Douar Abou Brar, Parc National Talassemtane	Chefchaouen	1254	35°10.977'N/005°08.005'W	
14. Oued Achekrade	Douar Aouighen	Tétouan	642	35°22.931'N/005°20.364'W	
15. Oued El Koub	Souk El Had	Chefchaouen	124	35°01.298'N/005°25.333'W	
16. Oued Izelfane	Beni Boufrah	Al Hoceima	206	35°07.272'N/004°12.555'W	
17. Oued Majjou (Hafa meqlouba)	Majjou	Chefchaouen	825	35°06.175'N/005°10.836'W	
18. Oued Zandoula	15 km au Nord de Ouazzane	Ouazzane	108	34°55.707'N/005°31.942'W	
19. Ruisseau Maison forestière	Parc National Talassemtane	Chefchaouen	1674	35°08.076'N/005°08.262'W	
20. Taghbalout	Larbaa Beni Hassan	Tétouan	379	35°15.323'N/005°20.887'W	
Middle Atlas Mountains					
21. Cascade Aïn Vittel	Aïn Vittel	Ifrane	1555	33°33.682'N/005°07.463'W	
22. Mchacha Aïn Vittel	Aïn Vittel	Ifrane	1585	33°33.206'N/005°06.722'W	
Anti Atlas Mountains					
23. Village Massa	Village Massa	Massa	24	29°59.353'N/009°38.708'W	



Figures 1–4. *Nemotelus danielsoni* female and male terminalia: **1** Genital furca **2** Subgenital plate **3** Genital capsule with Aedeagal complex (dorsal view) **4** Genital capsule with Median process (lateral view).

equipment described by (Nagatomi and Iwata 1976), then transferred to 70% alcohol where the internal tissues were removed with fine forceps. After washing, the dissected genital parts were preserved in (1:10) a mixture of glycerine and ethyl alcohol for observation.

Preparations of the male and/or female terminalia, as well as the illustrations are given here (Figs 1–4). Species were recognised according to the identification keys of Séguay (1930, 1953), Rozkošný (1982, 1983) and Rozkošný (1977). A list of 23 sampling sites, with coordinates and altitudes, is presented in Table 1, and the locations of the sites are shown in Map 1, done using the logiciel GisArc (Geographic Informa-



Map 1. Map showing all sampling sites for soldier flies collected in this study in Morocco.

tion System, version 9.3). Photographs of the sampling localities showing Moroccan habitats of the species newly recorded are given (by DY and BB). All the material is deposited in the insect collection of the department of Biology, Faculty of Sciences, University Abdelmalek Essaâdi, Tétouan, Morocco.

The nomenclature and the list of the species known to North Africa follow the world Catalogue of the Stratiomyidae (Woodley 2001).

Table 2 summarises the species presently known from North Africa.

Table 2. The following checklist summarises the species of solider fly presently known from North Africa. Taxa are listed according to the classification scheme of Woodley (2001). Those species which are new records for North Africa are noted with (^) symbol, those new for Morocco are marked with (+), and the species which represent the new site in Morocco are with one asterisk (*).

Specimens	Morocco	Algeria	Tunisia	Libya	Egypt
Subfamily Beridinae					
<i>Beris rozkosnyi</i> Kassebeer, 1996	X				
<i>Chorisops tunisiae</i> (Becker, 1915)	X	X	X		
Subfamily Pachygastrinae					
<i>Aspidacantha atra</i> Kertesz, 1916					X
<i>Pachygaster atra</i> Panzer, 1798	X^				
<i>Pachygaster maura</i> Lindner, 1939	X				
Subfamily Clitellariinae					
<i>Adoxomyia flauipes</i> (Fabricius, 1798)		X			
<i>Pycnomalla aterrima</i> Sack, 1912	X	X			
<i>Pycnomalla auriflua</i> (Erichson, 1841)	X	X			
<i>Pycnomalla splendens</i> (Fabricius, 1787)	X	X			
Subfamily Sarginae					
<i>Chloromyia formosa</i> (Scopoli, 1763)	X*	X	X		
<i>Sargus bipunctatus</i> (Scopoli, 1763)			X		
Subfamily Stratiomyinae					
<i>Dicorymbomyia annulatus</i> (Becker, 1906)				X	
<i>Oxycera germanica</i> (Szilady, 1932)		X			
<i>Oxycera morrisii</i> (Curtis, 1833)		X			
<i>Oxycera ochracea</i> (Vaillant, 1950)		X			
<i>Oxycera orientalis</i> (Lindner, 1974)					X
<i>Oxycera pardalina</i> (Meigen, 1822)	X^				
<i>Oxycera rara</i> (Scopoli, 1763)		X			
<i>Oxycera tenebricosa</i> (Vaillant, 1952)		X			
<i>Oxycera torrentium</i> (Vaillant, 1950)		X			
<i>Oxycera terminata</i> Meigen, 1822	X^				
<i>Oxycera trilineata</i> (Linnaeus, 1767)	X*	X			
<i>Peritta melichlora</i> Becker, 1906		X			
<i>Vanoyia tenuicornis</i> Macquart, 1834	X				
<i>Odontomyia discoclara</i> (Séguy, 1929)		X			
<i>Odontomyia aolena</i> (Séguy, 1930)	X				
<i>Odontomyia angulata</i> (Panzer, 1798)	X	X			X
<i>Odontomyia discolor</i> (Loew, 1846)	X	X			
<i>Odontomyia flavissima</i> (Rossi 1790)	X	X	X		
<i>Odontomyia limbata</i> (Wiedemann, 1822)	X*	X	X		
<i>Odontomyia megacephala</i> Olivier, 1812					X
<i>Odontomyia microcera</i> (Séguy, 1930)	X				
<i>Odontomyia xanthopus</i> Bezzi, 1906					X
<i>Oplodontha minuta</i> Fabricius, 1794					X
<i>Oplodontha viridula</i> (Fabricius, 1775)		X			
<i>Stratiomyia africana</i> Szilady, 1941		X			
<i>Stratiomys cenisia</i> Meigen, 1822	X	X	X		X

Specimens	Morocco	Algeria	Tunisia	Libya	Egypt
<i>Stratiomys deserticolor</i> Lindner, 1930					X
<i>Stratiomys hispanica</i> Pleske, 1901		X			
<i>Stratiomys longicornis</i> (Scopoli, 1763)	X	X	X		X
<i>Stratiomys singularior</i> (Harris, 1776)					X
Subfamily Nemotelinae					
<i>Lasiopa benoisti</i> Séguy, 1930	X	X			
<i>Lasiopa manni</i> Mik, 1882			X		
<i>Lasiopa pantherina</i> Séguy, 1930	X				
<i>Nemotelus anchora</i> Loew, 1846		X	X		X
<i>Nemotelus atriceps</i> Loew, 1856	X+	X	X		
<i>Nemotelus beckeri</i> Hauser, 1998		X	X		
<i>Nemotelus candidus</i> Becker, 1906		X			X
<i>Nemotelus carthaginis</i> Becker, 1906			X		
<i>Nemotelus cingulatus</i> Dufour, 1852	X*	X	X		
<i>Nemotelus danielsoni</i> Mason, 1989	X^				
<i>Nemotelus dentatus</i> Becker, 1902					X
<i>Nemotelus lasiops</i> Loew, 1846			X		
<i>Nemotelus latiusculus</i> Loew, 1871	X*	X	X		
<i>Nemotelus longirostris</i> Wiedemann, 1824	X	X	X		
<i>Nemotelus maculiventris</i> Bigot, 1861	X+	X			
<i>Nemotelus marinus</i> Becker, 1902					X
<i>Nemotelus matrouensis</i> Mohammad, Fadl, Gadalla & Badrawy, 2009					X
<i>Nemotelus nigrifrons</i> Loew, 1846	X*	X	X	X	
<i>Nemotelus niloticus</i> Olivier, 1811					X
<i>Nemotelus notatus</i> Zetterstedt, 1842					X
<i>Nemotelus pantherinus</i> (Linnaeus, 1758)	X				
<i>Nemotelus proboscideus</i> Loew, 1846		X	X	X	
<i>Nemotelus punctiventris</i> Becker, 1902					X
<i>Nemotelus subuliginosus</i> Rozkošný, 1974	X				
<i>Nemotelus ventralis</i> Meigen, 1830	X				
<i>Nemotelus nigrinus</i> Fallen, 1817	X				

Results

Faunistic records

Subfamily BERIDINAE

Genus *BERIS* Latreille, 1802

Beris rozkosnyi Kassebeer, 1996

North African literature records. Morocco: Middle Atlas, Meknès, Ifrane (Kasebber 1996: 155; Woodley 2001: 66).

World distribution. Spain (Woodley 2001: 66).

Genus *CHORISOPS* Rondani, 1856***Chorisops tunisiae* (Becker, 1915)**

= *Beris tunisiae* Becker, 1915

North African literature records. Morocco, Algeria, Tunisia: La Calle (Woodley 2001: 68).

World distribution. Portugal, Spain (Woodley 2001: 68); Sardinia (Mason et al. 2009: 509–510).

Subfamily PACHYGASTRINAE Loew, 1856**Genus *ASPIDACANTHA* Kertesz, 1916*****Aspidacantha atra* Kertesz, 1916**

North African literature record. Egypt (Lindner 1936: 211; Woodley 2001: 86).

World distribution. Palaearctic: Israel, Turkmenistan. Afrotropical: Ethiopia, Tanzania, Uganda, Zaire, Zimbabwe (Woodley 2001: 86); United Arab Emirates (Hauser 2008: 591–592).

Genus *PACHYGASTER* Meigen, 1803***Pachygaster atra* Panzer, 1798**

= *Nemotelus ater* Panzer, 179

= *Sargus pachygaster* Fallen, 1817

New record. Morocco, Rif: Daya Mezine (Fig. 5), 1♀, 11/VI/2013, sweep net, Coll. Yimlahi and Belqat.

World distribution. Austria, Belgium, Bulgaria, Czech Republic, England, France, Georgia, Germany, Greece, Hungary, Ireland, Israel, Italy, Netherlands, Poland, Portugal, Romania, Russia, Scotland, Slovakia, Spain, Sweden, Switzerland, Turkey, Ukraine, Wales, Yugoslavia (Woodley 2001: 125).

***Pachygaster maura* Lindner, 1939**

North African literature record. Morocco (Lindner 1939: 314).

World distribution. Portugal, Spain (Woodley 2001: 126).



Figure 5. Habitat of *Pachygaster atra*: Daya Mezine.

Subfamily CLITELLARIINAE
Genus *ADOXOMYIA* Kertesz, 1907

***Adoxomyia flauipes* (Fabricius, 1798)**

= *Stratiomys flauipes* Fabricius, 1798
= *Euparyphus kabylinus* Bigot, 1879b

World distribution. Only known in North African from Algeria: Oran, Tebessa (Lindner 1936: 158–159); Oran (Woodley 2001: 151).

Genus *PYCNOMALLA* Gerstaecker, 1857

***Pycnomalla aterrima* Sack, 1912**

World distribution. This species is known in North Africa only from Morocco: Middle Atlas and Algeria: Tizi s'Tkrine (Séguy 1930: 59); Morocco: Daiet Ahoua (Séguy 1953: 78); Morocco and Algeria (Woodley 2001: 172).

***Pycnomalla auriflua* (Erichson, 1841)**

= *Stratiomys auriflua* Erichson, 1841

World distribution. Known in North Africa only from Morocco: Middle Atlas and Algeria: Soufouloud (Séguy 1930: 59); Morocco, Algeria (Woodley 2001: 172).

***Pycnomalla splendens* (Fabricius, 1787)**

= *Stratiomys splendens* Fabricius, 1787

= *Ephippium rufitarse* Macquart, 1838

= *Pycnomalla splendens* ssp. *jordanica* Lindner, 1974

North African literature records. Morocco (Séguy 1930: 59, 1953: 78); Morocco, Tunisia (Lindner 1936: 149–150); Morocco, Algeria, Tunisia (Woodley 2001: 172).

World distribution. Armenia, Israel, Portugal, Spain (Woodley 2001: 172); Turkey (Üstüner et al. 2002: 23).

Subfamily SARGINAE**Genus *CHLOROMYIA* Duncan, 1837*****Chloromyia formosa* (Scopoli, 1763)**

= *Musca formosa* Scopoli, 1763

= *Nemotelus flavogeniculatus* De Geer, 1776

= *Musca cicur* M. Harris, 1778

= *Musca aurata* Fabricius, 1787

= *Sargus aeneus* Walckenaer, 1802

= *Sargus xanthopterus* Meigen, 1804

= *Sargus azureus* Loew, 1840

New localities. Morocco, Rif: Taghbalout, 1♂ 2♀, 5/IV/2014, sweep net, Coll. Yimlahi and Belqat; Lac Ametrasse (Fig. 6), 1♂, 24/V/2013, Coll. Yimlahi and Belqat, sweep net; Douar Kitane, 1♂, 10/IV/2014, Malaise trap, 1♀, 2/V/2014, sweep net, Coll. Yimlahi and Belqat.

North African literature records. Morocco: Tangier (Becker and Stein 1912: 64), Algeria: M’Rassine (Séguy 1930: 66), Merja Zerga (Pârvu et al. 2006); Morocco, Algeria, Tunisia (Woodley 2001: 189).

World distribution. Nearctic: USA (New York, introduced). Palaearctic, Austria, Bulgaria, Czech Republic, England, France, Germany, Greece, Italy, Poland, Portugal, Romania, Russia, Slovakia, Slovenija, Spain, Sweden, Switzerland, Turkey, Yugoslavia (Woodley 2001: 189); Sardinia (Mason et al. 2009: 508–509).



Figure 6. Habitat of *Chloromyia formosa*: Lac Ametrasse.

Genus *SARGUS* Fabricius, 1798

Sargus bipunctatus (Scopoli, 1763)

- = *Musca bipunctata* Scopoli, 1763
- = *Sargus reaumuri* Meigen, 1804
- = *Sargus reaumurii* Fabricius, 1805
- = *Sargus sulphureus* Meigen, 1822
- = *Sargus bipunctatus* O. Costa, 1844
- = *Chrysochroma fasciatus* Szilady, 1929
- = *Geosargus perlulcher* James, 1936

North African literature record. Tunisia (Woodley 2001: 221).

World distribution. Nearctic: Canada (British Columbia), USA (Oregon, Washington). Palaearctic: Albania, Austria, Belgium, Bulgaria, Croatia, Czech Republic, England, France, Georgia, Germany, Greece, Hungary, Ireland, Italy, Netherlands, Poland, Romania, Serbia, Slovakia, Slovenia, Switzerland (Woodley 2001: 221); Sardinia (Mason et al. 2009: 524).

Subfamily STRATIOMYINAE Latreille, 1802**Tribe OXYCERINI Enderlein, 1914****Genus *DICORYMBIMYIA* Woodley 2001*****Dicorymbomyia annulatus* (Becker, 1906)**

= *Oxycera annulata* Becker, 1906

World distribution. Tunisia: Zaghouan, Tunis (Woodley 2001: 240).

Genus *OXYCERA* Meigen, 1803

= *Hermione* Meigen, 1800

***Oxycera germanica* (Szilday, 1932)**

= *Hermione germanica* Szilady, 1932

= *Hermione dorieri* Vaillant, 1950

= *Hermione dorieri* var. *barbarica* Vaillant, 1950

North African literature record. Algeria: Aurès Mountains, Arris (Woodley 2001: 247).

World distribution. France, Germany, Switzerland (Woodley 2001: 247).

***Oxycera morrisii* (Curtis, 1833)**

= *Oxycera ranzonii* Schiner, 1857

= *Hermione muscaria ronzonii* Vaillant, 1950

= *Hermione morrisii* var. *auresi* Vaillant, 1950

= *Hermione morrisii* var. *minuta* Vaillant, 1950

North African literature records. Algeria: vicinity of Alger (Woodley 2001: 249).

World distribution. Austria, Belgium, Denmark, England, France, Germany, Greece, Ireland, Israel, Italy, Spain, Switzerland, Yugoslavia (Woodley 2001: 249).

***Oxycera ochracea* (Vaillant, 1950)**

= *Hermione ochracea* Vaillant, 1950

World distribution. Only known in North Africa from Algeria: Aurès Mountains, Arris, Constantine (Woodley 2001: 249).

Oxycera orientalis (Lindner, 1974)

- = *Heraclina orientalis* Lindner, 1974
- = *Heraclina stigmosa orientalis* (Lindner & Freidberg, 1978)

North African literature record. Egypt (Badrawy 2006: 252).

World distribution. Israel (Woodley 2001: 249).

Oxycera pardalina (Meigen, 1822)

- = *Oxycera amoena* Loew, 1857
- = *Oxycera engadinica* Jaennicke, 1866
- = *Oxycera calceata* Loew, 1871
- = *Hermione sahunica* Séguy, 1934
- = *Hermione pardalina* var. *oldenbergi* Lindner, 1938
- = *Hermione pardalina* var. *nigrifrons* Szilady, 1941
- = *Hermione morrisi* var. *bohemica* Hrbacek, 1945
- = *Hermione armata* Vaillant, 1950
- = *Hermione pardalina* var. *depressa* Vaillant & Delhom, 1956
- = *Hermione pardalina* var. *alticola* Vaillant & Delhom, 1956

New records. Oued Abou Bnar (Fig. 7a, b), 17/V/2014, sweep net, 1♀, 18/V/2014-12/VI/2014, reared, 2♂♂1♀; Oued Majjou (Hafa meqlouba) (Fig. 8a, b), 1♀, 9/IV/2015-15/VI/2015, reared; Oued Achekrade (Fig. 9), 1♀, 31/V/2014-8/VI/2014, reared; Ruisseau Maison forestière (Fig. 10), 2♀♀, 24/IV/2015, sweep net; Cascade Aïn Vittel (Fig. 11), 2♂♂1♀, 17/II/2016-20/IV/2016, reared; Mchacha Aïn Vittel (Fig. 12), 1♂, 17/II/2016-20/IV/2016, reared, Coll. Yimlahi and Belqat.

World distribution. Albania, Armenia, Austria, Belgium, Czech Republic, England, France, Georgia, Germany, Hungary, Ireland, Italy, Netherlands, Poland, Romania, Russia, Slovakia, Spain, Sweden, Switzerland, Yugoslavia (Woodley 2001: 250).

Oxycera rara (Scopoli, 1763)

- = *Musca rara* Scopoli, 1763
- = *Musca tardigradus* M. Harris, 1778
- = *Stratiomys maculata* Geoffroy in Fourcroy, 1785
- = *Oxycera pulchella* Meigen, 1822
- = *Hermione pulchella* var. *similis* (Vaillant, 1950)

North African literature record. Algeria: Aurès Mountains, Arris (Woodley 2001: 251)



Figure 7. **a** Habitat of *Oxycera pardalina*: Oued Abou Bnar (from where the substrate is taken for rearing adults) **b** Habitat of *Oxycera pardalina*: Oued Abou Bnar environment.



Figure 8. **a** Habitat of *Oxycera pardalina*: Oued Majjou (Hafa meqlouba) environment **b** Habitat of *Oxycera pardalina*: Oued Majjou (Hafa meqlouba).




Figure 9. Habitat of *Oxycera pardalina*: Oued Achekrade.



Figure 10. Habitat of *Oxycera pardalina*: Ruisseau Maison forestière environment.



Figure 11. Habitat of *Oxycera pardalina*: Cascade AïnVittel (photograph taken on 17/II/2016, when the material was collected).



Figure 12. Habitat of *Oxycera pardalina*: Mchacha Aïn Vittel (photograph taken on 17/II/2016, when the material was collected).

World distribution. Andorra, Austria, Belgium, Czech Republic, England, France, Germany, Hungary, Italy, Netherlands, Poland, Romania, Slovakia, Slovenia, Spain, Switzerland, Wales, Yugoslavia (Woodley 2001: 251); Sardinia (Mason et al. 2009: 524).

Oxycera tenebricosa (Vaillant, 1952)

= *Hermione tenebricosa* Vaillant, 1952

World distribution. Only known in North Africa from Algeria: Atlas de Blida (Woodley 2001: 252).

Oxycera terminata Meigen, 1822

New records. Morocco, Rif: Cascade Chrafate (Fig. 13a, b), 2♂♂ 2♀♀, 18/III/2015-9/V/2015, 1♂, 18/III/2015-20/IV/2015, reared, Coll. Yimlahi and Belqat.

World distribution. Austria, Czech Republic, Denmark, England, France, Greece, Hungary, Poland, Romania, Slovakia, Switzerland, Yugoslavia (Woodley 2001: 252).



Figure 13. **a** Habitat of *Oxycera terminata*: Cascade Chrafate environment **b** Habitat of *Oxycera terminata*: Cascade Chrafate (extraction site of the substrate for rearing adults).

Oxycera torrentium (Vaillant, 1950)

= *Hermione torrentium* Vaillant, 1950

World distribution. Only known in North Africa from Algeria: Atlas of Bilda, La Chiffa, Aurès Mountains (Woodley 2001: 252).

Oxycera trilineata (Linnaeus, 1767)

- = *Musca graeca* Pontoppidan, 1763
- = *Musca trilineata* Linnaeus, 1767
- = *Musca hypoleon* Linnaeus, 1767
- = *Stratiomys fasciata* Geoffroy in Fourcroy, 1785
- = *Oxycera proxima* Loew, 1873
- = *Oxycera trilineata* var. *collaris* Brunetti, 1889
- = *Hermione trilineata* ssp. *transfasciata* Pleske, 1925
- = *Hermione trilineata* ssp. *ferghanensis* Pleske, 1925
- = *Hermione ucrainica* Paramonov, 1926
- = *Hermione bucheti* Séguy, 1930
- = *Hermione trilineata* ssp. *angustistomata* Lindner, 1938



Figure 14. Habitat of *Oxycera trilineata* : Daya Aïn Jdioui.

- = *Hermione trilineata* var. *biroi* Szilady, 1941
- = *Hermione trilineata* var. *sajoi* Szilady, 1941
- = *Hermione trilineata* var. *algira* Vaillant, 1950

New locality. Morocco, Rif: Daya Aïn Jdioui (Fig. 14), 1♂, 28/III/2015-27/V/2015, reared, Coll. Yimlahi and Belqat.

North African literature records. Morocco: Tangier (Becker and Stein 1912: 63, Séguy 1930: 62); Tangier, Algeria: Aurès Mountains, Arris (Woodley 2001: 252).

World distribution. Austria, Belgium, Bulgaria, China, Czech Republic, Denmark, England, Finland, France, Georgia, Germany, Greece, Hungary, Ireland, Israel, Italy, Kyrgyzstan, Latvia, Lithuania, Mongolia, Netherlands, Norway, Poland, Portugal, Romania, Russia, Scotland, Slovakia, Spain, Sweden, Switzerland, Turkey, Ukraine, Uzbekistan, Wales, Yugoslavia (Woodley 2001: 252); Sardinia (Mason et al. 2009: 524); United Arab Emirates (Hauser 2014: 695-697).

Genus *PERITTA* Becker, 1906

Peritta melichlora Becker, 1906

World distribution. Only known in North Africa from Algeria: Biskra, spring at Hammam-Salahin (Woodley 2001: 254).

Genus **VANOYZA** Villeneuve, 1908

***Vanoyia tenuicornis* Macquart, 1834**

- = *Oxycera nigra* Macquart, 1834
- = *Oxycera longicornis* Dale, 1848
- = *Oxycera longicornis* Walker, 1851
- = *Vanoyia scutellata* Villeneuve, 1908
- = *Vanoyea separata* Kertesz, 1921

North African literature record. Morocco: Tangier (Lindner 1936: 194, Woodley 2001: 254).

World distribution. Belgium, England, France, Ireland, Spain (Woodley 2001: 254).

Tribe STRATIOMYINI Latreille, 1802

Genus ***ODONTOMYIA*** Meigen, 1803

***Odontomyia alolena* (Séguy, 1930)**

- = *Eulalia alolena* Séguy, 1930

World distribution. Only known in North Africa from Morocco: Casablanca, Tangier, Mahaidja, Aïn Leuh (Séguy 1930: 65, Woodley 2001: 269).

***Odontomyia angulata* (Panzer, 1798)**

- = *Eulalia angulata* Panzer, 1798
- = *Stratiomys angulata* Panzer, 1798
- = *Stratiomys vulpina* Panzer, 1798
- = *Stratiomys hydropota* Meigen, 1822
- = *Odontomyia latifaciata* Macquart, 1834
- = *Stratiomys brevicornis* Loew, 1840
- = *Stratiomys brevicornis* Loew, 1840
- = *Stratiomys ruficornis* Zetterstedt, 1842
- = *Odontomyia hydrophila* Loew, 1846

North African literature records. Morocco: Tangier (Becker and Stein 1912: 62); Morocco, Algeria, Egypt (Woodley 2001: 269); Egypt (Badrawy 2006: 253).

World distribution. Afghanistan, Albania, Austria, Belgium, Bulgaria, China, Czech Republic, Denmark, England, Estonia, Finland, France, Germany, Greece,

Hungary, Iran, Israel, Italy, Kazakhstan, Netherlands, Poland, Romania, Russia, Slovakia, Spain, Sweden, Switzerland, Turkey, Yugoslavia (Woodley 2001: 269); Sardinia (Mason et al. 2009: 521–522).

***Odontomyia disciclara* (Séguy, 1929)**

= *Eulalia disciclara* Séguy, 1929

World distribution. Only known in North African from Algeria: Touggourt (Lindner 1936: 87; Woodley 2001: 273).

***Odontomyia discolor* (Loew, 1846a)**

= *Eulalia (Odontomyia) discolor* Loew, 1897
= *Odontomyia limbata* Macquart in Lucas, 1849

North African literature records. Morocco: Tangier (Becker and Stein 1912: 62); Morocco, Algeria: Constantine (Woodley 2001: 274).

World distribution. Afghanistan, France, Greece, Israel, Italy, Kazakhstan, Kirgizia, Romania, Russia, Spain, Tajikistan, Turkey, Turkmenistan (Woodley 2001: 274); Sardinia (Mason et al. 2009: 522).

***Odontomyia flavissima* (Rossi 1790)**

= *Stratiomys flavissima* Rossi, 1790
= *Stratiomys decora* Wiedemann in Meigen, 1822
= *Stratiomys infoscata* Meigen, 1830
= *Odontomyia semiviolacea* Brullé, 1833
= *Odontomyia nigripes* Macquart, 1847
= *Odontomyia limbipennis* Macquart, 1847
= *Odontomyia laufferi* Strobl in Czemy & Strobl, 1909

North African literature records. Morocco, Algeria, Tunisia (Woodley 2001: 275).

World distribution. Albania, Austria, Azerbaijan, Bulgaria, Czech Republic, France, Georgia, Greece, Hungary, Israel, Italy, Kazakhstan, Portugal, Romania, Russia, Slovakia, Spain, Switzerland, Syria, Turkey, Yugoslavia (Woodley 2001: 275).

***Odontomyia limbata* (Wiedemann, 1822)**

- = *Stratiomys limbata* Wiedemann in Meigen, 1822
= *Clitellaria pacifica* Wiedemann in Meigen, 1822
= *Opseogymnus flavo signata* A. Costa, 1857

New locality. Morocco, Rif: Lac Ametrasse, 2♂♂1♀, 28/IV/2015; Aïn Sidi Brahim Ben Arrif, 4♂♂, 23/IV/2015; Daya Afrate (Fig. 15), 1♀, 18/IV/2015; Ruisseau Maisonn forestière, 11♂1♀, 8/V/2014; Aïn El Malâab (Fig. 16a, b), 1♂2♀♀, 17/V/2014; Daya Rmali El Malâab, 1♀, 17/V/2014, Daya Tazia, 2♂♂1♀, 12/V/2015, sweep net, Coll. Yimlahi and Belqat.

North African literature records. Morocco: Tangier (Becker and Stein 1912: 62); Middle Atlas, Tangier, Algeria (Séguy 1930: 65); Morocco, Algeria, Tunisia (Woodley 2001: 278).

World distribution. France, Italy, Portugal, Spain (Woodley 2001: 278).

***Odontomyia megacephala* Olivier, 1812**

World distribution. Only known in North Africa from Egypt (Lindner 1936: 92; Badrawy 2006: 253), Nil river (Woodley 2001: 279).



Figure 15. Habitat of *Odontomyia limbata* and *Nemotelus cingulatus*: Daya Afrate.




Figure 16. **a** Habitat of *Odontomyia limbata*: Aïn El Malâab **b** Habitat of *Odontomyia limbata*: Aïn El Malâab environment.

Odontomyia microcera (Séguy, 1930)

= *Eulalia microcera* Séguy, 1930

World distribution. Only known in North Africa from Morocco: Meknès (Séguy 1930: 65; Woodley 2001: 279).

Odontomyia xanthopus Bezzi, 1906

North African literature records. Egypt (Woodley 2001: 285; Badrawy 2006: 253).

World distribution. Palaearctic: Israel. Afrotropical: Ethiopia, Malawi (Woodley 2001: 285).

Genus *OPLODONTHA* Rondani, 1863

Oplodontha minuta Fabricius, 1794

= *Oplodontha oasisina* Lindner, 1925

= *Eulalia oasisina* Lindner, 1925

North African literature records. Egypt: Kharga Oasis, Dakhla Oasis (Lindner 1936: 93–94; Woodley 2001: 287); Egypt (Badrawy 2006: 253).

World distribution. Socotra Island (Tkoč and Rozkošný 2014: 431–437).

***Oplodontha viridula* (Fabricius, 1775)**

- = *Stratiomys viridula* Fabricius, 1775
- = *Stratiomys canina* Panzer, 1798
- = *Stratiomys jejuna* Schrank, 1803
- = *Musca jejuna* Schrank in Gistl, 1837
- = *Odontomyia dentata* Meigen, 1804
- = *Odontomyia holosericea* Olivier, 1811
- = *Odontomyia lunata* Olivier, 1811
- = *Stratiomys subvittata* Meigen, 1822
- = *Stratiomys bimaculata* Meigen, 1835
- = *Stratiomys bimaculata* Meigen, 1838
- = *Odontomyia personata* Loew, 1846
- = *Odontomyia interrupta* Loew, 1846
- = *Odontomyia heydenii* Jaennicke, 1866
- = *Odontomyia atrata* Verrall, 1909

North African literature record. Algeria (Woodley 2001: 288).

World distribution. Albania, Austria, Belgium, Bulgaria, China, Czech Republic, Denmark, England, France, Finland, Germany, Greece, Hungary, Iraq, Ireland, Israel, Italy, Kazakhstan, Kyrgyzstan, Mongolia, Netherlands, Norway, Poland, Romania, Russia, Slovakia, Spain, Sweden, Switzerland, Turkey (Woodley 2001: 288); Sardinia (Mason et al. 2009: 522).

Genus *STRATIOMYS* Geoffroy, 1762***Stratiomyia africana* Szilady, 1941**

World distribution. Only known in North Africa from Algeria: Biskra (Woodley 2001: 296).

***Stratiomys cenisia* Meigen, 1822**

- = *Stratiomys flaviventris* Loew, 1846
- = *Stratiomyia ahngeri* Pleske, 1901
- = *Stratiomyia cypria* Pleske, 1902
- = *Stratiomyia kervillei* Villeneuve, 1911
- = *Stratiomys hispanica* ssp. *planes* James, 1941

North African literature records. Morocco: Middle Atlas, Meknès, Rabat, Timahdit, Tangier (Séguy 1930: 64); Algeria: Surcouf (Séguy 1930: 64); Algeria, Egypt, Morocco, Tunisia (Woodley 2001: 297); Egypt (Badrawy 2006: 253).

World distribution. Armenia, Austria, Bulgaria, Cyprus, Czech Republic, France, Germany, Hungary, Iran, Israel, Italy, Kazakhstan, Poland, Romania, Russia, Slovakia, Spain, Syria, Turkey, Turkmenistan, Ukraine, Yugoslavia (Woodley 2001: 297).

Stratiomys deserticolor Lindner, 1930

World distribution. Only known in North Africa from Egypt: Siwa Oasis (Woodley 2001: 298); (Badrawy 2006: 254).

Stratiomys hispanica Pleske, 1901

North African literature record. Algeria (Woodley 2001: 300).

World distribution. France, Spain (Woodley 2001: 300).

Stratiomys longicornis (Scopoli, 1763)

- = *Hirtea longicornis* Scopoli, 1763
- = *Musca tenebricus* M. Harris, 1778
- = *Stratiomys strigata* Fabricius, 1781
- = *Stratiomys tomentosa* Schrank, 1803
- = *Stratiomys villosa* Meigen, 1804
- = *Stratiomys nubeculosa* Meigen, 1804
- = *Stratiomys thoracica* Fabricius, 1805
- = *Stratiomys hirtuosa* Meigen, 1830
- = *Stratiomys anubis* (Wiedemann, 1830)
- = *Stratiomyia flavifrons* Macquart, 1838
- = *Stratiomys strigata* var. *pallida* Loew, 1840
- = *Stratiomys lambessiana* (Bigot, 1879) (Lindner 1936: 62)
- = *Stratiomys flavo limbata* (A. Costa, 1893) (Lindner 1936: 60)
- = *Stratiomyia segnis* (Becker, 1906)
- = *Hirtea efflatouri* (Lindner, 1925) (Lindner 1936: 58-59)
- = *Stratiomyia* (*Hirtea*) *surcoufi* (Séguy, 1930) (Lindner 1936: 64)
- = *Hirtea surcoufi* (Séguy, 1932)
- = *Stratiomyia longicornis* ssp. *palaestinensis* Lindner, 1937
- = *Stratiomyia* (*Hirtea*) *longicornis* ssp. *flavoscutellata* Lindner, 1940

North African literature records. Morocco: Casablanca (Séguy 1930: 63); Morocco, Algeria: Lambessa, Tunisia (Lindner 1936: 60, 62, 64); Algeria, Egypt, Morocco, Tunisia (Woodley 2001: 301); Egypt (Badrawy 2006: 254).

World distribution. Afghanistan, Albania, Armenia, Austria, Azerbaijan, Belgium, Bulgaria, China, Cyprus, Czech Republic, Denmark, England, France, Germany, Greece, Hungary, Iran, Israel, Italy, Korea, Lithuania, Malta, Mongolia, Netherlands, Poland, Portugal, Romania, Russia, Scotland, Slovakia, Slovenia, Spain, Sweden, Switzerland, Tunisia, Turkey, Yugoslavia (Woodley 2001: 301); Sardinia (Mason et al. 2009: 524).

Stratiomys singularior (Harris, 1776)

- = *Musca singularius* Harris, 1776
- = *Stratiomys furcata* Fabricius, 1794
- = *Stratiomys panthaleon* Fallen, 1817
- = *Stratiomys riparia* Meigen, 1822
- = *Stratiomys paludosa* Siebke, 1863

North African literature record. Egypt (Badrawy 2006: 254).

World distribution. Armenia, Austria, Belgium, Bulgaria, China, Czech Republic, Denmark, England, Estonia, Finland, France, Germany, Hungary, Iran, Ireland, Italy, Kazakhstan, Lithuania, Mongolia, Netherlands, Norway, Poland, Romania, Russia, Slovakia, Spain, Sweden, Switzerland, Ukraine, Yugoslavia (Woodley 2001: 301).

Subfamily NEMOTELINAE Kertesz, 1912

Genus *LASIOPA* Brulle, 1832

Lasiopa benoisti Séguay, 1930

World distribution. Only known in North Africa from Morocco: Meknès and Algeria: Boghari (Séguay 1930: 60); Morocco: Meknès, Algeria: Boghari, Bougie (Woodley 2001: 309).

Lasiopa manni Mik, 1882

North African literature records. Algeria: Constantine (Becker and Stein 1912: 63); Tunisia (Woodley 2001: 309).

World distribution. Italy, Turkey (Woodley 2001: 309).

Lasiopa pantherina Séguay, 1930

World distribution. Only known in North African from Morocco: Maharidja (Séguay 1930: 62; Woodley 2001: 309).

Genus *NEMOTELUS* Geoffroy, 1762**Subgenus *Nemotelus* Geoffroy, 1762*****Nemotelus anchora* Loew, 1846**

= *Nemotelus siculus* Jaennicke, 1866

= *Nemotelus persicus* Pleske in Lindner, 1937

North African literature records. Algeria, Tunisia (Woodley 2001: 311); Egypt (Badrawy 2006: 252); Egypt: Fayoum (Mohammad et al. 2009: 99)

World distribution. Iran, Israel, Italy, Malta, Russia (Woodley 2001: 311); Sardinia (Mason et al. 2009: 510).

***Nemotelus atriceps* Loew, 1856**

= *Nemotelus longicornis* Lindner, 1937

New record. Morocco: Village Massa (Fig. 17), 11.V.2015, 1♂, sweep net, Coll. Yimlahi and Belqat.

North African literature records. Algeria, Tunisia (Woodley 2001: 312).

World distribution. France, Portugal, Spain (Woodley 2001: 312).

***Nemotelus beckeri* Hauser, 1998**

World distribution. Known in North Africa only from Algeria and Tunisia: Tabaraka, Khathairie (Hauser 1997: 453; Woodley 2001: 313).

***Nemotelus candidus* Becker, 1906**

World distribution. Known in North Africa only from Algeria: Hammam-Salahin (Lindner 1936: 122–123), Biskra, Hammam-Salahin (Woodley 2001: 314) and Egypt (Badrawy 2006: 252); Egypt: Dekhela Mariout, Wadi El Natroun (Mohammad et al. 2009: 99–100).

***Nemotelus carthaginis* Becker, 1906**

World distribution. Only known in North Africa from Tunisia (Lindner 1936: 123), Tunisia: Karthago (Woodley 2001: 314).



Figure 17. Habitat of *Nemotelus atriceps*: Village Massa.

***Nemotelus cingulatus* Dufour, 1852**

- = *Nemotelus lateralis* Dufour, 1852
- = *Nemotelus pulcher* Loew, 1871
- = *Nemotelus aemulus* Loew, 1871
- = *Nemotelus consimilis* (Becker, 1915)

New locality. Morocco, Rif: Daya Afrate (Fig. 15), 2♂♂, 18/IV/2015, sweep net; Oued El Koub (Figs 18, 19), 1♀, 06/V/2015, Coll. Yimlahi and Belqat.

North African literature records. Algeria, Tunisia (Lindner 1936: 123–124); Algeria, Morocco, Tunisia (Woodley 2001: 314).

World distribution. France, Spain (Woodley 2001: 314).

***Nemotelus danielsoni* Mason, 1989 (Figs 20, 21, 22, 23)**

World distribution. Greece (Woodley 2001: 315).

New record. Morocco: Oued Izelfane (Fig. 24), 8♂♂ 12♀♀, 25/VI/2013, sweep net, Coll. Yimlahi and Belqat.



Figure 18. Habitat of *Nemotelus cingulatus*: Oued El Koub environment.




Figure 19. Habitat of *Nemotelus cingulatus*: Oued El Koub.



Figure 20. *Nemotelus danielsoni*: Male adult (dorsal view).



Figure 21. *Nemotelus danielsoni*: Male adult (lateral view).




Figure 22. *Nemotelus danielsoni*: Female adult (dorsal view).



Figure 23. *Nemotelus danielsoni*: Female adult (lateral view).

Nemotelus danielsoni was described as a new species only from the male specimen (from Crete Island, Greece). Recently, Mason and Rozkošný (2003) have described the female.

The species is recorded from Izelfane in Morocco. This finding is very interesting, as it represents the first record from North Africa of a supposed endemic species of Greece.

Nemotelus dentatus Becker, 1902

World distribution. Only known in North Africa from Egypt: El Alagto Marg (Lindner 1925), Birket-el-Karun (Woodley 2001: 315); (Badrawy 2006: 252); Coastal strip, Lower Nile, Western desert (Mohammad et al. 2009: 100).

***Nemotelus lasiops* Loew, 1846**

= *Nemotelus striativentris* Lindner, 1937

North African literature record. Tunisia (Woodley 2001: 317).

World distribution. Italy (Woodley 2001: 317); Sardinia (Mason et al. 2009: 512).

***Nemotelus latiusculus* Loew, 1871**

= *Nemotelus cothurnatus* Bigot, 1879

= *Nemotelus cardinalii* Bezzii, 1898

= *Nemotelus perplexus* Becker, 1915

= *Nemotelus freidbergi* Lindner, 1975

New locality. Morocco, Rif: Barrage Moulay Bouchta (Fig. 25), 1♂, 05/IV/2014, sweep net, Coll. Yimlahi and Belqat.

North African literature records. Tunisia (Lindner 1936: 137); Algeria, Morocco, Tunisia (Woodley 2001: 318).

World distribution. Israel, Italy, Spain (Woodley 2001: 318).



Figure 24. Habitat of *Nemotelus danielsoni*: Oued Izelfane.



Figure 25. Habitat of *Nemotelus latiusculus*: Barrage Moulay Bouchta.



Figure 26. Habitat of *Nemotelus maculiventris* (Bigot, 1861): Oued Zandoula.

***Nemotelus longirostris* Wiedemann, 1824**

- = *Nemotelus gadensis* Schiner, 1868
- = *Nemotelus pilosus* Loew, 1871
- = *Nemotelus fuscinervis* Loew, 1871

North African literature records. Morocco: Tangier (Becker and Stein 1912: 62; Séguay 1930: 59); Morocco, Algeria (Lindner 1936: 131); Algeria, Morocco, Tunisia (Woodley 2001: 318).

World distribution. France, Spain (Woodley 2001: 318).

***Nemotelus maculiventris* Bigot, 1861**

- = *Nemotelus andalusiacus* Lindner, 1937

New record. Morocco, Rif: Oued Zandoula (Fig. 26), 1♂ 2♀, 06/V/2015, sweep net, coll. Yimlahi and Belqat.

North African literature record. Algeria (Woodley 2001: 318).

World distribution. Italy, Spain (Woodley 2001: 318).

***Nemotelus marinus* Becker, 1902**

World distribution. Only known in North Africa from Egypt (Badrawy 2006: 254); Egypt: Suez (Woodley 2001: 318), Fayed, Ferdan, Ismailia, Wadi Hoff, Wadi El Natroun (Mohammad et al. 2009: 100–101).

***Nemotelus matrouhensis* Mohammad, Fadl, Gadalla & Badrawy, 2009**

World distribution. Only known in North Africa from Egypt (Mohammad et al. 2009: 100–101).

***Nemotelus nigrifrons* Loew, 1846**

- = *Nemotelus tomentosus* Becker, 1906

New locality. Affluent Tarmast (Fig. 27), 2♂♂ 1♀, 26/VI/2013, sweep net, coll. Yimlahi and Belqat.



Figure 27. Habitat of *Nemotelus nigrifrons*: Affluent Oued Tarmast.

North African literature records. Morocco: Tangier (Becker and Stein 1912: 62); Morocco, Algeria, Tunisia (Lindner 1936); Algeria, Libya, Morocco, Tunisia (Woodley 2001: 319).

World distribution. Israel, Italy (Woodley 2001: 319); Sardinia (Mason et al. 2009: 512).

Nemotelus niloticus Olivier, 1811

- = *Nemotelus albifacies* Becker, 1902 syn. n.
- = *Nemotelus duofasciatus* Woodley 2001
- = *Nemotelus fasciatus* Olivier, 1811
- = *Nemotelus oasis* Becker, 1906
- = *Nemotelus theodori* Lindner, 1974

North African literature records. Algeria: Biskra (Lindner 1936: 117); Egypt: Alexandria (Woodley 2001: 311), Egypt (Badrawy 2006: 254), Dekhela Mariout, Fayid, Fayoum, Gabal Asfer, Dekhela, Ismailia, Mallaha Mariout, Ramleh, Sherbin, Zaranik protectorate (Mohammad et al. 2009: 98–101).

World distribution. United Arab Emirates (Hauser 2008: 598); Sardinia (Mason et al. 2009: 512).

Nemotelus notatus Zetterstedt, 1842

- = *Nemotelus brachystomus* Loew, 1846
- = *Nemotelus nigroaeneus* Verhoeff, 1891
- = *Nemotelus punctiventralis* Becker, 1902
- = *Nemotelus brachystomus* form *aegyptiacus* (Lindner, 1925)
- = *Nemotelus balearicus* Lindner, 1937
- = *Nemotelus zernyi* Lindner, 1937

North African literature record. Egypt (Badrawy 2006: 254), Egypt: Coastal strip (Mohammad et al. 2009: 102–103).

World distribution. Austria, Belgium, Denmark, England, Finland, France, Germany, Ireland, Netherlands, Norway, Poland, Spain, Sweden (Woodley 2001: 313); Sardinia (Mason et al. 2009: 515–520); Turkey (Üstüner et al. 2002: 21).

Nemotelus pantherinus (Linnaeus, 1758)

- = *Musca pantherina* Linnaeus, 1758
- = *Stratiomys albipes* Geoffroy in Fourcroy, 1785
- = *Stratiomys marginellus* Thunberg, 1789
- = *Musca marginella* Gmelin, 1790
- = *Nemotelus nigritus* Meigen, 1804
- = *Nemotelus marginellus* Fallen, 1817
- = *Nemotelus nigritus* Meigen, 1822
- = *Nemotelus Jraternus* Loew, 1846
- = *Nemotelus gracilis* Loew, 1846
- = *Nemotelus satunini* Pleske in Lindner, 1937
- = *Nemotelus zelleri* Pleske in Lindner, 1937
- = *Nemotelus albirostris* Szilady, 1941
- = *Nemotelus caucasicus* Nartshuk, 1969
- = *Nemotelus punctirostris* Lindner, 1974

North African literature record. Morocco: Tangier (Séguy 1930: 59), Morocco (Woodley 2001: 320).

World distribution. Albania, Armenia, Austria, Azerbaijan, Belgium, Bulgaria, Cyprus, Czech Republic, Denmark, England, Estonia, France, Germany, Greece, Hungary, Ireland, Israel, Italy, Latvia, Netherlands, Norway, Poland, Romania, Russia, Slovakia, Spain, Sweden, Switzerland, Tajikistan, Turkey, Yugoslavia (Woodley 2001: 320); Sardinia (Mason et al. 2009: 521).

***Nemotelus proboscideus* Loew, 1846**

- = *Nemotelus punctatus* Fabricius, 1794
= *Nemotelus algericus* Jaennicke, 1866

North African literature records. Morocco, Tunisia (Lindner 1936: 139); Algeria, Libya, Tunisia, omitted in Morocco by Woodley (2001: 321).

World distribution. Italy (Woodley 2001: 321).

***Nemotelus punctiventris* Becker 1902**

World distribution. Only known in North Africa from Egypt (Mohammad et al. 2009: 100–101).

***Nemotelus subuliginosus* Rozkošný, 1974**

World distribution. Only known in North Africa from Morocco: Tangier (Woodley 2001: 322).

***Nemotelus ventralis* Meigen, 1830**

World distribution. Only known in North Africa from Morocco (Lindner 1936: 146), Morocco: Essaouira (Woodley 2001: 323).

Subgenus *Camptopelta* (Williston, 1917)

***Nemotelus nigrinus* Fallen, 1817**

- = *Nemotelus carneus* Walker, 1849
= *Nemotelus crassus* Loew, 1863
= *Nemotelus unicolor* Loew, 1863
= *Nemotelus carbonarius* Loew, 1869

North African literature record. Morocco (Woodley 2001: 326).

World distribution. Nearctic: Canada, USA. Neotropical: Mexico. Palaearctic: Afghanistan, Austria, Azerbaijan, Belgium, Bulgaria, China, Czech Republic, Denmark, England, Estonia, Finland, Germany, Hungary, Ireland, Latvia, Lithuania, Mongolia, Netherlands, Norway, Poland, Romania, Russia, Slovakia, Spain, Sweden, Switzerland, Tibet, Ukraine, Yugoslavia (Woodley 2001: 326), Turkey (Üstüner 2010: 110).

Acknowledgments

It is our pleasure to thank warmly Dr. Martin Hauser, Plant Pest Diagnostics, California Department of Food & Agriculture, for sending important references. We also wish to thank Professor Haitham Badrawy, Lecturer of Entomology, Faculty of Science Ain Shams University, Cairo, for his generous help and kindness.

References

- Afzan H, Belqat B (2016) Faunistic and bibliographical inventory of the Psychodinae moth-flies of North Africa (Diptera, Psychodidae). *ZooKeys* 558: 119–145. <http://doi.org/10.3897/zookeys.558.6593>
- Badrawy HBM (2006) Taxonomic studies including numerical taxonomy of family: Stratiomyidae (Diptera: Brachycera) from Egypt, Thesis submitted in Partial Fulfilment of the Requirements for the Degree of M.Sc. in Entomology Science (Taxonomy), Department of Entomology, Faculty of Science, Ain Shams University, 252–254.
- Becker T, Stein P (1912) Dipteren aus Marokko. *Bulletin de l'Académie Impériale des Sciences de St.-Pétersbourg VI* série 6: 9, 602–603.
- Hauser M (1997) A new species of *Nemotelus* from North-Afrika (Diptera, Stratiomyidae). *Studia dipterologica* 4(2): 453–456.
- Hauser M (2008) Order Diptera, family Stratiomyidae. In: van Harten A (Ed.) A. Arthropod Fauna of the United Arab Emirates. Volume 1. Dar Al Ummah Publishers, Abu Dhabi, 754 pp.
- Hauser M (2014) Order Diptera, family Stratiomyidae. Further records and the description of a new species. In: van Harten A (Ed.) Arthropod Fauna of the United Arab Emirates 5: 695–703.
- Kassebeer CF (1996) A new species of the genus *Beris* Latreille, 1802 from Morocco (Diptera, Stratiomyidae). *Studia dipterologica* 3(1): 155–159.
- Lindner E (1925) Neue ägyptische Stratiomyidae (Dipt.). *Bulletin de la Société Royale Entomologique d'Egypte* 9: 145–151.
- Lindner E (1939) Neue westpaläarktische Stratiomyiden (Diptera). *Zoologischer Anzeiger* 127: 312–315.
- Lindner E (1936) 18. Stratiomyiidae [part]. Lieferung 104. In: Lindner E (Ed.) Die Fliegen der palaearktischen Region. Band IV. E. Schweizerbart'sche Verlagsbuchhandlung (Erwin Nägele), Stuttgart, 218 pp.
- Mason F (1989) A new species of *Nemotelus* from Crete Island (Greece) (Diptera, Stratiomyidae). *Boletín de la Asociación española de Entomología* 121(2): 147–150.
- Mason F, Rozkošný R (2003) Interesting records of European Stratiomyidae including description of the female of *Nemotelus danielsoni* (Diptera, Stratiomyidae). *Bollettino della Società Entomologica Italiana* 134(3): 253–264.
- Mason F, Rozkošný R, Hauser M (2009) A review of the soldier flies (Diptera: Stratiomyidae) of Sardinia. In: Cerretti P, Mason F, Minelli A, Nardi G, Whitmore D (Eds) Research on the Terrestrial Arthropods of Sardinia (Italy). *Zootaxa* 2318: 507–530.

- Mohammad SK, Fadl HH, Gadalla SMM, Badrawy HBM (2009) A revision of the genus *Nemotelus* Geoffroy (Diptera: Stratiomyidae) from Egypt, with one new species and three new records. Egyptian Academic Journal of Biological Sciences 2(2): 95–110.
- Nagatomi A, Iwata K (1976) Female terminalia of lower Brachycera—I (Diptera). Beitrage Zur Entomologie, Berlin 26: 5–47.
- Pârvu C, Răzvan PM, Răzvan Z (2006) Faunistic Data on some Dipteran Families (Insecta: Diptera) from Morocco. Travaux du Muséum d'Histoire Naturelle "Grigore Antipa" 274: 271–281.
- Roháček J, Hora M (2013) A northernmost European record of the alien black soldierfly *Hermetia illucens* (Linnaeus, 1758) (Diptera: Stratiomyidae). Casopis Slezskoho Zemskeho Muzea 62(2): 101–106.
- Rozkošný R (1977) The West Palaearctic species of *Nemotelus* Geoffroy (Diptera, Stratiomyidae). Folia Facultatis Scientiarum Naturalium Universitatis Purkynianae Brunensis 51(3): 1–105.
- Rozkošný R (1982) A Biosystematic study of the European Stratiomyidae (Diptera). Vol 1. W. Junk, The Hague, Boston, London, 401 pp.
- Rozkošný R (1983) A Biosystematic study of the European Stratiomyidae (Diptera). Vol. 2. W. Junk, The Hague, Boston, London, 431 pp.
- Rozkošný R, Nartshuk EP (1988) Family Stratiomyidae. In: Soós Á, Papp L (Eds) Catalogue of Palaearctic Diptera 5: 42–96.
- Rozkošný R (1997) Diptera Stratiomyidae, Soldier Flies. In: Nilsson A (Ed.) Aquatic Insects of North Europe— A taxonomic Handbook. Vol. 2. Odonata – Diptera. Apollo Books, Stenstrup, 321–332.
- Rozkošný R (1998) Chapter 24. Family Stratiomyidae. Manual of Palaearctic Diptera 2: 387–411.
- Rozkošný R (2004) Fauna Europaea: Stratiomyidae. – In: Pape T (Ed.) Fauna Europaea: Diptera Brachycera. Fauna Europaea version 1.1, <http://www.faunaeur.org> [accessed September 2009 as version 1.3 of April 19th 2007]
- Séguy E (1930) Contribution à l'étude des Diptères du Maroc. Mémoires de la Société des Sciences Naturelles du Maroc 24: 5–206.
- Séguy E (1953) Diptères du Maroc. - Encyclopédie Entomologique, Série (B) II. Diptera., 11: 77–92.
- Üstüner T, Hasbenli A, Aktümsek A (2002) Contribution to Subfamily Clitellariinae (Diptera, Stratiomyidae) Fauna of Turkey. Journal of the Entomological Research Society 4(1): 19–24.
- Üstüner T (2010) New records of Stratiomyidae (Diptera) from Turkey. Israel Journal of Entomology 40: 109–111.
- Tkoč M, Rozkošný R (2014) New synonymy of *Oplodontha minuta* (Diptera: Stratiomyidae) with its first record from Socotra Island. Acta entomologica Musei nationalis Pragae 54: 429–439.
- Woodley NE (2001) A World Catalog of the Stratiomyidae (Insecta: Diptera), Backhuys Publishers. Leiden, The Netherlands, 473 pp.
- Woodley NE (2011) A World Catalogue of the Stratiomyidae (Insecta: Diptera): A Supplement with Revisionary Notes and Erata. Myia 12: 485–521.