

Annotated catalogue of Iranian burrower bugs (Heteroptera, Pentatomoidea, Cydnidae)

Hassan Ghahari¹, Frédéric Chérot^{2,5}, Rauno E. Linnauori³, Hadi Ostovan⁴

1 Department of Agriculture, Islamic Azad University, Share Rey Branch, Tehran, Iran **2** Laboratoire de Systématique et d'Ecologie animales, CP 160/13, Av F. D. Roosevelt, 50, B-1050 Bruxelles, Belgique **3** Saukkokuja 10, FIN-21220 Raisio, Finland **4** Department of Entomology, Fars Science and Research Branch, Islamic Azad University, Marvdasht, Iran **5** Service Public de Wallonie, DG03, DEMNA, av. Maréchal Juin, 23, B-5030 Gembloux, Belgique

Corresponding author: Frédéric Chérot (frederic.cherot@spw.wallonie.be)

Academic editor: Thomas Henry | Received 21 June 2009 | Accepted 26 October 2009 | Published 30 October 2009

Citation: Ghahari H, Chérot F, Linnauori RE, Ostovan H (2009) Annotated catalogue of Iranian burrower bugs (Heteroptera, Pentatomoidea, Cydnidae). ZooKeys 26: 1–31. doi: 10.3897/zookeys.26.214

Abstract

A catalogue of burrower bugs (Heteroptera, Pentatomoidea, Cydnidae) of Iran is provided. A total of 58 species from 5 subfamilies, 6 tribes and 22 genera is listed in this paper. Of these, 14 species are newly recorded from Iran: *Byrsinus fossor* (Mulsant & Rey, 1866), *Byrsinus nigroscutellatus* (Montandon, 1900), *Byrsinus penicillatus* Wagner, 1964, *Canthophorus wagneri* Asanova, 1964, *Crocistethus waltianus* (Fieber, 1837), *Geotomus antennatus* Signoret, 1883, *Sehirus cypriacus* Dohrn, 1860, *Sehirus dissimilis* Horváth, 1919, *Sehirus luctuosus* Mulsant & Rey, 1866, *Sehirus ovatus* (Herrich-Schaeffer, 1840), *Sehirus parens* Mulsant & Rey, 1866, *Sehirus planiceps* Horváth, 1895, *Stibaropus henkei* (Jakovlev, 1874) and *Tritomegas delagrangei* (Puton, 1888). Additional Iranian records are provided for *Byrsinus laeviceps* (Kerzhner, 1972), *Exosehirus marginatus* (Signoret, 1881b), *Fromundus pygmaeus* (Dallas, 1851), *Geotomus elongatus* (Herrich-Schaeffer, 1840), *Geotomus punctulatus* (Costa, 1847), *Ochetostethus sahlbergi* Wagner, 1952 and *Tritomegas bicolor* (Linnaeus, 1758).

Keywords

Heteroptera, Cydnidae, Fauna, Iran

Introduction

The burrower or burrowing bugs (Insecta: Heteroptera: Cydnidae) include about 680 species worldwide, mainly in tropical parts of the both Old and New Worlds (Lis 2006). As suggested by their common name, a majority of cydnids are adapted to burrowing in the ground, feeding on roots of plants. The family does not include major agricultural pests (Lis *op. cit.*; Lis et al. 2000), although an outbreak in populations of some species could cause serious damage (Froeschner 1960, 1988).

Even if the monophyly and the delimitation of family Cydnidae has been questioned recently (Grazia et al. 2008; Pluot-Sigwalt and Lis, 2008), these insects are undoubtedly closely related to the Pentatomidae. To such an extent that they were classified as a subfamily of the Pentatomidae in early literature.

The Cydnidae can be separated from the Pentatomidae by a combination of:

- (a) their ovoid and convex body, frequently shinning, black or brown;
- (b) their broad, flattened head and legs adapted for digging;
- (c) their antenna generally 5-segmented and their tarsi generally 3-segmented and
- (d) their tibial armament. Pentatomidae have tibial hairs, but lack stout, hardened tibial spines that are characteristic of the Cydnidae.

The fauna of Iranian Cydnidae has been poorly studied. The main works concerning Cydnidae from Iran are the following: Hoberlandt (1955), Safavi (1959), Baroughi (1978), Modarres Awal (1987), Hoberlandt (1995), Modarres Awal (1996a, b, 1997), Lis (1998, 1999), Linnauori (2004), Lis (2006), Magnien (2006), Linnauori (2007), Modarres Awal (2008), Askari et al. (2009), Hassanzadeh et al. (2009), and Sadeghi et al. (2009). In a recent checklist of Iranian agricultural pests (Modarres Awal, 1997: 71), only 18 species of Cydnidae were listed. Magnien (2006) described one new species that occurs in Iran and Linnauori (2004, 2007) and Modarres Awal (2008) reported several other new species records for Iran. So, prior to our study, the total number of cydnids species recorded from Iran was 44.

We provide hereafter for the first time a catalog of Iranian Cydnidae, including a total of 58 species, 14 of which constitute new records for Iran. The new and additional records for Iran were analyzed by the third author and the specimens are preserved in his collection.

The arrangement of synonymies is adapted from Lis (1999, 2006); the distributional data including distribution outside Iran for each species were taken from Lis (2006). The classification of subfamilies and tribes follows in part Froeschner (1960), Lis (1994, 1999, 2006) and Pluot-Sigwalt and Lis (2008).

Catalogue

Subfamily Amaurocorinae Wagner, 1963

Tribe Amaurocorini Wagner, 1963

Genus *Amaurocoris* Stål, 1865

1) *Amaurocoris curtus* (Brullé, 1839)

Cydnus curtus Brullé, 1839: 81 (as new species).

Amaurocoris curtus: Lethierry and Severin 1893: 76 (new combination).

Iran records: Hoberlandt 1955: 122, 1995: 187–188; Linnauvori 2004: 285, 2007: 89; Lis 1999: 218–219, 2006: 138; Modarres Awal 1996a: 127, 1997: 71; Safavi 1959: 31; Wagner 1968: 438.

Distribution in Iran: Abbassi, Bushehr, Fars, Hormozgan, Kerman, Khorasan, Khuzestan, Sistan & Balouchestan provinces.

Distribution outside Iran: Afghanistan, Algeria, Armenia, Azerbaidjan, Canary Islands, Cape Verde Islands, Cyprus, Egypt, Eritrea, Ethiopia, Greece, India (?), Iraq, Israel, Kenya, Jordan, Libya, Mauretania, Morocco, Pakistan, Saudi Arabia, Senegal, Somalia, Sudan, Syria, Tadzhikistan, Tunisia, Turkey, Turkmenistan, Uzbekistan, Yemen.

Genus *Linospa* Signoret, 1881a

2) *Linospa candida* (Horváth, 1889)

Amaurocoris candidus Horváth, 1889: 173 [name only].

Linospa candida: Lis 1993a: 70 (new combination).

Iran records: Hoberlandt 1995: 188–189 [under the original combination]; Linnauvori 2007: 89; Lis 1999: 219, 2006: 138.

Distribution in Iran: Khorasan province.

Distribution outside Iran: Afghanistan, Kazakhstan, Turkmenistan, Uzbekistan.

3) *Linospa orbicularis* (Jakovlev, 1885)

Amaurocoris orbicularis Jakovlev, 1885: 110 (as new species).

Linospa orbicularis: Lis 1993a: 72 (new combination).

Iran records: Hoberlandt 1995: 188; Lis 1999: 219, 2006: 139; Seidenstücker 1957: 73 [under the name *A. aspericollis* Puton, 1886 a junior synonym]; Wagner 1968: 438 [under the combination *A. orbicularis* Jakovlev, 1885].

Distribution in Iran: Abbassi, Kerman, Khorasan, Khuzestan, Sistan & Baluchestan provinces.

Distribution outside Iran: Afghanistan, Algeria, Canary Islands, Egypt, Eritrea, Iraq, Israel, Libya, Morocco, Pakistan, Saudi Arabia, Sudan, Tadzhikistan, Tunisia, Turkmenistan, Uzbekistan, Yemen.

Subfamily Amnestinae Hart, 1919

Tribe Amnestini Hart, 1919

Genus *Amnestus* Dallas, 1851

4) *Amnestus raunoi* Lis, 1998

Amnestus raunoi Lis, 1998: 5–7 (as new species).

Iran records: Linnavuori 2007: 88; Lis 1998: 5–7, 2006: 119.

Distribution in Iran: Gilan province.

Distribution outside Iran: Unknown, apparently endemic to Iran.

Comment: *Amnestus raunoi* is very close to the species *Amnestus pusillus* Uhler. These two species were studied and compared taxonomically by Lis (1998).

Subfamily Cephalocheinae Mulsant & Rey, 1866

Tribe Scaptocorini Froeschner, 1960

Genus *Stibaropus* Dallas, 1851

5) *Stibaropus henkei* (Jakovlev, 1874)

Pachycnemis henkei Jakovlev, 1874a: 56 (as new species).

Stibaropus henkei: Signoret 1881a: 46 (new combination).

Material examined: East Azarbaijan province, Arasbaran (835m), 1♂, 3♀♀, September 2005. **New record for Iran.**

Distribution outside Iran: Armenia, Azerbaidjan, Bulgaria, Hungary, Kazakhstan, Kirgizia, Romania, Russia (Central and South European Territory), Tadzhikistan, Turkmenistan, Ukraine, Uzbekistan.

Subfamily Cydninae Billberg, 1820

Tribe Cydnini Billberg, 1820

Genus *Cydnus* Fabricius, 1803

6) *Cydnus aterrimus* (Forster 1771)

Cimex aterrimus Forster, 1771: 71 (as new species).

Cydnus aterrimus: Dallas 1851: 121 (new combination).

Cydnus infernalis Kiritshenko 1966: 799 (as new species ; the synonymy by Lis 1996: 405).

Iran records: Askari et al. 2009: 562; Baroughi 1978: 22; Hassanzadeh et al. 2009: 22; Hassanzadeh et al. 2009: 529; Hoberlandt 1955: 122, 1995: 187; Lindberg 1938: 89 ; Linnauvori 2007: 88; Lis 1999: 180, 2006: 124; Modarres Awal 1987: 19, 1996a: 127, 1997: 71, 2008: 246; Sadeghi et al. 2009: 499; Safavi 1959: 31; Wagner 1968: 438.

Distribution in Iran: Bushehr, East Azarbaijan, Fars, Golestan, Gorgan, Kerman, Khorasan, Mazandaran, Sistan & Baluchestan, Tehran, Zanjan provinces.

Distribution outside Iran: Afghanistan, Albania, Algeria, Armenia, Austria, Azerbaijan, Bangladesh, Belgium, Bosnia Herzegovina, Bulgaria, Canary Islands, China (Neimenggu, Ningxia), Congo (?), Crimea, Croatia, Cyprus, Czech Republic, Egypt, France, Georgia, Germany, Greece, Hungary, Iraq, India, Israel, Italy, Jordan, Kazakhstan, Kirgizia, Lebanon, Libya, Lithuania, Luxembourg, Macedonia, Madeira, Malta, Malaya (?), Malta, Moldavia, Morocco, Netherlands, Pakistan, Poland, Portugal, Romania, Russia (South European Territory, West Siberia), Senegal, Serbia and Montenegro, Slovakia, Slovenia, South Africa (introduction), Spain, Sri Lanka, Switzerland, Syria, Tadzhikistan, Tunisia, Turkey, Turkmenistan, Ukraine, USA (introduction), Uzbekistan, West Indies (introduction), Yemen.

Genus *Nishadana* Distant, 1899b

7) *Nishadana arabica* Horváth, 1919

Nishadana arabica Horváth, 1919: 265 (as new species).

Iran records: Hoberlandt 1995: 190; Lis 2006: 120.

Distribution in Iran: Lorestan province.

Distribution outside Iran: Chad, Gambia, Ghana, Senegal, Somalia, Yemen.

Remark: the genus *Nishadana* Distant, 1899 was reclassified in the tribe Cydnini of the subfamily Cydninae by Pluot-Sigwalt and Lis (2008) on the basis of spermatheca structure.

Tribe Geotomini Wagner, 1963**Genus *Aethus* Dallas, 1851****8) *Aethus hispidulus* (Klug, 1845)**

Cydnus hispidulus Klug, 1845: [10] (as new species).

Aethus hispidulus: Walker 1867: 154 (new combination).

Iran records: Hoberlandt 1995: 191; Lis 1999: 186, 2006: 126; Putshkov 1965: 35.

Distribution in Iran: North Iran (Putshkov 1965, without exact locality).

Distribution outside Iran: Afghanistan, Armenia, Azerbaidjan, Cyprus, Egypt, Greece, Iraq, Israel, Jordan, Saudi Arabia, Somalia, Syria, Turkey, Turkmenistan (Kugitang-Tau Mountains), Uzbekistan.

9) *Aethus ostiolatus* Signoret, 1882

Aethus ostiolatus Signoret, 1882: 433 (as new species).

Iran records: Hoberlandt 1995: 183 [under the name *Aethus macropthalmus* Wagner, 1951, a junior synonym]; Lis 1999: 187, 2006: 126; Seidenstücker, 1957: 73 [under the name *Aethus macropthalmus* Wagner, 1951].

Distribution in Iran: Abbassi, Baluchestan, Bushehr, Khuzestan provinces.

Distribution outside Iran: Algeria, Chad, Egypt, Kenya, Saudi Arabia, Somalia, Sudan, Yemen.

10) *Aethus pilosus* (Herrick-Schaeffer, 1834)

Cydnus pilosus Herrich-Schaeffer, 1834: plate 22 and legend.

Aethus pilosus: Walker 1867: 148 (new combination).

Iran records: Hoberlandt 1955: 121, 1995: 191; Lis 1999: 188, 2006: 126 ; Modarres-Awal 1997: 71; Safavi 1959: 31.

Distribution in Iran: Sistan & Balouchestan province.

Distribution outside Iran: Afghanistan, Algeria, Cameroon, Croatia, Egypt, Ethiopia, France, Greece, Israel, Italy, Jordan, Libya, Malta, Portugal, Saudi Arabia, Senegal, Serbia and Montenegro, Spain, Syria, Tunisia, Turkey.

Genus *Alonips* Signoret, 1881b

11) *Alonips acrostictus* (Distant, 1918)

Geotomus acrostictus Distant, 1918: 118 (as new species).

Alonips acrostictus: Lis 1994: 156 (new combination).

Iran records: Linnavuori 2004: 286; Lis 2006: 127.

Distribution in Iran: Hormozgan province.

Distribution outside Iran: China (Hainan), India, Laos, Pakistan, Sri-Lanka, Thailand, Vietnam.

12) *Alonips intrusus* (Wagner, 1953)

Geotomus intrusus Wagner, 1953: 467 (as new species).

Alonips intrusus: Lis 1994: 155 (new combination).

Iran records: Linnavuori 2004: 285; Lis 2006: 127.

Distribution in Iran: Hormozgan province.

Distribution outside Iran: Algeria, Burkina Faso, Canary Islands, Egypt, Eritrea, Iraq, Mali, Saudi Arabia, Senegal, Sudan, Tanzania, Tunisia, Yemen.

Genus *Byrsinus* Fieber, 1860

13) *Byrsinus comaroffii* (Jakovlev, 1879)

Cydnus comaroffii Jakovlev, 1879: 21 (as new species).

Byrsinus komaroffi: Wagner 1965: 50 (new combination).

Byrsinus comaroffii: Lis 1993b: 85 (revision, spelling).

Iran records: Hoberlandt 1995: 185 [under the name *Aethus comaroffi* (Jakovlev, 1879)]; Lis 1999: 191, 2006: 128.

Distribution in Iran: Abbassi, Baluchestan, Kerman, Khorasan.

Distribution outside Iran: Armenia, Afghanistan, Azerbaijan, China (Neimenggu, Xinjiang), Kazakhstan, Kirgizia, Mongolia, Russia (South European Territory), Tadzhikistan, Turkmenistan, Uzbekistan.

***Byrsinus discus* Jakovlev, 1906**

Byrsinus discus Jakovlev, 1906: 52 (as new species).

Iran records: Putshkov 1965: 41 [under the combination *Peribyssus discus* (Jak., 1906)]; Hoberlandt 1995: 185 [under the combination *Peribyssus discus* (Jakovlev, 1906)].

Distribution in Iran: The mention of *Byrsinus discus* Jakovlev, 1906 from North Iran, without exact locality information, by Putshkov (1965: 41) was hypothetical. Putshkov wrote “Described from Repetek (Turkmenistan), mostly it occurs in the upper layer of sandy soil near tufts of grasses. Recorded also from Uzbekistan and certainly will be recorded from grasslands in other Central Asian countries, including (...) northern Iran...” (Translated by D. Redei, our italicization). Hoberlandt (1995: 191) recorded the species from Iran referring only to Putshkov’s work but as a “published records”. Because Lis (2006: 128) did not list *B. discus* from Iran, its presence in Iran needs confirmation.

Distribution outside Iran: Turkmenistan, Uzbekistan.

14) *Byrsinus flavidornis* (Fabricius, 1794)

Cimex flavidornis Fabricius, 1794: 124 (as new species).

Byrsinus flavidornis: Lis 1993b: 86 (new combination).

Iran records: Hoberlandt 1955: 122, 1995: 190; Lis 1999: 192, 2006: 128; Modarres-Awal, 1997: 71; Safavi 1959: 31 [under the combination *Aethus flavidornis*, doubtful data].

Distribution in Iran: Khuzestan.

Distribution outside Iran: Afghanistan, Algeria, Armenia (?), Azerbaijan (?), Belgium, Bulgaria, Byelorussia (?), Canary Islands, China (Neimenggu, Xinjiang), Croatia, Cyprus, Czech Republic, Egypt, France, Germany, Great Britain (as migrant), Hungary, Italy, Kazakhstan, Latvia (?), Libya, Lithuania (?), Macedonia, Madeira, Malta, Moldavia, Mongolia, Morocco, Netherlands, Poland, Portugal, Romania, Russia [Central and South European Territory, East and West Siberia, Far East (?)], Serbia and Montenegro, Slovakia, Spain, Tunisia, Turkey, Turkmenistan, Ukraine.

15) *Byrsinus fossor* (Mulsant & Rey, 1866)

Cydnus (Byrsinus) fossor Mulsant and Rey, 1866: 18 (as new species).

Byrsinus fossor: Walker 1867: 164 (new combination).

Material examined: Mazandaran province, Beshahr (57m), 1♀, September 2006. **New record for Iran.**

Distribution in Iran: Mazandaran province.

Distribution outside Iran: Armenia, Azerbaidjan, Bulgaria, China (Xinjiang), France, Hungary, Kazakhstan, Romania, Russia (South European Territory), Serbia and Montenegro, Spain, Ukraine.

Comment: The records of *Byrsinus fossor* from Afghanistan, Egypt, Mongolia, Tadzhikistan and Turkmenistan pertain to other species of the genus, according to Lis (2006: 19).

16) *Byrsinus laeviceps* (Kerzhner, 1972)

Aethus laeviceps Kerzhner, 1972: 363 (as new species).

Byrsinus laeviceps: Lis 1993b: 87 (new combination).

Iran records: Hoberlandt 1995: 184–185 [under the original combination]; Linnavuori 2007: 88.

Additional data: Khorasan province, Serakhs (330m), 2♂♂, June 2001.

Distribution in Iran: Baluchestan, Kerman, Khorasan and Tehran provinces.

Distribution outside Iran: China (Xinjiang), Kazakhstan, Mongolia, Turkmenistan, Uzbekistan.

17) *Byrsinus laticollis* (Wagner, 1954)

Aethus (Cydnus) laticollis Wagner, 1954: 1 (as new species).

Byrsinus laticollis: Lis 1993b: 88 (new combination).

Iran records: Hoberlandt 1995: 183 (under the name *Aethus laticollis*).

Distribution in Iran: Tehran province.

Distribution outside Iran: Canary Islands, Italy, Libya, Madeira, Morocco, Spain, Tunisia.

***Byrsinus minor* Wagner, 1964.** The record of *Byrsinus minor* Wagner, 1964 from Gilan province (Assalem, 8 km NW Bandar-Pahlavi) by Wagner (1968: 437), also quoted by Hoberlandt (1995: 191), is erroneous, according to Lis (2006: 129). Consequently, this species is still not known from Iran.

18) *Byrsinus nigroscutellatus* (Montandon, 1900)

Byrsinocoris nigroscutellatus Montandon, 1900: 157 (as new species).

Byrsinus nigroscutellatus: Lis 2003: 40 (new synonymy and combination).

Material examined: Ilam province, Ilam (1430m), 1♂, 3♀♀, May 2004. **New record for Iran.**

Distribution outside Iran: Algeria, Chad, Egypt, Libya, Mauretania, Oman.

19) *Byrsinus pallidus* (Puton, 1887)

Cydnus pallidus Puton in Puton and Lethierry, 1887: 298 (as new species).
Byrsinus pallidus: Lis 1993b: 89 (new combination).

Iran records: Hoberlandt 1995: 184 [under the name *Aethus pallidus* (Puton, 1887)]; Linnavuori 2004: 285; Lis 1999: 193, 2006: 130; Wagner 1968: 437 [under the name *A. pallidus* (Puton, 1887)].

Distribution in Iran: Abbassi, Baluchestan, Golestan, Hormozgan provinces.

Distribution outside Iran: Algeria, Canary Islands, Djibouti, Egypt, Iran, Israel, Italy (?), Libya, Mauretania, Morocco, Pakistan, Saudi Arabia, Sudan, Tunisia, Yemen.

20) *Byrsinus penicillatus* Wagner, 1964

Byrsinus penicillatus Wagner, 1964: 81 (as new species).

Material examined: Khorasan province, Torbat-Jam (881m), 1♂, 2♀, September 2003. **New record for Iran**.

Distribution outside Iran: Afghanistan, China (Xinjiang, Xizang), Kazakhstan, Mongolia, Turkmenistan, Uzbekistan.

21) *Byrsinus pilosulus* (Klug, 1845)

Cydnus pilosulus Klug, 1845: [10] (as new species).
Byrsinus pilosulus: Lis 1993b: 90 (new combination).

Iran records: Hoberlandt 1995: 183 [under the name *Aethus pilosulus* (Klug, 1845), an incorrect spelling]; Linnavuori 2004: 285, 2007: 88; Lis 1999: 193, 2006: 130; Seidenstücker, 1957: 73 [under the combination *Aethus pilosulus* Klug].

Distribution in Iran: Abbassi, Baluchestan, Fars, Gilan, Hormozgan, Kerman, Khorasan, Sistan, Tehran, Zanjan provinces.

Distribution outside Iran: Afghanistan, Algeria, Bulgaria, Canary Islands, Cape Verde Islands, Egypt, France, Germany (?), Greece, Hungary, India, Iraq, Israel, Italy, Libya, Madeira, Malta, Mauretania, Morocco, Pakistan, Romania, Saudi Arabia, Socotra, Serbia and Montenegro, Spain, Sudan, Syria, Tadzhikistan, Tunisia, Turkey, Turkmenistan, Uzbekistan, Yemen.

22) *Byrsinus rugosus* (Jakovlev, 1874)

Cydnus rugosus Jakovlev, 1874b: 235 (as new species).
Byrsinus rugosus: Lis 1993b : 91 (new combination).

Iran records: Hoberlandt 1995 : 190 [under the combination *Aethus rugosus* (Jakovlev, 1874)]; Lis 1999: 194, 2006: 130; Modarres-Awal, 1996a: 126, 1997: 71 [both as *Aethus rugosus* (Jakovlev, 1874)]; Wagner, 1968: 437.

Distribution in Iran: Abbassi, Hormozgan, Khorasan provinces.

Distribution outside Iran: Afghanistan, Armenia (?), Azerbaijan, China (Neimenggu, Xinjiang), Egypt, Iraq, Kazakhstan, Mongolia, Russia (South European Territory), Saudi Arabia, Tadzhikistan, Turkmenistan, Uzbekistan.

Comment: Safavi (1959) reported *Aethus flavigornis* (Fabricius) from Iran (Khuzestan province). Because specimen(s) were not available to us, we do not know if the involved species is *Byrsinus flavigornis* (Fabricius, 1794) or *B. rugosus* (Jakovlev, 1874).

Genus *Fromundiellus* Lis, 1994

23) *Fromundiellus latiusculus* (Horváth, 1882)

Geotomus latiusculus Horváth, 1882: 217 (as new species).
Fromundiellus latiusculus: Lis 1997: 4 (new combination).

Iran record: Modarres Awal 2008: 247.

Distribution in Iran: Khorasan province.

Distribution outside Iran: Afghanistan, Azerbaijan, India (North), Kirgizia, Pakistan, Tadzhikistan, Turkmenistan, Uzbekistan.

Genus *Fromundus* Distant, 1901

24) *Fromundus pygmaeus* (Dallas, 1851)

Aethus pygmaeus Dallas, 1851: 120 (as new species).
Fromundus pygmaeus: Lis 1994: 181 (new combination).

Iran records: Linnauori 2004: 286; Lis 2006: 132.

Additional data: Kermanshah province, Sar-pole-zahab (558), 2♀♀, October 2001.

Distribution in Iran: Hormzegan and Kermanshah provinces.

Distribution outside Iran: Australia, Bismarck Archipelago (Manus, Mussau, New Ireland, New Britain), Burma, Brunei, Cambodia, Chagos Archipelago, China (Sichuan, Guangdong, Hainan, Guangxi, Yunnan, Jiangxi, Hubei, Taiwan), Christmas Islands, Cocos-Keel-

ing Islands, Fiji, Hawaii, Hong Kong, India (incl. Andamans and Nicobars), Indonesia, Iraq, Israel, Japan, Jordan, Korea, Laos, Malaysia, Maldives, Marquesas, Micronesia, Nepal, New Caledonia, New Hebrides, Pakistan, Papua New Guinea, Philippines, Samoa, Saudi Arabia, Singapore, Sri Lanka, Solomon Islands, Society Islands, Thailand, Vietnam, Yemen.

Genus *Geotomus* Mulsant & Rey, 1866

25) *Geotomus angustus* Wagner, 1953

Geotomus elongatus angustus Wagner, 1953: 471 [as new subspecies of *G. elongatus* (Herrich-Schaeffer, 1840)].

Geotomus angustus: Stichel 1961: 692 (upgrading to species rank).

Iran records: Hoberlandt 1995: 186 (under the original combination).

Distribution in Iran: Abbasi province.

Distribution outside Iran: Algeria, Cape Verde Islands, Chad, Libya, Morocco, Spain, Sudan, Tunisia, Turkey.

26) *Geotomus antennatus* Signoret, 1883

Geotomus antennatus Signoret, 1883: 56 (as new species).

Material examined: Khorasan province, Torbat-Jam (880m), 2♂♂, 2♀♀, July 2001.

New record for Iran.

Distribution outside Iran: Afghanistan, Israel, Syria, Turkey.

27) *Geotomus brunnipennis* Wagner, 1953

Geotomus brunnipennis Wagner, 1953: 466 (as new species).

Iran records: Modarres Awal 1996a: 127, 1996b: 106, 1997: 71.

Distribution in Iran: Ardabil, Khorasan provinces.

Distribution outside Iran: Albania, Austria, Bulgaria, Croatia, France, Greece, Italy, Macedonia, Serbia and Montenegro, Turkey.

28) *Geotomus ciliatitylus*, Signoret, 1881

Geotomus ciliatitylus Signoret, 1881b: 652–653 [the synonymy with *Geotomus caucasicus* (Kolenati, 1846), now *Geotomus elongatus* (Herrich-Schaeffer, 1840), by

Horváth 1918: 323, incorrect according to Lis 2006: 133; Hoberlandt's (1995: 187) remark about *G. caucasicus* subspecies distribution, based on Wagner (1953: 472–473), unjustified].

Iran records: Safavi 1959: 31 (under the combination *Geotomus caucasicus ciliatitulus*); Modarres-Awal 1997: 71; Signoret 1881b: 652–653 (type from Tehran).

Distribution in Iran: Tehran.

Distribution outside Iran: Albania, Azerbaijan, Bosnia Herzegovina, Bulgaria, Croatia, Georgia, Greece, Israel, Macedonia, Romania, Russia [South European Territory (?)], Serbia and Montenegro, Syria, Turkey, Turkmenistan, Uzbekistan.

29) *Geotomus elongatus* (Herrich-Schaeffer, 1840)

Cydnus elongatus Herrich-Schaeffer, 1840: 96 (as new species).

Cydnus caucasicus Kolenati 1846: 70 (as new species, the new combination by Horváth 1917: 371 and the synonymy by Kerzhner 2003: 95).

Geotomus elongatus: Mulsant and Rey 1866: 38 (new combination).

Iran records: Hoberlandt 1995: 186–187 (under the name *G. elongatus* and the junior synonym *G. caucasicus*); Lis 2006: 133; Wagner 1953: 471–472, 475.

Additional data: Golestan province, National Park (185m), 1♂, 2♀♀, October 2003.

Distribution in Iran: Abbassi, Baluchestan, Bushehr, Golestan, Kerman, Tehran provinces.

Distribution outside Iran: Algeria, Armenia, Azerbaidjan, Belgium, Bosnia Herzegovina, Bulgaria, Canary Islands, China (Neimenggu), Croatia, France, Georgia, Germany, Greece, Hungary, Israel, Iraq, Italy, Kazakhstan, Kirgizia, Macedonia, Morocco, Portugal, Romania, Russia (South European Territory), Serbia and Montenegro, Spain, Switzerland, Syria, Tadzhikistan, Tunisia, Turkey, Turkmenistan, Ukraine, Uzbekistan.

30) *Geotomus punctulatus* (Costa, 1847)

Cydnus punctulatus Costa, 1847: 30.

Geotomus punctulatus: Mulsant and Rey 1866: 35 (new combination).

Iran records: Hoberlandt 1995: 187; Lis 2006: 133.

Additional data: East Azarbajian province, Arasbaran (792m), 1♀, September 2005.

Distribution in Iran: East Azarbajian, Khorasan, Tehran provinces.

Distribution outside Iran: Albania, Algeria, Armenia, Azerbaidjan, Azores Islands, Belgium, Bosnia Herzegovina, Bulgaria, Canary Islands, Cyprus, France, Georgia, Germany, Great Britain, Greece, Hungary, Israel, Italy, Kazakhstan, Libya, Macedonia, Malta, Moldavia, Morocco, Portugal, Romania, Russia (South

European Territory), Serbia and Montenegro, Spain, Syria, Tunisia, Turkey, Turkmenistan, Ukraine.

Genus *Macroscytus* Fieber, 1860

31) *Macroscytus brunneus* (Fabricius, 1803)

Cydnus brunneus Fabricius, 1803: 185 (as new species).

Macroscytus brunneus: Fieber 1861: 362 (new combination).

Iran records: Hoberlandt 1955: 122, 1995: 185–186; Linnauvori 2004: 285, 2007: 88–89; Lis 1999: 208, 2000: 402–404, 2006: 135; Modarres Awal 1996a: 127, 1997: 71, 2008: 247; Safavi 1959: 31; Seidenstücker, 1957: 73, 1958: 5; Wagner 1968: 437.

Distribution in Iran: Abbassi, Fars, Gilan, Hormozgan, Kerman, Khorasan, Lorestan, Sistan & Baluchestan, Tehran, Zanjan provinces.

Distribution outside Iran: Afghanistan, Albania, Algeria, Azerbaidjan, Bosnia-Herzegovina, Botswana, Cameroon, Canary Islands, Cape Verde Islands, Central African Republic, China (North ?), Congo, Croatia, Cyprus, Egypt, Ethiopia, France, Gambia, Ghana, Greece, Guinea, Iraq, Israel, Italy, Ivory Coast, Jordan, Kazakhstan, Kenya, Lebanon, Libya, Macedonia, Mali, Malta, Morocco, Mozambique, Namibia, Nigeria, Oman, Pakistan, Portugal, Saudi Arabia, Senegal, Socotra, Somalia, South Africa, Serbia, Spain, Sudan, Switzerland, Syria, Tadzhikistan, Tanzania (incl. Zanzibar), Turkey, Turkmenistan, Tunisia, Uganda, Uzbekistan, Yemen, Zaire, Zambia, Zimbabwe.

Comment: Lis (2000) redescribed and illustrated *M. brunneus*.

Genus *Microporus* Uhler, 1872

32) *Microporus nigrita* (Fabricius, 1794)

Cimex nigrita Fabricius, 1794: 123 (as new species).

Microporus nigrita: Horváth 1917: 369 (new combination).

Iran records: Lis 1993b: 94, 1999: 212, 2006: 137.

Distribution in Iran: Gilan Province.

Distribution outside Iran: Albania, Armenia, Austria, Azerbaidjan, Belgium, Bulgaria, Byelorussia, Canary Islands, China (Beijing, Guangdong, Neimenggu, Shandong, Shanghai, Tianjin, Xinjiang, Xizang, Yunnan), Croatia, Czech Republic, Denmark (?), Ethiopia, France, Georgia, Germany, Greece, Hungary, India (Kashmir), Iran, Italy, Japan, Kazakhstan, Kirgizia, Korea, Latvia, Lithuania, Luxembourg, Macedonia, Moldavia, Mongolia, Netherlands, Poland, Romania, Russia (European Territory, Siberia, Far East), Senegal (?), Serbia and Montenegro, Slovakia, Slovenia, Spain, Sweden,

Switzerland, Tadzhikistan, Tunisia, Turkey, Turkmenistan, Ukraine, United States of America (introduced), Uzbekistan.

Subfamily Sehirinae Amyot & Serville, 1843

Tribe Sehirini Amyot & Serville, 1843

Genus *Adomerus* Mulsant & Rey, 1866

33) *Adomerus congener* (Jakovlev, 1879).

Schirus [sic!] *congener* Jakovlev, 1879: 25 (as new species).

Adomerus congener: Kerzhner and Jaczewski 1964: 825 (new combination).

Iran records: Lis 1999: 220, 2006: 139; Modarres-Awal 1987: 19 [under the combination *Sehirus congener* Jak], 1997: 71 [under the subsequent incorrect spelling *Sehirus congenes* (sic)].

Distribution in Iran: East Azarbaijan.

Distribution outside Iran: Armenia, Azerbaidjan, Georgia, Lebanon, Russia (South European Territory), Turkey.

Genus *Canthophorus* Mulsant & Rey, 1866

34) *Canthophorus coeruleus* (Reuter, 1902)

Sehirus coeruleus Reuter, 1902: 144 (as new species).

Canthophorus coeruleus: Asanova 1964: 143 (new combination).

Iran records: Brown 1966: 44; Hoberlandt 1955: 122 [under the combination *Sehirus coeruleus*], 1995: 191; Lis 1999: 220–221, 2006: 140; Modarres-Awal 1997: 71; Safavi 1959: 31 [under the combination *Sehirus coeruleus* (Reuter, 1902)]; Wagner 1968: 438.

Distribution in Iran: Tehran province.

Distribution outside Iran: Kirgizia, Tadzhikistan (?), Turkmenistan, Uzbekistan.

35) *Canthophorus dubius* (Scopoli, 1763)

Cimex dubius Scopoli, 1763: 121 (as new species).

Canthophorus (Canthophorus) dubius: Mulsant and Rey 1866: 63 (new combination).

Iran records: Brown, 1966: 44 [under the combination “*Sehirus dubius* Scop.”]; Hoberlandt 1995: 191; Lindberg 1938: 89 [under the combination “*Sehirus dubius* Scop.”];

Lis 1999: 221, 2006: 140; Modarres-Awal 1997: 71; Safavi 1959: 31 [under the combination “*Sehirus dubius* (Scopoli, 1763)”].

Distribution in Iran: Hamadan, Kerman, Tehran provinces.

Distribution outside Iran: Albania, Algeria (?), Austria, Bosnia Herzegovina, Bulgaria, Croatia, Cyprus (?), Czech Republic, Egypt (?), Finland, France, Germany, Great Britain, Greece, Hungary, Israel, Italy, Lebanon, Liechtenstein, Lithuania (?), Macedonia, Madeira, Moldavia, Poland, Romania, Russia (NW Caucasus), Serbia and Montenegro, Slovakia, Slovenia, Spain, Sweden, Switzerland, Syria (?), Turkey, Ukraine.

36) *Canthophorus melanopterus* (Herrich-Schaeffer, 1835)

Cydnus melanopterus Herrich-Schaeffer, 1835: 55 (as new species).

Canthophorus melanopterus: Asanova 1964: 143 (new combination).

Iran records: Hoberlandt 1995: 189 [under the spelling *Camptoporus melanopterus*]; Linnauori 2007: 89; Lis 1999: 221–222, 2006: 141; Modarres Awal 1996b: 107 [under the combination *Sehirus melanopterus*], 1997: 71 [under the combinations “*Sehirus dubius* var. *melanoptera* H-S” et “*S. melanopterus* H-S”, treated as the names of two different taxa]; Safavi 1959: 31 [as *Sehirus dubius* var. *melanoptera* (Herrich-Schaeffer, 1835)]; Wagner 1961: 157 [under the combination *Sehirus melanopterus*].

Distribution in Iran: Ardabil, Azarbaijan, Fars, Gilan, Golestan, Kurdistan, Tehran provinces.

Distribution outside Iran: Albania, Algeria, Armenia, Austria, Azerbaijan, Bosnia Herzegovina, Bulgaria, Canary Islands, China (Beijing ?), Croatia, Cyprus, Czech Republic, Egypt, Georgia, Great Britain, Greece, Hungary, Iraq, Israel, Italy, Jordan, Kazakhstan, Libya, Macedonia, Malta, Moldavia, Morocco, Portugal, Romania, Russia (South European Territory), Serbia and Montenegro, Slovakia, Slovenia, Spain, Syria, Tadzhikistan, Tunisia, Turkey, Ukraine.

37) *Canthophorus wagneri* Asanova, 1964

Canthophorus wagneri Asanova, 1964: 139.

Material examined: West Azerbaijan province, Seroo (1612m), 2♀♀, August 2003.

New record for Iran.

Distribution outside Iran: Armenia, Azerbaijan, Georgia, Russia (South European Territory), Turkmenistan, Turkey.

Genus *Crocistethus* Fieber, 1860

38) *Crocistethus waltlianus* (Fieber, 1837)

Cydnus waltlianus Fieber, 1837: 352 (as new species).

Crocistethus waltlianus: Stichel 1961: 678 (new combination).

Material examined: West Azerbaijan province, Piranshahr (1465m), 2♂♂, August 2004. **New record for Iran.**

Distribution outside Iran: Algeria, Canary Islands, Cyprus, Egypt, France, Greece, Iraq, Israel, Italy, Jordan, Malta, Morocco, Portugal, Serbia, Spain, Syria, Tunisia, Turkey.

Genus *Exosehirus* Wagner, 1963

39) *Exosehirus marginatus* (Signoret, 1881b)

Adomerus marginatus Signoret, 1881b: 656.

Exosehirus marginatus: Linnauori 1984: 2 (new combination, in the framework of *E. sargon* description).

Iran records: Modarres Awal 1996a: 127 [under the combination *Sehirus marginatus*], 1997: 71.

Additional data: West Azarbaijan province, Maco (1730m), 1♀, September 2000.

Distribution in Iran: Khorasan and West Azerbaijan provinces.

Distribution outside Iran: Syria, Turkey (type locality: Taurus Mt.).

40) *Exosehirus validus* (Jakovlev, 1877)

Gnathoconus validus Jakovlev, 1877: 76 (as new species).

Exosehirus validus: Wagner 1963: 107 (new combination).

Iran records: Hoberlandt 2005: 189; Lis 1999: 223, 2006: 142; Modarres Awal 2008: 247.

Distribution in Iran: Khorasan and Lorestan provinces; Semnan province: Shahkuh, near Shahrud (type locality).

Distribution outside Iran: Afghanistan, Iraq, Kazakhstan, Kirgizia, Tadzhikistan, Turkmenistan, Uzbekistan.

Genus *Legnotus* Schiødte, 1848**41) *Legnotus limbosus* (Geoffroy, 1785)**

Cimex limbosus Geoffroy in Fourcroy, 1785: 217 (as new species).

Legnotus limbosus: Stichel 1935: 362 (new combination).

Iran records: Hoberlandt 1995: 189; Lis 2006: 143; Linnauori 2007: 89.

Distribution in Iran: Khorasan and Tehran provinces.

Distribution outside Iran: Albania, Algeria, Armenia, Austria, Azerbaidjan, Belgium, Bosnia Herzegovina, Bulgaria, Croatia, Czech Republic, Denmark, France, Germany, Great Britain, Greece, Hungary, Israel, Italy, Kazakhstan, Lithuania, Luxembourg, Macedonia, Moldavia, Morocco, Netherlands, Poland, Portugal, Romania, Russia, Serbia and Montenegro, Slovakia, Slovenia, Spain, Sweden, Switzerland, Syria, Turkey, Ukraine, Uzbekistan.

42) *Legnotus picipes* (Fallén, 1807)

Cydnus picipes Fallén, 1807: 54 (as new species).

Legnotus picipes: Horváth 1917: 372 (new combination).

Iran records: Hoberlandt 1995: 189; Lis 2006: 143.

Distribution in Iran: Khorasan and Tehran provinces.

Distribution outside Iran: Albania, Algeria, Armenia, Austria, Azerbaidjan, Belgium, Bosnia Herzegovina, Bulgaria, Byelorussia, Croatia, Czech Republic, Denmark, Estonia, Finland, France, Germany, Great Britain, Greece, Hungary, Israel, Italy, Kazakhstan, Latvia, Liechtenstein, Lithuania, Luxembourg, Macedonia, Moldavia, Morocco, Netherlands, Poland, Portugal, Romania, Russia, Serbia and Montenegro, Slovakia, Slovenia, Spain, Sweden, Switzerland, Syria, Tunisia, Turkey, Ukraine, Uzbekistan.

Genus *Ochetostethus* Fieber, 1860**43) *Ochetostethus heissi* Magnien, 2006**

Ochetostethus heissi Magnien, 2006: 513–516 (as new species).

Iran records: Magnien 2006: 513–514; Linnauori 2007: 89.

Distribution in Iran: Gilan and Zanjan provinces.

Distribution outside Iran: Cyprus, Israel, Greece, Turkey.

44) *Ochetostethus nanus* (Herrich-Schaeffer, 1834)

Cydnus nanus Herrich-Schaeffer, 1834: plate 24 (as new species).

Ochetostethus nanus: Mulsant and Rey 1866: 76 (new combination).

Iran records: Modarres Awal 1996a: 127, 1997: 71.

Distribution in Iran: Khorasan province.

Distribution outside Iran: Belgium, France, Portugal (?), Spain.

Comments: This is a doubtful record. The report by Modarres Awal (1996) probably refers to one of the other species of *Ochetostethus*. This would also be the case for all records from North Africa and Central or Eastern Europe (Lis 2006: 144).

45) *Ochetostethus opacus* (Scholtz, 1847)

Cydnus opacus Scholtz, 1847: 158 (as new species).

Ochetostethus opacus: Kerzhner 1976: 36 (new combination).

Iran records: Hoberlandt 1995: 191; Kerzhner 1976: 36 (without exact locality); Linnavuori 2007: 89; Lis 1999: 226, 2006: 144; Magnien 2006: 515 (without exact locality); Modarres Awal 1996b: 107, 1997: 71.

Distribution in Iran: Ardabil, Gilan and Tehran provinces.

Distribution outside Iran: Afghanistan, Albania, Armenia, Austria, Azerbaijan, Bosnia Herzegovina, Bulgaria, Canary Islands, China (North), Croatia, Cyprus, Czech Republic, Finland, Georgia, Germany, Greece, Hungary, Iraq, Israel, Italy, Kazakhstan, Kirgizia, Latvia, Lebanon, Liechtenstein, Lithuania, Macedonia, Moldavia, Norway, Poland, Romania, Russia, Serbia and Montenegro, Slovakia, Slovenia, Switzerland, Syria, Tadzhikistan, Turkey, Ukraine, Uzbekistan.

46) *Ochetostethus sahlbergi* Wagner, 1952

Ochetostethus sahlbergi Wagner, 1952: 42 (as new species).

Iran records: Hoberlandt 1995: 190.

Additional data: West Azerbaijan, Ourmieh (1370m), 1♂, 3♀, August 2003.

Distribution in Iran: Tehran and West Azerbaijan provinces.

Distribution outside Iran: Afghanistan (?), Egypt, Israel, Jordan, Lebanon, Syria, Turkey.

47) *Ochetostethus tarsalis* Mulsant & Rey, 1852

Cydnus tarsalis Mulsant & Rey, 1852: 79 (as new species).

Ochetostethus tarsalis: Lethierry & Severin 1893: 80 (new combination).

Iran records: Hoberlandt 1995: 190.

Distribution in Iran: Fars and Tehran provinces.

Distribution outside Iran: Algeria, Canary Islands, France, Italy, Libya, Morocco, Saudi Arabia, Spain, Tunisia.

Genus *Sehirus* Amyot & Serville, 1843**48) *Sehirus cypriacus* Dohrn, 1860**

Sehirus cypriacus Dohrn, 1860: 101 (as new species).

Material examined: West Azerbaijan province, Naghadeh (1364m), 1♀, June 2002.

New record for Iran.

Distribution outside Iran: Cyprus, Israel, Jordan, Syria, Turkey.

49) *Sehirus dissimilis* Horváth, 1919

Sehirus dissimilis Horváth, 1919: 267 (as new species).

Material examined: West Azerbaijan province, Oshnavieh (1454m), 1♂, 2♀♀, June 2002. **New record for Iran**.

Distribution outside Iran: Turkey.

50) *Sehirus luctuosus* Mulsant & Rey, 1866

?*Cimex niger spinipes* De Geer, 1773: 269 [unavailable non binominal name; the synonymy with *S. luctuosus* by Reuter (1888); the synonymy with *Cydnus aterrimus* (Forster, 1771) by Amyot, 1845 seems doubtful since Reuter (1888) did examine De Geer's types (Lis 2006 contra 1999)].

Sehirus luctuosus Mulsant and Rey 1866: 52 (new combination).

Material examined: East Azerbaijan province, Arasbaran (826m), 2♂♂, 4♀♀, July 2005. **New record for Iran**.

Distribution outside Iran: Albania, Algeria, Armenia, Austria, Azerbaijan, Belgium, Bosnia Herzegovina, Bulgaria, Byelorussia, Croatia, Czech Republic, Den-

mark, Estonia, Finland, France, Georgia, Germany, Great Britain, Greece, Hungary, Ireland, Italy, Kazakhstan, Kirgizia, Latvia, Lithuania, Luxembourg, Macedonia, Moldavia, Mongolia, Netherlands, Norway, Poland, Romania, Russia, Serbia and Montenegro, Slovakia, Slovenia, Spain, Sweden, Switzerland, Tadzhikistan, Turkey, Ukraine.

51) *Sehirus morio* (Linnaeus, 1761)

Cimex morio Linnaeus, 1761: 250 (as new species).

Sehirus morio: Amyot and Serville 1843: 96 (new combination).

Iran records: Modarres Awal 1987: 19, 1997: 71; Linnavuori 2007: 89.

Distribution in Iran: Ardabil and East Azerbaijan provinces.

Distribution outside Iran: Albania, Algeria, Armenia, Austria, Azerbaidjan, Belgium, Bosnia Herzegovina, Bulgaria, Byelorussia, Canary Islands, Croatia, Cyprus, Czech Republic, Denmark, France, Georgia, Germany, Greece, Hungary, Italy, Kazakhstan, Kirgizia, Latvia (?), Libya, Lithuania, Luxembourg, Macedonia, Malta, Moldavia, Morocco, Netherlands, Norway, Poland, Portugal, Romania, Russia, Slovakia, Spain, Sweden, Switzerland, Tadzhikistan, Tunisia, Turkey, Turkmenistan, Ukraine, Uzbekistan.

52) *Sehirus ovatus* (Herrich-Schaeffer, 1840)

Cydnus ovatus Herrich-Schaeffer, 1840: 96 (as new species).

Sehirus ovatus: Fieber 1861: 367 (new combination).

Material examined: Ardabil province, Germi (764m), 1♂, 3♀♀, September 2004.

New Record for Iran.

Distribution outside Iran: Armenia, Austria, Azerbaijan, Bulgaria, Croatia, Czech Republic, Georgia, Greece, Hungary, Israel, Italy, Kazakhstan, Moldavia, Romania, Russia (Central and South European Territory), Slovakia, Syria, Turkey, Ukraine.

53) *Sehirus parens* Mulsant & Rey, 1866

Sehirus parens Mulsant and Rey, 1866: 52 (as new species).

Material examined: Khorasan province, Birjand (1434m), 2♀♀, April 2003.

New Record for Iran.

Distribution outside Iran: Afghanistan, Azerbaijan, China (Xinjiang), Czech Republic, Hungary, Kazakhstan, Kirgizia, Romania, Russia (South European Territory, West and East Siberia), Slovakia, Syria, Tadzhikistan, Ukraine.

54) *Sebirus planiceps* Horváth, 1895

Sebirus planiceps Horváth, 1895: 128 (as new species; in key).

Material examined: Mazandaran province, Galogah (82m), 1♂, 1♀, October 2001.

New Record for Iran.

Distribution outside Iran: Caucasus: Georgia and Russia.

55) *Sebirus robustus* Horváth, 1895

Sebirus robustus Horváth, 1895: 129, 130 (as new species; in key).

Iran records: Hoberlandt 1995: 191; Lis 1999: 228, 2006: 146; Wagner 1968: 438.

Distribution in Iran: Gorgan and Mazandaran provinces.

Distribution outside Iran: Armenia, Azerbaijan, Georgia, Iraq, Turkey.

Genus *Tritomegas* Amyot and Serville, 1843

56) *Tritomegas bicolor* (Linnaeus, 1758)

Cimex bicolor Linnaeus, 1758: 446 (as new species).

Tritomegas bicolor: Amyot & Serville 1843: 98 (new combination).

Iran record: Linnauori 2007: 90.

Additional data: Ardabil province, Pars-Abad (110m), 3♀♀, September 2004.

Distribution in Iran: Ardabil and Gilan provinces.

Distribution outside Iran: Algeria, Armenia, Austria, Azerbaidjan, Belgium, Bosnia Herzegovina, Bulgaria, Byelorussia, China (Liaoning), Croatia, Czech Republic, Denmark, Estonia, Finland, France, Georgia, Germany, Great Britain, Greece, Hungary, Israel, Italy, Kazakhstan, Kirgizia, Latvia, Liechtenstein, Lithuania, Luxembourg, Macedonia, Moldavia, Mongolia, Morocco, Netherlands, Norway, Poland, Portugal, Romania, Russia, Serbia and Montenegro, Slovakia, Slovenia, Spain, Sweden, Switzerland, Syria, Turkey, Ukraine, Uzbekistan.

57) *Tritomegas delagrangei* (Puton, 1888)

Sehirus bicolor var. *delagrangei* Puton, 1888: 256 (as new var.).

Sehirus delagrangei: Horváth 1901: 471 (upgrading to species rank).

Tritomegas delagrangei: Hoberlandt 1956: 212 (new combination).

Material examined: East Azarbaijan province, Maragheh (1495m), 1♂, June 1999.

New record for Iran.

Distribution outside Iran: Azerbaidjan, Israel, Lebanon, Syria, Turkey.

58) *Tritomegas sexmaculatus* Rambur, 1839

Cydnus sexmaculatus Rambur, 1839: 110 (as new species).

Canthophorus (*Tritomegas*) *sexmaculatus*: Mulsant and Rey 1866: 55 (new combination).

Iran records: Hassanzadeh et al. 2009: 529; Linnauori 2007: 90; Lis 1999: 229, 2006: 147; Modarres Awal 1996a: 128, 1997: 71.

Distribution in Iran: Khorasan and Tehran provinces.

Distribution outside Iran: Albania, Armenia, Austria, Azerbaidjan, Bosnia Herzegovina, Bulgaria, Croatia, Czech Republic, France, Georgia, Germany, Greece, Hungary, Italy, Liechtenstein, Macedonia, Moldavia, Netherlands, Poland, Portugal, Romania, Russia (Central and South European Territory), Serbia and Montenegro, Slovakia, Slovenia, Spain, Switzerland, Turkey, Ukraine.

Discussion

Fifty-eight species of burrower bugs are reported from Iran in the present checklist: one Amnestinae, one Cephalocheinae, and fifty-six Cydninae (three Amaurocorini, two Cydnini, twenty-five Geotomini and twenty-six Sehirini), i.e. about 9% of the World fauna. Of those, fourteen are newly recorded from the country:

- (a) one Cephalocheinae Scaptocorini: *Stibaropus henkei* (Jakovlev, 1874);
- (b) four Cydninae Geotomini: *Byrsinus fassor* (Mulsant & Rey, 1866), *Byrsinus nigroscutellatus* (Montandon, 1900), *Byrsinus penicillatus* Wagner, 1964 and *Geotomus antennatus* Signoret, 1883;
- (c) and nine Cydninae Sehirini: *Canthophorus wagneri* Asanova, 1964, *Crocistethus waltlianus* (Fieber, 1837), *Sehirus cypriacus* Dohrn, 1860, *Sehirus dissimilis* Horváth, 1919, *Sehirus luctuosus* Mulsant & Rey, 1866, *Sehirus ovatus* (Herrich-Schaeffer, 1840), *Sehirus parens* Mulsant & Rey, 1866, *Sehirus planiceps* Horváth, 1895 and *Tritomegas delagrangei* (Puton, 1888).



Figure 1. Map of Iran with boundaries of provinces.

One species is removed of the Iranian faunal list: *Byrsinus minor* Wagner, 1964, an erroneous mention according to Lis (2006: 129). The presence in Iran of *Byrsinus discus* Jakovlev, 1906 needs confirmation.

The number of species probably will increase in the future because Iran is a large, poorly explored, country, incorporating various geographical regions (Fig. 1), climates, and vegetation zones. We are convinced that additional field work will provide numerous new and interesting records.

Acknowledgments

The authors are indebted to late I. M. Kerzhner (Zoological Institute, Russian Academy of Sciences, St-Petersburg, Russia), P. Kment (National Museum of Natural History, Prague, Czech Republic), D. Pluot-Sigwalt (Muséum national d'Histoire naturelle, Paris, France), D. Redei (Hungarian Natural History Museum, Budapest, Hungary), D. Rider (North Dakota State University, Fargo, United States of America), H. Sakenin (Ghaemshahr Islamic Azad University, Mazandaran, Iran), M. Tabari (Iran Rice Research Institute, Mazandaran, Iran) and M. Havaskary (Tehran Islamic Azad University, Iran) for invaluable helps in progress of the project. The research was supported by Shahre Rey Islamic Azad University, Iran. Dr T. Henry (United States National Museum, Washington D.C., United States of America) and two anonymous reviewers provided constructive criticism on the manuscript.

References

- Amyot CJB (1845) Entomologie Française. Rhynchotes. Hémiptères. Homoptères. Phthiroptères. Pseudoptères. Méthode mononymique. Annales de la Société entomologique de France 2 (3): 369–492.
- Amyot CJB, Serville JGA (1843) Histoire naturelle des insectes. Hémiptères. Roret, Paris, i-lxxvi + 675pp + 6pp.
- Asanova RB (1964) [Genus *Canthophorus* Muls. and Rey (Heteroptera, Cydnidae) in the fauna of the USSR]. Entomologicheskoe Obozrenie 43 (1): 138–144 [in Russian].
- Askari O, Pourabad RF, Khaganinia S (2009) Faunistic study of Heteroptera of Zanjanroud Region in Zanjan Province of Iran. Munis Entomology and Zoology 4 (2): 560–563.
- Baroughi H (1978) Etude de la faune entomologique (Tabriz environs), 2, order des heteropteres (sic). Journal of the Entomological Society of Iran 12: 1–33.
- Billberg GJ (1820) Enumeratio insectorum in Museo Gust. Joh. Billberg. Gedalianis, Stockholm, 138 pp.
- Brown ES (1966) An account of the fauna associated with *Eurygaster intergiceps* Put. and *Aelia* species (Hem., Pentatomoidea) in their overwintering areas in the Middle East. Entomologist's Monthly Magazine 102: 29–46.
- Brullé A (1839) Insectes. In: Webb PB, Berthelot S (Eds) Histoire naturelle des îles Canaries 2 (2). Mellier, Paris, 55–119.

- Costa A (1847) Cimicum regni Neapolitani centuria secunda decas sexta, septima, octava, nona et decima. Napoli: 1–41 (separatum).
- Dallas WS (1851) List of the specimens of hemipterous insects in the collection of British Museum. Part I. Taylor, London, 368 pp.
- De Geer C (1773) Mémoires pour servir à l'histoire des Insectes 3. Hasselberg, Stockholm, i–viii, + 696 pp.
- Distant WL (1899a) Rhynchotal notes - Heteroptera: Plataspinae, Thyreocorinae and Cydninae. Annals and Magazine of Natural History (7) 4: 213–227.
- Distant WL (1899b). Correction. Annals and Magazine of Natural History (7) 4: 442.
- Distant WL (1901) Contributions to the knowledge of the Rhynchota. Transactions of the Entomological Society of London 1901: 581–592.
- Distant WL (1918) The fauna of British India, including Ceylan and Burma. *Rhynchota* vol. 7. (Homoptera: appendix; Heteroptera: addenda). Taylor and Francis, London, i–vii + 210pp.
- Dohrn A (1860) Hemipterologische Miscellaneen. Stettiner Entomologische Zeitung 21: 99–109.
- Fabricius JC (1794) Entomologia systematica emendata et aucta secundum classes, ordines, genera, species adjectis synonymis, locis, observationibus, descriptionibus. 4. Proft, Hafniae, i–vi + 472pp.
- Fabricius JC (1803) Systema Rhyngotorum secundum ordines, genera, species adjectis synonymis, locis, observationibus, descriptionibus. Reichard, Brunsvigae, i–vi + 314 pp.
- Fallén CF (1807) Monographia Cimicum Sueciae. Proft, Hafniae, 123pp.
- Fieber FX (1837) Beiträge zur Kenntnis der Schnabelkerfe (Rhynchota). In: Weitenweber WR (Ed). Beiträge zur gesammten Natur- und Heilwissenschaft 1: 337–355.
- Fieber FX (1860) Die europäischen Hemiptera. Halbflügler (Rhynchota Heteroptera). Gerold's Sohn, Wien, i–iv + 112 pp.
- Fieber FX (1861) Die europäischen Hemiptera. Halbflügler (Rhynchota Heteroptera). Gerold's Sohn, Wien, 331 pp.
- Forster JR (1771) Novae species insectorum. Centuria I. London, i–viii + 100 pp.
- Fourcroy AF (1785) Entomologia Parisiensis, sive catalogus insectorum quae in agro Parisiensi reperiuntur. Via et Aedibus Serpentineis, Paris, i–viii + 544 pp.
- Froeschner RC (1960) Cydnidae of the Western Hemisphere. Proceedings of the United States National Museum 111 (3430): 337–680.
- Froeschner RC (1988). Family Cydnidae Billberg, 1820 pp. 119–129. In: Henry TJ and Froeschner RC (Eds) Catalog of the Heteroptera, or True Bugs, of Canada and the Continental United States. E.J. Brill, Leiden, i–xix + 958 pp.
- Grazia J, Schuh RT, Wheeler WC (2008) Phylogenetic relationships of family groups in Pentatomidea based on morphology and DNA sequences (Insecta: Heteroptera). Cladistics 24: 1–45.
- Hart CA (1919) The Pentatomidae of Illinois, with keys of the Nearctic genera. Bulletin of the Illinois State Natural History Survey 13: i–ii, 157–223.
- Hassazadeh M, Pourabad RF, Shayesteh N (2009) An investigation on some Heteroptera in Marand region (Iran). Munis Entomology and Zoology 4 (1): 19–24.
- Hassazadeh M, Pourabad RF, Gharaat MA, Beykpor AR (2009) A study of the Heteroptera fauna of Shend Abad Region and environ (Iran). Munis Entomology and Zoology 4 (2): 527–530.

- Herrich-Schaeffer GA (1834) Fauna Insectorum Germanicae initia oder Deutschlands Insekten. 126. Regensburg, 24 plates.
- Herrich-Schaeffer GA (1835) Nomenclator entomologicus. Verzeichniss der europäischen Insecten; zur Erleichterung des Tauschverkehrs mit Preisen versehen. Erstes Heft. Lepidoptera und Hemiptera, leztere synoptisch bearbeitet und mit vollständiger Synonymie. Pustet, Regensburg, i-iv + 116 pp.
- Herrich-Schaeffer GA (1840) Die Wanzenartigen Insekten getreu nach der Natur abgebildet und beschrieben. 5. Nürnberg, 108 pp.
- Hoberlandt L (1955) Hemiptera – Heteroptera from Iran. I. Acta Entomologica Musei Nationalis Pragae 29 (1954): 121–148.
- Hoberlandt L (1956) Results of the zoological scientific expedition of the National Museum in Praha to Turkey. 18. Hemiptera IV. Terrestrial Hemiptera-Heteroptera of Turkey. Acta Entomologica Musei Nationalis Pragae, Suppl. 3 [1955]: 1–264.
- Hoberlandt L (1995) Results of the Czechoslovak-Iranian entomological expeditions to Iran 1970, 1973 and 1977; Heteroptera: Acanthosomatidae, Cydnidae, Scutelleridae, Pentatomidae. Acta Entomologica Musei Nationalis Pragae 44: 181–293.
- Horváth G (1882) Hemiptera nova vel minus cognitae. Természetrajzi Füzetek 5: 217–225.
- Horváth G (1889) Beitrag zur Hemipteren-fauna von Turkmenien. Wiener Entomologische Zeitung 8: 169–174.
- Horváth G (1895) Révision des *Sehirus* noirs. Revue d'Entomologie 14: 128–130.
- Horváth G (1901) Hémiptères du voyage de M. Martinez Escalera dans l'Asie-Mineure (1898). Természetrajzi Füzetek 24: 469–471.
- Horváth G (1917) Hemiptera palaearctica nova vel minus cognita. I. Annales Historico-Naturales Musei Nationalis Hungarici 15: 365–381.
- Horváth G (1918) Ad cognitionem faunae hemipterorum balcanicae. Annales Historico-Naturales Musei Nationalis Hungarici 16: 321–340.
- Horváth G (1919) Analecta ad cognitionem Cydnidarum. Annales Historico-Naturales Musei Nationalis Hungarici 17: 205–273.
- Jakovlev BE (1874a) [Contributions to the entomological fauna of the european Russia. IV]. Trudy Russkago Entomologicheskago Obshchestva 8: 46–82 [in Russian].
- Jakovlev BE (1874b) [Hemiptera Heteroptera of the Astrakhan' Region]. Bulletin de la Société Impériale des Naturalistes de Moscou 48 (2): 218–277 [in Russian and German].
- Jakovlev BE (1877) Bugs (Hemiptera Heteroptera) of northern Persia. Trudy Russkago Entomologicheskago Obshchestva 10: 67–98 [in Russian and German].
- Jakovlev BE (1879) [Bugs (Hemiptera Heteroptera) of the Caucasian region. I]. Trudy Russkago Entomologicheskago Obshchestva 12: 3–176 [in Russian and German].
- Jakovlev BE (1885) Hemiptera-Heteroptera aus Achal-Tekke. Horae Societatis Entomologicae Rossicae 19: 98–129 [in Russian and German].
- Jakovlev BE (1906) [Review of species of the genus *Byrsinus* Fieb. (Hemiptera-Heteroptera, Pentatomidae) of the Russian-Asiatic fauna]. Revue Russe d'Entomologie 6: 52 [in Russian (description of *B. discus* n. sp. and title) and French (key of species)].
- Kerzhner IM (1972) [New and little-known Heteroptera from Mongolia and from adjacent regions of the USSR. I]. Insects of Mongolia 1: 349–379 [in Russian].

- Kerzhner IM (1976) [New and little known Heteroptera from Mongolia and from adjacent regions of the USSR. III]. Insects of Mongolia 4: 30–86 [in Russian].
- Kerzhner IM (2003) Type specimens of Coreoidea and Pentatomoidea described by F. A. Kolenati (Heteroptera). Zoosystematica Rossica 12: 93–98.
- Kerzhner IM, Jaczewski TL (1964) Order Hemiptera (Heteroptera). Keys to the insects of the European USSR, 1, 655–845 [in Russian].
- Kiritschenko AN (1966) [Hemiptera-Heteroptera collected by D.M. Steinberg in Iran in 1955]. Entomologicheskoe Obozrenie 45 (4): 798–805 [in Russian].
- Klug JCF (1845) Symbolae physicae, seu icones et descriptiones insectorum, quae ex itinere Africam borealem et Asiam occidentalem F. G. Hemprich and C. H. Ehrenberg studio novae aut illustratae redierunt 5. Mittler, Berlin: a-l.
- Kolenati FA (1846) Meletemata Entomologica IV. Hemiptera Caucasi. Pentatomidae monographic dispositae. Academia Scientiarum, Petropoli, 72pp.
- Lethierry L, Severin G (1893) Catalogue général des Hémiptères. 1. Hétéroptères Pentatomidae. Hayez, Bruxelles, i-xi + 286 pp.
- Linnaeus C (1758) Systema Naturae per regna tria naturae, secundum classes, ordines, genera, species, cum characteribus, differentiis, synonymis, locis. Editio decima, reformata, Holmiae, i-iv + 824 pp.
- Linnaeus C (1761) Fauna Svecica sistens animalia Sveciae regni: Mammalia, Aves, Amphibia, Pisces, Insecta, Vermes, distributa per classes et ordines, genera et species. Salvii, Stockholmiae, i-xlvi + 578 pp.
- Lindberg H (1938) Einige Heteropterenfunde aus Nord-Persien. Notulae Entomologicae 18: 88–91.
- Linnavouri RE (1984) New species of Hemiptera Heteroptera from Iraq and the adjacent countries. Acta Entomologica Fennica 44: 1–59.
- Linnavuori RE (2004) Heteroptera of the Hormozgan province in Iran. III. Cimicomorpha (Anthocoridae, Cantacaderidae, Tingidae), Pentatomorpha (*sic*). Acta Universitatis Carolina, Biologica 48: 273–286.
- Linnavuori RE (2007) Studies on the Piesmatidae, Berytidae, Pyrrhocoridae, Stenocephalidae, Coreidae, Rhopalidae, Alydidae, Cydnidae and Plataspidae (Heteroptera) of Gilan and the adjacent provinces in Northern Iran. Acta Entomologica Musei Nationalis Praegae 47: 77–91.
- Lis JA (1993a) A review of the tribe Amaurocorini E. Wagner with remarks on its systematic position within the family (Heteroptera: Cydnidae). Annals of the Upper Silesian Museum Entomolgy 4: 59–78.
- Lis JA (1993b) On *Byrsinus* Fieber and *Microporus* Uhler, two allied cydnid genera (Heteroptera: Cydnidae). Annals of the Upper Silesian Museum Entomolgy 4: 79–98.
- Lis JA (1994) A revision of Oriental burrower bugs (Heteroptera: Cydnidae). Upper Silesian Museum, Bytom, 349 pp.
- Lis JA (1996) Taxonomy and phylogeny of the genus *Cydnus* F., 1803 (Hemiptera: Heteroptera: Cydnidae). Annales de la Société Entomologique de France 32 (4): 403–409.
- Lis JA (1997) Some remarks on the Palaearctic Cydnidae (Hemiptera: Heteroptera). Acta Entomologica Silesiana 4 [1996]: 3–7.

- Lis JA (1998) *Amnestus raunoi* n. sp. from Iran, the first Old World representative of the New World subfamily Amnestinae (Hemiptera: Heteroptera: Cydnidae). *Entomologica Fennica* 9: 5–7.
- Lis JA (1999) Burrower bugs of the Old World - a catalogue (Hemiptera: Heteroptera: Cydnidae). *Genus* 10 (2): 165–249.
- Lis JA (2000). A revision of the burrower-bug genus *Macroscytus* Fieber, 1860 (Hemiptera: Heteroptera: Cydnidae). *Genus* 11 (3): 359–509.
- Lis JA (2003) A redescription of *Byrsinus nigroscutellatus* (Montandon) comb. n. (Hemiptera: Heteroptera: Cydnidae), with remarks on the synonymy of *Byrsinocoris* Montandon, 1900 and *Byrsinus* Fieber, 1860. *Polskie Pismo Entomologiczne* 71: 43–71.
- Lis JA (2006) Family Cydnidae Billberg, 1820 – burrowing bugs (burrower bugs). In: Aukema B, Rieger E (Eds) Catalogue of the Heteroptera of the Palearctic Region. Volume 5, Pentatomomorpha II. Netherlands Entomological Society, Ponsen and Looijen, Wageningen, pp. 119–147.
- Lis JA, Becker M, Schaefer CW (2000) Burrower Bugs (Cydnidae). In: Schaefer CW, Panizzi, AR (Eds) Heteroptera of Economic Importance. CRC Press, Boca Raton, pp. 405–419.
- Magnien P (2006) A new species from the Near East for the genus *Ochetostethus* Fieber, 1860 (Heteroptera, Cydnidae). In: Rabitsch W (Ed) Hug the Bug – For the love of true bugs. *Festschrift zum 70. Geburtstag von Ernst Heiss. Denesia* 19 [Zugleich Kataloge der Oberösterreichischen Landesmuseen, N.S. 50]: 513–516.
- Modarres Awal M (1987) Collecting and determining of the fauna of Heteroptera in Ardabil area and research Station of University of Tabriz in Moghan. *Journal of the Entomological Society of Iran* 9 (1–2): 15–28.
- Modarres Awal M (1996a) Studies on some Pentatomorpha (Heteroptera) fauna in north of Khorasan province. *Journal of Agriculture and Science Technology, Mashhad University* 9 (2): 121–144.
- Modarres Awal M (1996b) Studies on some Cimicomorpha and Pentatomorpha (Heteroptera) fauna in Ardabil province. *Journal of Agriculture and Science Technology, Mashhad University* 10 (1): 102–112.
- Modarres Awal M (1997) List of agricultural pests and their natural enemies in Iran. Ferdowsi University press, 429 pp.
- Modarres Awal M (2008) Contribution to Heteroptera fauna of Khorasan Razavi province of Iran. *Turkiye Entomoloji Dergisi* 32 (4): 243–253.
- Montandon AL (1900) Hemiptera-Heteroptera. Espèces nouvelles de la faune paléarctique. *Buletinul Societății de Științe din Bucurescii* 9: 155–160.
- Mulsant E, Rey C (1852) Description de quelques Hémiptères Hétéroptères nouveaux ou peu connus. *Annles de la Société. Linnéenne de Lyon* 1852: 76–141.
- Mulsant E, Rey C (1866) Histoire naturelle des Punaises de France, Pentatomides. Savy and Deyrolle, Paris, 372pp.
- Pluot-Sigwalt D, Lis JA (2008) Morphology of the spermatheca in the Cydnidae (Hemiptera: Heteroptera): Bearing of its diversity on classification and phylogeny. *European Journal of Entomology* 105: 279–312.
- Putshkov VG (1965) [*Shield-bugs of Middle Asia* (Hemiptera, Pentatomoidea)]. Illim, Frunze, 331 pp.

- Puton A (1886) Enumération des Hémiptères récoltés en Tunisie en 1883 et 1884 par MM. Valéry Mayet et Maurice Sédillot, suivie de la description des espèces nouvelles. Exploration scientifique de la Tunisie. Imprimerie Nationale, Paris, 24 pp.
- Puton A (1888) Un genre nouveau d'Hémiptères et notes diverses. Revue d'Entomologie, 7: 255–257.
- Puton A, Lethierry L (1887) Hémiptères nouveaux de l'Algérie. Revue d'Entomologie 6: 298–311.
- Rambur P (1839) Faune entomologique de l'Andalousie. Arthus Bertrand, Paris, 336 pp.
- Reuter OM (1888) Revisio synonymica Heteropterorum Palaearcticorum quae descripserunt auctores vetustiores (Linnaeus 1758 – Latreille 1806). Acta Societatis Scientiarum Fennicae 15: 241–315, 443–812.
- Reuter OM (1902) Miscellanea Hemipterologica. Öfversigt af Finska Vetenskaps-Societatens Förhandlingar 44: 141–188.
- Sadeghi R, Pourabad RF, Kazemi H, Hassanzadeh M (2009) Introduction of some Heteroptera of Qurigol (Iran). Munis Entomology and Zoology 4 (2):498–500.
- Safavi M (1959) Monographie des Hémiptères-Hétéroptères de l'Iran. Entomologie et Phytopathologie appliquées 18: 31–40.
- Schiødte JC (1848) Om en gruppe af gravende Cimicies. Naturhistorisk Tidsskrift (2) 2 [1847]: 447–464.
- Scholtz H (1847) Prodromus zu einer Rhynchten-Fauna von Schlesien. Übersicht der Arbeiten und Veränderungen der Schlesischen Gesellschaft für Vaterländische Kultur [1846]: 104–164.
- Scopoli JA (1763) Entomologia carniolica exhibens Insecta Carnioliae indigena et distributa in ordines, genera, species, varietatis. Methodo Linnaeana. Trattner, Vindobonae, 420pp
- Seidenstücker G (1957) Heteropteren aus Iran 1954. I. Teil Hemiptera-Heteroptera (ohne Fam. Miridae). Jahreshefte des Vereins für Vaterländische Naturkunde in Würtemberg, 112: 66–73.
- Seidenstücker G (1958) Heteropteren aus Iran 1956. I. Hemiptera-Heteroptera (ohne Familie Miridae). Ergebnisse der Entomologischen Reisen Willi Richter, Stuttgart, in Iran, 1954 und 1956. Nr 17. Stuttgarter Beiträge zur Naturkunde aus dem Staatlichen Museum für Naturkunde in Stuttgart 11: 1–5.
- Signoret V (1881a). Révision du groupe des Cydnides de la famille des Pentatomides. 1^{ère} partie. Annales de la Société Entomologique de France (6) 1: 25–52.
- Signoret V (1881b) Revue des Cydnides contenus dans la collection du Musée Civique d'Histoire Naturelle de Gênes. Annali del Museo Civico di Storia Naturale di Genova 16: 621–657.
- Signoret V (1882) Révision du Groupe des Cydnides de la Famille des Pentatomides. 6^{ème} partie (1). Annales de la Société Entomologique de France (6) 2: 145–168.
- Signoret V (1883) Révision du Groupe des Cydnides de la Famille des Pentatomides. 9^{ème} partie. Annales de la Société entomologique de France (6) 3: 33–60.
- Stål C (1865) Hemiptera Africana 1. Norstedt, Homiae, i-iv 1+ 256 pp.
- Stichel W (1935) Illustrierte Bestimmungstabellen der Deutschen Wanzen. 12 (Hemiptera – Heteroptera). Stichel, Berlin-Hermsdorf: 331–362.
- Stichel W (1961) Illustrierte Bestimmungstabellen der Wanzen. II. Europa (Hemiptera -Heteroptera Europae), 4. 22. Heft. Pentatomomorpha, Cydnidae (2), Plataspidae. Stichel, Berlin-Hermsdorf, 667–700.

- Uhler H (1872) Notices of the Hemiptera of the western territories of the United States, chiefly from the surveys of Dr F.V. Hayden. In: Hayden FV (Ed). Preliminary report of the United States Geological Survey of Montana and portions of adjacent territories. 5. Government Printing Office, Washington D.C., 392–423.
- Wagner E (1951) *Aethus macropthalmus* nov. spec., eine neue Cydnidenart aus Aegypten (Hemiptera Heteroptera). Bulletin de la Société Fouad Ier d'Entomologie 35: 163–165.
- Wagner E (1952) Beitrag zur Systematik der Cydnidae (Hem. Heteropt.). Acta Entomologica Musei Nationalis Pragae 27 [1951]: 35–46.
- Wagner E (1953) Beitrag zur Systematik der Gattung *Geotomus* Muls.-Rey (Hemiptera Heteroptera: Cydnidae). Bulletin de la Société Fouad Ier d'Entomologie 37: 459–476.
- Wagner E (1954) Neue Heteropteren von den Kanarischen Inseln. Commentationes Biologicae 14 (2): 1–28.
- Wagner E (1961) Beitrag zur Heteropteren-Fauna von Iran. Ergebnisse der Österreichischen Iran-expeditionen 1949/1950 und 1956. Anzeiger der österreichische Akademie für Wissenschaften, Mathematisch-Naturwissenschaftliche Klass 10: 156–165.
- Wagner E (1963) Untersuchungen über den taxonomischen Wert des Baues der Gentalien bei den Cydnidae (Hem. Het.). Acta Entomologica Musei Nationalis Pragae 35: 73–115.
- Wagner E (1964) Die Gattung *Byrsinus* Fieber, 1860 und einige verwandte Gattungen (Heteroptera, Cydnidae). Reichenbachia 3 (3): 75–96.
- Wagner E (1965) Nachtrag zum *Byrsinus*-Komplex (Hem. Het., Cydnidae). Mitteilungen der Deutschen Entomologischen Gesellschaft 24 (3): 48–51.
- Wagner E (1968) Contribution à la faune de l'Iran. 7. Hémiptères Hétéroptères (*pro parte*). Annales de la Société Entomologique de France 4 (2): 437–453.
- Walker F (1867) Catalogue of the specimens of Hemiptera Heteroptera in the collection of the British Museum, vol. 1. London, 240 pp.