

Taxonomic study of the genus *Neurotettix* Matsumura (Hemiptera, Cicadellidae, Deltocephalinae) with a description of a new species from China

Renhuai Dai[†], Jichun Xing[‡], Zizhong Li[§]

Institute of Entomology, Guizhou University; The Provincial Key Laboratory for Agricultural Pest Management of Mountainous Region, Guiyang, Guizhou, P.R. China, 550025

[†] urn:lsid:zoobank.org:author:BCA15BC1-8DD9-4E25-99DE-EC94636C117E

[‡] urn:lsid:zoobank.org:author:1CD0E0C8-D09A-41EE-B9C0-478889BED58A

[§] urn:lsid:zoobank.org:author:9BA8A6EF-F7C3-41F8-AD7D-485FB93859F2

Corresponding author: Renhuai Dai (dairenhui@yahoo.com.cn)

Academic editor: Michael Wilson | Received 1 November 2010 | Accepted 2 February 2011 | Published 23 February 2011

urn:lsid:zoobank.org:pub:7EC687EC-72ED-4E54-8413-D0F4F67B30A3

Citation: Dai R, Xing J, Li Z (2011) Taxonomic study of the genus *Neurotettix* Matsumura (Hemiptera, Cicadellidae, Deltocephalinae) with a description of a new species from China. ZooKeys 82: 67–72. doi: 10.3897/zookeys.82.1158

Abstract

This paper treats all four known species of the genus *Neurotettix* Matsumura, 1914 (Hemiptera, Cicadellidae, Deltocephalinae), including one new species: *Neurotettix truncatus* **sp. n.** from China. A key is given to distinguish all species of the genus, and illustrations of genitalia are provided.

Keywords

Hemiptera, morphology, taxonomy, distribution

Introduction

The leafhopper genus *Neurotettix*, belonging to tribe Athysanini of subfamily Deltocephalinae (Hemiptera: Cicadellidae), was established by Matsumura (1914) for a single species, *Neurotettix horishanus* Matsumura from Taiwan. Later, Ishihara (1963)

redescribed and illustrated this genus. Cai and Shen (1999) described a second species, *Neurotettix bifurcatus*, from China, and recently, Shen et Dai (2002) described a third species, *Neurotettix flangenus*, from China. All the species of this genus are described from China.

Here we described and illustrated a new species from Guizhou Province, China. The type specimens of new species are deposited in the Institute of Entomology, Guizhou University, Guiyang, China (GUGC). The genus *Neurotettix* now contains four species. A key is given to separate all species.

Taxonomy

Neurotettix Matsumura

Neurotettix Matsumura 1914: 192; Esaki and Ito 1954: 128; Ishihara 1963: 223.

Type species: *Neurotettix horishanus* Matsumura, 1914

Description. Medium sized leafhoppers, body elongate, vertex triangular and produced. Head including eyes clearly narrower than pronotum. Eyes black, large. Ocelli located on anterior margin of vertex, near eyes. Face with dark, transverse streaks. Frontoclypeus long and narrow. Pronotum longer than vertex, its length of lateral carina 1/3 basal width of eye, anterior margin roundly protruded and posterior margin concave. Scutellum triangular, slightly shorter than pronotum, with transverse suture curved and depressed. Forewings with four apical cells and three subapical cells, apical cells short, anteapical cells with reticulate veins, clavus irregularly reticulated with many extra veins, appendix small.

Male pygofer side with about 10 stout setae, its ventro-posterior margin with a long appendage. Valve triangular. Subgenital plate with many setae in lateral margin. Aedeagus asymmetrical or symmetrical, base robust, aedeagal shaft slender or robust, with or without processes, gonopore apical or subapical. Connective nearly X-shaped. Style slender, elongate.

Diagnosis. *Neurotettix* may be distinguished from other Athysanini by the following combination of features: forewings with apical cells short, anteapical cells with reticulate veins, clavus irregularly reticulated with many extra veins; connective nearly X-shaped.

Distribution. Oriental Region and Palearctic Region.

Discussion. The aedeagus is asymmetrical in the type species in original description (Ishihara 1963). But, the aedeagus is symmetrical in the other two species (*Neurotettix bifurcatus* and *Neurotettix flangenus*) and the new species *Neurotettix truncatus* sp. n. According to the external characters and other male genitalia features, we confirmed the other two species and the new species belong to the genus *Neurotettix*.

Key to species (male) of *Neurotettix*

- | | | |
|---|---|--|
| 1 | Aedeagal shaft short and robust (Figs 5, 6)..... | <i>N. flangenus</i> Shen & Dai |
| – | Aedeagal shaft long and slender | 2 |
| 2 | Aedeagal with two paired apical processes (Figs 3, 4) | <i>N. bifurcatus</i> Cai & Shen |
| – | Aedeagal without paired apical process..... | 3 |
| 3 | Gonopore subapical, about one-third from apex (Figs 1, 2) | <i>N. horishanus</i> Matsumura |
| – | Gonopore at apex (Figs 10, 11)..... | <i>N. truncatus</i> Dai, Xing & Li, sp. n. |

– Aedeagal shaft long and slender **2**

2 Aedeagal with two paired apical processes (Figs 3, 4)

.....*N. bifurcatus* Cai & Shen

– Aedeagal without paired apical process.....**3**

3 Gonopore subapical, about one-third from apex (Figs 1, 2)

..... *N. horishanus* Matsumura

– Gonopore at apex (Figs 10, 11).....*N. truncatus* Dai, Xing & Li, sp. n.

Neurotettix horishanus Matsumura

Figs 1–2

Neurotettix horishanus Matsumura 1914: 193; Ishihara 1963: 224.

Distribution: China (Taiwan).

Neurotettix bifurcatus Cai & Shen

Figs 3–4

Neurotettix bifurcatus Cai and Shen 1999: 41–42.

Distribution: China (Henan).

Neurotettix flangenus Shen & Dai

Figs 5–6

Neurotettix flangenus Shen and Dai 2002: 89–91.


Distribution: China (Hunan).

Neurotettix truncatus Dai, Xing & Li, sp. n.

urn:lsid:zoobank.org:act:41C31A94-D0A4-4E56-96CC-BBC3D49AB654

Figs 7–19


Description. Body yellow-brown, vertex with black spots along anterior margin, and with two irregular orange markings behind the spots, eyes dark-brown (Fig. 16). Face



Figures 1–15. *Neurotettix horishanus* Matsumura **1** Aedeagus, dorsal view **2** Aedeagus, lateral view (after Ishihara 1963) *Neurotettix bifurcatus* Cai & Shen **3** Aedeagus, ventral view **4** Aedeagus, lateral view (after Cai and Shen 1999) *Neurotettix flangenus* Shen & Dai **5** Aedeagus, ventral view **6** Aedeagus, lateral view (after Shen & Dai 2002) *Neurotettix truncatus* sp. n. **7** Male pygofer side, lateral view **8** Valve, ventral view **9** Subgenital plate, ventral view **10** Aedeagus, ventral view **11** Aedeagus, lateral view **12** Connective **13** Style, dorsal view **14** female 1st valvula, lateral view **15** female 2nd valvula, lateral view.

brown, transverse streaks and a longitudinal band yellow, anteclypeus yellow with apex dark-brown (Fig.18). Pronotum and scutellum yellowish-brown. Forewings yellowish-brown, with irregular fuscous markings, viens yellowish-white, apical part of forewings pale brown (Fig. 17). Female abdominal genital segment pale brown (Fig. 19).

External features as in generic description.



Figures 16–19. *Neurotettix truncatus* sp. n. **16** ♂, dorsal view **17** ♂, lateral view **18** ♂, face **19** ♀ abdominal sternum VII, ventral view.

Male genitalia. Pygofer side short, with eleven stout setae along dorso-caudal margin, its ventro-posterior margin with a long appendage (Fig. 7). Valve triangle (Fig. 8). Subgenital plate short and broad, with many setae in lateral margin, distally truncate, with 2 to 3 irregular rows of setae from lateral margin to middle of subgenital plate (Fig. 9). Aedeagus symmetrical, base robust, aedeagal shaft slender without processes, gonopore apical (Figs 10, 11). Connective nearly X-shaped, its arms longer than stem (Fig. 12). Style slender, elongate, with apex of apophysis curved laterally (Fig. 13).

Female seventh sternum concaved medially on posterior margin. First valvula of ovipositor sculpture irregularly (Fig. 14), second valvula with teeth, tapered toward apex in lateral view (Fig. 15).

Measurement. Length (including tegmen): ♂4.8–5.3mm, ♀4.9–5.3mm.

Type Material. Holotype ♂, China: Guizhou Prov., Kuankuoshui, 16 August 2010, coll. Jichun Xing (GUGC). Paratypes: 1♂2♀♀, Guizhou Prov., Kuankuoshui, 12 August 2010, coll. Renhuai Dai (GUGC), 1♂1♀, China: Hubei Prov., Lichuan city, Pingba, 1 August 2010, coll. Junqiang Ni (GUGC).

Host. Bamboo.

Remarks. This species is similar to *Neurotettix horishanus* Matsumura, but can be distinguished from the latter by the symmetrical aedeagus and gonopore at apex, subgenital plate distally truncate and style apex curved.

Etymology. The new species name is derived from the Latin words “*truncatus*”, indicating that the subgenital plate distally truncate.

Acknowledgements

The project was supported by the Natural Science Foundation of China (31000952) and key project for social development of Guizhou (SZ[2009]3041).

References

- Cai P, Shen XC (1999) Six new species of Cicadellidae from Mt. Dabie in Henan (Homoptera: Cicadellidae). In: Shen XC, Pei HC (Eds) The Fauna and Taxonomy of Insects in Henan. Vol. 4, Insects of the Mountains Funiu and Dabie Regions. Beijing, China Agricultural Sciencetech Press, 36–45.
- Ishihara T (1963) Some genera, especially “*Eutettix*” of Japan and Formosa (Homoptera: Deltocephalinae). Transactions of the Shikoku Entomological Society 7(4): 119–224.
- Matsumura S (1914) Die Jassinen und einige neue Acocephalinen Japans. Journal of the College of Agriculture, Tohoku Imperial University of Tokyo, Saporro 5: 165–240.
- Shen L, Dai W (2002) A new species of the genus *Neurotettix* Matsumura (Homoptera: Cicadellidae: Deltocephalinae) from China. Entomotaxonomia 24(2): 89–92.
- Esaki T, Ito S (1954) A tentative catalogue of Jassoidea of Japan and her adjacent territories. Japan society for the promotion of science, Ueno Park, Tokyo, 315pp.