

# New Coleoptera records from New Brunswick, Canada: Elateridae

Reginald P. Webster<sup>1</sup>, Jon D. Sweeney<sup>1</sup>, Ian DeMerchant<sup>1</sup>

<sup>1</sup> Natural Resources Canada, Canadian Forest Service - Atlantic Forestry Centre, 1350 Regent St., P.O. Box 4000, Fredericton, NB, Canada E3B 5P7

Corresponding author: Reginald P. Webster (reginaldwebster@rogers.com)

Academic editor: R. Anderson | Received 28 December 2011 | Accepted 27 February 2012 | Published 4 April 2012

**Citation:** Webster RP, Sweeney JD, DeMerchant I (2012) New Coleoptera records from New Brunswick, Canada: Elateridae. In: Anderson R, Klimaszewski J (Eds) Biodiversity and Ecology of the Coleoptera of New Brunswick, Canada. ZooKeys 179: 93–113. doi: 10.3897/zookeys.179.2603

## Abstract

Twenty-two species of Elateridae are newly reported for New Brunswick, Canada. *Negastrius exiguus* (Randal) is removed from the faunal list and *Agriotes pubescens* Melsheimer is re-instated as a member of the New Brunswick fauna. *Agriotes pubescens* Melsheimer, *Dalopius brevicornis* W. J. Brown, *Danosoma obtectum* (Say) and *Megapenthes solitarius* Fall are newly reported for the Maritime provinces. Collection data, bionomic data, and distribution maps are presented for all these species.

## Keywords

Elateridae, new records, Canada, New Brunswick

## Introduction

The Elateridae (click beetles) is a species-rich family of beetles with about 965 named species in North America (Johnson 2002) and 369 species and subspecies from Canada and Alaska (Bousquet 1991). Although some groups are fairly well known taxonomically, genera such as *Ampedus* and *Dalopius* are in need of revisionary study and include a number of undescribed species (Johnson 2002). Larvae of some species of Elateridae are rhizophagous and are important agricultural pests; larvae of other species are predaceous, often living in soil, subcortical habitats, or rotten logs (Johnson 2002). However, little is known about the biology of most species.

The Elateridae of the Maritime provinces (New Brunswick, Nova Scotia, Prince Edward Island) was reviewed by Majka and Johnson (2008). They provided a detailed historical overview the collection of the Elateridae and other families of beetles in the Maritime provinces and a taxonomic review of the genus *Ctenicera*, which was in need of taxonomic review and generic re-assignment. Ninety-eight species were reported for New Brunswick, 13 as new provincial records; *Agriotes pubescens* Melsheimer, *Athous campyloides* Newman, and *Cardiophorus cardisce* (Say) were removed from the faunal list of the province by Majka and Johnson (2008). Later, Douglas (2011) newly reported *Pseudanstirus nigricollis* (Bland) and the adventive *Hemicrepidius niger* (Linnaeus) from New Brunswick. *Hemicrepidius niger* was also reported from Ontario and these represented the first records of this Eurasian species from North America. Here, we newly report 22 elaterid species from New Brunswick.

## Methods and conventions

The following records are based on specimens collected during a general survey by the first author to document the Coleoptera fauna of New Brunswick and from by-catch samples obtained during a study to develop a general attractant for the detection of invasive species of Cerambycidae. Additional provincial records were obtained from specimens contained in the collection belonging to Natural Resources Canada, Canadian Forest Service - Atlantic Forestry Centre, Fredericton, New Brunswick.

## Collection methods

Various methods were employed to collect the species reported in this study. Details are outlined in Webster et al. (2009, Appendix). Many specimens were also collected from 12-unit Lindgren funnel traps set in various forest habitats in New Brunswick between 2008 and 2011. These traps mimic tree trunks and are often effective for sampling species of Coleoptera that live in microhabitats associated with standing trees (Lindgren 1983). See Webster et al. (in press) for details of the methods used to deploy Lindgren funnel traps and for sample collection. A description of the habitat was recorded for all specimens collected during this survey. Locality and habitat data are presented exactly as on labels for each record. This information, as well as additional collecting notes, is summarized and discussed in the collection and habitat data section for each species.

## Specimen Preparation

Males of some species of Elateridae were dissected to confirm their identity. The genital structures were dehydrated in absolute alcohol and mounted in Canada balsam on celluloid microslides and pinned with the specimens they originated from.

## Distribution

Distribution maps, created using ArcMap and ArcGIS, are presented for each species in New Brunswick. Every species is cited with current distribution in Canada and Alaska, using abbreviations for the state, provinces, and territories. New records for New Brunswick are indicated in bold under Distribution in Canada and Alaska. The following abbreviations are used in the text:

|           |                       |                    |                           |
|-----------|-----------------------|--------------------|---------------------------|
| <b>AK</b> | Alaska                | <b>MB</b>          | Manitoba                  |
| <b>YT</b> | Yukon Territory       | <b>ON</b>          | Ontario                   |
| <b>NT</b> | Northwest Territories | <b>QC</b>          | Quebec                    |
| <b>NU</b> | Nunavut               | <b>NB</b>          | New Brunswick             |
| <b>BC</b> | British Columbia      | <b>PE</b>          | Prince Edward Island      |
| <b>AB</b> | Alberta               | <b>NS</b>          | Nova Scotia               |
| <b>SK</b> | Saskatchewan          | <b>NF &amp; LB</b> | Newfoundland and Labrador |

Acronyms of collections examined or where specimens reside referred to in this study are as follows:

|            |   |
|------------|---|
| <b>AFC</b> | Atlantic Forestry Centre, Natural Resources Canada, Canadian Forest Service, Fredericton, New Brunswick, Canada             |
| <b>CNC</b> | Canadian National Collection of Insects, Arachnids and Nematodes, Agriculture and Agri-Food Canada, Ottawa, Ontario, Canada |
| <b>NBM</b> | New Brunswick Museum, Saint John, New Brunswick, Canada   |
| <b>RWC</b> | Reginald Webster Collection, Charters Settlement, New Brunswick, Canada   |

## Results

Twenty-two species of Elateridae are newly reported for New Brunswick, *Negastrius exiguus* (Randall) is removed from the faunal list, and *Agriotes pubescens* Melsheimer reinstated as a member of the New Brunswick fauna, bringing the total number of species known from the province to 122. *Agriotes pubescens* Melsheimer, *Dalopius brevicornis*, *Danosoma obtectum* (Say), and *Megapenthes solitarius* Fall are newly reported for the Maritime provinces. Several apparently undescribed *Ampedus* sp. have also been found in New Brunswick but these are not reported here.

## Species accounts

All records below are species newly recorded for New Brunswick, Canada unless noted otherwise (additional records). Species followed by \*\* are newly recorded from the Maritime provinces of Canada.

The classification of the Elateridae follows Bouchard et al. (2011).

**Table 1.** Species of Elateridae recorded from New Brunswick, Canada.

|   |  |
|---|--|
| <b>Family Elateridae Leach</b>                | <i>Metanomus insidiosus</i> (LeConte)            |
| <b>Subfamily Agrypninae Candèze</b>           | <i>Nitidolimonius resplendens</i> (Eschscholtz)  |
| <b>Tribe Agrypnini Candèze</b>                | <i>Oxygonus montanus</i> Schaeffer               |
| <i>Danosoma brevicornis</i> (LeConte)         | <i>Oxygonus obesus</i> Say**                     |
| <i>Danosoma obiectum</i> (Say)**              | <i>Paractenicera fulvipes</i> (Bland)            |
| <i>Lacon auroratus</i> (Say)                  | <i>Prosternon medianum</i> (Germar)              |
| <b>Subfamily Lissominae Laporte</b>           | <i>Pseudanostirus hamatus</i> (Say)              |
| <i>Oestodes tenuicollis</i> (Randall)         | <i>Pseudanostirus hieroglyphicus</i> (Say)       |
| <b>Subfamily Pityobiinae Hyslop</b>           | <i>Pseudanostirus nigricollis</i> (Bland)        |
| <i>Pityobius anguinius</i> LeConte            | <i>Pseudanostirus propolus</i> (LeConte)         |
| <b>Subfamily Dendrometrinae Gistel</b>        | <i>Pseudanostirus triundulatus</i> (Randall)     |
| <b>Tribe Dendrometrini Gistel</b>             | <i>Selatosomus appropinquans</i> (Randall)       |
| <i>Athous acanthus</i> (Say)                  | <i>Selatosomus pulcher</i> (LeConte)             |
| <i>Athous brightwelli</i> (Kirby)             | <i>Selatosomus splendens</i> (Ziegler)           |
| <i>Athous fossularis</i> (LeConte)            | <i>Setasomus atratus</i> (LeConte)               |
| <i>Athous orvus</i> Becker                    | <i>Setasomus nitidulus</i> (LeConte)             |
| <i>Athous posticus</i> (Melsheimer)*          | <i>Setasomus rufopleuralis</i> (Fall)            |
| <i>Athous productus</i> (Randall)             | <i>Sylvanelater cylindriciformis</i> (Herbst)    |
| <i>Athous rufifrons</i> (Randall)             | <b>Tribe Hypnoidini Schwarz</b>                  |
| <i>Athous scapularis</i> (Say)*               | <i>Hypnoidus abbreviatus</i> (Say)               |
| <i>Denticollis denticornis</i> (Kirby)        | <i>Hypnoidus bicolor</i> (Eschscholtz)           |
| <i>Elathous discalceatus</i> (Say)*           | <i>Ligmargus lecontei</i> (Leng)*                |
| <i>Hemicrepidius brevicollis</i> (Candèze)    | <i>Margaistus grandicollis</i> (LeConte)         |
| <i>Hemicrepidius hemipodus</i> (Say)          | <b>Subfamily Negastrinae Nakane &amp; Kishii</b> |
| <i>Hemicrepidius memnonius</i> (Herbst)*      | <i>Microhypnus striatulus</i> (LeConte)          |
| <i>Hemicrepidius niger</i> (Linnaeus)         | <i>Negastrius arnetti</i> Stibick                |
| <i>Limonius aeger</i> LeConte                 | <i>Negastrius delumbis</i> (Horn)                |
| <i>Limonius anceps</i> LeConte                | <i>Negastrius atrosus</i> Wells**                |
| <i>Limonius confusus</i> LeConte              | <i>Neohypdonus tumescens</i> (LeConte)           |
| <i>Limonius pectoralis</i> LeConte            | <i>Oedostethus femoralis</i> LeConte             |
| <b>Tribe Prosternini Gistel</b>               | <i>Paradonus oliverea</i> Stibick                |
| <i>Actenicerus cuprascens</i> (LeConte)       | <i>Paradonus pectoralis</i> (Say)*               |
| <i>Anostirus vernalis</i> (Hentz)             | <i>Zorocheus melsheimeri</i> (Horn)              |
| <i>Beckerus appressus</i> (Randall)           | <b>Subfamily Elaterinae Leach</b>                |
| <i>Corymbitodes elongaticollis</i> (Hamilton) | <b>Tribe Agriotini Laporte</b>                   |
| <i>Corymbitodes pygmaeus</i> (Van Dyke)       | <i>Agriotes collaris</i> (LeConte)               |
| <i>Corymbitodes tarsalis</i> (Melsheimer)     | <i>Agriotes fuscus</i> (LeConte)                 |
| <i>Ctenicera kendalli</i> (Kirby)             | <i>Agriotes limosus</i> (LeConte)                |
| <i>Eanus estriatus</i> (LeConte)              | <i>Agriotes mancus</i> (Say)                     |
| <i>Eanus maculipennis</i> LeConte             | <i>Agriotes quebecensis</i> Brown*               |
| <i>Hypoganus sulcicollis</i> (Say)*           | <i>Agriotes sputator</i> (Linnaeus)              |
| <i>Hypoganus rotundicollis</i> (Say)**        | <i>Agriotes pubescens</i> Melsheimer*            |
| <i>Liotrichus falsificus</i> (LeConte)        | <i>Agriotes stabilis</i> (LeConte)               |
| <i>Liotrichus spinosus</i> (LeConte)          | <i>Dalopius cognatus</i> Brown                   |
| <i>Liotrichus vulneratus</i> (LeConte)        | <i>Dalopius fuscipes</i> Brown                   |

|                                      |   |
|--------------------------------------|---|
| <i>Dalopius pallidus</i> Brown       | <i>Ampedus subtilis</i> (LeConte)         |
| <i>Dalopius vagus</i> Brown          | <i>Ampedus virtuosus</i> (LeConte)        |
| <i>Dalopius brevicornis</i> Brown**  | <b>Tribe Elaterini Leach</b>              |
| <b>Tribe Ampedini Gistel</b>         | <i>Elater abruptus</i> Say*               |
| <i>Ampedus apicatus</i> (Say)        | <i>Sericus honestus</i> (Randall)         |
| <i>Ampedus areolatus</i> (Say)*      | <i>Sericus incongruus</i> (LeConte)       |
| <i>Ampedus collaris</i> (Say)        | <i>Sericus viridanus</i> (Say)*           |
| <i>Ampedus deletus</i> (LeConte)     | <b>Tribe Megapenthini Gurjeva</b>         |
| <i>Ampedus evansi</i> Brown          | <i>Megapenthes rogersi</i> Horn           |
| <i>Ampedus fuscus</i> (LeConte)      | <i>Megapenthes stigmatus</i> (LeConte)    |
| <i>Ampedus laurentinus</i> Brown     | <i>Megapenthes solitarius</i> Fall**      |
| <i>Ampedus luctuosus</i> (LeConte)   | <b>Tribe Melontini Candèze</b>            |
| <i>Ampedus minipennis</i> (LeConte)  | <i>Melanotus castanipes</i> (Paykull)     |
| <i>Ampedus mixtus</i> (Herbst)       | <i>Melanotus decumanus</i> (Erichson)     |
| <i>Ampedus molestus</i> (LeConte)    | <i>Melanotus leonardi</i> (LeConte)**     |
| <i>Ampedus nigricans</i> (Germar)    | <i>Melanotus similis</i> (Kirby)          |
| <i>Ampedus nigricollis</i> (Herbst)* | <i>Melanotus sagittarius</i> (LeConte)**  |
| <i>Ampedus nigrinus</i> (Herbst)     | <b>Tribe Pomachiliini Candèze</b>         |
| <i>Ampedus oblectus</i> (Say)        | <i>Agriotella bigeminata</i> (Randall)    |
| <i>Ampedus protervus</i> (LeConte)*  | <i>Agriotella debilis</i> (LeConte)       |
| <i>Ampedus pullus</i> Germar         | <b>Subfamily Cardiophorinae Candèze</b>   |
| <i>Ampedus rubricus</i> (Say)        | <i>Cardiophorus convexus</i> LeConte      |
| <i>Ampedus sanguinipennis</i> (Say)  | <i>Cardiophorus gagates</i> Erichson      |
| <i>Ampedus sayi</i> (LeConte)        | <i>Cardiophorus propinquus</i> Lanchester |
| <i>Ampedus semicinctus</i> (Randall) |   |

**Notes:** \*New to province, \*\*New to Maritime provinces.

## Family Elateridae Leach, 1815

### Subfamily Agrypninae Candèze, 1857

#### Tribe Agrypnini Candèze, 1857

##### *Danosoma obtectum* (Say, 1839)\*\*

[http://species-id.net/wiki/Danosoma\\_obtectum](http://species-id.net/wiki/Danosoma_obtectum)

Map 1

**Material examined.** New Brunswick, York Co., 14 km WSW of Tracy, S of Rt. 645, 45.6741°N, 66.8661°W, 13–27.VII.2010, R. Webster & C. MacKay, old mixed forest with red and white spruce, red and white pine, balsam fir, eastern white cedar, red maple, and *Populus* sp., Lindgren funnel trap (1, RWC).

**Collection and habitat data.** The single specimen from New Brunswick was captured during July in a Lindgren funnel trap deployed in an old mixed forest.

**Distribution in Canada and Alaska.** YK, NT, BC, AB, SK, MB, ON, PQ, NB (Bousquet 1991). Majka and Johnson (2008) removed *Danosoma obtectum* from the faunal list of Nova Scotia due to a lack of a supporting voucher specimen.

**Subfamily Dendrometrinae Gistel, 1848****Tribe Dendrometrini Gistel, 1848*****Athous posticus* (Melsheimer, 1846)**

[http://species-id.net/wiki/Athous\\_posticus](http://species-id.net/wiki/Athous_posticus)

Map 2

**Material examined.** **New Brunswick, Carleton Co.,** Meduxnekeag Valley Nature Preserve, 46.1957°N, 67.6803°W, 22.VII.2004, J. Edsall & R. P. Webster, mixed forest, u.v. light (1, RWC); Jackson Falls, Bell Forest, 46.2200°N, 67.7231°W, 5–12.VII.2008, 19–28.VII.2008, R. P. Webster, mature hardwood forest, Lindgren funnel traps (5, AFC, RWC). **Queens Co.,** Cranberry Lake P.N.A. (Protected Natural Area), 46.1125°N, 65.6075°W, 1–10.VII.2009, 15–21.VII.2009, 21–28.VII.2009, R. Webster & M.-A. Giguère, old red oak forest, Lindgren funnel traps (5, AFC); same locality data and forest type, 20.VII–4.VIII.2011, M. Roy & V. Webster, Lindgren funnel traps in forest canopy (3, NBM, RWC). **York Co.,** 15 km W of Tracy off Rt. 645, 45.6848°N, 66.8821°W, 14–20.VII.2009, R. Webster & M.-A. Giguère, old red pine forest, Lindgren funnel trap (1, RWC); same locality and habitat data but 30.VI–13.VII.2010, R. Webster & K. Burgess, Lindgren funnel trap (1, RWC); 14 km WSW of Tracy, S of Rt. 645, 45.6741°N, 66.8661°W, 16–30.VI.2010, 30.VI–13.VII.2010, R. Webster, C. MacKay, & K. Burgess, old mixed forest with red and white spruce, red and white pine, balsam fir, eastern white cedar, red maple, and *Populus* sp., Lindgren funnel traps (3, AFC, RWC).

**Collection and habitat data.** One adult was collected at an ultraviolet light, but most individuals were captured in Lindgren funnel traps in mixed and old mixed forests, a mature hardwood forest, an old red oak (*Quercus rubra* L.) forest, and an old red pine (*Pinus resinosa* Ait.) forest. Adults were collected during June, July, and August.

**Distribution in Canada and Alaska.** ON, QC, NB, NS (Bousquet 1991; Majka and Johnson 2008).

***Athous scapularis* (Say, 1839)**


[http://species-id.net/wiki/Athous\\_scapularis](http://species-id.net/wiki/Athous_scapularis)

Map 3


**Material examined.** **New Brunswick, Carleton Co.,** Jackson Falls, Bell Forest, 46.2208°N, 67.7211°W, 28.VI.2005, R. P. Webster, mature hardwood forest, u.v. light (1, RWC); same locality but 46.2200°N, 67.7231°W, 5–12.VII.2008, R. P. Webster, mature hardwood forest, Lindgren funnel trap (1, RWC). **Queens Co.,** Cranberry Lake P.N.A., 46.1125°N, 65.6075°W, 18–31.VIII.2011, M. Roy & V. Webster, old red oak forest, Lindgren funnel trap (1, AFC).

**Collection and habitat data.** Adults of this species were found in a mature hardwood forest with American beech (*Fagus grandifolia* Ehrh.), sugar maple (*Acer saccharum* Marsh.), and white ash (*Fraxinus americana* L.) and in an old red oak forest.







**Map 1.** Collection localities in New Brunswick, Canada of *Danosoma obrectum*.




**Map 2.** Collection localities in New Brunswick, Canada of *Athous posticus*.




**Map 3.** Collection localities in New Brunswick, Canada of *Athous scapularis*.



**Map 4.** Collection localities in New Brunswick, Canada of *Elathous discalceatus*.



**Map 5.** Collection localities in New Brunswick, Canada of *Hemicrepidius memnonius*.



**Map 6.** Collection localities in New Brunswick, Canada of *Hypoganus sulcicollis*.

Adults were captured at an ultraviolet light and in Lindgren funnel traps. This species was captured during June, July, and August. Becker (1974) reported larvae of this species from forest litter and decaying logs.

**Distribution in Canada and Alaska.** ON, QC, NB, NS (Bousquet 1991; Majka and Johnson 2008).

***Elathous discalceatus* (Say, 1839)**

[http://species-id.net/wiki/Elathous\\_discalceatus](http://species-id.net/wiki/Elathous_discalceatus)

Map 4

**Material examined.** New Brunswick, Carleton Co., Jackson Falls, Bell Forest, 46.2200°N, 67.7231°W, 19–28.VII.2008, R. P. Webster, mature hardwood forest, Lindgren funnel trap (1, RWC). York Co., 15 km W of Tracy off Rt. 645, 45.6848°N, 66.8821°W, 4–11.VIII.2009, R. Webster & M.-A. Giguère, old red pine forest, Lindgren funnel trap (1, RWC); same locality data, 27.VII–10.VIII.2010, R. Webster & C. Hughes, Lindgren funnel traps (3, AFC, RWC).

**Collection and habitat data.** Adults were captured during late July and August in Lindgren funnel traps in a mature hardwood forest with American beech, sugar maple, and white ash, and in an old red pine forest.

**Distribution in Canada and Alaska.** ON, QC, NB, NS (Bousquet 1991).

***Hemicrepidius memnonius* (Herbst, 1806)**

[http://species-id.net/wiki/Hemicrepidius\\_memnonius](http://species-id.net/wiki/Hemicrepidius_memnonius)

Map 5

**Material examined.** New Brunswick, York Co., Fredericton, 27.VII.1929, R. P. Gorham, (2, AFC); Charters Settlement, 45.8395°N, 66.7391°W, 1.VIII.2007, R. P. Webster, mixed forest, u.v. light (1, RWC); 15 km W of Tracy off Rt. 645, 45.6848°N, 66.8821°W, 20–29.VII.2009, 29.VII–4.VIII.2009, 11–18.VIII.2009, R. Webster & M.-A. Giguère, old red pine forest, Lindgren funnel traps (8, AFC, RWC); same locality data but 13–27.VII.2010, R. Webster & C. MacKay, Lindgren funnel traps (2, AFC, RWC).

**Collection and habitat data.** *Hemicrepidius memnonius* was collected at an ultraviolet light in a mixed forest and from Lindgren funnel traps in an old red pine forest. Adults were captured during July and August.

**Distribution in Canada and Alaska.** AB, SK, MB, ON, QC, NB, NS, PE (Bousquet 1991).

**Tribe Prosternini Gistel, 1856*****Hypoganus sulcicollis* (Say, 1834)**

[http://species-id.net/wiki/Hypoganus\\_sulcicollis](http://species-id.net/wiki/Hypoganus_sulcicollis)

Map 6

**Material examined.** New Brunswick, Carleton Co., Jackson Falls, Bell Forest, 46.2199°N, 67.7231°W, 9.IX.2006, 6.V.2007, R. P. Webster, mature hardwood forest, under bark of fallen beech logs (2, RWC); same locality but 20–26.V.2009, R. Webster & M.-A. Giguère, mature hardwood forest, Lindgren funnel trap (1, RWC).



**Queens Co.**, Grand Lake near Scotchtown, 45.8762°N, 66.1816°W, 25.IV.2004, R. Webster & M.-A. Giguère, oak and maple forest, under bark of oak (1, RWC); Cranberry Lake P.N.A., 46.1125°N, 65.6075°W, 21–27.V.2009, 10–15.VII.2009, R. Webster & M.-A. Giguère, old red oak forest, Lindgren funnel traps (2, RWC); Grand Lake Meadows P.N.A., 45.8227°N, 66.1209°W, 3–21.VI.2011, 21.VI–5.VII.2011, 5–19.VII.2011, 5–17.VIII.2011, M. Roy & V. Webster, old silver maple forest with green ash and seasonally flooded marsh, Lindgren funnel traps in forest canopy (4, NBM, RWC). **Sunbury Co.**, Burton near Sunpoke Lake, 45.7663°N, 66.5550°W, 20.VII.2006, R. P. Webster, oak forest, under loose bark of red oak (1, RWC).

**Collection and habitat data.** *Hypogonus sulcicollis* (Say) was collected in a mature hardwood forest with American beech, sugar maple, and white ash, in a red oak and red maple (*Acer rubrum* L.) forest, an old silver maple (*Acer saccharinum* L.) forest, and in an old red oak forest. Adults were collected from under bark of fallen American beech, under bark of red oak, and from Lindgren funnel traps. Adults were captured during April, May, June, July, August, and September.

**Distribution in Canada and Alaska.** MB, ON, QC, **NB**, NS (Bousquet 1991; Majka and Johnson 2008).

*Hypogonus rotundicollis* (Say, 1825)\*\*

[http://species-id.net/wiki/Hypogonus\\_rotundicollis](http://species-id.net/wiki/Hypogonus_rotundicollis)

Map 7

**Material examined.** New Brunswick, **Queens Co.**, near “Trout Creek”, 45.8237°N, 66.1225°W, 6.IX.2007, R. P. Webster, silver maple swamp, sweeping foliage on margin of marsh (1, RWC).

**Collection and habitat data.** The sole New Brunswick specimen of this species was collected during September by sweeping marsh vegetation on the margin of a silver maple swamp.

**Distribution in Canada and Alaska.** ON, **NB** (Bousquet 1991).

*Oxygonus obesus* (Say, 1823)\*\*

[http://species-id.net/wiki/Oxygonus\\_obesus](http://species-id.net/wiki/Oxygonus_obesus)

Map 8

**Material examined.** New Brunswick, **York Co.**, Canterbury, Browns Mountain Fen, 45.8967°N, 67.6343°W, 1.VI.2005, R. Webster & M.-A. Giguère, calcareous fen with shrubby cinquefoil, sweeping (2, RWC).

**Collection and habitat data.** Specimens of this species were swept from vegetation in an open calcareous cedar fen with shrubby cinquefoil (*Pentaphylloides floribunda* (Pursh) A. Löve) during early June.

**Distribution in Canada and Alaska.** AB, MB, ON, QC, **NB** (Bousquet 1991).

**Tribe Hypnoidini Schwarz, 1906 (1860)*****Ligmargus lecontei* (Leng, 1918)**

[http://species-id.net/wiki/Ligmargus\\_lecontei](http://species-id.net/wiki/Ligmargus_lecontei)

Map 9

**Material examined.** New Brunswick, Restigouche Co., Jacquet River Gorge P.N.A. near Jacquet R., 47.8897°N, 66.0835°W, 23.VI.2008, 26.VI.2008, R. P. Webster, river margin, among cobblestones (2, RWC); same locality but 47.8204°N, 66.0833°W, 14.VI.2009, R. P. Webster, river margin, among cobblestones (1, RWC); same locality but 47.8357°N, 66.0779°W, 14.V.2010, 24.V.2010, R. P. Webster, partially shaded gravel bar near confluence of brook and river, among cobblestones (2, RWC).

**Collection and habitat data.** *Ligmargus lecontei* adults were collected from under cobblestones along the margin of a fast-flowing, clear (cool water), rocky, river during May and June.

**Distribution in Canada and Alaska.** ON, QC, NB, NS (Bousquet 1991).

**Subfamily Negastrinae Nakane and Kishii, 1956*****Negastrius exiguus* (Randall, 1838)**

The record of *N. exiguus* in Majka and Johnson (2008) was based on a misidentification by C.G. Majka and was *N. atrosus* Wells (determined by Serge Laplante). In view of this, *N. exiguus* is removed from the faunal list of New Brunswick.

***Negastrius atrosus* Wells, 1996\*\***

[http://species-id.net/wiki/Negastrius\\_atrosus](http://species-id.net/wiki/Negastrius_atrosus)


Map 10

**Material examined.** New Brunswick, Queens Co., Bayard, at Nerepis River, 45.4426°N, 66.3280°W, 30.V.2008, R. P. Webster, river margin, under small rocks embedded in gravel (2, RWC). Restigouche Co., confluence of Restigouche River and Stillwater Brook, 26.VI.2000, R. Webster, F. Roy, & P. Poitras, in gravel on river margin (1, RWC). York Co., Rt. 105 at Nashwaaksis River, 45.9853°N, 66.6910°W, 9.V.2006, R. P. Webster, river margin, splashing water onto sand bar (2, RWC); 1.5 km S of Taymouth at the Nashwaak River, 46.1582°N, 66.6134°W, 15.VI.2008, R. P. Webster, on sand bar under drift material on sand (5, RWC).


**Collection and habitat data.** *Negastrius atrosus* adults were collected along river margins from under small rocks, in gravel, and under drift material on sand bars. Two adults were collected by splashing water onto sand on a sand bar. Adults were collected




**Map 7.** Collection localities in New Brunswick, Canada of *Hypogonus rotundicollis*.




**Map 8.** Collection localities in New Brunswick, Canada of *Oxygonus obesus*.




**Map 9.** Collection localities in New Brunswick, Canada of *Ligmargus lecontei*.



**Map 10.** Collection localities in New Brunswick, Canada of *Negastris atrosus*.



**Map 11.** Collection localities in New Brunswick, Canada of *Paradonus pectoralis*.



**Map 12.** Collection localities in New Brunswick, Canada of *Agriotes quebecensis*.

during May and June. No details on the habitat requirements of this species were given in Wells (1996). Wells (1996) reported that other species of *Negastris* were associated with riparian habitats and inhabit sandy-to-rocky stream and river margins.

**Distribution in Canada and Alaska.** ON, QC, NB (Wells 1996).

***Paradonus pectoralis* (Say, 1839)**[http://species-id.net/wiki/Paradonus\\_pectoralis](http://species-id.net/wiki/Paradonus_pectoralis)

Map 11

**Material examined.** **New Brunswick, Charlotte Co.,** St. Andrews, 45.0751°N, 67.0374°W, 25.VIII.2006, R. P. Webster, sea beach, sweeping foliage (1, RWC). **Queens Co.,** Grand Lake near Scotchtown, 45.8762°N, 66.1816°W, 5.VI.2004, R. P. Webster, in gravel near shoreline of lake (2, RWC); Grand Lake at Stony Point, 46.0031°N, 66.0337°W, 17.VIII.2004, D. Sabine & R. Webster, lakeshore, cobblestone beach, among cobblestones (3, RWC). **York Co.,** Charters Settlement, 45.8395°N, 66.7391°W, 19.VII.2005, 9.VII.2006, 17.VII.2008, R. P. Webster, mixed forest, u.v. light (3, RWC).

**Collection and habitat data.** Adults of this species were collected in gravel and among cobblestones along lakeshores, by sweeping foliage on a sea beach, and at an ultraviolet light in a mixed forest. Adults were collected during June, July, and August.

**Distribution in Canada and Alaska.** BC, AB, SK, MB, ON, QC, NB, NS (Bousquet 1991).

**Subfamily Elaterinae Leach, 1815****Tribe Agriotini Laporte, 1840*****Agriotes quebecensis* Brown, 1933**[http://species-id.net/wiki/Agriotes\\_quebecensis](http://species-id.net/wiki/Agriotes_quebecensis)

Map 12

**Material examined.** **New Brunswick, Restigouche Co.,** Jacquet River Gorge P.N.A., 47.7235°N, 66.1278°W, 16.VI.2009, K. A. A. Vandenbroeck (1, NBM). **York Co.,** Charters Settlement, 45.8380°N, 66.7310°W, 14.V.2004, R. P. Webster, beating foliage (1, RWC); 14 km WSW of Tracy, S of Rt. 645, 45.6741°N, 66.8661°W, 17–31.V.2010, R. Webster & C. MacKay, old mixed forest with red and white spruce, red and white pine, balsam fir, eastern white cedar, red maple, and *Populus* sp., Lindgren funnel trap (1, AFC).

**Collection and habitat data.** One individual was beaten from foliage in a mixed forest, another was captured in a Lindgren funnel trap deployed in an old mixed forest. Adults were captured during May and June.

**Distribution in Canada and Alaska.** MB, ON, QC, NB, PE, NS (Bousquet 1991; Majka and Johnson 2008).

***Agriotes pubescens* Melsheimer, 1845**[http://species-id.net/wiki/Agriotes\\_pubescens](http://species-id.net/wiki/Agriotes_pubescens)

Map 13

**Material examined.** New Brunswick, **Queens Co.**, Grand Lake Meadows P.N.A., 45.8227°N, 66.1209°W, 31.V–15.VI.2010, 15–29.VI.2010, R. Webster & C. MacKay, old silver maple forest with green ash and seasonally flooded marsh, Lindgren funnel traps (5, AFC, RWC). **Sunbury Co.**, Burton near Sunpoke Lake, 45.7658°N, 66.5546°W, 20.VI.2007, R. P. Webster, red oak and red maple forest, on foliage of *Quercus rubra* (1, RWC).

**Collection and habitat data.** Adults were collected during June from Lindgren funnel traps in an old silver maple forest (swamp) and from foliage of red oak in a red oak and red maple stand. Both forest sites were near seasonally flooded marshes.

**Distribution in Canada and Alaska.** MB, ON, QC, **NB** (Bousquet 1991). Bousquet (1991) reported *Agriotes pubescens* Melsheimer from New Brunswick. Majka and Johnson (2008) were unable to locate voucher specimens to support the record and, thus, they removed it from the faunal list of New Brunswick. The records above establish the presence of this species in the province.

***Dalopius brevicornis* Brown, 1934\*\***[http://species-id.net/wiki/Dalopius\\_brevicornis](http://species-id.net/wiki/Dalopius_brevicornis)

Map 14

**Material examined.** New Brunswick, **Carleton Co.**, Jackson Falls, Bell Forest, 46.2152°N, 67.7190°W, 1.VI.2005, M.-A. Giguère & R. P. Webster, upper river margin near floodplain forest, sweeping foliage (4, RWC); Meduxnekeag Valley Nature Preserve, 46.1931°N, 67.6825°W, 8. VI.2005, M.-A. Giguère & R. P. Webster, margin of floodplain forest with butternut, sweeping (3, RWC).

**Collection and habitat data.** Adults were collected during early June by sweeping foliage near floodplain forests adjacent to rivers.


**Distribution in Canada and Alaska.** ON, QC, **NB** (Bousquet 1991).

**Tribe Ampedini Gistel, 1848*****Ampedus areolatus* (Say, 1823)**[http://species-id.net/wiki/Ampedus\\_areolatus](http://species-id.net/wiki/Ampedus_areolatus)

Map 15

**Material examined.** New Brunswick, **Carleton Co.**, Jackson Falls, Bell Forest, 46.2152°N, 67.7190°W, 12.VI.2008, R. P. Webster, river margin, treading vegetation in seepage area (1, RWC); Meduxnekeag Valley Nature Preserve, 46.1931°N,







**Map 13.** Collection localities in New Brunswick, Canada of *Agriotes pubescens*.




**Map 14.** Collection localities in New Brunswick, Canada of *Dalpius brevicornis*.




**Map 15.** Collection localities in New Brunswick, Canada of *Ampedus areolatus*.



**Map 16.** Collection localities in New Brunswick, Canada of *Ampedus nigricollis*.



**Map 17.** Collection localities in New Brunswick, Canada of *Ampedus oblesus*.



**Map 18.** Collection localities in New Brunswick, Canada of *Ampedus protervus*.

67.6825°W, 8.VI.2005, M.-A. Giguère & R. P. Webster, floodplain forest with butternut, sweeping (1, RWC). **Queens Co.**, Grand Lake Meadows P.N.A., 45.8227°N, 66.1209°W, 19–31.V.2010, 31.V–15.VI.2010, 15–29.VI.2010, 29.VI–12.VII.2010, R. Webster, C. MacKay, M. Laity, & R. Johns, old silver maple forest with green ash and seasonally flooded marsh, Lindgren funnel traps (18,



AFC, RWC); same locality data and forest type, 17–30.VIII.2011, C. Hughes & R. P. Webster, Lindgren funnel traps (2, NBM).

**Collection and habitat data.** Adults were collected by treading vegetation in a seepage area along a river margin, by sweeping vegetation in a floodplain forest, and from Lindgren funnel traps deployed in an old silver maple forest near a seasonally flooded marsh. Adults were collected during May, June, July, and August.

**Distribution in Canada and Alaska.** ON, QC, NB, NS (Bousquet 1991; Majka and Johnson 2008).

***Ampedus nigricollis* (Herbst, 1801)**

[http://species-id.net/wiki/Ampedus\\_nigricollis](http://species-id.net/wiki/Ampedus_nigricollis)

Map 16

**Material examined.** New Brunswick, Queens Co., Grand Lake Meadows P.N.A., 45.8227°N, 66.1209°W, 21.VI–5.VII.2011, M. Roy & V. Webster, old silver maple forest with green ash and seasonally flooded marsh, Lindgren funnel traps in forest canopy (1, RWC). Sunbury Co., Mauderville, Portobello Creek N.W.A., 45.8990°N, 66.4200°W, 28.VI.2004, R. P. Webster, silver maple swamp, under bark of silver maple (1, RWC).

**Collection and habitat data.** One specimen was found under bark of a silver maple during late June in a silver maple swamp, another was captured between late June and early July in a Lindgren funnel trap deployed in the canopy of a silver maple in a silver maple swamp. Majka and Johnson (2008) reported this species from rotten wood of poplar, a spruce stump, and reared from an apple log in Nova Scotia.

**Distribution in Canada and Alaska.** ON, QC, NB, NS (Bousquet 1991; Majka and Johnson 2008).

***Ampedus oblessus* (Say, 1833)**

[http://species-id.net/wiki/Ampedus\\_oblessus](http://species-id.net/wiki/Ampedus_oblessus)

Map 17

**Material examined.** Additional New Brunswick records, Queens Co., Cranberry Lake P.N.A., 46.1125°N, 65.6075°W, 5–11.VI.2009, 18–25.VI.2009, R. Webster & M.-A. Giguère, old red oak forest, Lindgren funnel traps (3, AFC, RWC); same locality data and forest type, 13–25.V.2011, 25.V–7.VI.2011, 7–22.VI.2011, 29.VI–7.VII.2011, M. Roy & V. Webster, Lindgren traps in forest canopy (68, AFC, CNC, NBM, RWC); Grand Lake Meadows P.N.A., 45.8227°N, 66.1209°W, 1–3.VI.2011, 3–21.VI.2011, 21.VI–5.VII.2011, 5–19.VII.2011, M. Roy & V. Webster, old silver maple forest with green ash and seasonally flooded marsh, Lindgren funnel traps in forest canopy (13, AFC, NBM, RWC).

**Collection and habitat data.** Adults were captured during May, June, and July in Lindgren funnel traps in an old red oak stand and an old silver maple swamp.

Most (77 out of 81) individuals were captured in traps deployed in the forest canopy (mid crown).

**Distribution in Canada and Alaska.** AB, SK, MB, ON, QC, NB (Bousquet 1991). *Ampedus oblessus* (Say) was reported for New Brunswick in Bousquet (1991) but was not listed as a member of the fauna by Majka and Johnson (2008). The above record confirms the presence of this species for the province.

***Ampedus protervus* (LeConte, 1853)**

[http://species-id.net/wiki/Ampedus\\_protervus](http://species-id.net/wiki/Ampedus_protervus)

Map 18

**Material examined.** **New Brunswick, Carleton Co.,** Jackson Falls, Bell Forest, 46.2200°N, 67.7231°W, 9.IX.2006, 6.V.2007, R. P. Webster, mature hardwood forest, under bark of fallen beech log (3, RWC); same locality but 4–12.VI.2008, 12–19.VI.2008, 19–27.VI.2008, R. P. Webster, mature hardwood forest, Lindgren funnel traps (3, AFC, RWC); same locality and habitat data, 14–20.V.2009, M.-A. Giguère & R. Webster, Lindgren funnel traps (2, RWC). **Queens Co.,** Cranberry Lake P.N.A, 46.1125°N, 65.6075°W, 12–21.V.2009, 1–10.VII.2009, R. Webster & M.-A. Giguère, old red oak forest, Lindgren funnel traps (2, AFC, RWC); same locality data and forest type, 25.V–7.VI.2011, 7–22.VI.2011, 29.VI–7.VII.2011, M. Roy & V. Webster, Lindgren funnel traps (3, AFC, NBM). **York Co.,** Charters Settlement, 45.8331°N, 66.7410°W, 27.VII.2005, R. P. Webster, mixed forest, on foliage of *Alnus incana* (1, RWC); 15 km W of Tracy off Rt. 645, 45.6848°N, 66.8821°W, 14–20.VII.2009, R. Webster & M.-A. Giguère, old red pine forest, Lindgren funnel trap (1, RWC).

**Collection and habitat data.** Adults of this species were captured in Lindgren funnel traps deployed in mature hardwood forests with American beech, sugar maple, and white ash, an old red oak forest, and an old red pine forest. Adults were also collected from under bark of a fallen beech log in mature hardwood forest and from alder (*Alnus incana* (L.) Moench) foliage in a mixed forest. Adults were collected during May, June, July, and September.

**Distribution in Canada and Alaska.** ON, QC, NB, NS (Bousquet 1991).

**Tribe Elaterini Leach, 1915**

***Elater abruptus* Say, 1825**

[http://species-id.net/wiki/Elater\\_abruptus](http://species-id.net/wiki/Elater_abruptus)

Map 19

**Material examined.** **New Brunswick, Carleton Co.,** Jackson Falls, Bell Forest, 46.2200°N, 67.7231°W, 19–28.VII.2008, R. P. Webster, mature hardwood forest,

Lindgren funnel trap (1, RWC). **Queens Co.**, Cranberry Lake P.N.A., 46.1125°N, 65.6075°W, 28.VII–6.VIII.2009, R. Webster & M.-A. Giguère, old red oak forest, Lindgren funnel trap (1, AFC); same locality data and forest type, 20.VII–4.VIII.2011, 4–18.VIII.2011, M. Roy & V. Webster, Lindgren funnel traps in forest canopy (5, AFC, NBM, RWC); Grand Lake Meadows P.N.A., 45.8227°N, 66.1209°W, 19.VII–5.VIII.2011, 5–17.VII.2011, M. Roy & V. Webster, old silver maple forest and seasonally flooded marsh, Lindgren funnel traps in forest canopy (8, AFC, NBM, RWC).

**Collection and habitat data.** Adults were captured in Lindgren funnel traps in a mature hardwood forest with American beech, sugar maple, and white ash, in an old silver maple forest, and in an old red oak forest. Most adults were captured in traps deployed in the forest canopy. Adults were captured during July and August.

**Distribution in Canada and Alaska.** ON, QC, NB, NS (Bousquet 1991).

### *Sericus viridanus* (Say, 1825)

[http://species-id.net/wiki/Sericus\\_viridanus](http://species-id.net/wiki/Sericus_viridanus)

Map 20

**Material examined.** New Brunswick, Carleton Co., Jackson Falls, Bell Forest, 46.2200°N, 67.7231°W, 12–29.VI.2008, R. P. Webster, mature hardwood forest, Lindgren funnel trap (1, RWC).

**Collection and habitat data.** The only specimen known from New Brunswick was captured during June in a Lindgren funnel trap deployed in a mature hardwood forest with American beech, sugar maple, and white ash.

**Distribution in Canada and Alaska.** ON, QC, NB, NS (Bousquet 1991).


## Tribe Megapenthini Gurjeva, 1973

### *Megapenthes rogersi* Horn, 1871

[http://species-id.net/wiki/Megapenthes\\_rogersi](http://species-id.net/wiki/Megapenthes_rogersi)

Map 21


**Material examined.** Additional New Brunswick records, **Kings Co.**, Hampton, Hampton Marsh, 45.4787°N, 65.9007°W, 13.VII.2005, R. P. Webster, floodplain forest, on foliage of silver maple (1, RWC). **Queens Co.**, Cranberry Lake P.N.A., 46.1125°N, 65.6075°W, 29.VI–7.VII.2011, 7–13.VII.2011, 13–20.VII.2011, 20.VII–4.VIII.2011, 4–18.VIII.2011, M. Roy & V. Webster, old red oak forest, Lindgren funnel traps in forest canopy (16, AFC, NBM, RWC); Grand Lake Meadows P.N.A., 45.8227°N, 66.1209°W, 5–19.VII.2011, M. Roy & V. Webster, old silver maple forest with green ash and seasonally flooded marsh, Lindgren funnel traps in forest canopy (2, NBM, RWC). **York Co.**, 15 km W of Tracy off Rt. 645, 45.6848°N, 66.8821°W,




**Map 19.** Collection localities in New Brunswick, Canada of *Elater abruptus*.




**Map 20.** Collection localities in New Brunswick, Canada of *Sericus viridanus*.



**Map 21.** Collection localities in New Brunswick, Canada of *Megapenthes rogersi*.



**Map 22.** Collection localities in New Brunswick, Canada of *Megapenthes solitarius*.



**Map 23.** Collection localities in New Brunswick, Canada of *Melanotus leonardi*.



**Map 24.** Collection localities in New Brunswick, Canada of *Melanotus sagittarius*.

30.VI–13.VII.2010, R. Webster & K. Burgess, old red pine forest, Lindgren funnel trap (in forest canopy) (1, RWC).

**Collection and habitat data.** One adult of this species was collected from foliage of silver maple near a river. All others (19 specimens) from New Brunswick

were captured in Lindgren funnel traps deployed in the canopy of an old red oak forest, an old silver maple forest, and an old red pine forest. No adults were captured in traps deployed near the forest floor at the above sites, indicating that this species may be most active in the forest canopy. Adults were captured during June, July, and August.

**Distribution in Canada and Alaska.** ON, QC, NB (Bousquet 1991).

***Megapenthes solitarius* Fall, 1934\*\***

[http://species-id.net/wiki/Megapenthes\\_solitarius](http://species-id.net/wiki/Megapenthes_solitarius)

Map 22

**Material examined.** New Brunswick, Queens Co., Cranberry Lake P.N.A., 46.1125°N, 65.6075°W, 21–27.V.2009, 5–11.VI.2009, R. Webster & M.-A. Giguère, old red oak forest, Lindgren funnel traps (3, NBM, RWC); same locality data and forest type, 13–25.V.2011, 25.V–7.VI.2011, 22–29.VI.2011, M. Roy & V. Webster, Lindgren funnel traps (3, RWC). Restigouche, Co., Dionne Brook P.N.A., 47.9064°N, 68.3441°W, 30.V–15.VI.2011, 27.VI–14.VII.2011, M. Roy & V. Webster, old-growth northern hardwood forest, Lindgren funnel traps (3, NBM, CNC).

**Collection and habitat data.** In Alberta, two adults of *M. solitarius* were collected in mixed boreal forests; one from a window trap, the other was flying in a forest when captured (Fuller 2008). In New Brunswick, adults (9 specimens) of this rare species were captured in Lindgren funnel traps in an old red oak forest and an old-growth northern hardwood forest with sugar maple and yellow birch (*Betula alleghaniensis* Britt.). Adults were captured during May, June, and July.

**Distribution in Canada and Alaska.** AB, QC, NB (Bousquet 1991; Fuller 2008).

**Tribe Melontini Candéze, 1859**

***Melanotus leonardi* (LeConte, 1853)\*\***

[http://species-id.net/wiki/Melanotus\\_leonardi](http://species-id.net/wiki/Melanotus_leonardi)

Map 23

**Material examined.** New Brunswick, Saint John Co., Saint John, Taylor's Island 12.VI.1999, R. P. Webster, sea beach, under seaweed (1, RWC).

**Collection and habitat data.** One adult was collected from under seaweed (drift material) on a sea beach during June along with many other Coleoptera species from other families.

**Distribution in Canada and Alaska.** MB, ON, QC, NB (Bousquet 1991).



***Melanotus sagittarius* (LeConte, 1853)\*\***

[http://species-id.net/wiki/Melanotus\\_sagittarius](http://species-id.net/wiki/Melanotus_sagittarius)

Map 24

**Material examined.** New Brunswick, York Co., Charters Settlement, 45.8395°N, 66.7391°W, 27.VI.2006, 20.VII.2006, 10.VI.2007, 25.VI.2009, R. P. Webster, mixed forest, u.v. light (6, NBM, RWC).

**Collection and habitat data.** Adults from New Brunswick were collected at an ultraviolet light in a mixed forest during June and July.

**Distribution in Canada and Alaska.** QC, NB (Bousquet 1991).

## Acknowledgments

We thank Caroline Simpson (AFC) for editing this manuscript and an anonymous reviewer for helpful comments that improved this manuscript. Serge Laplante (Agriculture and Agri-Food Canada (CNC), Ottawa) is thanked for determining many of the specimens and for other invaluable assistance, and Nichole Brawn, Katie Burgess, Marie-Andrée Giguère, Jim Edsall, Nancy Harn, Cory Hughes, Rob Johns, Marsella Laity, Colin MacKay, Wayne MacKay, P. Poitras, Jessica Price, F. Roy, Michelle Roy, Dwayne Sabine, Katelyn Vandenbroeck, and Vincent Webster are thanked for technical assistance and collecting specimens. Natural Resources Canada - Canadian Forest Service; the Canadian Food Inspection Agency; and the USDA APHIS are thanked for funding the study on early detection of invasive cerambycids, which provided many specimens collected in Lindgren funnel traps. We thank the New Brunswick Environmental Trust Fund and New Brunswick Wildlife Trust Fund for funding various insect surveys over the past 7 years, and the Meduxnekeag River Association for permission to sample beetles at the Meduxnekeag Valley Nature Preserve (which includes the Bell Forest). The Canadian Wildlife Service is thanked for funding insect surveys at the Portobello Creek National Wildlife Area. The New Brunswick Department of Natural Resources (Fish and Wildlife Branch) is thanked for issuing permits for sampling in the Protected Natural Areas and for providing logistical support. Survey work in the Jacques River Gorge Protected Natural Area was organized through the New Brunswick Museum with external funding from the New Brunswick Environmental Trust Fund, Salamander Foundation, and the New Brunswick Wildlife Trust Fund.

## References

Becker EC (1974) Revision of the Nearctic species of *Athous* (Coleoptera: Elateridae) east of the Rocky Mountains. The Canadian Entomologist 106: 711–758. doi: 10.4039/Ent106711-7



- Bouchard P, Bousquet Y, Davies AE, Alonso-Zarazaga MA, Lawrence JF, Lyal CHC, Newton AF, Reid CAM, Schmitt M, Ślipiński SA, Smith ABT (2011) Family- group names in Coleoptera (Insecta). ZooKeys 88: 1–972. doi: 10.3897/zookeys.88.807
- Bousquet Y (1991) Family Elateridae: click beetles. In: Bousquet Y (Ed) Checklist of Beetles of Canada and Alaska. Agriculture Canada, Ottawa, Ontario, Publication 1861/E, Agriculture Canada, Research Branch, Ottawa, Ontario, 175–185.
- Douglas H (2011) New records of European wireworm pests and other click beetles (Coleoptera: Elateridae) in Canada and USA. Journal of the Entomological Society of Ontario 142: 11–17.
- Fuller ER (2008) New records of *Megapenthes solitarius* Fall from Alberta, Canada (Coleoptera: Elateridae). The Coleopterists Bulletin 62: 535–536. doi: 10.1649/1076.1
- Johnson PJ (2002) Family 58. Elateridae Leach 1815. In: Arnett RH Jr., Thomas MC, Skelley PE, Frank JH (Eds) American Beetles. Volume 2. Polyphaga: Scarabaeoidea through Curculionoidea, CRC Press, Boca Raton, Florida, 160–173.
- Lindgren BS (1983) A multiple funnel trap for scolytid beetles (Coleoptera). The Canadian Entomologist 115: 299–302. doi: 10.4039/Ent115299-3
- Majka CG, Johnson PL (2008) The Elateridae (Coleoptera) of the Maritime provinces of Canada: faunal composition, new records, and taxonomic changes. Zootaxa 1811: 1–33.
- Webster RP, Klimaszewski J, Pelletier G, Savard K (2009) New Staphylinidae (Coleoptera) records with new collection data from New Brunswick, Canada. I. Aleocharinae. In: Majka CG, Klimaszewski J (Eds) Biodiversity, biosystematics, and ecology of Canadian Coleoptera II. ZooKeys 22: 171–248. doi: 10.3897/zookeys.22.152
- Webster RP, Smetana A, Sweeney JD, DeMerchant I (in press) New Staphylinidae (Coleoptera) records with new collection data from New Brunswick and an addition to the fauna of Quebec: Staphylininae. In: Klimaszewski J, Anderson R (Eds) Biodiversity, Biosystematics and Ecology of Canadian Staphylinidae (Coleoptera) II. ZooKeys.
- Wells SA (1996) Studies on Nearctic *Negastrius* (Coleoptera: Elateridae). Great Basin Naturalist 56(4): 308–318.

