

Description of *Alpheus cedrici* sp. n., a strikingly coloured snapping shrimp (Crustacea, Decapoda, Alpheidae) from Ascension Island, central Atlantic Ocean

Arthur Anker^{1,†}, Sammy De Grave^{2,‡}

1 Instituto de Ciências do Mar (Labomar), Universidade Federal do Ceará (UFC), Fortaleza, Ceará, Brazil
2 Oxford University Museum of Natural History, Oxford, United Kingdom

† [urn:lsid:zoobank.org:author:B03ACF5D-1F1B-4653-9A7F-781804A2ABAB](https://zoobank.org/urn:lsid:zoobank.org:author:B03ACF5D-1F1B-4653-9A7F-781804A2ABAB)

‡ [urn:lsid:zoobank.org:author:D009C42B-FB1F-4473-B6D4-D6C8389D0F74](https://zoobank.org/urn:lsid:zoobank.org:author:D009C42B-FB1F-4473-B6D4-D6C8389D0F74)

Corresponding author: Sammy De Grave (sammy.degrave@oum.ox.ac.uk)

Academic editor: Niel Bruce | Received 12 March 2012 | Accepted 3 April 2012 | Published 19 April 2012

[urn:lsid:zoobank.org:pub:7861E107-FB5E-48CE-9AA8-23CAE43C86BD](https://zoobank.org/urn:lsid:zoobank.org:pub:7861E107-FB5E-48CE-9AA8-23CAE43C86BD)

Citation: Anker A, De Grave S (2012) Description of *Alpheus cedrici* sp. n., a strikingly coloured snapping shrimp (Crustacea, Decapoda, Alpheidae) from Ascension Island, central Atlantic Ocean. ZooKeys 183: 1–15. doi: 10.3897/zookeys.183.3073

Abstract

Alpheus cedrici sp. n. is described based on two specimens collected under rocks while scuba diving off the coast of Ascension Island, central Atlantic Ocean. The new species belongs to the *Alpheus macrocheles* (Hailstone, 1835) species complex and appears to be most closely related to the eastern–central Atlantic *A. macrocheles*, the western Atlantic *A. amblyonyx* Chace, 1972, and the eastern Pacific *A. bellimanus* Lockington, 1877 and *A. rectus* Kim & Abele, 1988. However, it differs from all these species by a combination of morphological characters and by a diagnostic and striking colour pattern.

Keywords

Caridea, Alpheidae, new species, *Alpheus*, Ascension Island, Atlantic Ocean

Introduction

The current knowledge of the alpheid shrimp fauna of the isolated central Atlantic islands St. Helena and Ascension is mainly based on two accounts, Chace (1966) for St. Helena and Manning and Chace (1990) for Ascension. Chace (1966) reported only

three alpheid species from St. Helena, viz. *Alpheus macrocheles* (Hailstone, 1835), *Synalpheus fritzmuelleri* Coutière, 1909, and *Metalpheus paragracilis* (Coutière, 1897). Manning and Chace (1990) reported the same three alpheids from Ascension Island, and in addition *Alpheus bouvieri* A. Milne Edwards, 1878, *A. crockeri* (Armstrong, 1941) [with some doubts], *A. dentipes* Guérin, 1832, *A. holthuisi* Ribeiro, 1964, *A. paracrinitus* Miers, 1881, *Automate dolichognatha* De Man, 1888, *Metalpheus rostratipes* (Pocock, 1890), *Parabetaeus hummelincki* (Schmitt, 1936) [as *Neotalpheopsis euryone* (De Man, 1910)], *Salmonesus setosus* Manning & Chace, 1990, and *S. teres* Manning & Chace, 1990, resulting in a total of 13 species of Alpheidae known to occur in the Central Atlantic Ocean south of Equator. Most alpheid specimens reported in Chace (1966) and Manning and Chace (1990) were collected in intertidal and shallow subtidal habitats, in tide pools, under rocks, in crevices of rocks and conglomerates of coralline algae, or in buoy fouling.

In April 2008, while scuba diving in English Bay, Ascension Island, one of us (SDG) collected two strikingly coloured snapping shrimps, by flipping rocks at a depth range of 10–15 m. A closer examination of these specimens revealed that they belong to a hitherto unnamed species of *Alpheus* Fabricius, 1798. This species is here-with described as new. Type material is deposited in the collections of the Oxford University Museum of Natural History, Oxford, the United Kingdom (OUMNH.ZC). Abbreviations used in the text: cl, carapace length (measured from the tip of the rostrum to the posterior margin of the carapace); Mxp, maxilliped; P, pereopod; CA, central Atlantic; EA, eastern Atlantic; WA, western Atlantic; EP eastern Pacific.

Systematics

Family Alpheidae Rafinesque, 1815

Genus *Alpheus* Fabricius, 1798

Alpheus cedrici sp. n.

urn:lsid:zoobank.org:act:7887B4BB-52D9-4329-A8BC-F32D8FA19FFC

http://species-id.net/wiki/Alpheus_cedrici

Figs 1–3

Material examined. Holotype: male, cl 10.1 mm, OUMNH.ZC.2008-11-0017, Ascension Island, west side of English Bay, 07°53.675'S, 014°22.999'W, depth 10 m, under rocks, leg S. & H. De Grave, 16.04.2008. Paratype: ovigerous female, cl 11.8 mm, OUMNH.ZC.2008-11-0018, Ascension Island, west side of English Bay, 07°53.675'S 014°22.999'W, depth 15 m, under rocks, leg. S. & H. De Grave, 17.04.2008.

Comparative material examined. *Alpheus macrocheles* (Hailstone, 1835): 1 male, cl 9.0 mm, OUMNH.ZC.2003-36-0002, Madeira, Canico, depth 20 m, leg. P. Wirtz, 02.11.2003. *Alpheus amblyonyx* Chace, 1972: 1 male, cl 5.6 mm, OUMNH.ZC.2011-03-0070, Panama, Isla Grande, in coral rubble, 1–1.5 m, leg. A. Anker, 09.12.2006.




Figure 1. *Alpheus cedrici* sp. n. **A–M** holotype, male from Ascension Island (OUMNH.ZC. 2008-11-0017); **N** paratype, female, same locality (OUMNH.ZC. 2008-11-0018). **A** frontal region, dorsal **B** anterior carapace, lateral **C** tooth of ventromesial carina of first article of antennular peduncle, lateral **D** third maxilliped, lateral **E** second pereopod, lateral **F** third pereopod, lateral **G** fifth pereopod, lateral **H** second pleopod, lateral **I** same, appendix masculina and appendix interna, mesial **J** same, detail of spiniform setae on protopod, lateral **K** third pleopod, detail of spiniform setae on protopod, mesial **L** uropod, dorsal **M** telson, dorsal **N** anterior carapace, dorsal.

Diagnosis. Frontal margin of carapace with rostrum slightly flattened dorsally, tapering distally, with acute tip, much longer than wide, reaching half-length of first article of antennular peduncle; rostral carina not distinct; orbital teeth in marginal position, small, acute distally, shorter than rostrum; margin between orbital teeth and rostrum broadly V-shaped; rostro-orbital process present; pterygostomial angle rounded; antennular peduncle with stylocerite not reaching distal margin of first article, with acute tip; second article about 2.5 times as long as wide; antenna with basicerite terminating in sharp distoventral tooth; carpocerite slightly exceeding both scaphocerite and antennular peduncle; scaphocerite with well-developed blade, shallowly concave lateral margin and large, stout distolateral tooth, latter reaching far beyond distal margin of blade; male minor cheliped with ventromesial margin of merus ending in small, acute distomesial tooth, and with minute spiniform setae; palm strongly compressed, with sculpture on both lateral and mesial surfaces, consisting of low crests ending in sharp teeth distally; lateroventral surface with distinct, rounded shoulder; pollex shallowly excavated on cutting edge; dactylus somewhat flattened and twisted laterally, not conspicuously broadened, only slightly convex dorsally; male and female major chelipeds similar in shape and in proportions; ventromesial surface of ischium with small spiniform seta; ventromesial margin of merus straight, ending in stout, sharp distomesial tooth, and with small, widely spaced spiniform setae; palm somewhat compressed, with strong sculpture on lateral and mesial surfaces, consisting of low crests ending in sharp teeth distally; lateroventral surface with rounded, smooth, non-projecting shoulder adjacent to deep notch, latter continuing transversely to shallow groove on mesial surface; dorsal margin with subcylindrical elevation ending in large adhesive disk distally; distomesial surface with transversally deeply notched crest ending in sharp tooth; pollex shorter than dactylus, somewhat twisted and shallowly depressed laterally, cutting edge bluntly projecting laterally; dactylus flattened, twisted laterally, convex dorsally, bulbous distally, plunger reduced to broad, low tooth; second pereopod with five-articulated carpus, ratio of articles approximately equal to 4 : 2 : 1 : 1.5 : 2; third and fourth pereopods similar; ischium armed with spiniform seta on ventrolateral surface; merus about five times as long as wide, without distoventral tooth; propodus with about eight spiniform setae along ventral margin and additional pair of spiniform setae close to propodo-dactylar articulation; dactylus about 0.4 length of propodus, simple, conical, faintly curved, with acute tip; pleopods with protopods furnished with spiniform setae on lateral margin, some inserted in pairs; male second pleopod with appendix masculina subequal in length to appendix interna, not reaching distal margin of endopod; uropodal exopod with sinuous diaeresis and small distolateral spiniform seta; uropodal endopod with row of small spiniform setae along distolateral margin; telson subrectangular, tapering posteriorly, about twice as long as wide at base; dorsal surface with two pairs of strong spiniform setae, first pair anterior to telson mid-length, second pair at about 0.7 telson length; posterior margin broadly convex, with two pairs of posterolateral spiniform setae, mesial about twice as long as lateral; anal tubercles well developed; gill–exopod formula typical for genus.




Figure 2. *Alpheus cedrici* sp. n. Holotype, male from Ascension Island (OUMNH.ZC. 2008-11-0017).

A minor (right) cheliped, lateral **B** minor (right) chela and carpus, mesial **C** same, distal palm and fingers, lateral, setae omitted **D** major (left) cheliped, lateral **E** major (right) cheliped, mesial **F** same, distal palm and fingers, lateral, setae omitted.

Description. Body stout, laterally not compressed. Carapace glabrous; frontal margin with well-developed rostrum and orbital teeth; rostrum slightly flattened dorsally, tapering distally, with acute tip, much longer than wide; lateral margins without setae; tip reaching half-length of first article of antennular peduncle; rostral carina not distinct; orbital teeth in marginal position, relatively small, acute distally, shorter than rostrum (note: right orbital hood atypical, i.e. without tooth in male); margin between orbital teeth and rostrum broadly V-shaped; orbital hoods moderately swollen, enclos-

ing eyes from all sides (Fig. 1A, B); rostro-orbital process present. Pterygostomial angle rounded, not protruding anteriorly (Fig. 1B); cardiac notch deep. Abdominal somites with posteroventral margins broadly rounded, fifth slightly more angular; sixth somite without articulated flap, bluntly projecting posteriorly.

Eyes with well-developed corneas; anteromesial margin bluntly protruding. Ocellar beak projecting, acute, visible in lateral view. Epistomial sclerites not acutely projecting.

Antennule with moderately slender peduncle; stylocerite not reaching distal margin of first article, with acute tip; ventromesial carina with large, subtriangular tooth as illustrated (Fig. 1C); second article much longer than dorsally visible portion of first article, about 2.5 times as long as wide (Fig. 1A); lateral flagellum with groups of aesthetascs starting from 12th article. Antenna with basicerite terminating in sharp distoventral tooth; carpocerite slightly exceeding both scaphocerite and antennular peduncle; scaphocerite with shallowly concave lateral margin and large, stout distolateral tooth, latter reaching far beyond distal margin of blade (Fig. 1A).

Mouthparts (not dissected) not specific in external view. Third maxilliped rather slender; coxa with lateral plate somewhat truncate distally; exopod long, overreaching distal margin of antepenultimate article; antepenultimate article somewhat flattened, ventral margin densely setose; penultimate article no more than three times as long as greatest width, distally slightly widening, very setose; ultimate article slender, tapering distally, with rows of serrulate setae and long, simple setae, tip unarmed (Fig. 1D).

Male minor cheliped with short, stout ischium; merus broad, subtriangular in cross-section; ventrolateral margin smooth; ventromesial margin straight, ending in small, acute distomesial tooth, and with four minute spiniform setae roughly equidistantly spaced along 0.6–0.7 of merus margin, and with tips falling just short of margin (therefore invisible in lateral view); distodorsal angle blunt; carpus rounded, cup-shaped; chela strongly compressed, with palm sculptured distally; lateral surface with low crest starting at about mid-length of palm and ending in a sharp distolateral tooth; ventral margin with blunt, non-protruding shoulder and adjacent deep notch, latter continuing transversely forming a shallow depression on mesial surface; dorsal margin with subcylindrical elevation ending distally in small adhesive disk; distomesial surface with crest ending in stout sharp tooth; fingers as long as palm; pollex shallowly excavated on cutting edge; dactylus somewhat flattened and twisted laterally, slightly convex dorsally, proximally with small adhesive disk (Fig. 2A–C). Female minor cheliped unknown (missing in the paratype).

Male major cheliped with short, stout ischium, ventromesial surface with small spiniform seta; merus stout, short, broad, subtriangular in cross-section; ventrolateral margin smooth; ventromesial margin straight, ending in stout, sharp distomesial tooth, and with small, widely spaced spiniform setae; dorsal margin ending bluntly distally; carpus very short, cup-shaped; chela somewhat compressed; palm strongly sculptured; lateral surface with low crest starting at about 0.6 length of palm and ending in sharp distolateral tooth; ventral margin with rounded, smooth, non-projecting shoulder adjacent to deep notch, latter continuing transversely to shallow groove on mesial surface; dorsal margin with subcylindrical elevation ending in large adhesive disk dis-




Figure 3. *Alpheus cedrici* sp. n. Holotype, male from Ascension Island (OUMNH.ZC. 2008-11-0017). **A** dorsal view **B** lateral view (photographs by S. De Grave).

tally; distomesial surface with transversally deeply notched crest ending in sharp tooth; pollex shorter than dactylus, somewhat twisted and shallowly depressed laterally, cutting edge bluntly projecting laterally; dactylus flattened, twisted laterally, convex dorsally, bulbous distally, plunger reduced to broad, low tooth (Fig. 2D–F). Female major cheliped generally similar in shape and proportions to male major cheliped.

Second pereiopod elongate, slender; ischium slightly longer than merus; carpus with five articles with ratio approximately equal to 4 : 2 : 1 : 1.5 : 2; chela simple, fingers with scarce tufts of setae (Fig. 1E). Third and fourth pereiopods generally similar,

moderately slender; third pereopod with ischium armed with spiniform seta on ventrolateral surface; merus about five times as long as wide, without distoventral tooth; carpus more slender and about half as long as merus; propodus longer than carpus, with eight or so robust spiniform setae along ventral margin and one pair of longer spiniform setae adjacent to dactylus; dactylus about 0.4 length of propodus, simple, conical, faintly curved, with acute tip (Fig. 1F). Fifth pereopod much more slender than third and fourth pereopods; merus only slightly longer than carpus; ischium with spiniform seta; propodus with some spiniform setae along ventral margin and well-developed setal brush distolaterally (Fig. 1G).

Pleopods with protopods furnished with spiniform setae on lateral margin, some inserted in pairs (Fig. 1H, J, K), first pleopod with small endopod furnished with setae, male second pleopod with appendix masculina subequal in length to appendix interna, not reaching distal margin of endopod, furnished with numerous stiff setae (Fig. 1I); female second pleopod with appendix interna only. Uropod with lateral lobe of protopod ending in large, acute tooth; diaeresis sinuous, with blunt tooth adjacent to one (occasionally two) stout distolateral spiniform seta(e); endopod with row of small spiniform setae along distolateral margin (Fig. 1L).

Telson subrectangular, tapering towards posterior margin, about twice as long as wide at base; lateral margins slightly convex; dorsal surface with two pairs of strong spiniform setae inserted at some distance from lateral margin, first pair anterior to telson mid-length, second pair at about 0.7 telson length; posterior margin about 0.6 length of anterior margin, broadly convex, with two pairs of posterolateral spiniform setae, mesial about twice as long as lateral (Fig. 1M); anal tubercles well developed.

Gill–exopod formula typical for *Alpheus*: five pleurobranchs (above P1–5), one arthrobranch (Mxp3), two lobe-shaped epipods (Mxp1–2), five mastigobranchs (Mxp3, P1–4), five setobranchs (P1–5); three exopods (Mxp1–3).

Size. *Alpheus cedrici* sp. n. is a medium-sized species of *Alpheus*, with 10.1 mm cl for the male, and 11.8 mm for the ovigerous female.

Colour. Body ground colour bright red or red–orange; carapace mostly red with transverse white band along posterior margin and several colourless or whitish areas on flanks; abdomen mostly red with transverse, more or less oval-shaped, white bands, latter mainly dorsal and not extending to ventral margins of pleura, additional colourless or whitish patches present near ventral margin of each pleuron; major chelae orange–brown marbled with pale yellow on mesial side, and with a distinct, somewhat zigzag-shaped, transverse, white band on palm, extending ventrally and posteriorly; dactylus pale brown with white tip; minor chela similar to major chela, orange–brown with transverse white bands on palm, a broader, more diffuse distal band, and smaller, well-delimited, V-shaped, proximal band; second to fifth pereopods pale reddish to yellowish, with white articulations; pleopods red; uropods and telson mostly red except for white uropodal protopods and most proximal portion of telson (Fig. 3).

Etymology. Named after our friend and colleague, Dr. Cedric d’Udekem d’Acoz, in recognition of his important contribution to the taxonomy of caridean shrimp and other decapods, particularly in the Atlantic Ocean.

Habitat. Both specimens were collected by lifting large, shallowly buried rocks on a fine sandy substrate at depths between 10 and 15 m, where the marginal boulder talus meets the sand. Other decapods obtained in the same habitat were the alpheid *Automate* cf. *dolichognatha*, *Alpheus holthuisi*, *Metalpheus paragracilis*, and the axiid *Axiopsis* cf. *serratifrons* (A. Milne-Edwards, 1873).

Type locality. English Bay, Ascension Island, central Atlantic Ocean.

Distribution. Central Atlantic Ocean: currently known only from the type locality.

Remarks. *Alpheus cedrici* sp. n. belongs to the monophyletic *Alpheus macrocheles* species group, which is comprised of about 30 species worldwide, all sharing a unique sculpture of the major chela (e.g. Coutière 1905; De Man 1911; Crosnier and Forest 1966; Banner 1953; Banner and Banner 1982; Kim and Abele 1988). Within the *A. macrocheles* species group, *A. cedrici* sp. n. belongs to the *A. macrocheles* species complex, characterised by the presence of acuminate orbital teeth on the frontal margin of the carapace; the major cheliped bearing a well-developed ventral notch and a dorsomesial notch or constriction; and the third and fourth pereopods (P3–4) with unarmed meri and simple or minutely biunguiculate (not conspicuously biunguiculate) dactyli. The majority of species in the *A. macrocheles* complex are found in the Atlantic Ocean: *A. macrocheles* (EA, CA), *A. platydactylus* Coutière, 1897 (EA), *A. amblyonyx* Chace 1972 (WA), *A. lentiginosus* Anker & Nizinski, 2011 (WA), *A. puapeba* Christoffersen, 1979 (WA), *A. pouang* Christoffersen, 1979 (WA), and *A. cedrici* sp. n. (CA). Two species are distributed in the eastern Pacific: *A. bellimanus* Lockington, 1877 (EP), and *A. rectus* Kim & Abele, 1988 (EP). Finally, only one Indo-West Pacific species presents the above combination of characters: *A. albatrossae* (Banner, 1953). All these species are contrasted and compared with the new species below, in order of geographical proximity.

Alpheus macrocheles is a well-known, mostly shallow-water species (0–50 m, exceptionally to 185 m), ranging in the eastern Atlantic from the British Isles and Mediterranean Sea south to Gabon, and extending to the Central Atlantic islands of Ascension and St. Helena (Holthuis 1951; Crosnier and Forest 1966; Chace 1966; Manning and Chace 1990). *Alpheus cedrici* sp. n. can be separated from *A. macrocheles* by the presence of a row of spiniform setae on the protopods of the pleopods (absent in *A. macrocheles*); the scaphocerite with a better developed blade (cf. Fig. 1A and Crosnier and Forest 1966, fig. 2a); and the male minor chela being more slender and with the dactylus less expanded and less arched dorsally (cf. Figs. 2A–C and Crosnier and Forest 1966, fig. 2c). The two species also differ in their colour patterns: the white bands and patches on the abdomen of *A. cedrici* sp. n. are contrasting with the mostly uniform deep-red to bright or pale orange abdomen of *A. macrocheles* (Fig. 4A, B). All records of *A. macrocheles* from the western Atlantic, e.g. records from Brazil (Ramos-Porto 1979; Guterres et al. 2005), have to be treated with some caution as they may refer to *A. amblyonyx* or other species of the *A. macrocheles* complex.

Alpheus platydactylus is a poorly known deep-water species (50–600 m) restricted to the northeastern Atlantic (Mediterranean Sea to the Azores and Cape Verde). *Alpheus cedrici* sp. n. can be easily distinguished from *A. platydactylus* by the much stout-




Figure 4. Colour patterns of some species of the *Alpheus macrocheles* (Hailstone, 1835) complex. **A** *Alpheus macrocheles* from Cadaques, Mediterranean coast of Spain **B** *Alpheus macrocheles* from Madeira **C** *Alpheus amblyonyx* Chace, 1972 from Guadeloupe **D** *Alpheus amblyonyx* from Isla Grande, Panama **E** *Alpheus bellimanus* Lockington, 1877 from Santa Barbara, California **F** *Alpheus bellimanus* from Galapagos **G** *Alpheus* sp. ? *rectus* Kim & Abele, 1988, from the Gulf of California. Photographic credits: A, Josep Lluís Peralta; B, Peter Wirtz; C, Frédéric Fasquel; D, Arthur Anker; E, Gregory Jensen; F, Todd Zimmerman (courtesy of Cleveland Hickman); G, Alex Kerstitch (from Kerstitch 1988, courtesy of A. Kerstitch).

er, shorter antennular peduncles; the broader scaphocerite; the smooth ventral margin of the major chela palm (vs. rugose in *A. platydactylus*); the stouter minor chela, with the relatively shorter fingers; and the less slender second pereopod (cf. Figs. 1, 2 and Crosnier and Forest 1966, fig. 2e–h).

Alpheus amblyonyx, a species widespread in the western Atlantic from the Gulf of Mexico to Brazil (Chace 1972; Christoffersen 1998), is generally very similar to *A. macrocheles*. Chace (1972) separated *A. amblyonyx* from *A. macrocheles* mainly on the basis of four characters: (1) the more prominent rostrum; (2) the transverse notch on the mesiodorsal surface of the major chela palm broader and less sharply defined; (3) the major chela dactylus more strikingly bulbous distally; and (4) the minor chela dactylus without a high dorsal crest. With the exception of the last feature, *A. cedrici* sp. n. can be separated from *A. amblyonyx* using the same criteria as from *A. macrocheles* (see above). The colour pattern of *A. amblyonyx* (Fig. 4C, D) is much more similar to the colour pattern of *A. macrocheles* (Fig. 4A, B) than to that of *A. cedrici* sp. n. (Fig. 3).

Alpheus pouang and *A. puapeba* are two deep-water species presently known only from the southwestern Atlantic, off southern Brazil and Uruguay, at depth ranges of 120–268 m and 45–175 m, respectively (Christoffersen 1979, 1998). *Alpheus cedrici* sp. n. can be distinguished from *A. pouang* by the anterior margin of the carapace between the rostrum and the orbital teeth being shallowly and broadly concave (vs. much more deeply incised in *A. pouang*), and the minor chela with a non-protruding ventral shoulder and a distinctly less flattened and dorsally arched dactylus (cf. Figs. 1–2 and Christoffersen 1979, figs. 14–15). The new species differs even more from *A. puapeba*, for example, by the much shorter antennular peduncles and the less elongate, more swollen major chela (cf. Figs. 1–2 and Christoffersen 1979, figs. 16–17). In addition, the pleopodal protopods of both *A. pouang* and *A. puapeba* are not armed with rows of spiniform setae (Christoffersen 1979, figs. 15r, 17d, 18f), as is the case of *A. cedrici* sp. n. (Fig. 1j).

Alpheus lentiginosus is another deep-water western Atlantic species presently known only from the northern Gulf of Mexico, at depths of 336–438 m (Anker and Nizinski 2011). *Alpheus cedrici* sp. n. can be separated from *A. lentiginosus* by the less expanded, dorsally non-arched dactylus of the minor chela; the less slender third to fifth pereopods, with simple, conical dactyli (vs. with a minute accessory unguis on the flexor margin in *A. lentiginosus*); and the presence of spiniform setae on the pleopodal protopods (absent in *A. lentiginosus*) (cf. Figs 1–2 and Anker and Nizinski 2011, figs. 1–2). The colour patterns of *A. cedrici* sp. n. and *A. lentiginosus* are different as well (cf. Fig. 3 and Anker and Nizinski 2011, fig. 3).

The two eastern Pacific species of the *A. macrocheles* complex, *A. bellimanus* and *A. rectus*, are both morphologically very close to *A. cedrici* sp. n. *Alpheus bellimanus* is a relatively common species with a very wide depth range (0–300 m), and also with a wide geographic range, from California via Mexico, Panama and Galapagos to northern Chile (Kim and Abele 1988). *Alpheus rectus* is a much less common species from moderately deep-water (55–73 m); it is currently known only from the type locality in Panama and one locality in southern Baja California (Kim and Abele 1988). *Alpheus cedrici* sp. n. shares with *A. bellimanus* the presence of spiniform setae on the protopods of pleopods. The two species also have very similar frontal margins of the carapace, antennules and antennae, major chelipeds, and walking legs. However, *A. cedrici* sp. n. can be separated from *A. bellimanus* by the non-protruding ventral shoulder of the male minor chela (vs. protruding in *A. bellimanus*); the less expanded, dorsally non-arched

dactylus of the male minor chela (vs. more expanded and dorsally strongly convex in *A. bellimanus*); and the anteriorly rounded tooth on the ventromesial carina of the first article of the antennular peduncle (vs. with a subacute tooth in *A. bellimanus*) (cf. Figs. 1–2 and Kim and Abele 1988, fig. 5). The colour pattern of *A. bellimanus* (Fig. 4E, F) resembles more the uniform colour patterns of *A. macrocheles* (Fig. 4A, B) and *A. amblyonyx* (Fig. 4C, D) than the distinctly banded colour pattern of *A. cedrici* sp. n. (Fig. 3).

The new species from Ascension also differs from *A. rectus*, for example, by the less rectangular general shape of the major chela, with the ventral shoulder of the palm broadly rounded, not protruding anteriorly (vs. bluntly projecting in *A. rectus*); and the anteriorly rounded tooth on the ventromesial carina of the first article of the antennular peduncle (vs. with a small point in *A. rectus*). The colour pattern of *A. rectus* remains unconfirmed. However, a colour photograph of a snapping shrimp erroneously identified as “*Alpheus paracrinatus*” in Kerstitch (1988) matches *A. rectus*, especially in the shape of the major and minor chelipeds. The colour pattern of this individual (*Alpheus* sp. ? *rectus* in Fig. 4G), although characterised by a conspicuous transversal orange-white banding, is different from that of *A. cedrici* sp. n., especially in the clearly banded carapace and the abdominal bands extending ventrally to the pleural margins (cf. Figs. 3, 4G).

All other species of the *A. macrocheles* group present in the western and eastern Atlantic and in the eastern Pacific differ more markedly from *A. cedrici* sp. n. (see Coutière 1910; Armstrong 1940, 1941; Holthuis 1951; Crosnier and Forest 1966; Chace 1972; Wicksten and Méndez 1981; Kim and Abele 1988; Wicksten and McClure 2003; Anker et al. 2008). The shape of the frontal margin of the carapace separates the new species from *A. inca* Wicksten & Méndez, 1981, *A. grahami* Abele, 1975, *A. cylindricus* Kingsley, 1878, *A. vanderbilti* Boone, 1930, *A. clamator* Lockington, 1877, *A. peasei* (Armstrong, 1940), *A. dentipes* Guérin, 1832 and *A. candei* Guérin-Méneville, 1855). The shape of the major cheliped separates the new species from *A. crockeri* (Armstrong, 1941), *A. hortensis* Wicksten & McClure, 2003, *A. grahami*, *A. cylindricus* and *A. vanderbilti*. The shape of the minor cheliped separates the new species from *A. hoonsooi* Kim & Abele, 1988, *A. crockeri*, *A. hortensis*, *A. grahami*, *A. cylindricus* and *A. vanderbilti*. The presence of a distinct distoventral tooth on the merus of the third and fourth pereopod in *A. hoonsooi*, *A. clamator*, *A. peasei* and *A. dentipes* separates these species from *A. cedrici* sp. n. Finally, the strongly biunguiculate dactylus of the third to fifth pereopods in *A. clamator*, *A. peasei*, *A. dentipes* and *A. candei* separates these species from *A. cedrici* sp. n.

Alpheus cedrici sp. n. can be separated from the Indo-West Pacific *A. albatrossae* by the presence of a distinct shoulder on the ventrolateral surface of the minor chela palm (absent in *A. albatrossae*); the stouter fingers of the minor chela; and the absence of a small unguis on the dorsal margin of the dactylus of the third to fifth pereopods (present in *A. albatrossae*) (cf. Figs. 1, 2 and Banner, 1953, fig. 18). None of the other Indo-West Pacific species of the *A. macrocheles* group appears to be closely related to *A. cedrici* sp. n. (e.g. Coutière 1905; De Man 1911; Banner 1953; Kensley 1969; Banner and Banner 1982; Burukovsky 1990).

Acknowledgments

Financial support for SDG's fieldwork in Ascension Island was provided by the Oxford University John Fell Fund, under a permit issued by the Ascension Conservation Department. H. De Grave assisted with and made fieldwork more enjoyable. AA is grateful to CAPES (Coordenação de Aperfeiçoamento de Pessoal de Nível Superior) for financial support in the form of a postdoctoral fellowship. Additional photographs used in this study were taken by Josep Lluís Peralta (Mataró, Spain), Gregory Jensen (University of Washington, USA), Peter Wirtz (Universidade do Algarve, Portugal), Todd Zimmerman / Cleveland P. Hickman (Lexington, USA), and the late Alex Kerstitch.

References

- Abele LG (1975) The macruran decapod Crustacea of Malpelo Island. *Smithsonian Contributions to Zoology* 176: 69–85.
- Anker A, Hurt C, Jara JA, Knowlton N (2008) Revision of the *Alpheus cylindricus* Kingsley, 1878 species complex (Crustacea: Decapoda: Alpheidae), with revalidation of *A. vanderbilti* Boone, 1930. *Zootaxa* 1943: 53–68.
- Anker A, Nizinski M (2011) Description of a new deep-water species of *Alpheus* Fabricius, 1798 from the Gulf of Mexico (Crustacea, Decapoda, Alpheidae). *Zootaxa* 2925: 49–56.
- Armstrong JC (1940) New species of Caridea from the Bermudas. *American Museum Novitates* 1096: 1–10.
- Armstrong JC (1941) The Caridea and Stomatopoda of the second Templeton Crocker–American Museum expedition to the Pacific Ocean. *American Museum Novitates* 1137: 1–14.
- Banner AH (1953) The Crangonidae, or snapping shrimp, of Hawaii. *Pacific Science* 7: 3–147.
- Banner DM, Banner AH (1982). The alpheid shrimp of Australia. Part III: The remaining alpheids, principally the genus *Alpheus* and the family Ogyrididae. *Records of the Australian Museum* 34: 1–357. doi: 10.3853/j.0067-1975.34.1982.434
- Boone L (1930) Scientific results of the cruises of the yachts “Eagle” and “Ara”, 1921–1928, William K. Vanderbilt, commanding. Crustacea: Anomura, Macrura, Schizopoda, Isopoda, Amphipoda, Mysidacea, Cirripedia, and Copepoda. *Bulletin of the Vanderbilt Marine Museum* 3: 1–221.
- Burukovsky RN (1990) Shrimps from the Sala-y-Gomez and Nazca Ridges [in Russian]. *Trudy Instituta Okeanologii Akademii Nauk SSSR* 124: 187–217.
- Chace FAJr (1966) Decapod crustaceans from St. Helena, South Atlantic. *Proceedings of the United States National Museum* 118: 622–662.
- Chace FAJr (1972) The shrimps of the Smithsonian–Bredin Caribbean expeditions with a summary of the West Indian shallow-water species (Crustacea: Decapoda: Natantia). *Smithsonian Contributions to Zoology* 98: 1–179.
- Christoffersen ML (1979) Campagne de la Calypso au large des côtes Atlantiques de l’Amerique du Sud (1961–1962). I. Decapod Crustacea: Alpheoida. *Annales de l’Institut Océanographique* 55(Suppl.): 297–377.

- Christoffersen ML (1998) Malacostraca. Eucarida. Caridea, Crangonoidea and Alpheoidea (except Glyphocrangonidae and Crangonidae). In: Young PS (Ed) Catalogue of Crustacea of Brazil. Museu Nacional, Rio de Janeiro, 351–372.
- Coutière H (1897) Note sur quelques Alphées nouveaux. Bulletin du Muséum d'Histoire Naturelle, serie 3 7: 303–306.
- Coutière H (1905) Les Alpheidae. In: Gardiner JS The Fauna and Geography of the Maldive and Laccadive Archipelagoes. Being the account of the work carried on and of the Collections made by an Expedition during the years 1899 and 1900. University Press, Cambridge, 852–921, pls. 70–87.
- Coutière H (1909) The American species of snapping shrimps of the genus *Synalpheus*. Proceedings of the United States National Museum 36: 1–93. doi: 10.5479/si.00963801.36-1659.1
- Coutière H (1910) The snapping shrimps (Alpheidae) of the Dry Tortugas, Florida. Proceedings of the United States National Museum 37: 485–487. doi: 10.5479/si.00963801.37-1716.485
- Crosnier A, Forest J (1966) Résultats Scientifiques des Campagnes de la “Calypso”. Fascicule VII. Campagne de la Calypso dans la Golfe de Guinée et aux Iles Principe, São Tomé et Annobon (1956), et campagne aux Iles du Cap Vert (1959) (suite). 19. Crustacés décapodes: Alpheidae. Annales de l'Institut Océanographique 44: 199–314.
- Fabricius JC (1798) Supplementum Entomologiae Systematicae. Proft et Storch, Hafniae, 572 pp.
- Guérin FE (1832) I.^{re} Classe. Crustacés. In: Brullé A (Ed) Expédition Scientifique de Morée. Section des Sciences Physiques. Tome III. – 1ère Partie. Zoologie. Deuxième Section. – Des animaux articulés. Paris, 30–50, Plate 27.
- Guérin-Méneville FE (1855–1856) Animales articulados con piés articulados. In: de la Sagra R (Ed) Historia física política y natural de la isla de Cuba. Segunda Parte. Historia Natural. Tomo VII (Crustaceos, Aragnides é Insectos) [1856]; Tomo VIII (Atlas de Zoologia) [1855]. Paris, i–xxxii, 1–371, pls. 1–20.
- Guterres LFR, Melo GAS, Castro-Guterres P (2005) Novos registros de ocorrência de *Alpheus macrocheles* (Crustacea, Caridea, Alpheidae) na costa do Brasil. Biociências 13: 231–233.
- Hailstone S (1835) The species of crustaceous animals discovered and described by Mr. Hailstone, and illustrated and annotated upon by Mr. Westwood. The Magazine of Natural History and Journal of Zoology, Botany, Mineralogy, Geology, and Meteorology 8: 394–395.
- Holthuis LB (1951) The caridean Crustacea of tropical West Africa. Atlantide Report 2: 1–187.
- Kensley B (1969) Decapod Crustacea from the south-west Indian Ocean. Annals of the South African Museum 52: 149–181.
- Kerstitch A (1988) Sea of Cortez marine invertebrates. A guide for the Pacific coast, Mexico to Ecuador. Sea Challengers, Monterey, 113 pp.
- Kim W, Abele LG (1988) The snapping shrimp genus *Alpheus* from the Eastern Pacific (Decapoda: Caridea: Alpheidae). Smithsonian Contributions to Zoology 454: 1–119. doi: 10.5479/si.00810282.454
- Kingsley JS (1878) A synopsis of the North American species of the genus *Alpheus*. Bulletin of the United States Geological and Geographical Survey 4: 189–199.
- Lockington WN (1877) Remarks on the Crustacea of the Pacific Coast, with descriptions of some new species. Proceedings of the California Academy of Sciences 7 [for 1876]: 28–36.

- Man JG de (1888) Bericht über die von Herrn Dr. J. Brock im indischen Archipel gesammelten Decapoden und Stomatopoden. Archiv für Naturgeschichte 53: 289–600.
- Man JG de (1910) Diagnoses of new species of macrurous decapod Crustacea from the “Siboga-Expedition”. Tijdschrift der Nederlandse Dierkundige Vereniging, series 2 11: 287–319.
- Man JG de (1911) The Decapoda of the Siboga Expedition. Part II. Family Alpheidae. Siboga Expeditie 39a1: 133–465.
- Manning RB, Chace FAJr (1990) Decapod and stomatopod Crustacea from Ascension Island, South Atlantic Ocean. Smithsonian Contributions to Zoology 503: 1–91. doi: 10.5479/si.00810282.503
- Miers EJ (1881) On a collection of Crustacea made by Baron Hermann-Maltzan at Goree island, Senegambia. The Annals and Magazine of Natural History, series 5 8: 204–220, 259.
- Milne-Edwards A (1873) Description de quelques crustacés nouveaux ou peu connus provenant du Musée de M. C. Godeffroy. Journal des Museum Godeffroy 1: 77–88.
- Milne-Edwards A (1878) Description de quelques espèces nouvelles de Crustacés provenant du voyage aux îles du Cap-Vert de MM. Bouvier et de Cessac. Bulletin de la Société Philomathique de Paris, serie 7 2: 225–232.
- Pocock RI (1890) Crustacea. The Journal of the Linnean Society. Zoology 20: 506–526.
- Rafinesque CS (1815) Analyse de la Nature ou Tableau de l’Univers et des corps organisés. Palerme, 224 pp.
- Ramos-Porto M (1979) Ocorrência de *Alpheus macrocheles* (Hailstone) (Crustacea: Decapoda: Alpheidae) no Norte e Nordeste do Brasil. Trabalhos do Instituto Oceanográfico da Universidade Federal do Pernambuco 14: 117–130.
- Ribeiro A (1964) Sobre uma espécie nova de *Alpheus* Fabricius, 1798 do Arquipélago de Cabo Verde, *Alpheus holthuisi* n. sp. Notas Mimeografadas do Centro de Biologia Piscatória, Lisboa 42: 1–14.
- Schmitt WL (1936) Zoologische Ergebnisse einer Reise nach Bonaire, Curaçao und Aruba im Jahre 1930. No. 16. Macruran and anomuran Crustacea from Bonaire, Curaçao und Aruba. Zoologische Jahrbücher 67: 363–378.
- Wicksten MK, McClure MR (2003) A new species of *Alpheus* (Decapoda: Caridea: Alpheidae) from the Gulf of Mexico. Crustacean Research 32: 26–31.
- Wicksten MK, Méndez MG (1981) *Alpheus inca*: a new snapping shrimp (Caridea: Alpheidae) from western South America. Journal of Crustacean Biology 1: 137–142. doi: 10.2307/1548211

