

New Staphylinidae (Coleoptera) records with new collection data from New Brunswick, Canada: Paederinae

Reginald P. Webster¹, Ian DeMerchant¹

¹ Natural Resources Canada, Canadian Forest Service - Atlantic Forestry Centre, 1350 Regent St., P.O. Box 4000, Fredericton, NB, Canada E3B 5P7

Corresponding author: Reginald P. Webster (reginaldwebster@rogers.com)

Academic editor: J. Klimaszewski | Received 6 December 2011 | Accepted 26 January 2012 | Published 26 April 2012

Citation: Webster RP, DeMerchant I (2012) New Staphylinidae (Coleoptera) records with new collection data from New Brunswick, Canada: Paederinae. In: Klimaszewski J, Anderson R (Eds) Biosystematics and Ecology of Canadian Staphylinidae (Coleoptera) II. ZooKeys 186: 273–292. doi: 10.3897/zookeys.186.2504

Abstract

We report 17 species of Paederinae new for New Brunswick, Canada. Ten of these species, *Lathrobium othioides* LeConte, *Lathrobium amplipenne* Casey, *Lathrobium armatum* Say, *Lathrobium confusum* LeConte, *Lathrobium debile* LeConte, *Achenomorphus corticinus* (Gravenhorst), *Rugilus rufipes* Germar, *Homaeotarsus bicolor* (Gravenhorst), *Homaeotharsus cribratus* (LeConte), and *Homaeotarsus pallipes* (Gravenhorst) are newly recorded for the Maritime provinces. This brings the total number of Paederinae recorded from New Brunswick to 36 species. Additional records are presented for the recently reported *Lathrobium simile* LeConte and *Lathrobium washingtoni* Casey. Collection and habitat data are presented for all species.

Keywords

Paederinae, new records, Canada, New Brunswick

Introduction

This paper treats new Staphylinidae records from New Brunswick of the subfamily Paederinae. The most recent taxonomic treatments of the North American Paederinae fauna were by Casey (1905, 1910). More recently Watrous (1981) reviewed the *Tetaropeus* (as a subgenus of *Lathrobium*) and the subgenus *Eulathrobium* of *Lobrathium*. Herman (1965a, b) revised the genus *Orus*. Many other genera in North America are in need of a modern revision.

Many species of Paederinae in Canada occur in and near wetland habitats such as marshes, bogs, and pond and river margins (Watrous 1980, 1981; Newton et al. 2000; Brunke et al. 2011). A few species, such as *Sunius confluentus* (Say), occur in subcortical habitats (Brunke et al. 2011). However, relatively little has been published on the bionomics of species occurring in Canada or North America.

Campbell and Davies (1991) reported 15 species of Paederinae from New Brunswick. Klimaszewski et al. (2005) added *Lathrobium simile* LeConte, *Lathrobium. washingtoni* Casey, and *Ochtheophilum fracticorne* (Paykull) from their study on rove beetles in red spruce (*Picea rubens* Sarg.) stands. The adventive *Lathrobium fulvipenne* (Gravenhorst) was reported from New Brunswick by Majka and Klimaszewski (2008). Here, we report an additional 17 species of Paederinae for New Brunswick, bringing the total number of species known from the province to 36.

Methods and conventions

The following records are based on specimens collected as part of a general survey to document the Coleoptera fauna of New Brunswick and from by-catch samples from Lindgren 12-funnel traps (Lindgren 1983) obtained during a study to develop a general attractant for the detection of invasive species of Cerambycidae.

Collection methods

Various collection methods were employed to collect the species reported in this study. Details are outlined in Campbell (1973) and Webster et al. (2009, Appendix). See Webster et al. (2012) for details of the methods used for deployment of Lindgren traps and sample collection. A description of the habitat was recorded for all specimens collected during this survey. Locality and habitat data are presented exactly as recorded on labels for each specimen. This information, as well as additional collecting notes, is summarized and discussed in the collection and habitat data section for each species.

Specimen preparation

Males and some females of most species were dissected to confirm their identity. The genital structures were dehydrated in absolute alcohol, mounted in Canada balsam on celluloid microslides, and pinned with the specimens from which they originated.

Distribution

Distribution maps, created using ArcMap and ArcGIS, are presented for each species in New Brunswick. Every species is cited with current distribution in Canada and Alaska, using abbreviations for the state, provinces, and territories. New provincial

records are indicated in bold under Distribution in Canada and Alaska. The following abbreviations are used in the text.

AK	Alaska	MB	Manitoba
YT	Yukon Territory	ON	Ontario
NT	Northwest Territories	QC	Quebec
NU	Nunavut	NB	New Brunswick
BC	British Columbia	PE	Prince Edward Island
AB	Alberta	NS	Nova Scotia
SK	Saskatchewan	NF & LB	Newfoundland and Labrador*

*Newfoundland and Labrador are each treated separately under the current Distribution in Canada and Alaska.

Acronyms of collections examined and referred to in this study are as follows:

- AFC** Atlantic Forestry Centre, Natural Resources Canada, Canadian Forest Service, Fredericton, New Brunswick, Canada
CNC Canadian National Collection of Insects, Arachnids and Nematodes, Agriculture and Agri-Food Canada, Ottawa, Ontario, Canada
NBM New Brunswick Museum, Saint John, New Brunswick, Canada
RWC Reginald P. Webster Collection, Charters Settlement, New Brunswick, Canada

Results

Unless noted otherwise (additional records), all records below are species newly recorded for New Brunswick, Canada. Species followed by ** are newly recorded from the Maritime provinces (New Brunswick, Nova Scotia, Prince Edward Island) of Canada.

Species accounts

The classification of the Paederinae follows Bouchard et al. (2011).

Seventeen species of Paederinae are newly reported for the New Brunswick, Canada. Ten of these are newly recorded for the Maritime provinces. This brings the total number of Paederinae recorded from New Brunswick to 36 (Table 1). Additional records and bionomic data of the recently reported *Lathrobium simile* and *L. washingtoni* are presented.

Table 1. Species of Paederinae (Staphylinidae) recorded from New Brunswick, Canada.

Subfamily Paederinae Fleming	<i>Lathrobium (Lathrobioma) othioides</i> LeConte**
Tribe Paederini Fleming	<i>Lathrobium (Lathrobium) amplipenne</i> Casey**
Subtribe Lathrobiina Laporte	<i>Lathrobium (Lathrobium) armatum</i> Say**
<i>Lathrobium (Lathrobioma) scolopaceum</i> (Casey)	<i>Lathrobium (Lathrobium) confusum</i> LeConte**

<i>Lathrobium (Lathrobium) fauveli</i> Duvivier	<i>Pseudomedon thoracica</i> Casey
<i>Lathrobium (Lathrobium) fulvipenne</i> (Gravenhorst)	<i>Sunius (Trachysectus) confluentus</i> (Say)
<i>Lathrobium (Lathrobium) simile</i> LeConte	Subtribe Stilicina Casey
<i>Lathrobium (Lathrobium) sparsellum</i> Casey	<i>Pachystilicus hanhami</i> (Wickham)
<i>Lathrobium (Lathrobium) spissicorne</i> Casey*	<i>Rugilus angustatus</i> (Geoffrey)*
<i>Lathrobium (Lathrobium) washingtoni</i> Casey	<i>Rugilus biarmatus</i> (LeConte)
<i>Lathrobium (Lathrolepta) debile</i> LeConte**	<i>Rugilus rufipes</i> Germar**
<i>Lobrathium (Lobrathium) collar</i> (Erichson)*	Subtribe Astenina Hatch
<i>Lobrathium (Eulathrobium) grande</i> (LeConte)*	<i>Astenus discopunctatus</i> (Say)
<i>Tetartopeus angularis</i> (LeConte)	Subtribe Cryptobiina Casey
<i>Tetartopeus furvulus</i> Casey	<i>Homaeotarsus (Gastrolobium) bicolor</i> (Gravenhorst)**
<i>Tetartopeus lacustris</i> Casey	<i>Homaeotarsus (Hesperobium) cinctus</i> (Say)*
<i>Tetartopeus niger</i> (LeConte)	<i>Homaeotarsus (Hesperobium) cribratus</i> (LeConte)**
<i>Tetartopeus nitidulus</i> (LeConte)	<i>Homaeotarsus (Hesperobium) pallipes</i> (Gravenhorst)**
<i>Tetartopeus capitosus</i> Casey*	<i>Ochtheophilum fracticorne</i> (Paykull)
<i>Tetartopeus rubripennis</i> Casey*	Subtribe Paederina Fleming
Subtribe Medonina Casey	<i>Paederus littorarius</i> (Gravenhorst)
<i>Achenomorphus corticinus</i> (Gravenhorst)**	
<i>Lithocaris (Lithocharis) ochracea</i> (Gravenhorst)	

Notes: *New to province, **New to Maritime Provinces

Family Staphylinidae Latreille, 1802

Subfamily Paederinae Fleming, 1821

Tribe Paederini Fleming, 1821

Subtribe Lathrobiina Laporte, 1835

Lathrobium (Lathrobioma) othioides LeConte, 1880**

http://species-id.net/wiki/Lathrobium_othioides

Map 1

Material examined. **New Brunswick, Carleton Co.,** Richmond, Hovey Hill Protected (Natural) Area, 46.1115°N, 67.7770°W, 10.V.2005, R. P. Webster, hardwood forest, in moist leaf litter and moss near seepage area (3 ♂, RWC). **Sunbury Co.,** Acadia Research Forest, 45.9816°N, 66.3374°W, 18.VI.2007, R. P. Webster, 8.5 year-old regenerating mixed forest, sifting leaf litter (1 ♂, AFC). **York Co.,** Charters Settlement, 45.8428°N, 66.7279°W, 19.IV.2004, 24.IV.2004, 15.IV.2005, 20.IV.2005, R. P. Webster, mixed forest, sedge marsh in moist grass litter and sphagnum (6 ♂, RWC).

Collection and habitat data. *Lathrobium othioides* was found in an old hardwood forest, a regenerating mixed forest, and a sedge (*Carex*) marsh. Adults were sifted from leaf litter, moist leaf litter, and moss in a seepage area and from moist grass litter and sphagnum. This species was collected during April, May, and June.

Distribution in Canada and Alaska. ON, QC, NB (Campbell and Davies 1991).

Lathrobium (Lathrobium) amplipenne* Casey, 1905*

http://species-id.net/wiki/Lathrobium_amplipenne

Map 2

Material examined. **New Brunswick, Restigouche Co.,** Little Tobique River near Red Brook, 47.4465°N, 67.0689°W, 13.VI.2006, R. P. Webster, river margin in *Carex* hummock (1 ♂, 1 ♀, RWC). **Sunbury Co.,** Sheffield, Portobello Creek N.W.A., 45.8952°N, 66.2728°W, 7.V.2004, R. P. Webster, silver maple swamp, in leaf litter (1 ♀, RWC).

Collection and habitat data. Adults were sifted from a *Carex* hummock on a river margin and from leaf litter in a silver maple (*Acer saccharinum* L.) swamp. Adults were collected during May and June.

Distribution in Canada and Alaska. ON, NB (Campbell and Davies 1991).

Lathrobium (Lathrobium) armatum* Say, 1834*

http://species-id.net/wiki/Lathrobium_armatum

Map 3

Material examined. **New Brunswick, Carleton Co.** Jackson Falls, Bell Forest, 46.2150°N, 67.7201°W, 14.V.2006, R. P. Webster, river margin, in drift material near seepage area (1 ♂, 1 ♀, RWC). **York Co.,** Dumfries, Slagundy Dry Ponds, 45.8596°N, 67.1849°W, 8.VII.2006, R. P. Webster, large vernal pond, pond margin in moist leaves (1 ♂, 2 ♀, RWC); 8.5 km W of Tracy off Rt. 645, 45.6888°N, 66.8004°W, 22.V.2008, R. P. Webster, *Carex* marsh/flowage near slow flowing brook, in *Carex* hummocks (3 ♂, 1 ♀, RWC).

Collection and habitat data. *Lathrobium armatum* was sifted from drift material near a seepage area along a river margin, from *Carex* hummocks in a *Carex* marsh/flowage near a slow-flowing brook and from moist leaves on the margin of a large vernal pond. Adults were collected during May and July.

Distribution in Canada and Alaska. ON, QC, NB (Campbell and Davies 1991).

Lathrobium (Lathrobium) confusum* LeConte, 1880*

http://species-id.net/wiki/Lathrobium_confusum

Map 4

Material examined. **New Brunswick, Carleton Co.,** near Hovey Hill P.N.A., 46.1155°N, 67.7631°W, 10.V.2005, R. P. Webster, mixed forest, in (leaf) litter near small brook (1 ♂, RWC); Meduxnekeag Valley Nature Preserve, 46.1964°N, 67.6840°W, 31.V.2005, M.-A. Giguère & R. Webster, mixed forest, vernal pond margin in moist (leaf) litter (2 ♀, NBM, RWC); same locality and forest type but, 46.1976°N, 67.6850°W, margin of vernal pond, in moist leaves (1 ♂, NBM). **Queens**

Co., W of Jemseg at “Trout Creek”, 45.8227°N, 66.1240°W, 26.IV.2004, 9.V.2004, R. P. Webster, silver maple swamp, sifting (leaf) litter at bases of large trees (2 ♂, 1 ♀, RWC); same locality, forest type, and collector but 45.8231°N, 66.1245°W, 11.IV.2006, sifting litter from crotch of silver maple with multiple trunks (1 ♂, NBM). **Sunbury Co.**, Sheffield, Portobello Creek N.W.A., 45.8952°N, 66.2728°W, 7.V.2004, R. P. Webster, silver maple swamp, in leaf litter (1 ♀, RWC); Burton, Sunpoke Lake, 45.7575°N, 66.5736°W, 30.IV.2004, 10.IV.2006, R. P. Webster, red maple swamp, in leaf litter near slow stream (1 ♂, 3 ♀, NBM, RWC). **York Co.** Kelly’s Creek at Sears Rd., 45.8723°N, 66.8414°W, 7.VI.2008, R.P. Webster, alder swamp with red maples, in moist leaf and grass litter near (small) pool (1 ♂, RWC).

Collection and habitat data. Watrous (1980) reported *L. confusum* from the margin of an intermittent stream. In New Brunswick, adults were found in leaf litter or grass litter near small brooks and streams and in leaf litter along vernal pond and forest pool margins. Some were sifted from leaf litter at the base of large silver maples or from litter in the crotch of a silver maple with multiple trunks. Adults were found in mixed forests, silver maple swamps, red maple (*Acer rubrum* L.) swamps, and an alder (*Alnus* sp.) swamp with red maple. This species was collected during April, May, and June.

Distribution in Canada and Alaska. ON, QC, NB (Campbell and Davies 1991).

Lathrobium (Lathrobium) simile LeConte, 1863

http://species-id.net/wiki/Lathrobium_simile

Map 5

Material examined. Additional New Brunswick records, Carleton Co., Hovey Hill P.N.A., 46.1115°N, 67.7770°W, 10.V.2005, R. P. Webster, hardwood forest, in moist leaf litter and moss near forest pool (1 ♂, RWC); Meduxnekeag Valley Nature Preserve, 46.1956°N, 67.6803°W, 15.IX.2004, R. P. Webster, mixed forest, in decaying fungi (1 ♂, RWC); same locality, forest type and collector but 46.1976°N, 67.6850°W, 4.V.2006, R. P. Webster, margin of vernal pond, in moist leaf litter (1 ♂, RWC); Jackson Falls, Bell Forest, 46.2210°N, 67.7210°W, 11.V.2005, M.-A. Giguère & R. Webster, hardwood forest, in leaf litter near small brook (1 ♂, 2 ♀, RWC). **Sunbury Co.**, Acadia Research Forest, 30.VI.1999, G. Gesner, Strip Cut 8, Site 1, pitfall trap (1, AFC); Acadia Research Forest, 45.9799°N, 66.3394°W, 18.VI.2007, R. P. Webster, mature red spruce and red maple forest, sifting leaf litter (1 ♂, 1 ♀, RWC).

Collection and habitat data. Adults of this species were found in hardwood forests, mixed forests, and a mature red spruce forest. Adults were collected from moist leaf litter near forest pools or vernal ponds, near a small brook, and from the forest floor. One individual was collected from decaying fungi on the forest floor. This species was collected during May, June, and September.

Distribution in Canada and Alaska. MB, ON, QC, NB, NS (Campbell and Davies 1991; Klimaszewski et al. 2005). This species was first reported from New Brunswick by Klimaszewski et al. (2005) from the Acadia Research Forest.

***Lathrobium (Lathrobium) spissicorne* Casey, 1905**http://species-id.net/wiki/Lathrobium_spissicorne

Map 6

Material examined. **New Brunswick, Charlotte Co.,** near New River, 45.2118°N, 66.6179°W, 7.VII.2006, R. P. Webster, mixed forest, margin small pond, treading *Carex* hummock into water (1 ♀, RWC). **Queens Co.,** W of Jemseg at “Trout Creek”, 45.8240°N, 66.1220°W, 4.VI.2004, R. P. Webster, silver maple swamp, margin of vernal pond in moist leaf litter on muddy soil (1 ♀, RWC); Grand Lake near Scotchtown, 45.8762°N, 66.1816°W, 5.VI.2004, 1.VII.2004, 25.V.2006, R. P. Webster, lakeshore, old (sand) dune with oaks, under dead fish and under drift material (3 ♂, 1 ♀, NBM, RWC). **Sunbury Co.,** Burton, Sunpoke Lake, 45.7665°N, 66.5545°W, 15.V.2004, R. P. Webster, (red) oak and (red and silver) maple forest, in leaf litter (3 ♂, RWC).


Collection and habitat data. *Lathrobium spissicorne* was found in mixed forests, a silver maple swamp, along a lakeshore, and in a mature forest with red oak (*Quercus rubra* L.), red maple, and silver maple. Adults were found in a *Carex* hummock on a pond margin, in moist leaf litter on vernal pond margin, and in leaf litter on the forest floor. Some adults were found under drift material and under a dead fish on a lakeshore. This species was collected during May, June, and July.

Distribution in Canada and Alaska. ON, QC, NB, PE (Campbell and Davies 1991).


***Lathrobium (Lathrobium) washingtoni* Casey, 1905**http://species-id.net/wiki/Lathrobium_washingtoni

Map 7


Material examined. **Additional New Brunswick records, Queens Co.,** W of Jemseg at “Trout Creek”, 45.8227°N, 66.1240°W, 4.VI.2004, R. P. Webster, silver maple swamp, sifting litter at base of large tree (silver maple) (1 ♂, RWC). **Saint John Co.,** ca. 2 km NE of Maces Bay, 45.1168°N, 66.4552°W, 8.V.2006, R. P. Webster, eastern white cedar swamp, in sphagnum and leaf litter (1 ♀, RWC). **Sunbury Co.,** Burton, SW of Sunpoke Lake, 45.7575°N, 66.5726°W, 17.IV.2005, R. P. Webster, red maple swamp, in leaf litter near margin of slow stream (1 ♂, RWC); Acadia Research Forest, 45.9799°N, 66.3394°W, 14.V.2007, R. P. Webster, mature red spruce and red maple forest, sifting leaf litter (1 ♂, AFC). **York Co.,** Charters Settlement, 45.8331°N, 66.7410°W, 16.IV.2004, R. P. Webster, mature red spruce and eastern white cedar forest, in moss and litter near small brook (1 ♂, RWC); same locality, forest type, and collector but 45.8341°N, 66.7445°W, 27.IV.2005, margin of vernal pond in leaf litter (1 ♂, NBM); New Maryland, off Hwy 2, E of Baker Brook, 45.8760°N, 66.6252°W, 6.IV.2005, 26.IV.2005, 4.VI.2005, R. P. Webster, old growth eastern white cedar swamp, in moss and litter at base of cedar (3 ♂, NBM, RWC); Canterbury, Browns Mountain Fen, 45.8967°N, 67.6343°W, 1.VI.2005, M.-A. Giguère and R. Webster,




Map 1. Collection localities in New Brunswick, Canada of *Lathrobium othioides*.




Map 2. Collection localities in New Brunswick, Canada of *Lathrobium ampliipenne*.




Map 3. Collection localities in New Brunswick, Canada of *Lathrobium armatum*.



Map 4. Collection localities in New Brunswick, Canada of *Lathrobium confusum*.



Map 5. Collection localities in New Brunswick, Canada of *Lathrobium simile*.



Map 6. Collection localities in New Brunswick, Canada of *Lathrobium spissicorne*.

calcareous fen, in moist sphagnum (1 ♂, NBM); Mazerolle Settlement, 45.8729°N, 66.8311°W, 28.IV.2006, R. P. Webster, eastern white cedar swamp, margin of vernal pool in leaf litter (1 ♂, 1 ♀, RWC).

Collection and habitat data. *Lathrobium washingtoni* was found in eastern white cedar (*Thuja occidentalis* L.) swamps, a silver maple swamp, a red maple swamp, a

mature red spruce forest, and in an open calcareous cedar fen. Adults were sifted from sphagnum, sphagnum and leaf litter, moss and/or litter at bases of trees (silver maple, eastern white cedar), leaf litter near the margin of a slow stream and a brook, and from leaf litter on vernal pond margins. Adults were collected during April, May, and June.

Distribution in Canada and Alaska. AK, NT, BC, AB, SK, MB, ON, QC, NB, NS, NF (Campbell and Davies 1991; Klimaszewski et al. 2005). This species was first reported from New Brunswick by Klimaszewski et al. (2005) from the Acadia Research Forest.

Lathrobium (Lathrolepta) debile* LeConte, 1880*

http://species-id.net/wiki/Lathrobium_debile

Map 8

Material examined. New Brunswick, Charlotte Co., at New River, 45.2166°N, 66.5953°W, 2.VI.2006, R. P. Webster, river margin, under debris (1 ♂, RWC). Queens Co., W of Jemseg at “Trout Creek”, 45.8227°N, 66.1240°W, 4.VI.2004, 3.IV.2006, 11.IV.2006, R. P. Webster, silver maple swamp, sifting litter at base of large tree (silver maple) (2 ♂, 1 ♀, NBM, RWC); Grand Lake near Scotchtown, 45.8762°N, 66.1816°W, 25.IV.2004, 12.V.2004, R. P. Webster, (red) oak and (silver) maple forest, in leaf litter (2 ♂, 2 ♀, RWC). Sunbury Co., Burton, SW of Sunpoke Lake, 45.7575°N, 66.5726°W, 17.IV.2005, R. P. Webster, red maple swamp, in leaf litter near margin of slow stream (1 ♀, NBM). York Co., Charters Settlement, 45.8380°N, 66.7310°W, 18.IV.2004, R. P. Webster, mixed forest, in leaf litter near stream (1 ♀, RWC); 8.5 km W of Tracy off Rt. 645, 45.6821°N, 66.7894°W, 6.V.2008, R. P. Webster, alder swamp, in leaf litter and grass on hummocks (1, RWC); Fredericton, Odell Park, 45.9570°N, 66.6695°W, 19.VI.2005, R. P. Webster, old growth hemlock forest, in leaf litter (1 ♂, RWC).

Collection and habitat data. In New Brunswick, *L. debile* was found in a silver maple swamp, a red oak and silver maple forest, a red maple swamp, a mixed forest, an alder swamp, an old-growth eastern hemlock (*Tsuga canadensis* (L.) Carr.) forest, and along a river margin. Most adults were sifted from leaf litter on the forest floor or near stream margins. One individual was found under drift material on a river margin. Adults were collected during April, May, and June.

Distribution in Canada and Alaska. ON, QC, NB (Campbell and Davies 1991).

***Tetartopeus capitosus* Casey, 1905**

http://species-id.net/wiki/Tetartopeus_capitosus

Map 9

Material examined. New Brunswick, Queens Co., W of Jemseg at “Trout Creek”, 45.8240°N, 66.1220°W, 4.VI.2004, R. P. Webster, silver maple swamp, margin of ver-

nal pond in moist leaf litter on muddy soil (1 ♀, RWC); Grand Lake near Scotchtown, 45.8762°N, 66.1816°W, R. P. Webster, 5.VI.2004, lake margin, under drift material (1 ♀, RWC); same locality data and collector, 25.V.2006, oak and maple forest near lakeshore, in litter near vernal pond (1 ♂, 1 ♀, RWC); Upper Gagetown, bog adjacent to Hwy 2, 45.8316°N, 66.2346°W, 23.V.2006, R. P. Webster, tamarack bog, treading *Carex* into water (1 ♂, RWC). **Sunbury Co.**, Maugerville, Portobello Creek N.W.A., 45.8992°N, 66.4248°W, 5.VI.2004, R. P. Webster, silver maple swamp, margin of small (vernal) pond, in leaf litter (1 ♀, RWC); Sheffield, Portobello Creek N.W.A., 45.8952°N, 66.2728°W, 17.VII.2004, R. P. Webster, silver maple swamp, u.v. light (1 ♀, RWC).

Collection and habitat data. Nearly all *Tetartopeus* spp. have been collected in association with wetland habitats, usually in accumulations of damp leaf litter, moss, and other debris along streams, bogs, marshes, swamps, and ponds (Watrous 1980). Watrous (1980) noted that *T. capitosus* (as *Latrobium* (*Tetartopeus*) *punctulatum* LeConte) occurred mostly in the boreal forest region but gave no specific habitat data for this species. In New Brunswick, adults of this species were found in silver maple swamps (floodplain forests), a red oak and silver maple forest, along a lake margin, and in a tamarack (*Larix laricina* (Du Roi) K. Koch) bog. Adults were found in leaf litter near vernal ponds, under drift material on a lake margin, and by treading *Carex* in a tamarack bog, and one individual was captured at an ultraviolet light.

Distribution in Canada and Alaska. AK, NT, BC, AB, SK, MB, ON, QC, NB, NS (Watrous 1980; Campbell and Davies 1991).

Tetartopeus rubripennis Casey, 1905

http://species-id.net/wiki/Tetartopeus_rubripennis

Map 10

Material examined. **New Brunswick, Madawaska Co.**, at Green River, 47.6918°N, 68.3202°W, 21.VI.2010, M. Turgeon & R. Webster, river margin among gravel on gravel bar (1 ♀, RWC). **Restigouche Co.**, Jacquet River Gorge P.N.A., 47.8197°N, 66.0835°W, 26.VI.2008, R. P. Webster, margin of Jacquet River among cobblestones near water (1 ♂, RWC); same locality and collector but 47.7894°N, 66.1065°W, 14.V.2010, river margin (Jacquet River), under drift material (1 ♂, RWC); same locality and collector but 47.8257°N, 66.0779°W, 24.V.2010, partially shaded cobblestone bar near outflow of brook at Jacquet River, under cobblestones and gravel (1 ♀, RWC); Kedgwick Forks, 47.9085°N, 67.9057°W, 22.VI.2010, R. P. Webster, river margin, gravel bar among gravel and cobblestones (1 ♂, 2 ♀, RWC).

Collection and habitat data. Watrous (1980) reported *T. rubripennis* (as *Lathrobium* (*Tetartopeus*) *rubripenne*) primarily from thin litter in marshes at one locality. In northern New Brunswick, *T. rubripennis* was found along river margins among gravel

and cobblestones or under drift material, usually near water. Adults were collected during May and June.

Distribution in Canada and Alaska. ON, QC, NB, NS (Campbell and Davies 1991). In the description of *T. rubripennis*, Watrous (1980) noted that the elytra are usually entirely red and rarely black with only apices reddish. Only one of the seven specimens of this species from New Brunswick had entirely reddish elytra. However, the male genitalia of the dark specimens conform to the illustration (Fig. 51) given in Watrous (1980) for *T. rubripennis*. There are also specimens of the dark form from Nova Scotia and Quebec in the CNC (A. Davies, personal communication).

***Lobrathium (Eulathrobium) grande* (LeConte, 1863)**

http://species-id.net/wiki/Lobrathium_grande

Map 11

Material examined. **New Brunswick, Charlotte Co.,** near Clark Ridge, 45.3155°N, 67.4406°W, 27.V.2007, R. P. Webster, beaver pond, treading vegetation (1, NBM). **Queens Co.,** Scotchtown near Indian Point (at Grand Lake), 45.8762°N, 66.1816°W, R. P. Webster, 5.VI.2004, lake margin, under drift material (3, RWC); W of Jemseg near Jemseg River, 45.8255°N, 66.1174°W, 1.VII.2008, R. P. Webster, seasonally flooded marsh, treading vegetation on margin of pool 1, NBM); Grand Lake Meadows P.N.A., 45.8227°N, 66.1209°W, 14-19.V.2010, R. Webster & C. MacKay, old silver maple forest with green ash and seasonally flooded marsh, Lindgren funnel trap (1, AFC); same locality data and forest type, 5-19.VII.2011, 19.VII-5.VIII.2011, M. Roy & V. Webster, Lindgren funnel traps (2, AFC, NBM). **Sunbury Co.,** Maugerville, Portobello Creek N.W.A., 45.8992°N, 66.4248°W, 27.V.2004, 5.VI.2004, R. P. Webster, silver maple forest, margin of small (vernal) pond, in leaf litter (5, RWC); Sheffield, Portobello Creek N.W.A., 45.8952°N, 66.2728°W, 24.VI.2004, R. P. Webster, seasonally flood marsh, treading marsh vegetation (1, RWC). **York Co.,** Douglas, near Nashwaaksis River, 45.9845°N, 66.6908°W, 4.VI.2003, R. P. Webster, silver maple forest, margin of small pond in leaf litter (1, RWC); Charters Settlement, 45.8340°N, 66.7450°W, 27.IV.2006, R. P. Webster, mixed forest, margin of vernal pond in moist leaves (1 ♀, NBM).

Collection and habitat data. Watrous (1981) reported *Lobrathium grande* from a variety of riparian habitats. Adults occurred in moss roots, in leaf litter and sticks at a stream margin, in litter in a North American beaver (*Castor canadensis* Kuhl) lodge and in leaf litter at the margin of a swamp. Larvae were described by Watrous (1981). In New Brunswick, was adults were found in silver maple swamps, seasonally flooded marshes, a mixed forest, and beaver pond margins. Adults were usually found among moist leaves along pond and vernal pond margins or by treading vegetation in marshes. One individual was captured in a Lindgren funnel trap deployed in an old silver maple swamp. Adults were captured during April, May, June, July, and August.

Distribution in Canada and Alaska. ON, QC, NB, NS (Campbell and Davies 1991).

***Lobrathium (Lobrathium) collare* (Erichson, 1840)**

http://species-id.net/wiki/Lobrathium_collare

Map 12

Material examined. New Brunswick, Carleton Co., Meduxnekeag Valley Nature Preserve, 46.1931°N, 67.6825°W, 31.V.2005, M.-A. Giguère & R. Webster, mixed forest, river margin, under drift material (1, RWC). Queens Co., Scotchtown near Indian Point (at Grand Lake), 45.8762°N, 66.1816°W, R. P. Webster, 5.VI.2004, lake margin, under drift material (1, RWC); Bayard at Nerepis River, 45.4426°N, 66.3380°W, 30.V.2008, R. P. Webster, river margin, on sand bar in moist sand, collected by lightly splashing sand with water (1, RWC). Sunbury Co., Sheffield, Portobello Creek N.W.A., 45.8952°N, 66.2728°W, 18.VI.2004, R. P. Webster, silver maple forest (swamp), black light trap (1, RWC). York Co., Charters Settlement, 45.8395°N, 66.7391°W, 26.VI.2003, 1.VIII.2004, 10.VI.2005, 29.VI.2005, R. P. Webster, mixed forest, u.v. light (6, RWC).

Collection and habitat data. Two adults of this species were collected along river margins from under drift material and in sand (splashing), and at an ultraviolet light in a silver maple swamp, and near a mixed forest. Adults were collected during May, June, and August.

Distribution in Canada and Alaska. MB, ON, QC, NB, NS (Campbell and Davies 1991).

Subtribe Medonina Casey, 1905


Achenomorphus corticinus* (Gravenhorst, 1802)*

http://species-id.net/wiki/Achenomorphus_corticinus


Map 13

Material examined. New Brunswick, Sunbury Co., Maugerville, Portobello Creek N.W.A., 45.8992°N, 66.4248°W, 18.VI.2004, R. P. Webster, silver maple forest (swamp), black light trap (2 ♀, RWC). York Co., Fredericton, Odell Park, 45.9570°N, 66.6695°W, 19.VI.2005, R. P. Webster, compost (with) wood chips and decaying plant material (2 ♂, 7 ♀, RWC); Charters Settlement, 45.8395°N, 66.7391°W, 17.VII.2004, 9.VII.2008, R. P. Webster, mixed forest, u.v. light (2 ♀, RWC); same locality and collector but, 45.8456°N, 66.7267°W, 10.VI.2010, beaver dam, among sticks and debris near outflow area of dam (1 sex undetermined, RWC).


Collection and habitat data. In New Brunswick, *A. corticinus* adults were collected at ultraviolet light near a mixed forest and in a silver maple swamp. Adults were




Map 7. Collection localities in New Brunswick, Canada of *Lathrobium washingtoni*.




Map 8. Collection localities in New Brunswick, Canada of *Lathrobium debile*.




Map 9. Collection localities in New Brunswick, Canada of *Tetartopeus capitosus*.



Map 10. Collection localities in New Brunswick, Canada of *Tetartopeus rubripennis*.



Map 11. Collection localities in New Brunswick, Canada of *Loblathium grande*.



Map 12. Collection localities in New Brunswick, Canada of *Loblathium collar*.

common in compost with wood chips and decaying plant material. One individual was collected from among sticks and debris near the outflow area of a beaver dam.

Distribution in Canada and Alaska. MB, ON, QC, NB (Campbell and Davies 1991).

Subtribe Stilicina Casey, 1905***Rugilus angustatus* (Geoffroy, 1785)**

http://species-id.net/wiki/Rugilus_angustatus

Map 14

Material examined. **New Brunswick, Carleton Co.**, Jackson Falls, Bell Forest, 46.2152°N, 67.7190°W, 1.VI.2005, M.-A. Giguère & R. Webster, upper river margin, collected with aerial net between 16:00 and 18:00 h (2 ♀, RWC); **York Co.**, Canterbury, 45.8920°N, 67.6592°W, 8.VI.2004, D. Sabine & R. Webster, hardwood forest, wood pile, under bark (1 ♀, NBM); Charters Settlement, 45.8340°N, 66.7450°W, 27.IV.2005, 30.IV.2005, R. P. Webster, mixed forest, in wood pile, under (loose) bark of spruce (5 ♂, 2 ♀, RWC); Fredericton, at Saint John River, 45.9588°N, 66.6254°W, 7.VI.2005, R. P. Webster, river margin in flood debris (1 ♀, NBM); Fredericton, Odell Park, 45.9570°N, 66.6695°W, 19.VI.2005, R. P. Webster, compost (with) wood chips and decaying plant material (1 ♂, NBM); Fredericton, 45.9361°N, 66.6747°W, 17.VIII.2009, R. P. Webster, beaver dam, outer margin under over-hanging sticks near water (1 ♂, RWC).

Collection and habitat data. In the Palearctic region, *R. angustatus* occurs under decaying organic matter along forest borders and watercourses and in meadows (Hoebeke 1995). In New Brunswick, adults of this adventive species were found along river margins, in hardwood and mixed forests, and in a beaver dam. Adults were found under loose bark in wood piles, among composted wood chips and decaying plant material, under overhanging sticks on the outer margin of a beaver dam, and in flood debris on a river margin. Two individuals were collected with an aerial net during evening flight (16:00–18:00 h) on a river margin. Adults were collected during April, June, and August.

Distribution in Canada and Alaska. ON, QC, NB, NS (Campbell and Davies 1991; Hoebeke 1995).

Rugilus rufipes* Germar, 1836*

http://species-id.net/wiki/Rugilus_rufipes

Map 15

Material examined. **New Brunswick, Carleton Co.**, Jackson Falls, Bell Forest, 46.2152°N, 67.7190°W, 15.IX.2004, R. P. Webster, upper river margin, under litter on clay soil (2 ♂, 1 ♀, NBM, RWC); same locality and collector but 46.2246°N, 67.7206°W, 12.IV.2007, upper river margin, in drift material in area without snow cover, adults very active (2 ♂, 3 ♀, RWC); same locality but 46.2200°N, 67.7231°W, 20–26.V.2009, M.-A. Giguère & R. Webster, mature hardwood forest, Lindgren funnel traps (2, AFC); Meduxnekeag River Valley Nature Preserve, 46.1907°N, 67.6740°W, 20.VI.2006, R. P. Webster, mixed forest, in decaying gilled mushroom, (2 ♂, RWC); Jackson Falls, 46.2257°N, 67.7437°W,

18.VI.2010, R. P. Webster, water falls, splashing moss on rocks near fast flowing water (1, RWC). **York Co.**, Charters Settlement, 45.8395°N, 66.7391°W, 23.IX.2009, R. P. Webster, mixed forest, in decaying (moldy) corncobs and cornhusks (1, RWC).

Collection and habitat data. This adventive Palaearctic species lives in both dry and wet habitats in the Palaearctic region, including meadows, fields, heaths, forests, and hilly steppe (Hoebeke 1995). Adults occurred in decaying organic matter and compost, under stones, and among leaves. In New Brunswick, adults were collected from grass litter and drift material along river margins and from decaying gilled mushrooms and decaying (moldy) corncobs and cornhusks in mixed and hardwood forests. One individual was collected from wet moss on rocks adjacent to a waterfall. Adults become active very early in the season when a deep snow cover is still present, as a number of very active adults were collected from a sun-exposed bare patch of drift material on an upper river margin on 12 April when a 60-cm snow pack was still present. Adults were captured during April, May, June, and September.

Distribution in Canada and Alaska. ON, QC, NB, (Campbell and Davies 1991; Hoebeke 1995).

Subtribe Cryptobiina Casey, 1905

Homaeotarsus (Gastrolobium) bicolor (Gravenhorst, 1802)**

http://species-id.net/wiki/Homaeotarsus_bicolor

Map 16

Material examined. New Brunswick, Carleton Co., Belleville, Meduxnekeag River Valley Nature Preserve, 46.1944°N, 67.6832°W, 2.VI.2008, R. P. Webster, river margin, under cobblestone in sand/gravel among scattered grasses (1, RWC).

Collection and habitat data. *Homaeotarsus* are generally riparian and occur along river margins (Brunke et al. 2011). The single adult of *H. bicolor* from New Brunswick was collected from under a cobblestone along a river margin during early June.

Distribution in Canada and Alaska. ON, QC, NB (Campbell and Davies 1991).

Homaeotarsus (Hesperobium) cinctus (Say, 1830)

http://species-id.net/wiki/Homaeotarsus_cinctus

Map 17

Material examined. New Brunswick, Carleton Co., “Two Mile Brook Fen”, 46.3619°N, 67.6730°W, 6.V.2005, M.-A. Giguère & R. Webster, calcareous cedar fen, open area with sedges, in sphagnum hummock (3, RWC). **Charlotte Co.**, near New River, 45.1616°N, 66.6649°W, 7.VII.2006, R. P. Webster, mixed forest in sedge marsh, treading sedges (1, NBM). **Madawaska Co.**, Loon Lake, 236 m elev., 47.7839°N, 68.3943°W, 21.VI.2010, R. P. Webster, boreal forest, small lake surrounded by sedges,

treading sedges and grasses into water (1, NBM). **Restigouche Co.**, Jacquet River Gorge P.N.A., 47.8207°N, 65.9955°W, 12.VIII.2010, R. P. Webster, black spruce bog, treading vegetation (*Carex* & sphagnum) (1, NBM). **Saint John Co.**, Chance Harbour off Rt. 790, 45.1374°N, 66.3633°W, 25.VI.2010, R. P. Webster, saturated green sphagnum mat, treading (1, NBM). **York Co.**, Canterbury, Browns Mountain Fen, 45.8967°N, 67.6344°W, 21.VII.2004, D. Sabine, R. Webster, & J. Edsall, calcareous cedar fen, in moss and sphagnum among scattered sedges (1, RWC); same locality and habitat data, 2.V.2005, M.-A. Giguère & R. Webster, open area with sedges, in sphagnum hummock (3, RWC); Charters Settlement, 45.8267°N, 66.7343°W, 14.V.2005, 23.V.2005, R. P. Webster, margin of *Carex* marsh/fen, in sphagnum and leaf litter at base of tree (1 ♂, 2 sex undetermined, RWC); Upper Brockway, 45.5684°N, 67.0993°W, 23.IV.2006, R. P. Webster, forested black spruce bog, in sphagnum (1, NBM); Magundy, 45.8491°N, 67.1573°W, 8.VII.2006, R. P. Webster, kettle hole bog, treading bog margin (1, NBM).

Collection and habitat data. In New Brunswick, *H. cinctus* was found in *Carex* marshes, open calcareous cedar fens, a forested black spruce (*Picea mariana* (Mill.) B.S.P.) bog, and in a kettle hole bog with a floating bog mat. Adults were found in wet to saturated sphagnum with scattered sedges often in a floating mat, among emergent sedges and grasses, and in sphagnum hummocks. Adults were collected by treading these microhabitats. Some adults were sifted from sphagnum and leaf litter at bases of trees on a *Carex* marsh margin. This species was collected during April, May, June, July, and August.

Distribution in Canada and Alaska. BC, AB, ON, QC, NB, NS (Campbell and Davies 1991).


Homaeotarsus (Hesperobium) cribratus* (LeConte, 1863)*

http://species-id.net/wiki/Homaeotarsus_cribratus


Map 18

Material examined. **New Brunswick, Carleton Co.**, Jackson Falls, Bell Forest, 46.2208°N, 67.7211°W, 19.IV.2005, R. P. Webster, mature hardwood forest, in leaf litter at base of tree (2, RWC); same locality but, 46.2152°N, 67.7190°W, 11.V.2005, M.-A. Giguère & R. Webster, river margin, in drift material (2, RWC); same locality data and collectors, 1.VI.2005, upper river margin, collected with aerial net between 16:00 and 18:00 h (2, RWC); Belleville, Meduxnekeag River Valley Nature Preserve, 46.1888°N, 67.6762°W, 20.V.2005, R. P. Webster, river margin, in flood debris (2, RWC); same locality and collector but, 46.1942°N, 67.6832°W, 2.VI.2008, river margin, under cobblestones (1 ♂, RWC). **York Co.** Fredericton, at Saint John River, 45.9588°N, 66.6254°W, 4.VII.2004, R. P. Webster, river margin, in drift material (mostly maple seeds) (1, RWC).


Collection and habitat data. In New Brunswick, most adults of this species were collected along river margins. Adults were collected from flood debris and drift material (maple seeds), and from under a cobblestone. Two adults were collected with an aerial



Map 13. Collection localities in New Brunswick, Canada of *Achenomorphus corticinus*.




Map 14. Collection localities in New Brunswick, Canada of *Rugilus angustatus*.




Map 15. Collection localities in New Brunswick, Canada of *Rugilus rufipes*.



Map 16. Collection localities in New Brunswick, Canada of *Homaeotarsus bicolor*.



Map 17. Collection localities in New Brunswick, Canada of *Homaeotarsus cinctus*.



Map 18. Collection localities in New Brunswick, Canada of *Homaeotarsus cribratus*.

net during an evening flight (16:00–18:00 h) along a river margin, and two individuals were collected from leaf litter at the base of a tree in mature hardwood forest (0.5 km from a river margin) during late April when some snow was still present. This may have been an overwintering site. Adults were collected during April, May, June, and July.

Distribution in Canada and Alaska. ON, QC, NB (Campbell and Davies 1991).

Homaeotarsus (Hesperobium) pallipes* (Gravenhorst, 1802)*


http://species-id.net/wiki/Homaeotarsus_pallipes

Map 19

Material examined. New Brunswick, York Co., Charters Settlement, 45.8456°N, 66.7267°W, 16.V.2010, R. P. Webster, beaver dam, among sticks and debris near outflow area of dam (1 ♀, RWC).

Collection and habitat data. The only specimen from New Brunswick was collected during May from among sticks and debris in a beaver dam near an outflow area with flowing water.

Distribution in Canada and Alaska. ON, QC, NB (Campbell and Davies 1991).



Map 19. Collection localities in New Brunswick, Canada of *Homaeotarsus pallipes*.

Acknowledgments

Caroline Simpson is thanked for editing this manuscript. Two anonymous reviewers are thanked for their helpful suggestions that improved this manuscript. Anthony Davies (Agriculture and Agri-Food Canada (CNC), Ottawa) is thanked for determining specimens and other invaluable assistance. Jon Sweeney (AFC) revised the first draft of this manuscript and provided very useful comments. Jon Sweeney, Nichole Brawn, Katie Burgess, Marie-Andrée Giguère, Jim Edsall, Nancy Harn, Cory Hughes, Colin MacKay, Wayne MacKay, Jessica Price, Michelle Roy, and Vincent Webster are thanked for technical assistance and collecting specimens. Natural Resources Canada, Canadian Forest Service; the Canadian Food Inspection Agency; and the USDA APHIS funded the study on early detection of invasive cerambycids, which provided the records from the specimen captured in Lindgren funnel traps. The Canadian Wildlife Service is thanked for funding insect surveys at the Portobello Creek National Wildlife Area, the New Brunswick Environmental Trust Fund and New Brunswick Wildlife Trust Fund are thanked for funding various insect surveys over the past 7 years, and the Meduxnekeag River Association is thanked for permission to sample beetles at the Meduxnekeag Valley Nature Preserve (which includes the Bell Forest). The New Brunswick Department

of Natural Resources (Fish and Wildlife Branch) is thanked for issuing permits for sampling in the Protected Natural Areas and for providing logistical support. Survey work in the Jacquet River Gorge Protected Natural Area was organized through the New Brunswick Museum with external funding from the New Brunswick Environmental Trust Fund, Salamander Foundation, and the New Brunswick Wildlife Trust Fund.

References

- Bouchard P, Bousquet Y, Davies AE, Alonso-Zarazaga MA, Lawrence JF, Lyal CHC, Newton AF, Reid CAM, Schmitt M, Ślipiński SA, Smith ABT (2011) Family- group names in Coleoptera (Insecta). *ZooKeys* 88: 1–972. doi: 10.3897/zookeys.88.807
- Brunke A, Newton A, Klimaszewski J, Majka C, Marshall S (2011) Staphylinidae of eastern Canada and adjacent United States. Key to subfamilies: Staphylininae: tribes and subtribes, and species of Staphylinina. *Canadian Journal of Arthropod Identification* 12: 1–110.
- Campbell JM (1973) A revision of the genus *Tachinus* (Coleoptera: Staphylinidae) of North and Central America. *Memoirs of the Entomological Society of Canada* 90: 1–137. doi: 10.4039/entm10590fv
- Campbell JM, Davies A (1991) Family Staphylinidae: rove beetles. In: Bousquet Y (Ed) *Checklist of Beetles of Canada and Alaska*. Agriculture Canada, Research Branch, Ottawa, Ontario, Publication 1861/E, 86–124.
- Casey TL (1905) A revision of the American Paederini. *Transactions of the Academy of Science of St. Louis* 15: 17–248.
- Casey TL (1910) Synonymic and descriptive notes on the Paederini and Pinophilini, In: *Memoirs on the Coleoptera*, Vol. 1. New Era, Lancaster, Pennsylvania, 184–201.
- Herman LH (1965a) A revision of *Orus* Casey. I. Subgenus *Leucorus* Casey and a new subgenus (Coleoptera: Staphylinidae). *The Coleopterists Bulletin* 18: 112–121.
- Herman LH (1965b) A revision of *Orus* Casey. II. Subgenera *Orus*, *Pycnorus* and *Nivorus* (Coleoptera: Staphylinidae). *The Coleopterists Bulletin* 19: 73–90.
- Hoebeke RE (1995) Three Palearctic species of *Rugilus* Leach in North America (Coleoptera: Staphylinidae, Paederinae): redescription, new synonymy, and new records. *Insecta Mundi* 9: 69–80.
- Lindgren BS (1983) A multiple funnel trap for scolytid beetles (Coleoptera). *The Canadian Entomologist* 115: 299–302. doi: 10.4039/Ent115299-3
- Klimaszewski J, Sweeney J, Price J, Pelletier G (2005) Rove beetles (Coleoptera: Staphylinidae) in red spruce stands, eastern Canada: diversity, abundance, and descriptions of new species. *The Canadian Entomologist* 137: 1–48. doi: 10.4039/n03-123
- Majka CG, Klimaszewski J (2008) Adventive Staphylinidae (Coleoptera) of the Maritime provinces of Canada: further contributions. *ZooKeys* 2: 151–174. doi: 10.3897/zookeys.2.5
- Newton AF, Thayer MK, Ashe JS, Chandler DS (2000) [2001] Family 22. Staphylinidae Latreille, 1802. In: Arnett RH, Thomas MC (Eds) *American Beetles*. Volume 1. Archostemata, Myxophaga, Adephaga, Polyphaga. CRC Press, Boca Raton, Florida, xv + 272–418.

- Watrous LE (1980) Studies of *Lathrobium* (*Tetrartopeus*): natural history, revision, and phylogeny of Nearctic species. *Systematic Entomology* 5: 303–338. doi: 10.1111/j.1365-3113.1980.tb00418.x
- Watrous LE (1981) Studies of *Lathrobium* (*Lobrathium*): Revision of the *grande* species group (Coleoptera: Staphylinidae). *Annals of the Entomological Society of America*: 74: 144–150.
- Webster RP, Klimaszewski J, Pelletier G, Savard K (2009) New Staphylinidae (Coleoptera) records with new collection data from New Brunswick, Canada. I. Aleocharinae. In: Majka CG, Klimaszewski J (Eds) Biodiversity, biosystematics, and ecology of Canadian Coleoptera II. *ZooKeys* 22: 171–248. doi: 10.3897/zookeys.22.152
- Webster RP, Smetana A, Sweeney JD, DeMerchant I (2012) New Staphylinidae (Coleoptera) records with new collection data from New Brunswick and an addition to the fauna of Quebec: Staphylininae. In: Klimaszewski J, Anderson R (Eds) Biosystematics and Ecology of Canadian Staphylinidae (Coleoptera) II. *ZooKeys* 186: 293–348. doi: 10.3897/zookeys.186.2469