

# First report of *Xiphinema brevicolle* Lordello et Costa, 1961 (Nematoda, Longidoridae) in Japan

Hiromichi Sakai<sup>1,†</sup>, Ai Takeda<sup>2,‡</sup>, Takayuki Mizukubo<sup>1,§</sup>

**1** National Agriculture and Food Research Organization, Agricultural Research Center, Kannondai 3-1-1, Tsukuba, Ibaraki 305-8666, Japan **2** Chiba Prefectural Agriculture and Forestry Research Center, 808 Daizenno, Midori-Ku, Chiba 266-0006 Japan

Corresponding author: Hiromichi Sakai (sakaihm@affrc.go.jp)

Academic editor: Sergei Subbotin | Received 15 June 2011 | Accepted 23 August 2011 | Published 7 October 2011

**Citation:** Sakai H, Takeda A, Mizukubo T (2011) First report of *Xiphinema brevicolle* Lordello et Costa, 1961 (Nematoda, Longidoridae) in Japan. ZooKeys 135: 21–40. doi: 10.3897/zookeys.135.1716

## Abstract

Mixed populations of *Xiphinema americanum*-group species were detected from a root zone soil sample of Japanese holly, *Ilex crenata*, during a survey for plant-parasitic nematodes of commercial ornamental plant nurseries in Chiba Prefecture, Japan. From the result of the morphological study, the species were identified as *X. brevicolle* and *Xiphinema* sp. This is the first record of *X. brevicolle* in Japan. Morphometrics of *X. brevicolle* generally agree with those of the type specimens and the topotype specimens. *Xiphinema* sp. morphometrically resembles *X. paramonovi* except for tail length. The mitochondrial COI region, the nuclear 18S rDNA and the nuclear large subunit rDNA D2/D3 region of the species were sequenced and compared in the molecular study. For the COI region, PCR primers were newly designed to obtain longer sequences, ca. 900 bp, than previously used. Sequence identities of COI, 18S and D2/D3 regions between these two populations were 84.0–84.1%, 99.9% and 98.1–98.2%, respectively. Phylogenetic analyses of maximum likelihood trees were carried out to compare genetic relationships among the group and some suggestions were made on the *X. brevicolle*-subgroup.

## Keywords

COI, *Ilex crenata*, Japanese holly, rDNA, *Xiphinema americanum*-group, *Xiphinema brevicolle*-subgroup

## Introduction

During a survey for plant-parasitic nematodes of commercial ornamental plant nurseries in Chiba Prefecture, Japan, we detected mixed populations of *Xiphinema americanum*-group species from a root zone soil sample of Japanese holly, *Ilex crenata* Thunb.,

one of the major garden tree species in Japan. This study was conducted using morphological characters of females and DNA sequences of the mitochondrial COI region and the nuclear ribosomal RNA (rDNA) regions to identify and characterize the species in the mixed populations.

## Methods

### Collection of the nematode specimens

The soil sample containing the *Xiphinema* spp. (No. 001-001) was taken from the root zone of Japanese holly growing in a commercial plant nursery at Sosa City, Chiba, Japan. Nematodes were extracted with Cobb's wet-sieving technique. Material collected on a 75 µm mesh sieve was placed on a Baermann funnel and nematodes were collected after one day at room temperature.

### Morphological observation

Females of *Xiphinema* spp. were removed from the nematode suspension using a dissecting microscope and nematode pick. Female nematodes were transferred to a small amount of water then killed by either heating at 60°C for 2 min or by adding hot FP4:1 (Netscher and Seinhorst 1969). Killed nematodes were fixed with FG4:1 (De Grisse 1969). The specimens were fixed for more than one week, processed into glycerin by the ethanol/glycerin method (Seinhorst 1959, De Grisse 1969), and mounted in dehydrated glycerin supported with both minute glass beads and paraffin on glass slides. Morphological observations were made using a DIC microscope (BX51, Olympus Co., Japan). A digital camera, Olympus DP20 or DP21 was used for measuring and taking photo images. The body length and position of vulva were measured with a digital curvimeter (CV-9 Jr., Koizumi Sokki Mfg. Co. Ltd., Japan) on nematode line drawings prepared using a drawing tube. Illustrations of nematodes were sketched directly on highly transparent tracing film (No. 200Z-A4 (S): Tochiman Technical Paper Co. Ltd., Japan) by tracing DP21 digital camera images on a panel-protected LCD-display. Sketched images were converted to digital images using the software Adobe Illustrator CS4 (Adobe Systems Inc., USA) and a pen tablet (Intuos4: Wacom Co., Ltd., Japan).

### Molecular study

DNA extractions from single female specimens were carried out according to Sakai (2010), which yielded 200 µL lysate for each specimen. Before DNA extraction, the specimens were tagged, killed by gentle heat, prepared as temporary water mounts,

digital images of mounts were photographed, and then images were measured for body and odontostyle length, respectively.

DNA fragments of the mitochondrial COI region were amplified by PCR using the set of primers, COI-F1 and COI-R1 (Table 1), which were originally designed from sequence comparison of the COI region between *X. americanum* (He et al. 2005a: GenBank accession AY382608; NCBI Reference Sequence NC\_005928) and *Caenorhabditis elegans* (Howe and Denver 2008: GenBank accession EU407804). COI-R1 is virtually identical to COIR (He et al. 2005a), with the latter having only a difference in degeneracy. The PCR reaction mixture consisted of 0.2 mM dNTPs, 0.3  $\mu$ M of each primer, 0.5 U PrimeSTAR HS DNA Polymerase with PrimeSTAR Buffer (5 mM  $Mg_2^+$  plus) (Takara Bio Inc., Japan), and 10  $\mu$ L of DNA lysate as PCR template, in a total volume of 20  $\mu$ L. The reaction conditions were as follows: a single step of pre-denaturation at 94°C for 2 min, followed by 35 cycles of denaturation at 94°C for 30 s, annealing at 40°C for 15 s, 0.5°C/s ramp up to extension temperature, and extension at 72°C for 1 min.

DNA fragments of the nuclear 18S rDNA region were amplified using the set of primers, 18S 39 and 18S 1573R (Mullin et al. 2005). The PCR reaction mixture consisted of 0.2 mM dNTPs, 0.2  $\mu$ M of each primer, 0.5 U TaKaRa Ex Taq Hot Start Version with Ex Taq Buffer ( $Mg_2^+$  plus) (Takara Bio), and 2  $\mu$ L of DNA lysate as PCR template, in a total volume of 20  $\mu$ L. The reaction conditions were as follows: a single step of pre-denaturation at 94°C for 2 min, followed by 35 cycles of denaturation at 94°C for 30s, annealing at 55°C for 30s, and extension at 72°C for 1 min.

DNA fragments of the nuclear large subunit rDNA D2/D3 region were amplified using the set of primers, D2Ab (De Ley et al. 1999) and D3B (Thomas et al. 1997). The PCR reaction conditions were the same as those for 18S rDNA.

These PCR products were purified with QIAquick PCR Purification Kit (Qiagen K.K., Japan), subjected to cycle sequencing with BigDye Terminator v3.1 Cycle Sequencing Kit (Life Technologies Japan), purified with either DyeEx 2.0 Spin Kit (Qiagen) or Agencourt CleanSEQ (Beckman Coulter Inc., USA), and sequenced on the ABI PRISM 3100 Genetic Analyzer (Life Technologies Japan Ltd., Japan). Primers 18S 599R, 18S 550, 18S 977R, 18S 965 (Mullin et al. 2005), and D3B (Thomas et al. 1997) were also used as inner primers for sequencing as well as those designed by us (Table 1). The DNA sequences were aligned by MUSCLE (Edgar 2004) and arranged using BioEdit (Hall 1999). Multiple alignments were manually refined where necessary.

Phylogenetic analyses including substitution model selections were performed using MEGA5 (Tamura et al. 2011) to compare the obtained sequences with homologous sequences of the *X. americanum*-group in GenBank searched via BLAST. A maximum likelihood (ML) tree was constructed using 1,578 sites for multiply aligned sequences of the 18S rDNA region, where the model K2 + G, the heuristic search with the Close-Neighbor-Interchange (CNI) method, bootstrapping with 500 replications, gaps treatment using all sites, and the neighbor-joining (NJ) tree as the initial tree were employed. A ML tree was constructed using 676 sites for multiply aligned sequences of the D2/D3 region with gaps and an ambiguous site deleted, where the model K2 +

**Table 1.** Primers designed and employed originally in this study for PCR amplification and sequencing of mitochondrial COI region and rDNA D2/D3 region.

primer name	Sequence	Note
CO1-F1	5'-ATAATTTTTTTTATGGTAATACC-3'	PCR, sequencing
CO1-R1	5'-ACTACATAATAAGTATCATG-3'	PCR, sequencing
CO1-F2	5'-TATATTTTAATTTTACCTGG-3'	sequencing
CO1-R2	5'-CCAGGTAAAATTAAAATATA-3'	sequencing
D3A-R	5'-AGACTCCTTGGTCCGTGTTTC-3'	sequencing

G, the CNI method, bootstrapping with 500 replications, and a maximum parsimony (MP) tree as the initial tree were employed. A ML tree was constructed using 342 sites for multiply aligned sequences of the COI region, where the model T92 + G + I, the CNI method, bootstrapping with 500 replications, and a MP tree as the initial tree were employed. Among COI homologous sequences of the *X. americanum*-group in the database, those including alignment gaps were excluded from the analysis because they possibly represent pseudogenes. Sequences of *Longidorus* species available in Gen-Bank were used to root those trees.

Results

The DNA sequence analysis of the mitochondrial COI region for 9 specimens suggested two different *X. americanum*-group species were present. Morphometric data were used to identify them as *Xiphinema brevicolle* Lordello et Costa, 1961, and an undescribed *X. americanum*-group species.

Morphological observations

*Xiphinema brevicolle* Lordello & Costa, 1961

[http://species-id.net/wiki/Xiphinema\\_brevicolle](http://species-id.net/wiki/Xiphinema_brevicolle)

Figs 1A, 1C, 1E, 1G, 2, 4A, 5A, 6A

**Measurements.** See Tables 2, 3.

**Remarks.** Morphometrics of the specimens obtained here generally agree with those of the type specimens and topotype specimens (Lamberti et al. 1992, Luc et al. 1998) of the species (Table 2). No male was detected.

**Nomenclatorial note.** The emended name of this species, *Xiphinema brevicollum*, was proposed by Luc et al. (1998) and used in many works to date. Monteiro (2010) claimed that the correct species name is *Xiphinema brevicolle* Lordello et Costa, 1961, and should have been preserved unaltered. We support the claim by Monteiro (2010) and *X. brevicolle* is used here.

***Xiphinema* sp.**

Figs 1B, 1D, 1F, 1H, 3, 4B, 5B, 6B

**Measurements.** See Tables 2, 3.

**Remarks.** These specimens morphometrically resemble *Xiphinema paramonovi* Romanenko, 1981, except for the clearly different tail length (Table 2). The morphometrics of the specimens partly overlap those of *X. brevicolle*. General morphology and DNA information addressed below suggest that these specimens belong to some species related to *X. brevicolle*, though a specific species accommodating them was not found. We finally regarded these specimens as an unidentified *X. americanum*-group species. Further information is required to identify the specimens as a new species or determine if they represent intra-specific variation of species previously described. No male was detected.

**Molecular study.** DNA sequences of 886 bp except for primer regions were obtained for the mitochondrial COI region. The five *X. brevicolle* specimens observed had identical sequences, whereas a single nucleotide in the sequence differed among the four specimens of *Xiphinema* sp. though this variation resulted in no difference between the translated amino acid sequences within the specimens. Sequence identity between these two species was 84.0–84.1%, whereas the sequences of *X. brevicolle* and *Xiphinema* sp. were 80.7 and 80.4–80.5% identical to that of *X. americanum* (He et al. 2005a) respectively, with no gap found among them (Fig. 7). Putative amino acid sequences were available without any stop codon when translations were made from the second base of the obtained sequences. DNA sequences of 1,566 bp except for primer regions were obtained for the 18S rDNA region from one specimen for each species. The difference between sequences of the two species studied here was only a single nucleotide, resulting in 99.9% identity without any gap. DNA sequences of 788–791 bp except primer regions were obtained for the D2/D3 region from four specimens for each species. A single nucleotide variation of sequence among four specimens of *X. brevicolle* was observed, whereas no variation of sequence was observed among four specimens of *Xiphinema* sp. Sequence identity between these two species was 98.1–98.2%, with gaps found.

ML trees inferred for the 18S rDNA and D2/D3 regions placed our specimens in similar clades, which include *X. brevicolle* and its junior synonyms by Luc et al. (1998) such as *X. diffusum* Lamberti et Bleve-Zacheo, 1979, *X. incognitum* Lamberti and Bleve-Zacheo, 1979, *X. taylori* Lamberti et al., 1992, as well as other different species like *X. inaequale* (Khan et Ahmad, 1975) and *X. lamberti* Bajaj et Jairajpuri, 1977 (Figs 8, 9). On the other hand, the ML tree inferred for the COI region didn't show strong support for such a clade because of a low bootstrap value though several subclades were strongly supported by high bootstrap values (Fig. 10).

**Discussion**

This study reports the occurrence of *X. brevicolle* in Japan for the first time. The only member species of the *Xiphinema americanum*-group recorded to date in Japan was *X.*

**Table 2.** Morphometrics of *Xiphinema brevicolle*, *Xiphinema* sp., *X. paramonovi* and *X. diffusum* (female). Mean  $\pm$  standard deviation (range) in  $\mu\text{m}$ , except for L in mm, and ratios.

	<i>Xiphinema brevicolle</i>				<i>Xiphinema</i> sp. (this study)	<i>Xiphinema</i> <i>paramonovi</i> Paratypes (Romanenko 1981)	<i>Xiphinema diffusum</i> Paratypes (Lamberti and Bleve-Zacheo 1979)
	Population in this study	Topotypes (Luc et al. 1998)		Types (Lordello and Costa 1961)			
n	22	25	17	-	13	27	10
L	1.93 $\pm$ 0.11 (1.71-2.10)	1.92 $\pm$ 0.122 (1.7-2.16)	2.1 $\pm$ 0.1 (1.8-2.2)	(1.82-2.20)	2.30 $\pm$ 0.12 (2.08-2.47)	2.1 (2.0-2.3)	1.7 (1.6-1.8)
a	46.1 $\pm$ 2.6 (40.5-50.5)	46.1 $\pm$ 1.72 (44-51)	44.5 $\pm$ 2.3 (40.7-50.1)	(36.0-42.2)	48.8 $\pm$ 2.4 (45.0-52.3)	49.6 (44-54.2)	47 (46-51)
b	6.6 $\pm$ 0.6 (5.5-8.2)	5.92 $\pm$ 0.37 (5.1-6.7)	6.4 $\pm$ 0.6 (5.6-7.7)	(7.0-10.5)	7.0 $\pm$ 0.4 (6.5-8.1)	6.1 (4.8-7.1)	6.9 (5.3-8.9)
c	69.8 $\pm$ 4.4 (60.5-79.1)	76.9 $\pm$ 5.64 (67.6-89.9)	77.8 $\pm$ 0.6 (60.3-94.0)	(62.5-93.0)	79.6 $\pm$ 8.5 (67.5-94.5)	60.5 (49.1-68.5)	72 (63-84)
c'	1.0 $\pm$ 0.1 (0.9-1.2)	0.96 $\pm$ 0.06 (0.89-1.10)	1.0 $\pm$ 0.06 (0.9-1.1)	-	0.9 $\pm$ 0.1 (0.8-1.2)	1.1 (0.9-1.2)	0.9 (0.8-1.1)
V	50.5 $\pm$ 1.2 (48.1-53.6)	53 $\pm$ 1.9 (50-55)	53 $\pm$ 0.9 (51.0-54.0)	(50.0-54.0)	52.4 $\pm$ 1.1 (50.9-54.0)	52.1 (50.8-55.0)	50 (47-52)
Total stylet	149.0 $\pm$ 4.4 (144-161)	159 $\pm$ 8.05 (144-173)	-	(156.0-168.3)	166.5 $\pm$ 2.6 (161-171)	-	-
Odontostyle	94.1 $\pm$ 3.3 (88-102)	101 $\pm$ 6.14 (89-110)	101.9 $\pm$ 7.2 (84.7-108.2)	-	107.3 $\pm$ 2.7 (103-111)	103.5 (88.5-120.0)	87 (84-89)
Odontophore	54.9 $\pm$ 2.1 (51-59)	59 $\pm$ 3.43 (50-64)	57.0 $\pm$ 2.9 (48.8-60.0)	(61.2-62.7)	59.2 $\pm$ 1.6 (57-62)	56.7 (53.1-60.0)	50 (48-51)
Oral aperture to guide ring	76.5 $\pm$ 3.8 (67-83)	86.0 $\pm$ 4.23 (77-92)	86.3 $\pm$ 5.6 (72.3-92.3)	-	85.5 $\pm$ 5.0 (78-93)	79.6 (66.0-103.0)	62 (60-64)
Pharyngeal bulb length	79.3 $\pm$ 4.0 (73-86; n = 10)	-	-	(61-79)	84.9 $\pm$ 2.7 (82-90; n = 7)	94.2 (65.0-117)	-
Pharyngeal bulb width	21.2 $\pm$ 1.2 (18-23; n = 21)	-	-	(12-19)	22.9 $\pm$ 1.7 (20-26)	21.1 (18.0-24.0)	-

Tail	27.8 ± 2.0 (25-32)	26.0 ± 1.71 (23-28)	26.8 ± 2.0 (24.1-31.2)	(24.5-29.0)	29.2 ± 2.8 (24-34)	36.1 (33.0-47.0)	24 (21-28)
Hyaline portion of tail (I)	10.4 ± 1.5 (8-13)	-	8.0 ± 0.9 (5.9-9.4)	-	11.4 ± 1.5 (9-14)	9 (7-11)	12 (10-14)
Body diam. at lip region	12.5 ± 0.4 (12-13)	11.5 (11-13)	11.5 ± 0.5 (10.6-12.3)	-	12.4 ± 0.4 (11-13)	14.6 (13.5-15.0)	11 (10-12)
Body diam. at guide ring	30.4 ± 0.8 (29-32)	-	29.8 ± 1.5 (27.1-31.8)	-	32.6 ± 0.8 (31-34)	31.7 (30.0-36.0)	26 (26-27)
Body diam. at base of pharynx	37.9 ± 1.2 (35-40)	-	39.4 ± 2.9 (35.3-45.3)	-	41.3 ± 2.2 (38-45)	40.1 (36.0-42.0)	33 (31-35)
Body diam. at vulva	42.0 ± 2.0 (38-46)	-	46.6 ± 3.4 (39.4-50.0)	(49.0-59.7)	47.2 ± 2.5 (44-52)	43.4 (39.0-47.2)	36 (33-38)
Body diam. at anus	27.0 ± 1.6 (24-30)	26 (22-29)	26.6 ± 1.7 (21.8-29.4)	(30.6-36.7)	29.7 ± 1.1 (28-32)	32.4 (27.0-41.0)	25 (23-28)
Body diam. at beginning of J	16.2 ± 2.2 (12-20)	-	13.7 ± 1.1 (11.2-14.7)	-	18.4 ± 2.2 (15-22)	7 (6-8)	17 (15-20)

**Table 3.** Uterine and vaginal region lengths of *Xiphinema brevicolle* and *Xiphinema* sp. measured in this study (female). Mean  $\pm$  standard deviation (range; number of specimens) in  $\mu\text{m}$ , except where indicated.

	<i>Xiphinema brevicolle</i>	<i>Xiphinema</i> sp.
Anterior uterus length	44.3 $\pm$ 4.8 (34–54; n = 14)	35.1 $\pm$ 2.9 (32–40; n = 9)
Posterior uterus length	41.1 $\pm$ 2.9 (37–46; n = 9)	32.7 $\pm$ 2.7 (29–37; n = 9)
<i>Pars proximalis vaginae</i>	5.9 $\pm$ 0.7 (5–7; n = 22)	7.6 $\pm$ 1.5 (5–10; n = 13)
<i>Pars distalis vaginae</i>	10.5 $\pm$ 0.7 (9–12; n = 22)	11.3 $\pm$ 1.0 (9–13; n = 13)
Vagina length	16.4 $\pm$ 1.0 (14–18; n = 22)	18.9 $\pm$ 1.5 (17–22; n = 13)
Vagina length / body diam. at vulva (%)	39.2 $\pm$ 3.2 (31.3–44.6; n = 22)	40.1 $\pm$ 2.9 (34.8–44.2; n = 13)

*incognitum* Lamberti & Bleve-Zacheo, 1979, described as a new species for nematode specimens detected from soil of bonsai trees exported from Japan to England (Lamberti and Bleve-Zacheo 1979).

The specimens of this study were obtained from mixed populations of the *X. americanum*-group. Information on juveniles was not included because the coexistence of the two closely related nematodes requires special care to separate juvenile specimens of the different species. Situations like this will make matters worse if one is to identify the species, since it is difficult enough to identify even a single population of the group in many cases. Detection of mixed populations of *X. americanum* Cobb, 1913 and *X. rivesi* Dalmasso, 1969 were reported (Vrain et al. 1992, Vrain 1993), and such coexistence of multiple populations and/or species of the *X. americanum*-group may be common. Therefore, it is strongly recommended to check the genetic uniformity of a given population to be identified. The mitochondrial COI region has been recently used to examine variations in populations of *Xiphinema* species including members of the *X. americanum*-group (Lazarova et al. 2006, Kumari et al. 2010a, 2010b), where ca 400 bp sequences were examined. This genetic region has much more information to differentiate between populations than 18S rDNA and D2/D3 regions. As shown above, we developed new primers and examined longer sequences of the COI region than previously used. Examination of longer sequences provides not only more reliable comparative results but also another option to develop a more specific primer to the species to be tested since high variability of this region may contribute to some unfitness of reported primers. Among options available at present, sequences of the COI region can be efficiently used to examine the diversity of *Xiphinema* specimens.

ML phylogenetic analyses of 18S rDNA and D2/D3 regions moderately supported the clade including *X. brevicolle*, *X. diffusum*, *X. inaequale*, *X. incognitum*, *X. lamberti*, and *X. taylori* with our materials (Figs 8, 9), whereas such a clade was not so highly supported in the ML tree of COI (Fig. 10). Member species of *X. americanum*-group harbor endosymbionts (Vandekerckhove et al. 2000). The

difference in phylogenetic inference between nuclear and mitochondrial genetic regions may result from the symbionts' effect on mitochondrial DNA since unreliable results of phylogenetic inference based on mitochondrial DNA due to the presence of a symbiont are known (Hurst and Jiggins 2005). Furthermore, the ML tree of COI brought another problem to us. It showed closer relationship of our *X. brevicolle* population to *X. diffusum* than to other *X. brevicolle* populations. Our specimens identified as *X. brevicolle* were reasonably larger than type specimens of *X. diffusum* (Table 2), and our identification was made considering more overlapping morphometrics of type specimens of *X. brevicolle*. If COI sequences can differentiate species of *X. americanum*-group, our specimens may be identified as *X. diffusum* rather than *X. brevicolle*. In such a situation, morphological features used to identify species should be reconsidered since a single species like *X. diffusum* can have a wide range of morphometrics which may result in more difficult diagnoses without reducing the number of species by synonymizing them intensively. In any case, it should be desirable to collect much more COI sequences of other species and populations, such as the sequence data of topotype specimens of *X. diffusum*, which is unavailable at present. Our results of phylogenetic analyses, however, may be helpful to refine the concept of a *X. brevicolle*-subgroup, which was previously discussed in some works (Romanenko and Stegaresku 1985, Lamberti and Ciancio 1993, He et al. 2005b). Taking our results of phylogenetic analyses into consideration, we suggest that the *X. brevicolle*-subgroup includes at least the five species of *X. brevicolle*, *X. diffusum*, *X. inaequale*, *X. incognitum*, and *X. taylori*, and our specimens also belong to the subgroup, though the validity of respective species is a different matter. Appropriate establishment of subgroups within the *X. americanum*-group will contribute to a more feasible identification process of the member species and requires further research.

## Conclusion

In this study, specimens from mixed populations of the *X. americanum*-group, present in the root zone of Japanese holly in Japan, were identified as *X. brevicolle* and *Xiphinema* sp. This record of *X. brevicolle* is the first for Japan. PCR primers to amplify longer sequences of the mitochondrial COI regions were originally designed and used to efficiently differentiate the specimens. Phylogenetic analyses using 18S rDNA, D2/D3, and COI regions supported a close relationship among our specimens and species related to *X. brevicolle* or the *X. brevicolle*-subgroup.

## Acknowledgements

The first author thanks Tom Prior for useful suggestions and information and Dr. Nobuhiro Minaka for helpful comments on phylogenetic analyses. The authors thank Dr. Jerome


T. Gaspard for improving the manuscript. This study was supported by Research and development projects for application in promoting new policy of Agriculture Forestry and Fisheries (21043) from the Ministry of Agriculture, Forestry and Fisheries of Japan.

## References


- De Grisse AT (1969) Redescription ou modifications de quelques techniques utilisées dans l'étude des nématodes phytoparasitaires. Mededelingen van de Rijksfaculteit Landbouwetenschappen Gent 34: 351–369.
- De Ley P, Félix M-A, Frisse LM, Nadler SA, Sternberg PW, Thomas K (1999) Molecular and morphological characterisation of two reproductively isolated species with mirror-image anatomy (Nematoda: Cephalobidae). Nematology 1: 591–612.
- Edgar RC (2004) MUSCLE: multiple sequence alignment with high accuracy and high throughput. Nucleic Acids Research 32: 1792–1797. doi: 10.1093/nar/gkh340
- Hall TA (1999) BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. Nucleic Acids Symposium Series 41: 95–98.
- He Y, Jones J, Armstrong M, Lamberti F, Moens M (2005a) The mitochondrial genome of *Xiphinema americanum sensu stricto* (Nematoda: Enoplea): considerable economization in the length and structural features of encoded genes. Journal of Molecular Evolution 61: 819–833. doi: 10.1007/s00239-005-0102-7
- He Y, Subbotin SA, Rubtsova TV, Lamberti F, Brown DJF, Moens M (2005b) A molecular phylogenetic approach to Longidoridae (Nematoda: Dorylaimida). Nematology 7: 111–124. doi: 10.1163/1568541054192108
- Howe DK, Denver DR (2008) Muller's Ratchet and compensatory mutation in *Caenorhabditis briggsae* mitochondrial genome evolution. BMC Evolutionary Biology 8: 62. doi: 10.1186/1471-2148-8-62
- Hurst GDD, Jiggins FM (2005) Problems with mitochondrial DNA as a marker in population, phylogeographic and phylogenetic studies: the effects of inherited symbionts. Proceedings of the Royal Society B 272: 1525–1534. doi: 10.1098/rspb.2005.3056
- Kumari S, Decraemer W, De Luca F (2010a) Molecular characterization of *Xiphinema brevicollum* (Nematoda: Longidoridae) from the Czech Republic. European Journal of Plant Pathology 128: 243–250. doi: 10.1007/s10658-010-9651-8
- Kumari S, Decraemer W, De Luca F, Tiefenbrunner W (2010b) Cytochrome c oxidase subunit 1 analysis of *Xiphinema diversicaudatum*, *X. pachtaicum*, *X. simile* and *X. vuittenzei* (Nematoda, Dorylaimida). European Journal of Plant Pathology 127: 493–499. doi: 10.1007/s10658-010-9614-0
- Lamberti F, Bleve-Zacheo T (1979) Studies on *Xiphinema americanum sensu lato* with descriptions of fifteen new species (Nematoda, Longidoridae). Nematologia Mediteranea 7: 51–106.
- Lamberti F, Ciancio A (1993) Diversity of *Xiphinema americanum*-group species and hierarchical cluster analysis of morphometrics. Journal of Nematology 25: 332–343.

- Lamberti F, Ciancio A, Agostinelli A, Coiro MI (1992) Relationship between *Xiphinema brevicolle* and *X. diffusum* with a redescription of *X. brevicolle* and descriptions of three new species of *Xiphinema* (Nematoda: Dorylaimida). *Nematologia Mediterranea* 19 (1991): 311–326.
- Lazarova SS, Malloch G, Oliveira CMG, Hübschen J, Neilson R (2006) Ribosomal and mitochondrial DNA analyses of *Xiphinema americanum*-group populations. *Journal of Nematology* 38: 404–410.
- Lordello LGE, Costa CP (1961) A new nematode parasite of coffee roots in Brazil. *Revista Brasileira de Biologia*, 21: 363–366.
- Luc M, Coomans A, Loof PAA, Baujard P (1998) The *Xiphinema americanum*-group (Nematoda: Longidoridae). 2. Observations on *Xiphinema brevicollum* Lordello & da Costa, 1961 and comments on the group. *Fundamental and Applied Nematology* 21: 475–490.
- Monteiro AR (2010) Nomenclatorial note on *Xiphinema brevicolle* Lordello & Costa, 1961 (Nematoda: Longidoridae). *Nematologia Brasileira* 34: 1–2.
- Mullin PG, Harris TS, Powers TO (2005) Phylogenetic relationships of Nygolaimina and Dorylaimina (Nematoda: Dorylaimida) inferred from small subunit ribosomal DNA sequences. *Nematology* 7: 59–79. doi: 10.1163/1568541054192199
- Netscher C, Seinhorst JW (1969) Propionic acid better than acetic acid for killing nematodes. *Nematologica* 15: 286.
- Romanenko ND (1981) [Discovery of a new species of nematode *Xiphinema paramonovi* n. sp. (Nematoda: Longidoridae) in the territory of the soviet Union]. *Tezisy Dokladov, Pervaya Konferentsiya po Nematodam Rastenii, Nasekomykh, Pochvy i Vod*, 68–69.
- Romanenko ND, Stegaresku OP (1985) [Taxonomic problem in nematodes of the *Xiphinema americanum* group (Nematoda, Dorylaimoidea)]. *Parazitologiya* 19: 434–442.
- Sakai H (2010) A DNA extraction method with SDS from single nematodes for direct application to PCR amplification. *Nematological Research* 40: 13–14. doi: 10.3725/jjn.40.13
- Seinhorst JM (1959) A rapid method for the transfer of nematodes from fixative to anhydrous glycerin. *Nematologica* 4: 67–69. doi: 10.1163/187529259X00381
- Tamura K, Peterson D, Peterson N, Stecher G, Nei M, Kumar S (2011) MEGA5: Molecular Evolutionary Genetics Analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Molecular Biology and Evolution*. doi: 10.1093/molbev/msr121
- Thomas WK, Vida JT, Frisse LM, Mundo M, Baldwin JG (1997) DNA Sequences from formalin-fixed nematodes: integrating molecular and morphological approaches to taxonomy. *Journal of Nematology* 29: 250–254.
- Vandekerckhove TTM, Willems A, Gillis M, Coomans A (2000) Occurrence of novel verucomicrobial species, endosymbiotic and associated with parthenogenesis in *Xiphinema americanum*-group species (Nematoda, Longidoridae). *International Journal of Systematic and Evolutionary Microbiology* 50: 2197–2205. doi: 10.1099/00207713-50-6-2197
- Vrain TC (1993) Restriction fragment length polymorphism separates species of the *Xiphinema americanum* group. *Journal of Nematology* 25: 361–364.
- Vrain TC, Wakarchuk DA, Lévesque AC, Hamilton RI (1992) Intraspecific rDNA restriction fragment length polymorphism in the *Xiphinema americanum* group. *Fundamental and Applied Nematology* 15: 563–573.


# Figures




**Figure 1.** Female morphology of *Xiphinema brevicolle* (A, C, E, G) and *Xiphinema* sp. (B, D, F, H) A–B Anterior region C–D Tail E–F Reproductive system G–H Entire body.




**Figure 2.** *Xiphinema brevicolle* (female) **A–C** Anterior region **D–G** Tail **H–K** Vulval region **L–P** Entire body.




**Figure 3.** *Xiphinema* sp. (female) **A–C** Anterior region **D–F** Tail **G–I** Vulval region **J–M** Entire body.



**Figure 4.** Intra-population variation of lip region in females **A** *Xiphinema brevicolle* **B** *Xiphinema* sp.




**Figure 5.** Intra-population variation of tail in females **A** *Xiphinema brevicolle* **B** *Xiphinema* sp.




**Figure 6.** Intra-population variation of vagina in females **A** *Xiphinema brevicolle* **B** *Xiphinema* sp.

Xbre	1	AATTTAATTGGAGGATTGGTAATATTTATTGCCGCTGATACTAGGAGCCCAAGATATGTGTTTCCTCGTTTAAATA	80
Xsp1	1	G.....C.....A.A.A.A...T...A...C...A.....A.....	80
Xsp2	1	G.....C.....A.A.A.A...T...A...C...A.....A.....	80
Xame	9214	...C.....G.....A..CTC...T...A...C...C.C.C.C..G..C.	9293
Xbre	81	ATTTTAGATTTTGGCTTTTACCGATCAGAGGATTGTTTATATTAAGCTCTATATTTATAGTTGTGGCGCTGGAACCGGA	160
Xsp1	81	.....A.....AG.A.....A...T..A.....A.....A.....G..T..	160
Xsp2	81	.....C.....A.....AG.A.....A...T..A.....A.....A.....G..T..	160
Xame	9294	.C.....G...AT.A.....G...G..AC...A...T...C.C.G.A.C.T...T..T..T	9373
Xbre	161	TGAACATTTTATCCTCCGCTTAGAGGTATTCAGGTCATTCTAATTGAAGGTAGACTTAGTAATTTTATGCTACACTT	240
Xsp1	161	.....TA.A..CA...A..C..A..C...G.....AT.....	240
Xsp2	161	.....TA.A..CA...A..C..A..C...G.....AT.....	240
Xame	9374	.....C...T...A.....A.....G..C...A..G.....G.....T..G..TC.	9453
Xbre	241	AGCTGGAGTAAGATCGATTGCTGGATCCATTATTTTGTGTACTATTAATAATTTAAAAACAGTGCCTATTTTCATGAA	320
Xsp1	241	.....A.....T.....A.....T.....C.....G..	320
Xsp2	241	.....A.....T.....A.....T.....C.....G..	320
Xame	9454	.....T..T...A..C...T..T.....C.....CC.T..GTCT..AT.A..C.T...T...	9533
Xbre	321	TGTCATACCTTTATTTTGGATTAGGGTATGAGTAACCGCTTTTGTAGTTCTTAGATTACCAAGTATTTGGCAGGTGGA	400
Xsp1	321	.A....G.....CC.T...A..T..G..T..T....C.TC...T.A..C...T..GC.A.....G	400
Xsp2	321	.A....G.....CC.T...A..T..G..T..T....C.TC...T.A..C...T..GC.A.....G	400
Xame	9534	.A.GT...G..G.....A..T...T..T.....G...T.A.....T..T..A..T.....	9613
Xbre	401	ATTACGATGCTTTGTGTTGATCGAAATTTAAACACTTCTTTCTTTGATCCTTTAGGAGGGGGGATCCAATTTTATTC	480
Xsp1	401	.....T..AT.A..A.....C...T.....G..T.....A..T.....C.....T..	480
Xsp2	401	.....T..AT.A..A.....C...T.....G..T.....C.....A..T.....C.....T..	480
Xame	9614	.....T..A.....A..C.....C..G.....C.....C...AC...T.A.....C...T..	9693
Xbre	481	ACATTTATCTGATTTTGGGCACCCAGAGTATATATTTTAATTTTACCTGGGTTGGATTAGTTAGGCATGCTGTTA	560
Xsp1	481	...CC.C.....C...T..T.....A.....G..A.....A.....AG	560
Xsp2	481	...CC.C.....C...T..T.....A.....G..A.....A.....AG	560
Xame	9694	G...C..T..T...C...A.....G..G..G.....G.....A..A..C..C.....	9773
Xbre	561	TAGTTTCTAGAGGAAAGCCATCTCCGTTTGGTGTTCCAGGTATGTTTTAGCTATTACTAGGATTGGTGCTCTAGGTTGT	640
Xsp1	561	.....C.....AA.....T.....AA.C..G.....C.....A.....A.....A...	640
Xsp2	561	.....C.....AA.....T.....AA.C..G.....C.....A.....A.....A...	640
Xame	9774	..A...C.....T...A.....G..A...C..A..C.....A.....A..G..A..C	9853
Xbre	641	GTGGTCTGAGCGCATCATATATTTAGAGTAGGATTAGACATAGATACTCGACTGTATTTTACAGCGGCCACTATAATTAT	720
Xsp1	641	..A..T...T.....G.....T..T.....C..A...T.A.....T.....	720
Xsp2	641	..A..T...T.....G.....T..T.....C..A...T.A.....T.....	720
Xame	9854	..A..A...A.....C..G.....G..T.....T.....T.....A.....	9933
Xbre	721	TGCTGTTCCAACCTGGGATTAAGGTATTCAGATGATTGGTATCCTTTAGAGGAAGAAAAATATTATTGCCCGCTCCAAT	800
Xsp1	721	C....A..T..A..A.....GA.A..A..TACT...	800
Xsp2	721	C....A..T..A..A.....GA.A..A..TACT...	800
Xame	9934	.....A.....T.....G..A..C...T.....G.....G...A..ATA..A..A...	10013
Xbre	801	TGTGGATTTTAGGCTTTTGTGTTTTATTACAGTGGGAGGACTTACTGGCATTGTATTAGCCAACGGGACTTTAGATTTA	880
Xsp1	801	.A..AG.AC.T..T..C..A.....T.A...T...T...T..T..A.....	880
Xsp2	801	.A..AG.AC.T..T..C..A.....T.A...T...T...T..T..A.....	880
Xame	10014	.A..A...C..T..T..C..A.....G.....G..T..T..A...T...G...T..T.....	10093
Xbre	881	TTATAT	886
Xsp1	881	C.T...	886
Xsp2	881	C.T...	886
Xame	10094	..G..C	10099


**Figure 7.** Sequence comparison of mitochondrial COI region. Xbre: *X. brevicolle* (this study; GenBank accession AB604337); Xsp1, Xsp2: *Xiphinema* sp. (this study: AB604338 and AB604339 respectively); Xame: *X. americanum* (He et al. 2005: usAY382608).



**Figure 8.** Maximum likelihood tree for 18S rDNA region. Bootstrap values higher than 50 are shown. Arrows indicate specimens examined in this study: *Xiphinema brevicolle* GenBank accession AB604340; *Xiphinema* sp. AB604341. The tree is drawn to scale, with branch lengths measured in the number of substitutions per site.



**Figure 9.** Maximum likelihood tree for D2/D3 region. Bootstrap values higher than 50 are shown. Arrows indicate specimens examined in this study: *Xiphinema brevicolle* GenBank accession AB635401 and AB635402; *Xiphinema* sp. AB635403. The tree is drawn to scale, with branch lengths measured in the number of substitutions per site.



**Figure 10.** Maximum likelihood tree for COI region. Bootstrap values higher than 50 are shown. Arrows indicate specimens examined in this study. The tree is drawn to scale, with branch lengths measured in the number of substitutions per site.