

Calls of *Boana latistriata* (Caramaschi & Cruz, 2004) (Amphibia, Anura, Hylidae), an endemic tree frog from the State of Minas Gerais, Brazil

Cyro de Luna-Dias¹, Sergio P. de Carvalho-e-Silva¹

¹ *Laboratório de Anfíbios e Répteis. Departamento de Zoologia, Instituto de Biologia, Universidade Federal do Rio de Janeiro (UFRJ). Av. Carlos Chagas Filho, 373 – Prédio do CCS, Bloco A, Sala A1-111, Cidade Universitária, CEP 21941-902, Rio de Janeiro, RJ, Brazil*

Corresponding author: *Cyro de Luna Dias* (cyroluna@gmail.com)

Academic editor: *A. Crottini* | Received 20 October 2018 | Accepted 11 January 2019 | Published 29 January 2019

<http://zoobank.org/B6F99B28-E27D-43B8-9B86-E761B0AEAD32>

Citation: Luna-Dias C, Carvalho-e-Silva SP (2019) Calls of *Boana latistriata* (Caramaschi & Cruz, 2004) (Amphibia, Anura, Hylidae), an endemic tree frog from the State of Minas Gerais, Brazil. ZooKeys 820: 83–94. <https://doi.org/10.3897/zookeys.820.30711>

Abstract

Bioacoustical data are useful for studying amphibians, especially their conservation, taxonomy, and evolution, among others. Of the 12 species of the *Boana polytaenia* clade, only *B. buriti* and *B. latistriata* have no published information about their advertisement calls. We recorded four males of *B. latistriata* in its type locality at Parque Nacional do Itatiaia, south-eastern Brazil. We used a Roland R26 digital recorder with a Sennheiser ME-67 microphone and analysed the recordings using the Raven Pro 1.5 software. We recorded two different types of calls (call A and call B). Both were composed of one pulsed note and presented a slightly ascending-descending frequency modulation. Call A was more frequent, having durations between 0.042 and 0.093 s with the dominant frequency ranging from 3375.0 to 3937.5 Hz, and was composed of 11 to 21 pulses separated by intervals that were not fully silent. Call B had durations between 0.711 and 1.610 s, with dominant frequency from 3281.2 to 3750.0 Hz, and was composed of 11 to 29 pulses separated by fully silent intervals. Among the *B. polytaenia* clade, the calls of *B. latistriata* are more similar to those of *B. bandeirantes*, *B. beckeri*, *B. polytaenia*, and *B. aff. beckeri*. The calls of *B. latistriata* differ from these species in its lower dominant frequency. *Boana latistriata* present a short, single-note call with a lower pulse period (call A) and a long call with higher pulse period (call B), which differ from the other species of the clade. The coefficients of variation for the various bioacoustical attributes were calculated within- and between-males and these have been discussed. We also report a fight event between two males of *B. latistriata*. This is the first report of a fight in members of the *B. polytaenia* clade.

Keywords

Cophomantini, *Boana pulchella* species group, communication, bioacoustics, taxonomy

Introduction

Vocalization plays an essential role during the reproductive period of anurans, being species-specific and constituting a pre-zygotic mechanism of reproductive isolation (Duellman and Trueb 1994, Wells 2007). This makes bioacoustical data useful for studying topics such as conservation (Laiolo 2010, Forti et al. 2017), taxonomy (Hepp and Carvalho-e-Silva 2011, Carvalho and Giarretta 2013, Rivadeneira et al. 2018), social interaction (Giasson and Haddad 2006), and evolution (Robillard et al. 2006, Goicoechea et al. 2010). Amongst the different anuran call types, advertisement calls have the highest value in taxonomy (Köhler et al. 2017). Despite being the most studied call type, there is a need for further documentation (Toledo et al. 2014, Guerra et al. 2018a) and understanding of the importance of advertisement call variation both within- and between-individuals (Gerhardt 1991, Köhler et al. 2017).

The *Boana polytaenia* clade is composed of 12 species of tree frogs with a striped dorsal pattern (Caramaschi and Cruz 2013): *B. bandeirantes* (Caramaschi & Cruz, 2013); *B. beckeri* (Caramaschi & Cruz, 2004); *B. botumirim* (Caramaschi, Cruz & Nascimento, 2009); *B. buriti* (Caramaschi & Cruz, 1999); *B. cipoensis* (Lutz, 1968); *B. goiana* (Lutz, 1968); *B. jaguariaivensis* (Caramaschi, Cruz & Segalla, 2010); *B. latistriata* (Caramaschi & Cruz, 2004); *B. leptolineata* (Braun & Braun, 1977); *B. phaeopleura* (Caramaschi & Cruz, 2000); *B. polytaenia* (Cope, 1870); and *B. stenocephala* (Caramaschi & Cruz, 1999). Of these, there is a lack of information on the advertisement calls of *B. buriti* and *B. latistriata* alone, while the information for *B. cipoensis* and *B. leptolineata* is very limited (Kwet 2001, Batista et al. 2015).

Boana latistriata was described from Brejo da Lapa, an artificial pond in the Parque Nacional do Itatiaia (PNI), Itamonte Municipality, State of Minas Gerais, Brazil (Caramaschi and Cruz 2004). The type series of this species also includes five individuals from Marmelópolis, in the same state, which is the only record of the species outside the PNI. It is the largest species of the *B. polytaenia* clade, with the males measuring 34.9–40.6 mm and females 40.9–51.6 mm (Caramaschi and Cruz 2004). Tadpoles of this species have been described by Orrico et al. (2007). Toledo and Haddad (2009) have reported the distress call of one female of this species. The lack of information about *B. latistriata* led IUCN to list it as being “Data Deficient” (Stuart 2018), thus supporting the need to study this species. In this paper, we describe the advertisement call of *B. latistriata* from its type locality and report an event of a combat between males of this species.

Material and methods

Recordings were made at Brejo da Lapa (22.3589°S, 44.7372°W, 2140 m altitude) on 12 November 2014, from 20:30 to 23:00. Vocalizations from four males were recorded

with a Roland R26 digital recorder and a Sennheiser ME-67 shotgun microphone positioned between 20 and 50 cm from the calling males. After the calls were recorded, the specimens were collected, anesthetised and euthanized with 5% lidocaine, fixed in 10% formaldehyde, and subsequently preserved in 70% ethanol. These have been deposited in the amphibian collection of the Departamento de Zoologia, Instituto de Biologia, Universidade Federal do Rio de Janeiro (ZUFRJ), with collection numbers 15073, 15074, 15076, and 15077. Recordings were deposited at Fonoteca Neotropical Jacques Viellard (FNJV; <https://www2.ib.unicamp.br/fnjv/>) with the respective numbers 40238, 40239, 40240 and 40241. All procedures were conducted under licence No. 40371 issued by the Instituto Chico Mendes de Conservação da Biodiversidade – ICMBio.

Recordings were made using 24 bits of resolution and 48 kHz sampling rate. Sound analyses were performed using the Raven Pro 1.5 software (Bioacoustics Research Program 2014) using window type = Hann, size = 256 samples, and overlap = 99%. Measurements and terminology follow Köhler et al. (2017). Terminology used to describe calls of the species belonging to the *B. polytaenia* clade (calls, notes and pulses) is highly variable (for example, see Acioli and Toledo 2008, Pinheiro et al. 2012, Martins et al. 2016). To keep nomenclature stability, we opted to follow the most recent papers on this topic (e.g. Martins et al. 2016, Guerra et al. 2017) and adopted a call centred approach (Köhler et al. 2017).

The following parameters were measured or calculated: call duration (CD), pulse duration (PD), pulse period (PP), interval between “call A” and “call B” (ABI), call dominant frequency (DF), call fundamental frequency (FF; measured through the “Peak Frequency” function of Raven Pro), number of visible harmonics (NH; integer multiples of the fundamental frequency), first pulse dominant frequency (FPDF), central pulse dominant frequency (CPDF, measured in the central pulse with higher dominant frequency), last pulse dominant frequency (LPDF), pulse rise time (RT; measured through the “Max Time” function of Raven Pro), proportion of pulse rise time in relation to pulse duration (RTR), and pulse number (PN). Measurements are given as a range, followed by mean and standard deviation (SD). For DF, FPDF, CPDF, LPDF, and PN, the mode (Mo; the most frequent value among the measurements) is also presented.

To determine the variation in the bioacoustical attributes, we calculated the within-individual coefficient of variation (CV_w ; Gerhardt 1991) for each parameter of each male. Between-individual coefficient of variation (CV_b) was calculated for each parameter by pooling the measurements from all the males. Parameters with a coefficient of variation below 5.0% were considered static, whereas those with a coefficient of variation above 12.0% were considered dynamic (Gerhardt 1991).

Results

Males of *B. latistriata* called from dusk (about 18:00) to at least 02:00, perched on shrubs and grass, near or above the water. At the time of the recordings, we visually counted more than 20 males of *B. latistriata* and heard many more at a distance. Dur-

ing the fieldwork, the weather was rainy, air temperature between 12 and 14 °C, and humidity above 90%. Other amphibian species recorded at the site were *Aplastodiscus albosignatus* (Lutz & Lutz, 1938), *Bokermannohyla gouveai* (Peixoto & Cruz, 1992), *Rhinella icterica* (Spix, 1824), and *Scinax duartei* (Lutz, 1951).

We recorded two types of calls from the males of *B. latistriata*, herein called “call A” and “call B” (Fig. 1). Call A was emitted more frequently (103 of 142 recorded calls in the complete dataset), and we interpreted it as an advertisement call due to the social context in which it was emitted (frequently emitted even by isolated males, without aggressive interactions). Call B was always emitted after call A, with ABI varying from 0.545 to 1.622 s (mean = 0.782 s, SD = 0.231, n = 38 intervals), and possibly had some degree of territorial function.

Call A was composed of a single, short, pulsed note, with up to seven visible harmonics and duration from 0.042 to 0.093 s (mean = 0.073 s, SD = 0.012, n = 97 calls; Fig. 1C, D). On an FFT size of 256 (270 Hz 3 dB filter bandwidth), sidebands are visible between the harmonics, caused by the pulse rate of the call. The dominant frequency was equal to the fundamental frequency, varying from 3375.0 to 3937.5 Hz (mean = 3626.3 Hz, SD = 168.2, Mo = 3562.5 Hz, n = 97 calls). Each call was composed of 11 to 21 pulses (mean = 15.5 pulses, SD = 2.4, Mo = 15 pulses, n = 68 calls) separated by intervals that were not fully silent, resulting in pulse duration equal to pulse period, which varied from 0.001 to 0.010 s (mean = 0.005 s, SD = 0.001, n = 1049 pulses). Each pulse had a rise time from 0.001 to 0.005 s (mean = 0.002 s, SD = 0.001, n = 1049 pulses), corresponding to 13.8–89.2% of the pulse duration (mean = 36.2%, SD = 8.7, n = 1049 pulses). A slightly ascending-descending frequency modulation is visible from the first to the last pulses, with the first pulse dominant frequency from 3000.0 to 3937.5 Hz (mean = 3547.3 Hz, SD = 190.0, Mo = 3562.5 Hz, n = 68 pulses), central pulse dominant frequency from 3375.0 to 4031.2 Hz (mean = 3745.9 Hz, SD = 143.5, Mo = 3843.8 Hz, n = 68 pulses), and last pulse dominant frequency from 3000.0 to 3843.8 Hz (mean = 3438.4 Hz, SD = 218.1, Mo = 3375.0 Hz, n = 68 pulses). In one individual (ZUF RJ 15077), all calls of this type presented some pulses fused in a pulsatile (Fig. 2), which made it impossible to count pulse number and to measure pulse parameters properly. Table 1 shows the parameters for each recorded male.

Call B was composed of a single, long, pulsed note, with up to seven visible harmonics and call duration from 0.711 to 1.610 s (mean = 1.114 s, SD = 0.279, n = 37 calls; Fig. 1E, F). The dominant frequency was equal to the fundamental frequency, varying from 3281.2 to 3750.0 Hz (mean = 3435.8 Hz, SD = 140.1, Mo = 3468.8 Hz, n = 37 calls). Each call was composed of 11 to 29 pulses (mean = 17.8 pulses, SD = 4.5, Mo = 17, n = 37 calls) separated by a fully silent interval. Pulse duration varied from 0.003 to 0.013 s (mean = 0.008 s, SD = 0.002, n = 682 pulses) and pulse period varied from 0.009 to 0.196 s (mean = 0.065 s, SD = 0.027, n = 645 periods). Each pulse had a rise time from 0.001 to 0.004 s (mean = 0.002 s, SD = 0.000, n = 682 pulses), corresponding to 14.2–51.6% of the pulse duration (mean = 26.9, SD = 5.7, n = 682 pulses). A slightly ascending-descending frequency modulation is visible from the first to the last pulses, with first pulse dominant frequency from 3000.0 to 3562.5 Hz (mean = 3310.9, SD = 138.0, Mo = 3375.0 Hz, n = 38 pulses), central pulse domi-




Figure 1. Calls of *Boana latistriata* from its type locality. **A** Waveform and **B** spectrogram showing three instances of call A and one of call B emitted in sequence **C** waveform and **D** spectrogram of a call A, in detail, indicating seven visible harmonics numbered I to VII **E** waveform and **F** spectrogram of a call B, in detail. Images were obtained using Raven Pro 1.5 software. Spectrograms parameters: window type = Hann, size = 256 samples, overlap = 99%. Individual ZUFRJ 15076 (snout-vent length = 39.3 mm), recorded at a temperature between 12 and 14 °C.

nant frequency from 3187.5 to 3750.0 Hz (mean = 3473.6 Hz, SD = 178.6, Mo = 3468.8 Hz, n = 38 pulses), and last pulse dominant frequency from 2625.0 to 3656.2 Hz (mean = 3399.7 Hz, SD = 195.8, Mo = 3375.0 Hz, n = 38 pulses). Table 2 shows the parameters for each recorded male.




Figure 2. Calls of *Boana latistriata* from its type locality. Two examples of call A, emitted by the individual ZUFRJ 15077 (snout-vent length = 40.0 mm), showing the last pulses fused in a pulsatile (below the red lines).

In general, spectral parameters were static, while temporal parameters were dynamic. The most static parameter was the dominant frequency (mean CV_w and CV_b was 2.1% and 4.6% in call A and 1.5% and 4.1% in call B, respectively), while pulse period of call B was the most dynamic (CV_w and CV_b was 38.5% and 41.4%, respectively). Some parameters, including first pulse dominant frequency and last pulse dominant frequency in call A and central pulse dominant frequency and last pulse dominant frequency in call B, were intermediate in CV_b , but static in CV_w . All CV_w and CV_b values are shown in Tables 1 and 2.

A fight event between two males of *B. latistriata* was witnessed during the recordings. A male (M1) was calling from the marginal vegetation, perched at approximately 20 cm from the water surface. A second male (M2) was calling from a floating shrub at approximately 40 cm from M1. Without any previous alteration on vocalization, M1 jumped over M2 and started the fight, both grasping and kicking. This first round lasted less than a second and resulted in M2 moving approximately 20 cm away. After a few seconds, M1 pursued M2, starting a second round of fighting, which lasted about the same time as the first one. After this, M2 swam to the other side of the pond, where it started calling a few minutes later. M1 returned to its original calling site and started calling again immediately. M1 was then recorded and collected, and is one of the individuals included in this study (ZUFRJ 15076). Individual M2 was not recorded nor collected.

Table 1. Call parameters for the call A of each recorded individual of *Boana latistriata*. Values are given as range, mean, standard deviation (SD), mode (Mo; when applicable) and sample number (n). Abbreviations: SVL, snout-vent length; CV_w , within-individual coefficient of variation; CV_b , between-individual coefficient of variation; CD, call duration; DF, dominant frequency; FF, fundamental frequency; PD, pulse duration; PP, pulse period; RT, pulse rise time; RTR, proportion of pulse rise time in relation to pulse duration; FPDF, first pulse dominant frequency; CPDF, central pulse dominant frequency; LPDF, last pulse dominant frequency; PN, pulse number. * All instances of call A of this individual presented some pulses fused in a pulsatile, which made it impossible to measure pulse parameters properly.

Parameters (Call A)	Individuals				Mean $C_{y,w}$	$C_{y,b}$
	ZUF RJ 15073 SVL = 35.4 mm	ZUF RJ 15074 SVL = 39.2 mm	ZUF RJ 15076 SVL = 39.3 mm	ZUF RJ 15077 SVL = 40.0		
CD (s)	0.066–0.089 mean = 0.080 SD = 0.006; n = 31 CV_w = 7.1%	0.067–0.078 mean = 0.072 SD = 0.003; n = 15 CV_w = 4.7%	0.077–0.094 mean = 0.084 SD = 0.004; n = 22 CV_w = 5.2%	0.042–0.078 mean = 0.059 SD = 0.008; n = 29 CV_w = 13.6%	7.7%	16.2%
PD (s) (= PP)	0.003–0.011 mean = 0.006 SD = 0.001; n = 434 CV_w = 18.5%	0.001–0.011 mean = 0.005 SD = 0.001; n = 215 CV_w = 22.3%	0.001–0.008 mean = 0.005 SD = 0.001; n = 406 CV_w = 19.9%	*	20.2%	24.0%
PN	11–16 mean = 13.9 SD = 1.1 Mo = 15; n = 31 CV_w = 8.1%	11–17 mean = 14.3 SD = 1.5 Mo = 15; n = 15 CV_w = 10.8%	17–21 mean = 18.5 SD = 1.1 Mo = 18; n = 22 CV_w = 6.2%	*	8.4%	15.5%
RT (s)	0.001–0.004 mean = 0.002 SD = 0.001; n = 434 CV_w = 23.9%	0.001–0.005 mean = 0.002 SD = 0.000; n = 215 CV_w = 24.4%	0.001–0.003 mean = 0.002 SD = 0.000; n = 406 CV_w = 28.8%	*	25.7%	25.0%
RTR (%)	13.8–89.2 mean = 37.5 SD = 11.5; n = 434 CV_w = 30.8%	18.2–64.3 mean = 37.4 SD = 5.6; n = 215 CV_w = 15.1%	15.4–63.6 mean = 34.3 SD = 5.9; n = 406 CV_w = 17.1%	*	21.0%	24.1%
DF (Hz) (= FF)	3656.2–3937.5 mean = 3807.5 SD = 92.8 Mo = 3750.0; n = 31 CV_w = 2.5%	3656.2–3843.8 mean = 3712.5 SD = 69.1 Mo = 3656.2; n = 15 CV_w = 1.9%	3375.0–3656.2 mean = 3536.9 SD = 71.9 Mo = 3562.5; n = 22 CV_w = 2.0%	3375.0–3562.5 mean = 3455.8 SD = 69.6 Mo = 3468.8; n = 29 CV_w = 2.0%	2.1%	4.6%
FPDF (Hz)	3375.0–3937.5 mean = 3668.3 SD = 132.0 Mo = 3656.2; n = 31 CV_w = 3.6%	3562.5–3750.0 mean = 3618.7 SD = 69.1 Mo = 3562.5; n = 15 CV_w = 1.9%	3000.0–3468.8 mean = 3328.1 SD = 107.3 Mo = 3375.0; n = 22 CV_w = 3.2%	*	2.9%	5.4%
CPDF (Hz)	3656.2–4031.2 mean = 3861.9 SD = 85.3 Mo = 3843.8; n = 31 CV_w = 2.2%	3656.2–3843.8 mean = 3737.5 SD = 69.7 Mo = 3750.0; n = 15 CV_w = 1.9%	3375.0–3750.0 mean = 3588.1 SD = 77.5 Mo = 3562.5; n = 22 CV_w = 2.2%	*	2.1%	3.8%
LPDF (Hz)	3468.8–3843.8 mean = 3634.9 SD = 95.9 Mo = 3656.2; n = 31 CV_w = 2.6%	3375.0–3468.8 mean = 3393.8 SD = 38.8 Mo = 3375.0; n = 15 CV_w = 1.1%	3000.0–3468.8 mean = 3191.8 SD = 124.4 Mo = 3187.5; n = 22 CV_w = 3.9%	*	2.5%	6.3%

Table 2. Call parameters for the call B of each recorded individual of *Boana latistriata*. Values are given as range, mean, standard deviation (SD), mode (Mo; when applicable) and sample number (n). Abbreviations: SVL, snout-vent length; CV_w, within-individual coefficient of variation; CV_b, between-individual coefficient of variation; CD, call duration; DF, dominant frequency; FF, fundamental frequency; PD, pulse duration; PP, pulse period; RT, pulse rise time; RTR, proportion of pulse rise time in relation to pulse duration; FPDF, first pulse dominant frequency; CPDF, central pulse dominant frequency; LPDF, last pulse dominant frequency; PN, pulse number. Mode followed by an “-” indicates that no value was more frequent than the others.

Parameters (Call B)	Individuals				Mean CV _w	CV _b
	ZUF RJ 15073 SVL = 35.4 mm	ZUF RJ 15074 SVL = 39.2 mm	ZUF RJ 15076 SVL = 39.3 mm	ZUF RJ 15077 SVL = 40.0		
CD (s)	1.030–1.234 mean = 1.130 SD = 0.073; n = 12 CV _w = 6.5%	1.301–1.476 mean = 1.384 SD = 0.550; n = 8 CV _w = 4.0%	1.564–1.610 mean = 1.577 SD = 0.022; n = 4 CV _w = 1.4%	0.711–0.894 mean = 0.814 SD = 0.048; n = 14 CV _w = 5.9%	4.5%	25.1%
PD (s)	0.003–0.012 mean = 0.007 SD = 0.002; n = 214 CV _w = 26.4%	0.001–0.010 mean = 0.008 SD = 0.001; n = 170 CV _w = 14.6%	0.003–0.010 mean = 0.008 SD = 0.001; n = 106 CV _w = 13.0%	0.004–0.013 mean = 0.009 SD = 0.001; n = 192 CV _w = 13.4%	16.9%	21.2%
PP (s)	0.009–0.152 mean = 0.066 SD = 0.027; n = 203 CV _w = 40.7%	0.024–0.166 mean = 0.068 SD = 0.021; n = 162 CV _w = 31.1%	0.024–0.121 mean = 0.062 SD = 0.017; n = 102 CV _w = 27.2%	0.026–0.196 mean = 0.063 SD = 0.035; n = 178 CV _w = 54.9%	38.5%	41.4%
PN	15–24 mean = 17.4 SD = 2.3 Mo = 17; n = 12 CV _w = 13.5%	20–23 mean = 21.3 SD = 1.0 Mo = 21; n = 8 CV _w = 4.9%	24–29 mean = 26.5 SD = 2.4 Mo = -; n = 4 CV _w = 9.0%	11–16 mean = 13.7 SD = 1.2 Mo = 14; n = 14 CV _w = 8.8%	9.1%	25.3%
RT (s)	0.001–0.003 mean = 0.002 SD = 0.001; n = 214 CV _w = 24.2%	0.001–0.002 mean = 0.002 SD = 0.000; n = 170 CV _w = 12.7%	0.001–0.002 mean = 0.002 SD = 0.000; n = 106 CV _w = 14.7%	0.001–0.004 mean = 0.002 SD = 0.000; n = 192 CV _w = 13.1%	16.2%	20.0%
RTR (%)	18.8–51.6 mean = 31.9 SD = 5.6; n = 214 CV _w = 17.7%	14.8–50.0 mean = 24.7 SD = 4.1; n = 170 CV _w = 16.6%	16.3–50.0 mean = 25.5 SD = 4.0; n = 106 CV _w = 15.8%	14.2–44.2 mean = 24.0 SD = 3.9; n = 192 CV _w = 16.1%	16.5%	21.2%
DF (Hz) (= FF)	3281.2–3750.0 mean = 3579.5 SD = 137.9 Mo = 3656.2; n = 12 CV _w = 3.9%	3375.0–3468.8 mean = 3457.1 SD = 33.2 Mo = 3468.8; n = 8 CV _w = 1.0%	3468.8 min = max = mean = Mo SD = 0.0 n = 4 CV _w = 0.0%	3281.2–3375.0 mean = 3301.3 SD = 39.9 Mo = 3281.2; n = 14 CV _w = 1.2%	1.5%	4.1%
FPDF (Hz)	3187.5–3562.5 mean = 3414.1 SD = 123.0 Mo = 3562.5; n = 12 CV _w = 3.6%	3375.0–3468.8 mean = 3386.7 SD = 33.2 Mo = 3375.0; n = 8 CV _w = 1.0%	3093.8–3187.5 mean = 3140.7 SD = 54.1 Mo = 3187.5; n = 4 CV _w = 1.7%	3000.0–3375.0 mean = 3227.7 SD = 102.1 Mo = 3281.2; n = 14 CV _w = 3.2%	2.4%	4.2%
CPDF (Hz)	3562.5–3750.0 mean = 3703.1 SD = 63.2 Mo = 3750.0; n = 12 CV _w = 1.7%	3468.8 min = max = mean = Mo SD = 0.0 n = 8 CV _w = 0.0%	3468.8 min = max = mean = Mo SD = 0.0 n = 4 CV _w = 0.0%	3187.5–3375.0 mean = 3293.7 SD = 60.00 Mo = 3281.2; n = 14 CV _w = 1.8%	0.9%	5.1%
LPDF (Hz)	2625.0–3656.2 mean = 3507.8 SD = 283.9 Mo = 3562.5; n = 12 CV _w = 8.1%	3281.2–3468.8 mean = 3386.7 SD = 60.1 Mo = 3375.0; n = 8 CV _w = 1.8%	2906.2–3468.8 mean = 3210.9 SD = 234.4 Mo = -; n = 4 CV _w = 7.3%	3281.2–3468.8 mean = 3368.3 SD = 57.8 Mo = 3375.0; n = 14 CV _w = 1.7%	4.7%	5.8%

Discussion

In this study, we describe two calls of *B. latistriata* and the coefficient of variation of its parameters. A fight event between two males is also reported. Although males of the species of the *B. polytaenia* clade frequently present dorsum marks that suggest fighting (C. Luna-Dias pers. comm.), this is the first published record of a fight for the clade. Although both intraspecific (Giasson and Haddad 2006, Toledo et al. 2007, Nali and Prado 2014) and interspecific (Reichert and Gerhardt 2014, Guerra et al. 2018b) fighting events are broadly reported for anurans, the event reported here was short when compared to similar fights in other species (for example, Nali and Prado 2014, Fernandes et al. 2018).

The calls of *B. latistriata* are similar to those of other species of the *B. polytaenia* clade. By presenting a shorter, single-note call with lower pulse period (call A), and a longer call with higher pulse period (call B), it resembles the calls from *B. bandeirantes*, *B. beckeri*, *B. polytaenia*, and *B. aff. beckeri* (Acioli and Toledo 2008, Pinheiro et al. 2012). In these characteristics, it differs from the described calls of *B. botumirim* (call B absent; Caramaschi et al. 2009), *B. cipoensis* (call A composed of 1–3 notes; Batista et al. 2015), *B. goiana* (call A composed of two notes; Menin et al. 2004), *B. jaguariaivensis* (call A composed of 1–4 notes; Guerra et al. 2017), *B. leptolineata* (call B absent; Kwet 2001), *B. phaeopleura* (call A composed of 2–5 notes; Pinheiro et al. 2012), and *B. stenocephala* (call A composed of 2–4 notes; Martins et al. 2016). The call A of *B. latistriata* also differs from some of these species in the pulse number: 11–21 in *B. latistriata* (present study), 3–5 in *B. botumirim* (Caramaschi et al. 2009), 1–5 in *B. jaguariaivensis* (Guerra et al. 2017), and 2–5 in *B. stenocephala* (Martins et al. 2016).

The calls of *B. latistriata* are distinguished from those of the four species whose calls are most similar to it by the lower dominant frequency of both calls A and B: 3375.0–3937.5 and 3281.2–3750.0 Hz, respectively, in *B. latistriata* (present study); 5340.2–5857.0 and 5340.2–5512.5 Hz, respectively, in *B. bandeirantes* (Pinheiro et al. 2012); 3938.0–5063.0 and 3938.0–4875.0 Hz, respectively, in *B. beckeri* (Martins et al. 2016); 6890.0–7320.0 and 6460.0–7320.0 Hz, respectively, in *B. aff. beckeri* (Acioli and Toledo 2008). These species are also morphologically similar (Caramaschi and Cruz 2013), and hence, the use of calls for identifying species and clarifying phylogenetic relationships must be encouraged. Additionally, of all species of the *B. polytaenia* clade, frequency modulation was previously reported only for *B. goiana* (Menin et al. 2004). However, in this species the frequency ascends from the first to the last pulses, while in *B. latistriata* the frequency ascends from the first to the central pulses, thereafter decreasing to the last pulses.

Values of CV are linked to issues like recognition (at the species, population, and individual levels) and female preferences, and comparing those coefficients can be taxonomically informative (Köhler et al. 2017). Gerhardt (1991) stated that static CV_w parameters may be linked with recognition. All spectral parameters, as well as call B duration, were static on a within-individual basis. Furthermore, CV_b values much greater than CV_w indicates parameters useful for individual recognition (Gambale et al. 2014, Forti et al. 2014, Köhler et al. 2017). Call B duration had CV_b 5.6 times greater than CV_w and is the parameter that best fits this purpose. The finding that spectral parameters were static and temporal parameters were dynamic is congruent

with Köhler et al. (2017) and with the results reported by Guerra et al. (2017) for *B. jaguariaivensis*. However, call B duration being static at the within-individual level is a novelty for the *B. polytaenia* clade. As social context can influence call duration in some anurans (Gambale and Bastos 2014, Gambale et al. 2014), social interaction experiments involving *B. latistriata* will be useful to understand this variation.

Despite the similarities between the calls of the species of the *B. polytaenia* clade, different terminology used in call descriptions can lead to difficulties in comparing those descriptions if the terminology is not well explained. For example, structures of the call B herein defined as “pulses” were called as “notes” by Acioli and Toledo (2008) and by Pinheiro et al. (2012). In these cases, the terminology is well stated, and can be comprehended without ambiguity. However, the lack of clear definitions for the calls of *B. cipoensis* and *B. leptolineata* (Kwet 2001; Batista et al. 2015) resulted in few possible comparisons. The redescription of the calls of these species, as well as the description of the call of *B. buriti*, will make possible a full comparison of the calls in the *B. polytaenia* clade, serving as a powerful tool for the taxonomy of this clade.

Acknowledgments

We thank Dr. Vinicius Guerra and an anonymous reviewer for their suggestions on the manuscript and Dr. Fabio Hepp for his help with the call structure and sidebands. This study was partially funded by grants 312606/2017-6 and 311156/2013-4 provided by Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq). CLD received a fellowship from Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES/CNPq). ZUF RJ amphibian collection was partially funded by Fundação de Amparo à Pesquisa do Estado do Rio de Janeiro (FAPERJ), grant E-26/111.638/2012.

References

- Acioli ECS, Toledo LF (2008) Amphibia, Anura, Hylidae, *Hypsiboas beckeri*: filling gap and description of its advertisement call. Check List 4: 182–184. <https://doi.org/10.15560/4.2.182>
- Batista VG, Gambale PG, Lourenço-De-Moraes R, Campos RM, Bastos RP (2015) Vocalizations of two species of the *Hypsiboas pulchellus* group (Anura: Hylidae) with comments on this species group. North-Western Journal of Zoology 11: 253–261. http://biozoojournals.ro/nwjz/content/v11n2/nwjz_151503_Batista.pdf
- Bioacoustics Research Program (2014) Raven Pro: Interactive Sound Analysis Software. Version 1.5. <http://www.birds.cornell.edu/raven>
- Caramaschi U, Cruz CAG (2004) Duas novas espécies de *Hyla* do grupo de *H. polytaenia* Cope, 1870 do sudeste do Brasil (Amphibia, Anura, Hylidae). Arquivos do Museu Nacional 62: 247–254.
- Caramaschi U, Cruz CAG, Nascimento LB (2009) A New Species of *Hypsiboas* of the *H. polytaenius* Clade from Southeastern Brazil (Anura: Hylidae). South American Journal of Herpetology 4: 210–216. <https://doi.org/10.2994/057.005.0301>

- Caramaschi U, Cruz CAG (2013) A new species of the *Hypsiboas polytaenius* clade from south-eastern Brazil (Anura: Hylidae). South American Journal of Herpetology 8: 121–126. <https://doi.org/10.2994/SAJH-D-13-00009.1>
- Carvalho TR, Giaretta AA (2013) Bioacoustics reveals two new syntopic species of *Adenomera* Steindachner (Anura: Leptodactylidae: Leptodactylinae) in the Cerrado of central Brazil. Zootaxa 3731: 533–551. <https://doi.org/10.11646/zootaxa.3731.4.6>
- Duellman WE, Trueb L (1994) Biology of Amphibians. The Johns Hopkins University Press, Baltimore, 670 pp.
- Fernandes IYP, Cândido-Junior J, Kaefer IL (2018) Combat behaviour in the glassfrog *Vitreorana uranoscopa* (Müller, 1924). Herpetology Notes 11: 671–673. <https://biotaxa.org/hn/article/view/39009/34648>
- Forti LR, Costa WP, Martins LB, Nunes-de-Almeida CHL, Toledo LF (2014) Advertisement call and genetic structure conservatism: good news for an endangered Neotropical frog. PeerJ 4: e2014. <https://doi.org/10.7717/peerj.2014>
- Forti LR, da Silva TRA, Toledo LF (2017) The acoustic repertoire of the Atlantic Forest Rocket Frog and its consequences for taxonomy and conservation (Allobates, Aromobatidae). ZooKeys 692: 141–153. <https://doi.org/10.3897/zookeys.692.12187>
- Gambale PG, Bastos RP (2014) Vocal repertoire and bioacoustic analyses in *Physalaemus cuvieri* (Anura, Leptodactylidae) from southern Brazil. The Herpetological Journal 24: 31–40.
- Gambale PG, Signorelli L, Bastos RP (2014) Individual variation in the advertisement calls of a Neotropical treefrog (*Scinax constrictus*). Amphibia-Reptilia 3: 271–281. <https://doi.org/10.1163/15685381-00002949>
- Gerhardt HC (1991) Female mate choice in treefrogs: static and dynamic acoustic criteria. Animal Behaviour 42: 615–635. [https://doi.org/10.1016/S0003-3472\(05\)80245-3](https://doi.org/10.1016/S0003-3472(05)80245-3)
- Giasson LOM, Haddad CFB (2006) Social interactions in *Hypsiboas albomarginatus* (Anura: Hylidae) and the significance of acoustic and visual signals. Journal of Herpetology 40: 171–180. <https://doi.org/10.1670/205-05A.1>
- Goicoechea N, De La Riva I, Padial JM (2010) Recovering phylogenetic signal from frog mating calls. Zoologica Scripta 39: 141–154. <https://doi.org/10.1111/j.1463-6409.2009.00413.x>
- Guerra V, Lingnau R, Bastos RP (2017) Vocalizations and Bioacoustic Analysis of *Boana jaguariavensis* (Caramaschi, Cruz, and Segalla, 2010) (Anura: Hylidae). South American Journal of Herpetology 12: 34–41. <https://doi.org/10.2994/SAJH-D-16-00018.1>
- Guerra V, Llusia D, Gambale PG, Morais AR, Márquez R, Bastos RP (2018a) The advertisement calls of Brazilian anurans: Historical review, current knowledge and future directions. PloS ONE 13: e0191691. <https://doi.org/10.1371/journal.pone.0191691>
- Guerra V, Costa NQ, Bastos RP (2018b) Interspecific behaviour between two species of *Boana* (Anura: Hylidae) from central Brazil. Herpetology Notes 11: 489–493. <https://biotaxa.org/hn/article/view/34737/33663>
- Hepp FSFS, Carvalho-e-Silva SP (2011) Description and comparison of advertisement calls of *Euparkerella brasiliensis* (Parker, 1926) and *E. cochranae* Izecksohn, 1988 (Amphibia: Anura: Strabomantidae). Herpetology Notes 4: 45–51.
- Köhler J, Jansen M, Rodríguez A, Kok PJR, Toledo LF, Emmrich M, Glaw F, Haddad CFB, Rödel M, Vences M (2017) The use of bioacoustics in anuran taxonomy: theory, terminol-

- ogy, methods and recommendations for best practice. *Zootaxa* 4251: 1–124. <https://doi.org/10.11646/zootaxa.4251.1.1>
- Kwet A (2001) Frösche im brasilianischen Araukarienwald. Anurengemeinschaft des Araukarienwaldes von Rio Grande do Sul: Diversität, Reproduktion und Ressourcenaufteilung. Natur und Tier-Verlag, Münster, 192 pp.
- Laiolo P (2010) The emerging significance of bioacoustics in animal species conservation. *Biological Conservation* 143: 1635–1645. <https://doi.org/10.1016/j.biocon.2010.03.025>
- Martins LB, Giaretta AA, Carvalho TR, Miwa RY (2016) Vocalizations of *Hypsiboas beckeri* and *H. stenocephalus* (Anura: Hylidae), two species of the *H. polytaenius* group from southeastern Brazil. *Phyllomedusa: Journal of Herpetology* 15: 51–64. <https://doi.org/10.11606/issn.2316-9079.v15i1p51-64>
- Menin M, Silva RA, Giaretta AA (2004) Reproductive biology of *Hyla goiana* (Anura, Hylidae). *Iheringia, Série Zoologia* 94: 49–52. <https://doi.org/10.1590/S0073-47212004000100008>
- Nali RC, Prado CP (2014) The fight call of *Bokermannohyla ibitiguara* (Anura: Hylidae): first record for the genus. *Salamandra* 50: 181–184. <http://www.salamandra-journal.com/index.php/home/contents/2014-vol-50/374-nali-r-c-c-p-a-prado>
- Orrico VGD, Mongin MM, Carvalho-e-Silva AMPT (2007) The tadpole of *Hypsiboas latistriatus* (Caramashi & Cruz, 2004) a species of the *Hypsiboas polytaenius* (Cope, 1870) clade (Amphibia, Anura, Hylidae). *Zootaxa* 1531: 25–37.
- Pinheiro PDP, Pezzuti TL, Garcia PCA (2012) The tadpole and vocalizations of *Hypsiboas polytaenius* (Cope, 1870) (Anura, Hylidae, Hylineae). *South American Journal of Herpetology* 7: 123–133. <https://doi.org/10.2994/057.007.0202>
- Reichert MS, Gerhardt HC (2014) Behavioral strategies and signaling in interspecific aggressive interactions in gray tree frogs. *Behavioral Ecology* 25: 520–530. <https://doi.org/10.1093/beheco/aru016>
- Rivadeneira CD, Venegas PJ, Ron SR (2018) Species limits within the widespread Amazonian treefrog *Dendropsophus parviceps* with descriptions of two new species (Anura, Hylidae). *ZooKeys* 726: 25–77. <https://doi.org/10.3897/zookeys.726.13864>
- Robillard T, Höbel G, Gerhardt CH (2006) Evolution of advertisement signals in North American hylid frogs: vocalizations as end-products of calling behavior. *Cladistics* 22: 533–545. <https://doi.org/10.1111/j.1096-0031.2006.00118.x>
- Stuart S (2018) *Hypsiboas latistriatus*. The IUCN Red List of Threatened Species: e.T61778A12540948. www.iucnredlist.org [accessed on 05 October 2018]
- Toledo LF, Araújo OG, Guimarães LD, Lingnau R, Haddad CF (2007) Visual and acoustic signaling in three species of Brazilian nocturnal tree frogs (Anura, Hylidae). *Phyllomedusa* 6: 61–68. <https://doi.org/10.11606/issn.2316-9079.v6i1p61-68>
- Toledo LF, Haddad CFB (2009) Defensive vocalizations of Neotropical anurans. *South American Journal of Herpetology* 4: 25–42. <https://doi.org/10.2994/057.004.0104>
- Toledo LF, Martins IA, Bruschi DP, Passos MA, Alexandre C, Haddad CFB (2014) The anuran calling repertoire in the light of social context. *Acta Ethologica* 18: 87–99. <https://doi.org/10.1007/s10211-014-0194-4>
- Wells KD (2007) *The Ecology and Behavior of Amphibians*. University of Chicago Press, Chicago, 1400 pp. <https://doi.org/10.7208/chicago/9780226893334.001.0001>