

# First description of the male of *Barylestis saaristoi* Jäger, 2008 (Araneae, Sparassidae) from China

Yang Zhong<sup>1,2</sup>, Ye-Jie Lin<sup>3</sup>, Jie Liu<sup>2</sup>

**1** School of Nuclear Technology and Chemistry & Biology, Hubei University of Science and Technology, Xiangning 437100, Hubei, China **2** The State Key Laboratory of Biocatalysis and Enzyme Engineering of China, Centre for Behavioural Ecology and Evolution, College of Life Sciences, Hubei University, Wubao 430062, Hubei, China **3** Institute of Zoology, Chinese Academy of Sciences, Beijing 100101, China

Corresponding author: Jie Liu ([sparassidae@aliyun.com](mailto:sparassidae@aliyun.com))

---

Academic editor: G. Blagoev | Received 20 December 2018 | Accepted 1 March 2019 | Published 20 March 2019

<http://zoobank.org/5A46A734-C743-4AA2-9182-A32C28C532F3>

---

**Citation:** Zhong Y, Lin Y-J, Liu J (2019) First description of the male of *Barylestis saaristoi* Jäger, 2008 (Araneae, Sparassidae) from China. ZooKeys 832: 153–160. <https://doi.org/10.3897/zookeys.832.32569>

---

## Abstract

The male of *Barylestis saaristoi* Jäger, 2008 is described for the first time from Menglun Town, Yunnan Province, China. This is the first record of this genus from China. An illustrated morphological description of this species is given.

## Keywords

biodiversity, Yunnan, huntsman spiders, taxonomy

## Introduction

The genus *Barylestis* was established by Simon (1910). Currently, of all 11 known *Barylestis* species, 10 from tropical Africa (Cameroon, Congo, Central Africa, Equatorial Guinea, Gabon, Nigeria, Rwanda, Sudan, Uganda, West Africa) and one from South-east Asia (Thailand, Myanmar) have been recorded (Jäger 2002; World Spider Catalog 2018). This genus had long been recognized as a purely African genus by Jäger (2002) until *Barylestis saaristoi* Jäger, 2008 was first described based on female specimens from Thailand and Myanmar. Jäger and his colleagues tried to collect the male of *B. saaristoi* specifically because of its potential contribution to zoogeographic and phylogenetic rela-

tionships of this genus, but the search was not successful (Jäger 2008). Recently, the authors examined specimens collected from Yunnan Province and found three females and one male belonging to this species. This is also the first record of this genus from China.

## Materials and methods

Specimens were examined and measured with a Leica M205C stereomicroscope. Epigynes were examined and illustrated after dissection from the spider bodies. All photos were taken with a Leica DFC450 digital camera attached to a Leica M205C stereomicroscope, with 10–20 photographs taken in different focal planes and combined using image stacking software (Leica LAS). Photographic images were edited using Adobe Photoshop. Left palps are depicted. Most hairs and macrosetae are omitted in the palp drawings. All specimens examined in this study are deposited in the College of Life Sciences, Hubei University.

Leg measurements are shown as: total length (femur, patella, tibia, metatarsus, tarsus). Number of spines is listed for each segment in the following order: prolateral, dorsal, retrolateral, ventral (in femora and patellae ventral spines are absent and the fourth digit is omitted in the spination formula). Abbreviations follow Zhong et al. (2017, 2018):

<b>ALE</b>	anterior lateral eyes;	<b>Pa</b>	patella;
<b>AME</b>	anterior median eyes;	<b>PL</b>	prosoma length;
<b>AW</b>	anterior width of prosoma;	<b>PLE</b>	posterior lateral eyes;
<b>CH</b>	clypeus height;	<b>PME</b>	posterior median eyes;
<b>FE</b>	femur;	<b>Pp</b>	palp;
<b>Mt</b>	metatarsus;	<b>PW</b>	prosoma width;
<b>OL</b>	opisthosoma length;	<b>Ta</b>	tarsus;
<b>OW</b>	opisthosoma width;	<b>Ti</b>	tibia I, II, III, IV—legs I to IV.

Abbreviations for the collection depositories:

<b>HBU</b>	Hubei University, Wuhan, China;
<b>SMF</b>	Research Institute Senckenberg, Frankfurt, Germany

## Taxonomy

**Family Sparassidae Bertkau, 1872**

**Genus *Barylestis* Simon, 1910**

***Barylestis saaristoi* Jäger, 2008**


Figures 1–26

*Barylestis saaristoi* Jäger, 2008: 106, figs 1–14 (holotype female from Mae Hong Son Province, Thailand, deposited in SMF 58342).

**Material examined.** 1 male and 1 female (HBU), Mengxin Farm [21.89°N, 101.36°E, 736m], Dai Autonomous Prefecture of Xishuangbanna, China, 4 May 2018, Yiwu Zhu leg.; 2 females (HBU), Xishuangbanna Tropical Botanical Garden [21.96°N, 101.22°E, 757m] Dai Autonomous Prefecture of Xishuangbanna, China, 30 May 2015, Wancheng Li leg.


**Diagnosis.** Male of *B. saaristoi* can be separated from *B. montandoni* (Lessert, 1929) and *B. occidentalis* (Simon, 1887) by embolus arising from tegulum in an 11-o'clock-position (3-o'clock-position in *B. montandoni* and *B. occidentalis*, Figs 2, 5), separated from *B. fagei* (Lessert, 1929) and *B. variatus* (Pocock, 1900) by the long and slender dRNA (short and wide in *B. fagei* and *B. variatus*, Figs 3, 6), separated from *B. scutatus* (Pocock, 1903) by tegulum partly covered embolic base (wholly covered in *B. scutatus*). Females of this species can be recognised as this particular species by the following combination of characters: 1. Epigyne with V-shaped pit; 2. Vulva with tips of lateral coils pointing mediad and first part of copulatory ducts slender, running parallel (Jäger 2008).

**Description. Male.** PL 6.4, PW 7.5, AW 4.0, OL 5.4, OW 3.9. Eyes: AME 0.34, ALE 0.43, PME 0.26, PLE 0.44, AME–AME 0.26, AME–ALE 0.15, PME–PME 0.59, PME–PLE 1.03, AME–PME 0.55, ALE–PLE 0.97, CH AME 0.47, CH ALE




**Figures 1–3.** *Barylestis saaristoi* Jäger, 2008. 1–3 Left male palp (1 prolateral 2 ventral 3 retrolateral). Abbreviations: C—conductor; dRTA—dorsal retrolateral tibial apophysis; E—embolus; ST—subtegulum; T—tegulum, vRTA—ventral retrolateral tibial apophysis. Scale bar: 0.5 mm.





**Figures 4–8.** *Barylestis saaristoi* Jäger, 2008. **4–6** Left male palp (**4** prolateral **5** ventral **6** retrolateral) **7** epigyne, ventral **8** vulva, dorsal. Abbreviations: AB—anterior bands, CD—copulatory ducts, CO—copulatory opening, GS—glandular structures of internal duct system, LW—lateral winding of internal duct system, PP—posterior pits of lateral lobes. Scale bar: 0.5 mm.



**Figures 9–21.** *Barylestis saaristoi* Jäger, 2008. **9, 10** Cheliceral dentition, ventral view (**9** male **10** female) **11** palpal claw of female, retrolateral view **12–15** leg I prolateral and retrolateral view of two claws (**12, 13** male **14, 15** female) **16, 17** eye arrangement, dorsal view (**16** male **17** female) **18–21** Leg I–IV (male, right). Scale bars: 1 mm (**9, 10, 16, 17**); 0.2 mm (**11–15**); 2 mm (**18–21**).


0.62. Spination: Palp: 131, 101, 2021; Fe: I 333, II 000, III 333, IV 331; Pa: I 101, II 000, III 101, IV 000; Ti: I 2226, II 0004, III 2116, IV 2014; Mt: I 1014, II 0004, III 1014, IV 1016. Measurements of palp and legs: Palp 9.5 (2.9, 1.2, 1.6, –, 3.8), I 32.5 (8.6, 3.2, 8.7, 9.4, 2.6), II 27.1 (7.5, 2.3, 7.5, 7.1, 2.7), III 29.0 (9.1, 2.4, 8.4,





**Figures 22–25.** *Barylestis saaristoi* Jäger, 2008. **22, 23** Male (**22** dorsal **23** ventral) **24, 25** female (**24** dorsal **25** ventral). Scale bar: 2 mm.

7.1, 2.0), IV 29.1 (9.3, 2.5, 8.1, 6.8, 2.4). Leg formula: I-IV-III-II (second leg may have fractured before collection, as it is very tiny; Figs 18–21). Cheliceral furrow with 3 anterior and 4 posterior teeth, without denticles. Claws of leg I with long and slightly curved teeth in both male and female. Female palpal claws with seven long teeth, almost same size as those of leg I (Figs 9–15). Dorsal carapace reddish-brown, posterior margins dark. Chelicerae, sternum, gnathocoxae and labium deep reddish-brown to black. Legs reddish-brown without spots and patches. Dorsal opisthosoma covered by long and dense hairs. Ventral opisthosoma uniformly yellowish-brown (Figs 22, 23). Cymbium significantly longer than tibia. Conductor membranous, arising from direction of tegulum 11:30. Embolus running 1.25 coils around tegulum, with tip situated



**Figure 26.** Collection localities of *Barylestis saaristoi* in Yunnan Province, China.

near conductor. RTA arising medially from tibia, vRTA developed, almost rectangle-shaped and dRTA finger-shaped in ventral view (Figs 1–6).

**Female.** For details see Jäger (2008).

**Distribution.** China (Yunnan Province, new record) (Fig. 26), Thailand (Mae Hong Son Province), Myanmar (Karen, Kayin State).

## Acknowledgements

We thank Mr Wancheng Li and Yiwu Zhu for providing the Sparassidae specimens. This study was financially supported by the National Natural Sciences Foundation of China (NSFC- 31572236/31272268/31772420) and a PhD grant from Hubei University Science and Technology (BK201811).

## References

- Jäger P (2002) Heteropodinae: transfers and synonymies (Arachnida: Araneae: Sparassidae). *Acta Arachnologica* 51: 33–61. <https://doi.org/10.2476/asjaa.51.33>
- Jäger P (2008) *Barylestis* Simon, 1910 – first record in Asia with comments on its zoogeography (Araneae: Sparassidae: Heteropodinae). *Arthropoda Selecta* 17: 105–110.

- Simon E (1910) Arachnides recueillis par L. Fea sur la côte occidentale d'Afrique. 2<sup>e</sup> partie. Annali del Museo Civico di Storia Naturale di Genova 44: 335–449.
- World Spider Catalog (2018) World Spider Catalog. Version 19.5. Natural History Museum Bern. <http://wsc.nmbe.ch> [Accessed on 17 September 2018]
- Zhong Y, Cao XW, Liu J (2017) Six *Sinopoda* species (Araneae: Sparassidae) from Fujian and Yunnan Provinces in China. Zootaxa 4227: 151–172. <https://doi.org/10.11646/zootaxa.4227.2.1>
- Zhong Y, Jäger P, Chen J, Liu J (2018) Taxonomic review of the *Sinopoda okinawana*-group (Araneae: Sparassidae) in China. Zootaxa 4388(3): 328–346. <https://doi.org/10.11646/zootaxa.4388.3.2>