

ZooKeys

Table S1: Average individual snout-vent length (SVL) in *Pelobates* populations (in mm).

Taxon	Locality	Female			Male			Reference
		n	mean ± SE	Range	n	mean ± SE	Range	
<i>b. balcanicus</i>	Utrine, Serbia	30	60.8 ± 1.7	47.8–79.6	21	61.1 ± 2.9	47.6–75.6	Rot-Nikcevic et al. 2001
<i>b. balcanicus</i>	Deliblato sand, Serbia	32	60.6 ± 1.5	47.8–79.6	29	60.4 ± 1.4	47.6–75.6	Üğurtas et al. 2002
<i>b. balcanicus</i>	Ivanovo, Serbia	9	57.1 ± 1.8	48.1–66.2	4	53.9 ± 4.4	46.1–62.1	Üğurtas et al. 2002
<i>b. balcanicus</i>	Staro Laniste, Serbia	2	71.0 ± 0.9	70.2–71.9				Üğurtas et al. 2002
<i>b. balcanicus</i>	Macedonia + Bulgaria	11	70.5 ± 1.2	66.1–79.3	19	75.8 ± 0.7	68.8–79.7	Rot-Nikcevic et al. 2001
<i>b. balcanicus</i>	Ovce pole, Macedonia	3	72.3 ± 2.0	68.4–75.2	4	71.5 ± 1.9	66.6–75.5	Üğurtas et al. 2002
<i>b. balcanicus</i>	Monospitovo, Macedonia	24	63.4 ± 1.6	54.2–83.8	15	60.5 ± 1.4	53.7–76.1	Üğurtas et al. 2002
<i>b. balcanicus</i>	Predejci, Macedonia	6	70.2 ± 1.0	67.7–73.2	7	71.6 ± 3.4	59.1–85.4	Üğurtas et al. 2002
<i>b. balcanicus</i>	Djavato, Macedonia	13	71.5 ± 2.1	59.8–84.7	10	76.5 ± 1.6	70.0–88.2	Üğurtas et al. 2002
<i>b. balcanicus</i>	Ezerani, Macedonia	13	59.9 ± 1.0	52.1–68.3	11	60.6 ± 1.2	53.3–65.9	Üğurtas et al. 2002
<i>b. balcanicus</i>	Romania	5	73.0	67.0–79.0				Bacescu 1954
<i>b. balcanicus</i>	Kresna, Bulgaria	4	72.9 ± 3.0	66.1–79.3	14	76.9 ± 0.6	72.4–79.7	Üğurtas et al. 2002
<i>b. balcanicus</i>	Grindul Lupilor, Romania	39	74.4 ± 1.9	55.4–99.5	37	76.5 ± 1.8	59.3–93.7	Cogalniceanu et al. 2014
<i>b. balcanicus</i>	Durunkulak, Bulgaria	13	69.6 ± 1.5	62.8–79.4	17	67.8 ± 1.4	57.8–79.5	Üğurtas et al. 2002
<i>b. balcanicus</i>	Plovdiv, Bulgaria				1	90.0		Müller 1932
<i>b. balcanicus</i>	Greece	44	73.2	66.0–84.0	38	72.4	65.0–85.0	Sofianidou 2012
<i>b. balcanicus</i>	Greece	35	53.9	50.0–60.0	29	50.8	49.0–55.0	Sofianidou 2012
<i>cultripes</i>	Spain	?	78.0		?	70.1		Salvador et al. 1986
<i>cultripes</i>	Arrifa, Portugal	40	54.2	?-72	20	52.8	?-62	Leclair et al. 2005
<i>cultripes</i>	Porto Covo, Portugal	4	72.7	?-84	11	71.5	?-80	Talavera 1990
<i>cultripes</i>	Montalvos, Spain	66	74.6	?-86	76	72.0	?-85	Lizana et al. 1994
<i>cultripes</i>	Madrid, Spain	14	79.5	?-101	15	79.8	?-87	Talavera 1990
<i>cultripes</i>	Cadiz, Spain	30	72.1	?-99	30	65.8	?-84	Beukema et al. 2013
<i>cultripes</i>	Pedroso, Spain	18	90.4 ± 2.0		56	83.9 ± 1.1		Marangoni and Tejedo 2008
<i>cultripes</i>	Navas, Spain	5	91.8 ± 3.8		6	83.8 ± 3.5		Marangoni and Tejedo 2008
<i>cultripes</i>	Gerena, Spain				8	93.3 ± 3.0		Marangoni and Tejedo 2008
<i>cultripes</i>	Aznalcóllar, Spain	14	104.7 ± 2.3		31	88.1 ± 1.5		Marangoni and Tejedo 2008
<i>cultripes</i>	Lázaro, Spain	1	68.0		18	58.4 ± 2.0		Marangoni and Tejedo 2008
<i>cultripes</i>	Bodogones, Spain	40	76.2 ± 1.3		43	71.6 ± 1.3		Marangoni and Tejedo 2008
<i>cultripes</i>	Abalarío, Spain	69	71.8 ± 1.0		83	67.7 ± 0.9		Marangoni and Tejedo 2008
<i>cultripes</i>	Doñana, Spain	24	62.9 ± 1.7		47	66.1 ± 1.2		Marangoni and Tejedo 2008
<i>cultripes</i>	Doñana, Spain	8	60.8		8	55.9		Díaz-Paniagua 2005
<i>cultripes</i>	Mas de Melons, Spain	26	76.1		27	71.8		Pascual-Pons et al. 2017
<i>cultripes</i>	Spain?	12	44.9 ± 2.4	32.0–57.1	17	47.7 ± 2.6	33.9–64.3	Busack et al. 1985
<i>fucus</i>	Stagni di Belangero, Italy	2	52.3 ± 0.3		12	45.6 ± 0.7		Mercurio and Li Vigni 2007
<i>fucus</i>	Arsago-Seprio, Italy	26	52.1 ± 1.1	43.0–61.5	17	50.4 ± 1.6	44.5–57.5	Scali and Gentilli 2003
<i>fucus</i>	Ivrea, Italy	73	54.5 ± 3.7		70	45.7 ± 2.6		Andreone et al. 2004
<i>fucus</i>	Ivrea II, Italy	11	58.4 ± 2.6	55.0–63.0	22	50.2 ± 4.2	40.0–58.0	Andreone and Pavignano 1988
<i>fucus</i>	Varese, Italy	?	52.1 ± 5.4		?	50.4 ± 3.4		Andreone et al. 2004
<i>fucus</i>	Wien, Austria	199	45.0 ± 0.4		427	39.6 ± 0.2		Wiener 1997
<i>fucus</i>	Braunschweig, Germany	289	49.7		866	43.3		Tobias 2000
<i>fucus</i>	Mecklenburg, Germany	37	52.5 ± 0.9	40.0–67.0	61	46.0 ± 0.6	37.0–56.0	Szekely and Nemec 2002
<i>fucus</i>	Brandenburg, Germany	25	52.6	48.0–59.0	100	42.2	40.0–48.0	Szekely and Nemec 2002
<i>fucus</i>	Niedersachsen, Germany	223	63.0 ± 0.3	48.0–78.0	545	52.7 ± 0.2	38.0–65.0	Szekely and Nemec 2002
<i>fucus</i>	Sachsen-Anhalt, Germany	21	54.8 ± 1.2	46.0–62.0	29	47.8 ± 0.6	42.0–57.0	Szekely and Nemec 2002
<i>fucus</i>	Unteren Saaletal, Germany	40	56.9 ± 0.8	48.3–69.5	65	47.0 ± 0.8	38.9–55.6	Grosse 2008
<i>fucus</i>	Nordrhein-Westfalen, Germany	?		?-71.5	?		?-62.3	Chmela and Kronshage 2011
<i>fucus</i>	Nordrhein-Westfalen, Germany	2	69.5 ± 1.5	68.0–71.0	7	55.7 ± 1.3	50.0–61.0	Szekely and Nemec, 2002
<i>fucus</i>	Germany + Hungary	23	44.2 ± 4.0		12	42.6 ± 3.8		Andreone et al. 1993
<i>fucus</i>	Aleksandrow Kujawski, Poland	157	50.3 ± 0.3	43.0–57.6	161	44.4 ± 0.2	38.0–50.9	Adrzejewski et al. 1977
<i>fucus</i>	Poland	?	?	50.0–77.0	?	?	45.0–65.0	Juszczuk 1974
<i>fucus</i>	Utrine, Serbia	26	53.7 ± 1.0	46.5–65.2	5	44.8 ± 1.2	41.2–48.1	Rot-Nikcevic et al. 2001
<i>fucus</i>	Cavolj, Serbia	28	54.0 ± 0.8	47.1–62.2	39	46.0 ± 0.5	40.6–56.7	Rot-Nikcevic et al. 2001
<i>fucus</i>	Lesino Kopovo, Serbia	27	45.4 ± 1.8	41.4–57.9	16	43.9 ± 0.8	36.4–49.5	Rot-Nikcevic et al. 2001
<i>fucus</i>	Cluj-Napoca, Romania	15	64.1 ± 1.4	52.0–72.0	80	52.2 ± 0.4	48.0–63.0	Szekely and Nemec 2002
<i>fucus</i>	Cluj-Napoca, Romania	41	62.4 ± 0.8	50.0–73.0	117	54.3 ± 0.3	45.0–63.0	Szekely and Nemec 2002
<i>fucus</i>	Grindul Lupilor, Romania	34	47.7 ± 1.1	36.9–61.5	43	41.7 ± 0.4	37.6–47.3	Cogalnicanu et al. 2014
<i>syriacus boettgeri</i>	Edirne, Turkey	5	69.4 ± 2.0	61.7–73.4	21	68.6 ± 1.1	56.7–77.6	Üğurtas et al. 2002
<i>syriacus boettgeri</i>	Izmir, Turkey	9	62.2 ± 4.3	40.0–79.0	12	70.8 ± 2.0	57.0–79.0	Basoglu and Zaloglu 1964
<i>syriacus boettgeri</i>	Izmir, Turkey	20		64.0–92.0	13		66.0–72.0	Caydam 1974
<i>syriacus boettgeri</i>	Seydisehir, Turkey	9	58.2 ± 1.4	50.8–63.7	13	63.1 ± 1.2	57.2–70.8	Üğurtas et al. 2002
<i>syriacus boettgeri</i>	Tbilisi, Georgia	9	71.5	64.4–86.0	16	71.6	61.5–82.5	Delwig 1928
<i>syriacus boettgeri</i>	Belesuwar, Azerbaijan	1	78.0					Mertens 1923
<i>s. syriacus</i>	Haifa, Israel	1	78.0		1	73.5		Boettger 1889
<i>varaldii</i>	Morocco	9	50.7 ± 2.7	35.6–62.5	20	51.0 ± 1.6	33.2–64.6	Busack et al. 1985
<i>varaldii</i>	Foret Mamora, Morocco	20	52.8 ± 1.3	45.0–64.1	66	51.7 ± 0.4	43.9–60.6	Guarino et al. 2011
<i>varaldii</i>	Foret Mamora, Morocco	30	53.6	?-66	98	51.9	?-61	Beukema et al. 2013
<i>varaldii</i>	Foret Mamora, Morocco				1	49.5		Pasteur and Bons 1959
<i>varaldii</i>	Merja Samora, Morocco	1	55.0					Pasteur and Bons 1959
<i>vespertinus</i>	Karadagh NR, Russia				3	52.9 ± 4.8	44.5–61.3	Kukushkin 2004
<i>vespertinus</i>	Karadagh NR, Russia				5	56.3 ± 2.2	51.3–61.4	Kukushkin 2006
<i>vespertinus</i>	Opul NR, Russia				1	58.2		Kukushkin and Moroz 2007
<i>vespertinus</i>	Mordovia Republic, Russia	7	52.7 ± 1.5	48.1–58.0	12	42.1 ± 1.6	31.0–47.0	Ruchin 2014
<i>vespertinus</i>	Uritskoe, Russia	32	45.7 ± 0.7	37.8–55.1				Yermokhin et al. 2014
<i>vespertinus</i>	Uritskoe, Russia	?	40.0–45.4	29.3–59.2	?	35.7–43.2	29.0–56.0	Yermokhin et al. 2016
<i>vespertinus</i>	Uritskoe, Russia				15	48.0 ± 0.7	43.0–52.3	Yermokhin and Tabachishin 2011
<i>vespertinus</i>	Uritskoe, Russia				49	46.5 ± 0.5	39.7–54.8	Yermokhin and Tabachishin 2011
<i>vespertinus</i>	Uritskoe, Russia				21	48.9 ± 0.6	42.7–54.3	Yermokhin and Tabachishin 2011
<i>vespertinus</i>	Uritskoe, Russia				17	46.3 ± 0.8	40.2–52.5	Yermokhin and Tabachishin 2011
<i>vespertinus</i>	Uritskoe, Russia				18	46.1 ± 0.8	39.8–53.8	Yermokhin and Tabachishin 2011
<i>vespertinus</i>	Uritskoe, Russia				35	47.9 ± 0.5	41.3–53.1	Yermokhin and Tabachishin 2011
<i>vespertinus</i>	Uritskoe, Russia				15	46.4 ± 0.9	42.0–53.6	Yermokhin and Tabachishin 2011

Cited references

- Andreone E, Fortina R, Chiminello A (1993) Natural history, ecology and conservation of the Italian spadefoot toad, *Pelobates fuscus insubricus*. Scientific Reports of the Zoological Society "La Torbiera" 2: 1–96.
- Andreone E, Pavignano I (1988) Observations on the breeding migration of *Pelobates fuscus insubricus* Cornalia 1873 at a ditch in north western Italy (Amphibia, Anura, Pelobatidae). Bollettino della Museo Regionale di Scienze Naturali, Torino 1: 241–250.
- Andreone F, Bergo PA, Bovero S, Gazzaniga E (2004) On the edge of extinction? The spadefoot *Pelobates fuscus insubricus* in the Po Plain, and a glimpse at its conservation biology. Italian Journal of Zoology 71: 61–72.
- Andrzejewski H, Przystalski A, Wilczynska B (1977) The biometric structure of the common spadefoot toad (*Pelobates fuscus* Laur., Salientia, Amphibia) of the environs of Aleksandrow Kujawski Poland. Acta Biologica Cracoviensis Series Zoologia 20: 65–74.
- Bacescu M (1954) *Pelobates syriacus balcanicus* Karaman, a new frog for the fauna of Romania (in Romanian). Comunicări ale Academiei Republicii Populare Române 4: 483–490.
- Basoglu M, Zaloglu S (1964). Morphological and osteological studies in *Pelobates syriacus* from Izmir region, Western Anatolia. Senckenbergiana biologica 45: 233–242.
- Beukema W, De Pous P, Donaire-Barroso D, Bogaerts S, Garcí-Port J, Escoriza D, Arribas OJ, El Mouden EH, Carranza S (2013) Review of the systematics, distribution, biogeography and natural history of Moroccan amphibians. Zootaxa 3661: 1–60.
- Boettger O (1889). Ein neuer Pelobates aus Syrien. Zoologischer Anzeiger 12: 144–147.
- Busack SD, Maxson LR, Wilson MA (1985) *Pelobates varaldii* (Anura: Pelobatidae): A morphologically conservative species. Copeia 1985: 107–112.
- Caydam, O. 1974. Izmir'de bulunan anura türlerinden *Bufo bufo*, *Bufo viridis* (Bufonidae), *Rana ridibunda* (Ranidae), *Pelobates syriacus* (Pelobatidae) ve *Hyla arborea* (Hylidae)'nin üreme biyolojisi üzerinde arastırırmalar. Ege Üniversitesi Fen Fakültesi İlimi Raporlar Serisi 198: 1–22.
- Chmela C, Kronshage A (2011) Knoblauchkröte - *Pelobates fuscus*. In: Hachtel M, Schlüpmann M, Weddeling K, Thiesmeier B, Geiger A, Willigalla C (Eds) Handbuch der Amphibien und Reptilien Nordrhein-Westfalens Band 1. Laurenti Verlag, Bielefeld, 896 pp.
- Cogălniceanu D, Rosioru DM, Székely P, Székely D, Buhaciuc-Ionita E, Stanescu F, Miaud C (2014) Age and body size in populations of two syntopic spadefoot toads (Genus *Pelobates*) at the limit of their ranges. Journal of Herpetology 48: 537–545.
- Delwig W (1928). Eine neue Art der Gattung *Pelobates* Wagl. aus dem zentralen Transkaukasus. Zoologischer Anzeiger 75: 24–31.
- Díaz-Paniagua C, Gómez-Rodríguez C, Portheault A, de Vries W (2005) Los anfibios de Doñana. Naturaleza y parques nacionales. Serie técnica. Organismo Autónomo Parques Nacionales, Madrid, 181 pp.
- Yermokhin MV, Tabachishin VG (2011) Reproductive parameters of females *Pelobates fuscus* (Laurenti, 1768) as functions of size and weight characteristics. Current Studies in Herpetology, Saratov 11: 28–39.
- Grosse W-R. (2008) Dynamik und Struktur einer Population der Knoblauchkröte (*Pelobates fuscus*) im Naturpark Unteres Saaletal bei Zaschwitz. RANA 5: 141–160.

ZooKeys

Guarino FM, de Pous P, Crottini A, Mezzasalma M, Andreone F (2011) Age structure and growth in a population of *Pelobates varaldii* (Anura, Pelobatidae) from northwestern Morocco. *Amphibia-Reptilia* 32: 550–556.

Juszczyk W (1974) Płazy i gady krajowe. Państwowe Wydawn. Naukowe, Warszawa, 721 pp.

Kukushkin OV (2004) Materials to ecology and morphology of little studied amphibian species in the Crimea — the common spadefoot toad, *Pelobates fuscus* (Laurenti, 1768) (Amphibia, Anura, Pelobatidae). In: Letopis' Prirody. Karadagskiy Prirodnyi Zapovednik. T. 20. Simferopol, Sonat, pp. 152–174.

Kukushkin OV (2006) Materials to distribution, ecology and morphology of the common spadefoot toad, *Pelobates fuscus* (Laurenti, 1768), in southeastern coast of the Mountain Crimea. In: Letopis' Prirody. Karadagskiy Prirodnyi Zapovednik. T. 21. Simferopol, Sonat, pp. 179–200.

Kukushkin OV, Moroz PA (2007) Materials to study of herpetofauna of Opuk Nature Reserve and adjacent territories of the Kerch Black Sea coast. In: Letopis' Prirody. Opukskiy Prirodnyi Zapovednik. Simferopol, p. 2.

Leclair MH, Leclair JR, Gallant J (2005) Application of skeletochronology to a population of *Pelobates cultripes* (Anura: Pelobatidae) from Portugal. *Journal of Herpetology* 39: 199–207.

Lizana M, Marquez R, Martin-Sanchez R (1994) Reproductive biology of *Pelobates cultripes* (Anura: Pelobatidae) in Central Spain. *Journal of Herpetology* 28: 19–27.

Marangoni F, Tejedo M (2008) Variation in body size and metamorphic traits of Iberian spadefoot toads over a short geographic distance. *Journal of Zoology* 275: 97–105.

Mercurio V, Li Vigni F (2007) Rediscovery of *Pelobates fuscus insubricus* in the Asti Province, north-western Italy. *Acta Herpetologica* 2: 1–6.

Mertens R (1923) Beiträge zur Kenntnis der Gattung Pelobates Wagler. *Senckenbergiana Biologica* 5: 118–128.

Müller L (1932) Beiträge zur Herpetologie der südosteuropäischen Halbinsel. 1. Herpetologisch Neues aus Bulgarien. *Zoologischer Anzeiger* 100: 299–309.

Pasteur G, Bons J (1959) Les batraciens du Maroc. *Travaux de l'Institut scientifique Chérifien. Série Zoologie* 17, Rabat, 241 pp.

Pascual-Pons M, Oromi N, Pujol-Buxo E, Fibla M, Sanuy D, Montori A (2017) Life history traits of a spadefoot toad (*Pelobates cultripes*) population from a semiarid zone in the north east of the Iberian Peninsula. *Herpetological Journal* 27: 57–61.

Rot-Nikcevic I, Sidorovska V, Dzukic G, Kalezic ML (2001) Sexual size dimorphism and life history traits of two European spadefoot toads (*Pelobates fuscus* and *P. syriacus*) in allopatry and sympatry. *Annales (Koper), Series Historia Naturalis* 11: 107–112.

Ruchin AB (2014) Ecology of amphibians and reptiles of Mordovia. Communication 1. The Pallas's spadefoot toad, *Pelobates vespertinus* (Pallas 1771). *Trudy Mordovskogo Gosudarstvennogo Prirodnogo Zapovednika Imeni P.G. Smidovicha*, 12: 337–349.

Salvador A, Alvarez J, Garcia C (1986) Reproductive biology of a northern population of the western spadefoot *Pelobates cultripes* (Anura, Pelobatidae). In Z. Roček (Ed) *Studies in Herpetology. Societas Europaea Herpetologica*, Prague, pp. 403–408.

ZooKeys

Scali S, Gentilli A (2003) Biology aspects in a population of *Pelobates fuscus insubricus* Cornalia, 1873 (Anura: Pelobatidae). *Herpetozoa* 16: 51–60.

Skékely P, S Nemes (2002) Sex ratio and sexual dimorphism in a population of *Pelobates fuscus* from Transylvania, Romania. *Zeitschrift für Feldherpetologie* 9: 211–216.

Sofianidou T (2012) *Pelobates syriacus* (Boetger, 1899)—Syrische Schaufelkröte. In: Grossenbacher K (Ed) *Handbuch der Reptilien und Amphibien Europas* Bd 51 Froschlurche I. Aula, Wiebelsheim, Germany, pp. 563–620.

Talareva R (1990) Evolucion de Pelobatidos y Peloditidos (Amphibia: Anura): morfologia y desarrollo del sistema esqueletico. Unpublished Ph.D. dissertation, Universidad Complutense de Madrid, Madrid.

Tobias M (2000) Zur Populationsökologie von Knoblauchkröten (*Pelobates fuscus*) aus unterschiedlichen Agrarökosystemen. Dissertation. Technische Universität Braunschweig, 149 pp.

Uğurtas IH, Ljubusavljevic K, Sidorovska V, Kalezic ML, Džukić G. (2002). Morphological differentiation of eastern spadefoot toad (*Pelobates syriacus*) populations. *Israel Journal of Zoology* 48: 13–32.

Wiener K (1997) Struktur und Dynamik einer Knoblauchkrötenpopulation (*Pelobates fuscus fuscus*, Laurenti 1768) nördlich von Wien – ein Vergleich der Untersuchungsjahre 1986, 1987 und 1989 bis 1995. In: *Populationsbiologie von Amphibien: Eine Langzeitstudie auf der Wiener Donauinsel*, Stapfia, pp. 165–181.

Yermokhin MV, Tabachishin VG (2011) Reproductive parameters of females *Pelobates fuscus* (Laurenti, 1768) as functions of size and weight characteristics. *Current Studies in Herpetology*, Saratov, 11: 28–39.

Yermokhin MV, Tabachishin VG, Ivanov GA (2014) Results convergence of fecundity determination of *Pelobates fuscus* (Laurenti, 1768) by full and partial eggs counting methods. *Current Studies in Herpetology*, Saratov, 14: 14–18.

Yermokhin MV, Tabachishin VG, Ivanov GA (2016) Long-term dynamics of the size-weight and sexual structure in populations of *Pelobates fuscus* (Anura, Pelobatidae) in the Medveditsa river valley (Saratov region). *Current Studies in Herpetology*, Saratov, 16: 113–122.