

# Two new species of the stonefly genus *Amphinemura* (Insecta, Plecoptera, Nemouridae) from China

Xiao-Yu Ji<sup>1,†</sup>, Yu-Zhou Du<sup>1,‡</sup>, Zhi-Jie Wang<sup>1,2,§</sup>

**1** School of Horticulture and Plant Protection & Institute of Applied Entomology, Yangzhou University, Yangzhou, 225009, China **2** Suqian College, Suqian, Jiangsu, 223800, China

† <http://zoobank.org/8D9CA829-4EE5-421D-B282-1F0A8CB3AAFA>

‡ <http://zoobank.org/1FDB5781-C8A0-4088-8D18-DCD5BB01C548>

§ <http://zoobank.org/DEEC63FC-CCE1-4485-BC5B-2D7F15845328>

Corresponding author: Yu-Zhou Du (yzdu@yzu.edu.cn)

---

Academic editor: M. Gottardo | Received 18 January 2014 | Accepted 31 March 2014 | Published 22 April 2014

---

<http://zoobank.org/F1856766-0CEE-4F38-A285-FEA7E0BF4F20>

---

**Citation:** Ji X-Y, Du Y-Z, Wang Z-J (2014) Two new species of the stonefly genus *Amphinemura* (Insecta, Plecoptera, Nemouridae) from China. ZooKeys 404: 23–30. doi: 10.3897/zookeys.404.7067

---

## Abstract

Two new species of the genus *Amphinemura* Ris from China are described and illustrated, i.e. *A. annulata* Du & Ji, **sp. n.** from Zhejiang, Shanxi, Shaanxi and Guizhou Province, and *A. lingulata* Du & Wang, **sp. n.** from Shaanxi and Sichuan Province. *A. annulata* is similar to *A. tricintusidens* Wang & Zhu in having an apical cavity of the epiproct, but the epiproct ventral sclerite and the median paraproct lobe of the two species are different. *A. lingulata* is related to *A. didyma* Zhu & Yang in having the similar epiproct, but they differ mostly in paraproct median and outer lobes.

## Keywords

*Amphinemura*, Nemouridae, Plecoptera, new species, China

## Introduction

The subfamily Amphinemurinae includes seven genera, i.e. *Amphinemura* Ris, *Indonemoura* Baumann, *Mesonemoura* Baumann, *Protonemura* Kempny, *Sphaeronemoura* Shimizu & Sivec, *Malenka* Ricker and *Tominemoura* Sivec & Stark, and the first five genera were

found in China. The genus *Amphinemura* is the largest genus of Amphinemurinae with more than 170 species from the Oriental and Holarctic Regions (Baumann 1975; DeWalt et al. 2013). The *Amphinemura* in China is represented by at least 70 species (Du et al. 2007; Du and Wang 2007; Li et al. 2005; Li and Yang 2005, 2006, 2007, 2008a, b, c, d, e, 2011, 2013; Wang et al. 2006; Wang et al. 2007; Wu 1926, 1935, 1938, 1949, 1962, 1973; Yang et al. 2005; Yang et al. 2005 and Zhu and Yang 2002, 2003). Herein we describe two new Chinese species of *Amphinemura* based on male specimens.

## Materials and methods

All type specimens are preserved in 75% or 99% ethanol and are deposited at the School of Horticulture and Plant Protection & Institute of Applied Entomology, Yangzhou University, China. Specimens were examined and illustrated using a Leica stereomicroscope-MZAPO. Abdomens were cut from the bodies, then treated in 5% NaOH, slowly heated to 40–50 °C for 1–3 minutes, and then the specimens were cleared rinsing in clean water. The morphological terminology follows that of Baumann (1975).

## Results

### *Amphinemura annulata* Du & Ji, sp. n.

<http://zoobank.org/F99B5974-7245-4B96-A61C-147FA0A25B4E>


[http://species-id.net/wiki/Amphinemura\\_annulata](http://species-id.net/wiki/Amphinemura_annulata)

Figs 1–6

**Material examined.** Holotype ♂ from China, Zhejiang Province, Mt. Tianmu, 300m, 18–20 Mar. 2006, leg. Wang Zhi-Jie, Sun Yun. Paratypes 30♂♂, the same details as holotype; 1♂, Shanxi Province, Lishan Natural Reserve, 1222m, 24 Aug. 2012, leg. Shi Fu-Ming; 1♂, Shaanxi Province, Houzhenzi, Upriver of Hougou, 26 May 1995, leg. Du Yu-Zhou; 1♂, Shaanxi Province, Foping County, East River, 1240m, 25 Sept. 1996, leg. Xing Lian-Xi; 1♂, Guizhou Province, Maolan City, Dongkuang, 22 Oct. 1996, leg. Li Zi-Zhong.

**Adult habits.** Head and antennae brown, pronotum light brown, subquadrate, angles bluntly rounded, anterior margin wider than posterior margin, with lightly rugosities. Legs brown. Wings hyaline, light brown, veins brown.

**Male.** Forewing length 7.7–8.2 mm, hind wing length 6.6–6.8 mm. Tergum 9 weakly sclerotized, with a small mid-posterior indentation, and bearing a large bundle of tiny spines medially. Tergum 10 weakly sclerotized, with a rounded concavity below epiproct, bearing several spines on lateral margin, and a small triangular projections extruding from the base of the concavity, which is more distinct in lateral view (Figs 1 and 3). Hypoproct narrow basally, extending at midpoint and tapering with a blunt rounded tip, bulging before apex, which is more distinct in lateral view; vesicle slender, four times longer than



**Figures 1–6.** *Amphinemura annulata* male structures. **1** terminalia, dorsal aspect **2** terminalia, ventral aspect **3** terminalia, lateral aspect **4** paraproct (left) **5** epiproct, dorsal aspect **6** epiproct, ventral aspect.

wide (Figs 2 and 3). Paraproct divided into 3 lobes; inner lobe weakly sclerotized, thin and long, about half length of median lobe; median lobe broad basally, mostly membranous, with a long sclerotized band in its outer margin, apically curved inwards and forming an annular projection, bearing many tiny dense spines on the projection; outer lobe sclerotized, short, equal length to inner lobe (Fig. 4). Epiproct elongated, dorsal sclerite sclerotized basally, apical portion weakly sclerotized, taking up half of the dorsal sclerite, forming a small cavity at tip (Fig. 5); in lateral view, slender, well sclerotized pair of large lobes extending to ventral surface from midpoint (Fig. 3); ventral sclerite forming a triangular keel, which is more distinct in lateral view, bearing several black spines ventrally, tip of the ventral sclerite rising from the tip cavity of dorsal sclerite (Fig. 6).

**Female.** Unknown.

**Etymology.** The Latin “*annulata*” refers to median lobe of paraproct forming an annular projection pointing inwards.

**Diagnosis.** This new species is similar to *A. tricintusidens* Wang & Zhu, 2007 (in Wang et al. 2007). Both species having the dorsal sclerite of the epiproct with an apical cavity, but the new species can be separated from the latter by the form of the ventral sclerite of the epiproct and the median lobe of the paraproct. In *A. annulata* sp. n., the ventral sclerite forming a triangular keel, bearing several black spines ventrally, whereas in *A. tricintusidens*, the ventral sclerite of the epiproct forms two heavily sclerotized sclerites which fused at the apex, each bearing two rows of denticles on the surface of the sclerotized sclerites. The median paraproct lobe of *A. annulata* sp. n. is mostly membranous, its outer margin sclerotized to form a long sclerotized band, and forming an annular projection pointing inwards bearing many tiny dense spines on the projection, In *A. tricintusidens*, the median lobe is sclerotized, tapering medially, with a darkly sclerotized rounded tip at the apex, and 3 or 4 dentations surrounding one side of the tip.

***Amphinemura lingulata* Du & Wang, sp. n.**

<http://zoobank.org/DC92AC76-9A64-482E-887A-4C3D2F594D46>


[http://species-id.net/wiki/Amphinemura\\_lingulata](http://species-id.net/wiki/Amphinemura_lingulata)

Figs 7–12

**Material examined.** Holotype ♂ China, Shaanxi Province, Houzhenzi, Upriver of Hougou, 26 May 1995, leg. Du Yu-Zhou. Paratypes 6♂♂, Sichuan Province, Laohegou Natural Reserve, 1700m, 25 May 2012, leg. Ji Xiao-Yu, Tang Xiao-Tian; 2♂♂, Shaanxi Province, Qinling Mountain Range, Railway Station of Qinling, 15 May 1995, leg. Wang Min.

**Adult habitus.** Head brown, antennae light brown, pronotum brown, with light rugosities. Legs brown. Wings subhyaline, light brown, veins brown.

**Male.** Forewing length 6.5–6.8 mm, hind wing 5.4–5.8 mm. Tergum 9 weakly sclerotized, bearing a row of long hairs at distal margin. Tergum 10 weakly sclerotized, with a large flat area below the epiproct, with few spines each side of the epiproct (Fig. 7). Hypoproct broad basally and tapering toward blunt rounded tip, vesicle slightly constricted



**Figures 7–12.** *Amphinemura lingulata* male structures. **7** terminalia, dorsal aspect **8** terminalia, ventral aspect **9** terminalia, lateral aspect **10** paraproct (left) **11** epiproct, dorsal aspect **12** epiproct, ventral aspect.



basally, three times longer than wide (Fig. 8). Paraproct divided into 3 lobes; inner lobe weakly sclerotized, slender, with a short darkly sclerotized line medially; median lobe weakly sclerotized basally, bearing several long strong spines on the large membranous and strongly curved tip; outer lobe darkly sclerotized, slender, with 4 or 5 strong spines at the triangular tip (Fig. 10). Epiproct slender in the dorsal aspect; dorsal sclerite side mostly membranous, with a pair of sclerotized small triangular tongue-shaped projections encasing each side of the bifurcated tip (Fig. 11); lateral arms slender, darkly sclerotized, a pair of sclerotized lateral bands convergent apically (Fig. 9); ventral sclerite forming a median keel-shaped ridge, bearing many black spines ventrally (Fig. 12).

**Female.** Unknown.

**Etymology.** The Latin “*lingulata*” refers to the pair of small triangular tongue-shaped projections encasing each side of the tip of epiproct.

**Diagnosis.** This new species is related to *A. didyma* Zhu & Yang (2002) in having the similar median and outer paraproct lobe. However, their epiprocts are markedly different. *A. lingulata* sp. n. is also similar to *A. zhoui* Li & Yang (2008b), *A. helan-shana* Li, Murányi & Yang (2013) and *A. tibetensis* Zhu & Yang (2003) in the epiproct with a pair of sclerotized small projections encasing each side of the tip, but their paraproct lobes are different obviously.

## Concluding remarks

Shimizu (1996), in a PhD study on East Asian Nemouridae, suggested seven species-groups for East Asian species of *Amphinemura*, i.e. the *flavicollis* group, the *clavigera* group, the *spinigera* group, the *sagittata* group, the *flavostigma* group, the *pentagona* group and the *megaloba* group, and arranged some Chinese species in these species-groups. However, only the *flavostigma* group and the *megaloba* group were subsequently published (Shimizu 1998a, b). A revision of the genus *Amphinemura* will be needed considering the high number of species included.

So far 74 species of *Amphinemura*, including the two new species documented in this paper, were recorded in China. It can be expected that more *Amphinemura* species will be found in the future because China is across the Palaearctic and Oriental Region and features not yet faunistically explored areas with suitable resources for stoneflies. Additionally, studies focused on the biogeography, phylogeny and evolution are also needed to further understand the species diversity of *Amphinemura*.

## Acknowledgements

We heartily thank the editor and reviewers for their critical review of an earlier version of this manuscript. We are very indebted to Qian Yu-Han for assistance with this paper. This research was supported by National Natural Science Foundation of China (No. 31071958).

## References

- Baumann RW (1975) Revision of the stonefly family Nemouridae (Plecoptera): A study of the world fauna at the generic level. *Smithsonian Contributions to Zoology* 211: 1–74. doi: 10.5479/si.00810282.211
- DeWalt RE, Neu-Becker U, Stueber G (2013) Plecoptera Species File Online. Version 5.0/5.0. [3.24.2013] <http://Plecoptera.SpeciesFile.org>
- Du Y-Z, Wang Z-J (2007) New species of the genus *Amphinemura* (Plecoptera: Nemouridae) from Yunnan, China. *Zootaxa* 1554: 57–62.
- Du Y-Z, Wang Z-J, Zhou P (2007) Four new species of the genus *Amphinemura* (Plecoptera: Nemouridae) from China. *Aquatic Insects* 29: 297–305. doi: 10.1080/01650420701552482
- Li W-H, Murányi D, Pan J-J, Yang D (2013) New and little known species of Nemouridae (Plecoptera) from Inner Mongolia of China. *Zootaxa* 3746: 473–480. doi: 10.11646/zootaxa.3746.3.6
- Li W-H, Yang D (2005) Two new species of *Amphinemura* (Plecoptera: Nemouridae) from Sichuan, China. *Zootaxa* 1083: 63–68.
- Li W-H, Yang D (2006) Three new species of *Amphinemura* (Plecoptera: Nemouridae) with a key to the species from Guizhou province, China. *Zootaxa* 1154: 41–48.
- Li W-H, Yang D (2007) Review of the genus *Amphinemura* (Plecoptera: Nemouridae) from Guangdong, China. *Zootaxa* 1511: 55–64.
- Li W-H, Yang D (2008a) Species of *Amphinemura* (Plecoptera: Nemouridae) from Tibet, China. *Zootaxa* 1688: 54–60.
- Li W-H, Yang D (2008b) New species of Nemouridae (Plecoptera) from China. *Aquatic Insects* 30: 205–221. doi: 10.1080/01650420802334038
- Li W-H, Yang D (2008c) A new species of *Amphinemura* (Plecoptera: Nemouridae) from China. *Zootaxa* 1892: 65–68.
- Li W-H, Yang D (2008d) Two new species of *Amphinemura* (Plecoptera: Nemouridae) from Yunnan, China, with the redescription of *A. triramia* (Wu, 1962). *Zootaxa* 1926: 61–67.
- Li W-H, Yang D (2008e) Two new species and two new records of stonefly family Nemouridae from Henan (Plecoptera: Nemouroidea). In: Shen X, Lu Ch (Ed) *The Fauna and Taxonomy of Insects in Henan*. China Agricultural Science and Technology Press, Beijing, 6: 11–16.
- Li W-H, Yang D (2011) Two new species of *Amphinemura* (Plecoptera: Nemouridae) from China. *Zootaxa* 2975: 29–34.
- Li W-H, Yang D, Sivec I (2005) A new species of *Amphinemura* (Plecoptera: Nemouridae) from China. *Entomological News* 116: 93–96.
- Shimizu T (1996) A Contribution to the Knowledge of the Family Nemouridae from East Asia (Insecta, Plecoptera). Ph.D thesis. Tokyo University of Agriculture, Tokyo, Japan.
- Shimizu T (1998a) The Group of *Amphinemura flavostigma* (Plecoptera, Nemouridae). *Aquatic Insects* 20: 203–208. doi: 10.1076/aqin.20.4.203.4469
- Shimizu T (1998b) The Group of *Amphinemura megaloba* (Plecoptera, Nemouridae). *Japanese Journal of Systematic Entomology* 42: 227–236.

- Wang Z-J, Du Y-Z, Sivec I, Li Z-Z (2006) Records and descriptions of some Nemouridae species (Order: Plecoptera) from Leigong Mountain, Guizhou province, China. *Illiesia* 2: 50–56.
- Wang Z-J, Zhang J-H, Zhu J-Y (2007) A new species of the genus *Amphinemura* (Plecoptera: Nemouridae) from Xinjiang, China. *Acta Zootaxonomica Sinica* 29: 13–16.
- Wu C.F. (1926) Two new species of stoneflies from Nanking. *The China Journal of Science and Arts* 5: 331–332.
- Wu C-F (1935) New species of stoneflies from East and South China. *Bulletin of the Peking Society of National History* 9: 227–243.
- Wu C-F (1938) *Plecopterorum sinensium*: A monograph of stoneflies of China (Order Plecoptera). Yenching University, 225 pp.
- Wu C.F. (1949) Sixth Supplement to the stoneflies of China (Order Plecoptera). *Bulletin of the Peking Natural History* 17: 251–256.
- Wu C-F (1962) Results of the Zoologico-Botanical expedition to Southwest China, 1955–1957 (Plecoptera). *Acta Entomologica Sinica* 11 (Supplement): 139–153.
- Wu C-F (1973) New species of Chinese stoneflies (Order Plecoptera). *Acta Entomologica Sinica* 16: 97–118.
- Yang D, Li W-H, Sivec I (2005) A new species of *Amphinemura* (Plecoptera: Nemouridae) from China. *Entomological News* 115: 226–228.
- Yang D, Li W-H, Zhu F (2005) A new species of *Amphinemura* from south China (Plecoptera: Nemouridae). *Zootaxa* 805: 1–4.
- Zhu F, Yang D (2002) Three new species of *Amphinemura* from China (Plecoptera: Nemouridae). *Acta Zootaxonomica Sinica* 27: 745–749.
- Zhu F, Yang D (2003) Three new species of *Amphinemura* (Plecoptera: Nemouridae) from Tibet. *Entomologia Sinica* 10: 51–56.