

A new species of *Metaeuchromius* (Lepidoptera, Crambidae) from the Tibetan glacier area of China

Wei-Chun Li¹, Dong Liu¹

¹ College of Agronomy, Jiangxi Agricultural University, Nanchang, 330045, China

Corresponding author: Wei-Chun Li (weichunlee@126.com)

Academic editor: Matthias Nuss | Received 16 October 2014 | Accepted 17 December 2014 | Published 22 January 2015

<http://zoobank.org/2ADA6539-7541-4C1C-B124-5F55CA974F37>

Citation: Li W-C, Liu D (2015) A new species of *Metaeuchromius* (Lepidoptera, Crambidae) from the Tibetan glacier area of China. ZooKeys 475: 113–118. doi: 10.3897/zookeys.475.8766

Abstract

Metaeuchromius glacialis Li, **sp. n.** is described from the Tibetan glacier area of China. The new species is similar to *M. circe* Bleszynski by the distal projection of costa exceeding the apex of valva, and the phallus with strong spine-like cornuti in the male genitalia. Images of male adult, tympanal and scent organs as well as genitalia of the new species are provided.

Keywords

Pyraloidea, Crambinae, taxonomy, Qinghai-Tibetan Plateau

Introduction

Metaeuchromius was established with *Eromene yuennanensis* Caradja, 1937 as the type species (Bleszynski 1960). The genus has fourteen species with a Palearctic and Oriental distribution (Bleszynski 1965; Park 1990; Schouten 1997; Nuss and Speidel 1999; Chen et al. 2002; Sasaki 2005; Li et al. 2009; Nuss et al. 2014). Prior to this study, eight species of *Metaeuchromius* were recorded in China (Li et al. 2009). However, the known species are rarely recorded around the monsoon temperate glaciers of the south-eastern part of the Qinghai-Tibetan Plateau in China. Moreover, it is becoming more and more critical to pay close attention to the species living around mountain glaciers because global warming is highly likely to cause changes in the biodiversity

of these regions (Li and Liu 2014). In the present paper, *Metaeuchromius glacialis*, collected from Galongla Mountain, Tibetan glacier area, is described as new to science.

Material and methods

Specimens were collected by a 250-W high-pressure mercury lamp. They were hand-collected alive and killed by ammonium hydroxide just prior to mounting and spreading (Landry and Landry 1994). The type specimens of *Metaeuchromius* species are deposited in the Natural History Museum of London (NHM) have been examined by corresponding author. All the type specimens of the new species are deposited in the Insect Museum, Jiangxi Agricultural University, Nanchang, China (JXAUM).

The terminology for the terminal dots formula of the forewings follows Schouten (1997), the terminology for the tympanal organs follows Maes (1985) and Nuss (2005).

Genitalia preparation followed the standard procedure, using boiling 10% KOH solutions to digest internal tissues; after careful cleaning and removal of scales, genitalia were examined, compared, and described before being mounted on microscope slides by corresponding author. The images of the adults were taken with a digital camera (Canon G12). The illustrations of the genitalia were prepared with a digital camera DV320 OPTPro2010_Ch5 attached to a digital microscope Optec BK-DM320.

Taxonomic account

Metaeuchromius glacialis Li, sp. n.


<http://zoobank.org/D38737BC-E4CF-432C-A1D4-ECFB14E07C8E>

Figs 1–3

Holotype. ♂, China, Tibet, Medog, the foot of Mt. Galongla (29°44.2947'N, 95°40.6068'E), 3415 m, 22.VII.2014, coll. Wei-Chun Li and Dong Liu et al.

Paratypes. 4 ♂♂, same data as the holotype except dated 20–23.VII.2014.


Diagnosis. In male genitalia, this new species is similar to *Metaeuchromius circe* Bleszynski, 1965 in the distal projection of costa exceeding the apex of valva, and the phallus with strong spine-like cornuti. Based on a comparison with the type specimens and additional specimens (2 ♂♂, China, Hunan Province, Shimen County, Mt. Huping, ca. 1200 m, 18.VII.2013, coll. Wei-Chun Li et al.) of *M. circe*, the new species can be distinguished by the forewing with a conspicuously convex medial fascia and terminal dots with formula 2-3-1 (Fig. 1) whereas the medial fascia of *M. circe* is straight and the terminal dots with formula 2-3-2 (Fig. 4). The new species has the saccus tympani of male tympanal organ extending to two thirds of the second sternite and the third sternite with two scent organs (Fig. 2) while the saccus tympani of *M. circe* exceeds the posterior margin of the second sternite and there is no scent organ (Fig. 5). The male genitalia of the new species have a strong spine-like projection at the



Figures 1–3. *Metaeuchromius glacialis* Li, sp. n. **1** Adult, Holotype **2** Abdomen, paratype **a** Tympanal and scent organs with scales **b** Tympanal and scent organs without scales **3** Male genitalia, paratype **a** Without phallus **b** Phallus **c** Cornuti.

end of costa of valva and the phallus with eight cornuti (Fig. 3) while *M. circe* only has a small pointed costal tip and four cornuti (Fig. 6).

Description. Adult (Fig. 1). Forewing length 9.0–10.0 mm. Frons pale brown. Vertex white except pale brown in middle. Labial palpus approximately one and half as long as compound eye's diameter, pale brown. Maxillary palpus pale brown, distally white. Antenna scapus white; flagellomere pale brown. Patagium and thorax white. Tegula blackish brown mixed with white, posterior margin with long and thin white scales. Forewing densely covered with blackish brown scales from basal one fifth to medial fascia, the other area sparsely suffused with pale brown scales; costa with longitudinal blackish brown stripe extending from base to near medial fascia; medial fascia conspicuously convex, incurved slightly near middle, running to before middle of dorsum, golden, edged with pale brown; discoidal cell with two brown spots; apex golden mixed with pale brown, with one white stripe; subterminal line golden mixed with pale brown; six terminal black dots running from middle of termen to tornus, formula 2-3-1, each group delimited by white, each dot of a group divided by golden; cilia pale brown. Hindwing white, suffused with gray scales; subterminal fascia pale brown, inconspicuous; cilia white. Fore- and midlegs pale brown, tarsi with white rings; hindleg yellowish white.



Figures 4–6. *Metaeuchromius circe* Bleszynski. **4** Adult, paratype, female **5** Tympanal organ **6** Male genitalia, additional specimen **a** Without phallus **b** Phallus.

Male abdomen (Fig. 2). Praecinctorium with a cluster of slender white scales; bulla tympani of tympanal organ open, bean-shaped, inner margin convex anteriorly and concave posteriorly; saccus tympani broad, rounded, extending to two thirds of second sternite; venula secunda present. Third sternite with two clusters of yellowish white scales on lateral side (Fig. 2a), bearing two oblong scent organs, opening towards lateral side, outside wall with pits, a cluster of slender scales attached to pits on inward tip (Fig. 2b).

Male genitalia (Fig. 3). Uncus curved downward, tapering to blunt apex. Gnathos curved upward slightly, distally rounded. Tegumen arms approximately twice as long as gnathos. Valva broad basally, narrowed towards apex; apex rounded. Costa strongly sclerotized, ending with a strong spine-like projection directed upward, reaching apex of valva. Juxta basally convex, gently broadened towards tip; distal margin incurve and forming two triangular lateral projections. Phallus nearly as long as valva; cornuti composed of eight strong spines.

Female unknown.

Distribution. This species is only known from Galongla Mountain, in Medog County, Tibet, China.

Natural history. Unknown except that the moths fly late July and are attracted to light. The habitat in which this species has been collected is located at an altitude of 3415 m, at the foot of Galongla Mountain. Most parts of the mountain are covered



Figure 7. Natural environment of collecting localitiy of *Metaeuchromius glacialis* Li, sp. n.

with snow; the vegetation on the south slope is a blend of alpine meadows, shrubs and conifer trees (Fig. 7).

Etymology. The specific name is derived from the Latin *glacialis* = glacier, in reference to the species occurance in the Tibetan glacier area.

Acknowledgments

We are grateful to Xiao-Ke Zhang and Dan-Dan He for their help during the fieldwork in the expedition to the Tibetan glacier area. Special thanks are given to Dr. Bernard Landry (Muséum d'histoire naturelle, Genève, Switzerland) and an anonymous reviewer for their critical reading of the manuscript and helpful comments. This project was supported by the National Natural Science Foundation of China (No. 31160428).

References

- Bleszynski S (1960) Studies of the Crambidae (Lepidoptera). Part XXIX. Species of the genus *Euchromius* Guenée. Acta Zoologica Cracoviensia 5: 203–247.
- Bleszynski S (1965) Crambinae. In: Amsel HG, Reisser H, Gregor F (Eds) Microlepidoptera Palaearctica 1 (1–2). Verlag Georg Fromme and Co., Wien, 553 pp.
- Chen TM, Song SM, Yuan DC (2002) A study on the genera *Metaeuchromius* Bleszynski, *Euchromius* Guenée and *Miyakea* Marumo from China, with descriptions of two new species

- of *Metaeuchromius* (Lepidoptera: Pyralidae, Crambinae). Acta Entomologica Sinica 45 (3): 365–370. doi: 10.3321/j.issn:0454-6296.2002.03.015
- Landry JF, Landry B (1994) A technique for setting and mounting Microlepidoptera. Journal of the Lepidopterists' Society 48: 205–227.
- Li WC, Li HH, Nuss M (2009) Notes on the genus *Metaeuchromius* Bleszynski (Lepidoptera: Crambidae: Crambinae) from China, with description of two new species. Zootaxa 2287: 55–63.
- Li WC, Liu D (2014) DNA barcoding and morphology reveal exceptional species diversity of *Scoparia* (Lepidoptera: Crambidae) from the Hailuoguo Glacier area, China. Zoological Journal of the Linnean Society 171: 732–752. doi: 10.1111/zoj.12154
- Maes KVN (1985) A comparative study of the abdominal tympanal organs in Pyralidae (Lepidoptera). I. Description, terminology, preparation technique. Nota lepidopterologica 8(4): 341–350.
- Nuss M (2005) Scopariinae. In: Goater B, Nuss M, Speidel W, Pyraloidea I, Huemer P, Karsholt O (Eds) Microlepidoptera of Europe 4. Stenstrup, Apollo Books, 127–180.
- Nuss M, Speidel W (1999) A new crambid moth species from the northeastern part of Turkey (Crambidae: Crambinae). Nota Lepidopterologica 22(2): 155–159.
- Nuss M, Landry B, Vegliante F, Tränkner A, Mally R, Hayden J, Segerer A, Li H, Schouten R, Solis MA, Trofimova T, De Prins J, Speidel W (2003–2014) Global information system on Pyraloidea. Available at: <http://www.pyraloidea.org/> [accessed 16 October 2014]
- Park KT (1990) Two new species of Pyralidae (Lepidoptera) from Korea. Korean Journal of Entomology 20(3): 139–144.
- Sasaki A (2005) A new species of genus *Metaeuchromius* Bleszynski (Crambinae, Crambidae) from Okinawa Is., Japan. Japan Heterocerists' Journal 236(10): 195–196.
- Schouten RTA (1997) Revision of the genus *Metaeuchromius* Bleszynski (Lepidoptera: Pyralidae: Crambinae). Tijdschrift voor Entomologie 140: 111–127.