

Annotated Checklist of the Terrestrial Gastropods of Nepal

Prem B. Budha^{1,4}, Fred Naggs², Thierry Backeljau^{3,4}

1 Tribhuvan University, Kirtipur, Kathmandu, Nepal **2** Natural History Museum, Cromwell Road, London, SW7 5BD, UK **3** Royal Belgian Institute of Natural Sciences, Vautierstraat 29, B-1000, Brussels, Belgium
4 University of Antwerp, Evolutionary Ecology Group, Groenenborgerlaan 171, B-2020, Antwerp, Belgium

Corresponding author: Prem B. Budha (pbbudha@cdtu.edu.np; prembudha@yahoo.com)

Academic editor: M. Schilthuizen | Received 26 December 2014 | Accepted 10 March 2015 | Published 30 March 2015

<http://zoobank.org/79C9A851-D8B3-4024-B5F1-C3EC84C6095E>

Citation: Budha PB, Naggs F, Backeljau T (2015) Annotated Checklist of the Terrestrial Gastropods of Nepal. ZooKeys 492: 1–48. doi: 10.3897/zookeys.492.9175

Abstract

This is the very first checklist of the terrestrial gastropods of Nepal. It includes 138 species and six subspecies, of which 22 species are endemic and four are introduced. It highlights 34 species recorded for the first time in Nepal and provides new distribution records for another 30 species.

Keywords

Gastropoda, checklist, endemic, Nepal

Introduction

The rich diversity of non-marine molluscs of the Indian subcontinent was explored by pioneering 19th century British malacologists. However, as foreigners were restricted from entering Nepal until 1951, the Nepalese malacofauna remained poorly known. *Pupa eurina* Benson, 1864 (now *Pupilla eurina*) may have been the first land snail recorded from Nepal (Budha 2005), but its type locality of ‘Tribeni Ghat’ has not been identified. The earliest confirmed records of terrestrial gastropods from Nepal were an unidentified *Nanina* species and *Anadenus* sp. [= *A. giganteus* Heynemann] from Kathmandu Valley (Nevill 1878: p. 27 and 65 respectively). No publications on Nepa-

lese snails appeared for the following 95 years until the description of two new species and two new subspecies of *Hemiphaedusa* by Nordsieck (1973) and a chromosomal study on some ariophantids (Kiauta and Butot 1973). Subsequently, Schileyko and Frank (1994) described a new species (*Laevozebrinus nepalensis*), a new genus (*Nepaliella*) and a new enid subfamily (Pseudonapaeinae) from hills surrounding Kathmandu Valley. In addition, they described the reproductive anatomy of *Oxytesta orobia* (Benson, 1848). Since then, several occasional papers on the Nepalese terrestrial gastropods have been published (Kuznetsov 1996, Kuznetsov and Schileyko 1997, 1999, Schileyko and Kuznetsov 1998a, 1998b, Raut 1999, Schileyko 1999, Wiktor 2001a, Subba and Ghosh 2001, 2008, Wiktor and Bössneck 2004, Budha 2005, Kuzminykh and Schileyko 2005, Bössneck 2006, Budha and Naggs 2008, Gerber and Bössneck 2009, Budha et al. 2012, Schileyko and Balashov 2012, Khanal and Budha 2013). Despite the fact that the study of Nepalese terrestrial gastropods is still in its infancy, there is a need for at least a provisional checklist as a starting point for further study. The present paper aims at providing such a list.

The data included here are based on published records and field investigations from 2006–2010 by Prem Budha. Collected material has been deposited in the Central Department Zoology Museum of Tribhuvan University (CDZMTU), Kirtipur, Kathmandu, Nepal. The list provides taxonomic notes where needed, as well as distribution ranges of genera and species. The original names of the type species of genera and subgenera are provided. An attempt was made to standardize the use of geographical place names and local features but, owing to the nature of this data, it was not always possible to do so. The district name is mentioned for all species from Nepal with particular locations such as hill, forest, and village names wherever data are available. National park or conservation areas are given without district names because most national parks extend across more than one district. Particular locations within national parks are given where known. Indian states are given with particular location(s) wherever data are available. The systematic arrangement at family and more inclusive levels is based on Bouchet and Rocroi (2005). Family names are arranged according to Bouchet and Rocroi (2005), while genus and species names are arranged alphabetically. The list includes 138 species and six subspecies, including 22 endemic species, four introduced species, 34 new species for Nepal, and new distribution records for 30 species.

Systematics

Class: Gastropoda Cuvier, 1795

Clade: Caenogastropoda Cox, 1960

Superfamily: Cyclophoroidea J.E. Gray, 1847

Family: Cyclophoridae J.E. Gray, 1847

Subfamily: Cyclophorinae J.E. Gray, 1847

Genus: *Cyclophorus* Montfort, 1810¹

Distribution: Subtropical and tropical Asia (Gude 1921, Benthem Jutting 1948, Zilch 1956, Kongim et al. 2006).

Type species: *Helix volvulus* O.F. Müller, 1774

Subgenus: *Glossostylus* Kobelt & Möllendorff, 1897

Distribution: India; Sri Lanka; Myanmar; Thailand; Vietnam; Taiwan; Malaysia; Philippines (Gude 1921).

Type species: *Cyclostoma validum* Sowerby, 1842

Cyclophorus (Glossostylus) fulguratus (L. Pfeiffer, 1852)²

Distribution: Myanmar; Thailand; Vietnam (Gude 1921).

Nepal: Ilam, Jhapa, Morang, Sunsari, Dharan, Udayapur and Gulmi Districts (Subba and Ghosh 2001).

Subgenus: *Kobeltostylus* Egorov, 2006³

Distribution: Bangladesh; India; Sri Lanka; Myanmar; Philippines (Gude 1921).

Type species: *Helix involvulus* O.F. Müller, 1774

Cyclophorus (Kobeltostylus) pyrotrema Benson, 1854

Distribution: Bangladesh; India; Myanmar (Gude 1921).

Nepal: Lalitpur District-Phulchowki Hill (Kuznetsov and Schileyko 1997).

Subgenus: *Annularia* Schumacher, 1817⁴

Distribution: India; Sri Lanka; Myanmar; Philippines (Gude 1921).

Type species: *Annularia aurantiaca* Schumacher, 1817

Cyclophorus (Annularia) aurantiacus (Schumacher, 1817)⁵

Distribution: Thailand; Myanmar; W Malaysia (Nevill 1878, Gude 1921).

Nepal: Ilam, Morang, Sunsari, Dharan and Udayapur Districts (Subba and Ghosh 2001).

Genus: *Theobaldius* Nevill, 1878

Distribution: Sri Lanka; S and NE India; Myanmar (Gude 1921).

Type species: *Cyclophorus annulatus* L. Pfeiffer, 1847

Theobaldius sp.

New species record for Nepal: Shivapuri-Nagarjun and Langtang National Parks.

Genus: *Scabrina* W.T. Blanford, 1863⁶

Distribution: S and SE Asia (Gude 1921, Maassen 2006).

Type species: *Cyclophorus calyx* Benson, 1847.

Scabrina phaenotopicus (Benson, 1851)

Distribution: India: W Bengal-Darjeeling, Sikkim (Gude 1921).

Nepal: Raheem et al. (2010).

New distribution records from Nepal: Chitwan National Park, Tanahu District-Shiddha Cave area and Lalitpur District-Phulchowki Hill.

Genus: *Pterocyclos* Benson, 1832

Distribution: India; Sri Lanka; SE Asia (Raheem and Naggs 2006, Ramakrishna et al. 2010, Kongim et al. 2013).

Type species: *Pterocyclos rupestris* Benson, 1832

***Pterocyclos cf. brahmakundensis* Godwin-Austen, 1915**

Distribution: India: Assam-Brahmakund (Gude 1921).

New species record for Nepal: Langtang National Park.

Subfamily: Alycaeinae W.T. Blanford, 1864⁷**Genus: *Alycaeus* J.E. Gray, 1850**

Distribution: India; Nepal; Myanmar; China; Japan; Taiwan; Korea; Thailand; Vietnam; Laos; Philippines; Indonesia; Malaysia; Australia (Gude 1921, Tarruella and Domènec 2011).

Type species: *Alycaeus eydouxi* Venmans, 1956⁸

Subgenus: *Alycaeus* J.E. Gray, 1850

Distribution: India; Myanmar; China; Malaysia; Japan (Gude 1921).

Type species: *Alycaeus eydouxi* Venmans, 1956

***Alycaeus (Alycaeus) burti* Godwin-Austen, 1874**

Distribution: India: Arunachal Pradesh, Assam, Mizoram-Akha Hills, Dihiri Parbat; Bhutan (Gude 1921, Ramakrishna et al. 2010).

Nepal: Solukhumbu District (Kuznetsov and Schileyko 1997).

New distribution records from Nepal: Kathmandu District-Champadevi Hill, Lalitpur District-Phulchowki Hill and Shivapuri-Nagarjun National Park.

***Alycaeus (Alycaeus) lohitensis* Godwin-Austen, 1914**

Distribution: India: Assam, Arunachal Pradesh (Gude 1921, Ramakrishna et al. 2010).

Nepal: Lalitpur District-Phulchowki Hill (Kuznetsov 1996).

***Alycaeus (Alycaeus) yamneyensis* Godwin-Austen, 1914**

Distribution: India: Arunachal Pradesh-Yamne Valley, Abor Hills (Gude 1921).

Nepal: Tarruella and Domènec (2011).

Genus: *Chamalycaeus* Kobelt & Möllendorff, 1897

Distribution: India; Nepal; Myanmar; China; Taiwan; Korea; Thailand; Vietnam; Laos; Philippines; Indonesia; Malaysia; Australia (Gude 1921, Tarruella and Domènec 2011).

Type species: *Alycaeus andamaniae* Benson, 1861

Subgenus: *Dicharax* Kobelt & Möllendorff, 1900

Distribution: NE India; Myanmar; China; Malaysia (Gude 1921).

Type species: *Alycaeus hebes* Benson, 1857

Chamalycaeus (Dicharax) birenatus (Godwin-Austen, 1874)

Distribution: NE India: Assam, Nagaland-Naga Hill (Ramakrishna et al. 2010).

Nepal: Lalitpur District-Phulchowki Hill (Kuznetsov 1996).

Chamalycaeus (Dicharax) digitatus (H.F. Blanford, 1871)

Distribution: NE India: W Bengal-Darjeeling, Sikkim-Richila Peak; W Bhutan (Gude 1921).

New species record for Nepal: Kathmandu District-Champadevi Hill, Lalitpur District-Phulchowki Hill, Shivapuri-Nagarjun National Park.

Chamalycaeus (Dicharax) inflatus (Godwin-Austen, 1874)

Distribution: NE India: Nagaland-Naga Hills (Ramakrishna et al. 2010).

Nepal: Shivapuri-Nagarjun National Park-Nagarjun Forest (Khanal and Budha 2013).

Chamalycaeus (Dicharax) notatus (Godwin-Austen, 1876)

Distribution: NE India: Nagaland-Naga Hills, Arunachal Pradesh-Dafta Hills (Gude 1921).

Nepal: Solukhumbu District (Kuznetsov and Schileyko 1997).

Chamalycaeus (Dicharax) plectochilus (Benson, 1859)

Distribution: NE India: W Bengal-Darjeeling, Sikkim-Damsang Peak; W Bhutan (Gude, 1921).

New species records for Nepal: Kathmandu District-Champadevi Hill, Shivapuri-Nagarjun and Langtang National Parks.

Chamalycaeus (Dicharax) strangulatus (L. Pfeiffer, 1846)⁹

Distribution: This is the only species of *Chamalycaeus (Dicharax)* recorded from the W Himalaya, NW India: Himachal Pradesh-Simla, Uttarakhand-Kumaon, Nainital (Ramakrishna et al. 2010).

New species record for Nepal: Shivapuri-Nagarjun National Park.

Chamalyceus (Dicharax) stylifer (Benson, 1857)

Distribution: NE India: W Bengal-Darjeeling and Sikkim; Bhutan (Gude 1921).
New species records for Nepal: Lalitpur District-Phulchowki Hill, Shivapuri-Nagarjun and Langtang National Parks.

Subgenus: *Cycloryx* Godwin-Austen, 1914

Distribution: NE India to Myanmar (Gude 1921).
Type species: *Alyceus constrictus* Benson, 1851

Chamalyceus (Cycloryx) otiphorus (Benson, 1858)

Distribution: NE India: W Bengal-Darjeeling, Sikkim-Pankhabari, Meghalaya, Nagaland (Gude 1921, Ramakrishna et al. 2010).
Nepal: Lalitpur District-Phulchowki Hill (Kuznetsov 1996).
New distribution record from Nepal: Shivapuri-Nagarjun National Park.

Chamalyceus (Cycloryx) summus (Godwin-Austen, 1914)

Distribution: NE India: Sikkim-Richila Peak; W Bhutan (Gude 1921).
Nepal: Solukhumbu District (Kuznetsov and Schileyko 1997).

Family: Diplommatinidae L. Pfeiffer, 1856

Subfamily: Diplommatininae L. Pfeiffer, 1856

Genus: *Diplommatina* Benson, 1849¹⁰

Distribution: India; Nepal; China; Indonesia; Vietnam; Singapore; Malaysia; Japan; Philippines; Taiwan; Papua New Guinea; Fiji (Gude 1921, Zilch 1953, Schileyko and Kuznetsov 1997, Maassen 2002, Panha and Burch 2005, Webster et al. 2012).
Type species: *Bulimus folliculus* L. Pfeiffer, 1846

Subgenus: *Diplommatina* Benson, 1849

Distribution: N India; Nepal; China; Malaysia; Philippines; Japan; Taiwan; Papua New Guinea; Fiji (Gude 1921, Zilch 1953, Schileyko and Kuznetsov 1997, Panha and Burch 2005, Webster et al. 2012).

Type species: *Bulimus folliculus* L. Pfeiffer, 1846

Diplommatina (Diplommatina) exserta Godwin-Austen, 1886

Distribution: Myanmar: Damotha Cave, etc., Moulmein, now Mawlamyine (Gude 1921).
New species record for Nepal: Tanahu District-Siddha Cave area.

Diplommatina (Diplommatina) folliculus (L. Pfeiffer, 1846)

Distribution: NW India: Himachal Pradesh-Landour, Simla, Uttarakhand-Nainital (Naggs 1997, Ramakrishna et al. 2010).
New species records for Nepal: Lalitpur District-Phulchowki Hill, Shivapuri-Nagarjun and Langtang National Parks.

Diplommatina (Diplommatina) munipurensis Godwin-Austen, 1892

Distribution: NE India: Manipur; Myanmar (Gude 1921).

New species records for Nepal: Lalitpur District-Phulchowki Hill, Langtang National Park.

Diplommatina (Diplommatina) oviformis Fulton, 1901

Distribution: India: W Bengal-Darjeeling (Gude 1921).

Nepal: Solukhumbu District (Kuznetsov and Schileyko 1997).

New distribution records from Nepal: Hills surrounding Lalitpur and Kathmandu Districts, Shivapuri-Nagarjun and Langtang National Parks.

Diplommatina (Diplommatina) pachychilus Benson, 1857

Distribution: NE India: W Bengal-Darjeeling (Gude 1921).

Nepal: Solukhumbu District (Kuznetsov and Schileyko 1997).

New distribution records from Nepal: Shivapuri-Nagarjun and Langtang National Parks.

Diplommatina (Diplommatina) regularis Fulton, 1901.

Distribution: NE India: W Bengal-Darjeeling (Ramakrishna et al. 2010).

New species record for Nepal: Shivapuri-Nagarjun National Park-Baghitar.

Diplommatina (Diplommatina) silvicola Godwin-Austen, 1886

Distribution: NE India: Assam-North Cachar, Jenta Hajuma Peak (Gude 1921, Ramakrishna et al. 2010).

New species record for Nepal: Shivapuri-Nagarjun National Park-Balaju, Pani Tanki.

Diplommatina (Diplommatina) sperata W.T. Blanford, 1862

Distribution: Myanmar (Gude 1921).

Nepal: Solukhumbu District (Kuznetsov and Schileyko 1997).

Subgenus: *Metadiancta* Möllendorff, 1898

Distribution: NE India: Assam, Manipur, Nagaland; Myanmar; Vietnam (Gude 1921).

Type species: *Diplommatina dohertyi* Godwin-Austen, 1892

Diplommatina (Metadiancta) miriensis Godwin-Austen, 1917

Distribution: NE India: Arunachal Pradesh-Miri Hills (Gude 1921).

New species records for Nepal: Shivapuri-Nagarjun and Langtang National Parks.

Subgenus: *Sinica* Möllendorff, 1885

Distribution: India; Nepal; Myanmar; China; Japan; Philippines; Indonesia; Malaysia; Papua New Guinea; Taiwan (Gude 1921, Zilch 1953, Kuznetsov and Schileyko 1997).

Type species: *Diplommatina collarifera* Schmacker and Boettger, 1877

Diplommatina (Sinica) canarica Beddome, 1875¹¹

Distribution: India: Western Ghats, Karnataka, Maharashtra (Ramakrishna et al. 2010, Raheem et al. 2014).

Nepal: Solukhumbu District (Kuznetsov and Schileyko 1997).

Family: Pupinidae L. Pfeiffer, 1853

Subfamily: Pupininae L. Pfeiffer, 1853

Genus: *Schistoloma* Kobelt, 1902

Distribution: Indian Himalaya; Nepal; China; Thailand; W Malaysia; Sumatra; Borneo; Philippines (Gude 1921, Bartsch 1915, Tumpeeswan and Panha 2008).

Type species: *Cyclostoma altum* Sowerby, 1842

Schistoloma cf. funiculatum (Benson, 1838)¹²

Distribution: India: W Bengal-Darjeeling (Gude 1921).

New species records for Nepal: Lalitpur District-Phulchowki Hill, Shivapuri-Nagarjun and Langtang National Parks.

Superfamily: Ellobioidea L. Pfeiffer, 1854 (1822)

Family: Ellobiidae L. Pfeiffer, 1854 (1822)

Subfamily: Carychiinae Jeffreys, 1830

Genus: *Carychium* O.F. Müller, 1773

Distribution: Very widely distributed from N and C America, Europe to S and SE Asia (Burch and Panha 2002, Thompson 2011).

Type species: *Carychium minimum* O.F. Müller, 1774

Carychium minusculum Gredler, 1888¹³

Distribution: China "aus Hope" (Gredler 1888).

Nepal: Langtang National Park-Syabru (Kuznetsov and Schileyko 1997), Kavre District-Chandeshwari (Nesemann et al. 2007).

Carychium sp.¹⁴

New species records for Nepal: Lalitpur District-Phulchowki Hill, Shivapuri-Nagarjun and Langtang National Parks.

Clade: Systellomatophora Pilsbry, 1948

Superfamily: Veronicelloidea J.E. Gray, 1840

Family: Veronicellidae J.E. Gray, 1840

Genus: *Laevicaulis* Simroth, 1913

Distribution: Pantropical (Stanisic 1998).

Type species: *Vaginulus alte* Féussac, 1822

Laevicaulis alte (Férussac, 1822)¹⁵

Distribution: The geographical origin of *L. alte* is uncertain, but it has been widely distributed in tropical and subtropical countries through human agency (Stanisic 1998). New distribution records from Nepal: Widely distributed throughout most districts of Tarai and inner Tarai.

Clade: Stylommatophora Schmidt, 1855

Superfamily: Succineoidea H. Beck, 1837

Family: Succineidae H. Beck, 1837

Subfamily: Catinellinae Odhner, 1950

Genus: *Quickia* Odhner, 1950

Distribution: W and E Africa; Mascarene Islands; Seychelles; Aldabra; India; Nepal (Patterson 1975, Schileyko 2007).

Type species: *Succinea concisa* Morelet, 1848

Quickia sp.

New species record for Nepal: Chitwan District-Sauraha (collected from flower vase at hotel).

Subfamily: Succineinae H. Beck, 1837

Genus: *Succinea* Draparnaud, 1801

Distribution: Nearly circumglobal (Schileyko 2007), the Northern Hemisphere; Australia; some Pacific islands (Thompson 2011).

Type species: *Succinea amphibia* Draparnaud, 1801 (= *Helix putris* Linnaeus, 1758)

Succinea sp.

New species records for Nepal: Kathmandu and Lalitpur Districts.

Superfamily: Pupilloidea Turton, 1831

Family: Pupillidae Turton, 1831

Subfamily: Pupillinae Turton, 1831

Genus: *Pupilla* Fleming, 1828

Distribution: Temperate N America; Europe; Africa; Asia; Australia (Gude 1914, Pokryszko et al. 2009).

Type Species: *Pupa marginata* Draparnaud, 1801 (= *Turbo muscorum* Linnaeus, 1758).

Pupilla annandalei Pilsbry, 1921¹⁶

Distribution: Pakistan.

Nepal: Pokryszko et al. (2009).

Pupilla eurina (Benson, 1864)¹⁷

Distribution: Endemic to Nepal.

Nepal: Tribeni Ghat (Blanford and Godwin-Austen 1908), Annapurna Conservation Area-Tukuche (Kuznetsov and Schileyko 1997).

New distribution record from Nepal: Langtang National Park-Gosainkund.

Pupilla triplicata (Studer, 1820)

Distribution: Europe and C Asia (Sysoev and Schileyko 2009).

Nepal: Annapurna Conservation Area-Tukuche (Kuznetsov and Schileyko 1997).

Family: Pyramidulidae Kennard & Woodward, 1914

Genus: *Pyramidula* Fitzinger, 1833

Distribution: Holarctic and S Asia (Gittenberger and Bank 1996, Schileyko and Balashov 2012).

Type species: *Helix rupestrис* Draparnaud, 1801

Pyramidula humilis (Hutton, 1838)¹⁸

Distribution: NW India: Himachal Pradesh, Punjab, Uttarakhand (Ramakrishna et al. 2010)

Nepal: Shivapuri-Nagarjun National Park-Nagarjun Forest (Khanal and Budha 2013).

Pyramidula kuznetsovi Schileyko & Balashov, 2012¹⁹

Distribution: Endemic to Nepal.

Nepal: Mustang District-Muktinath (Schileyko and Balashov 2012).

Family: Valloniidae Morse, 1864

Genus: *Vallonia* Risso, 1826

Distribution: Holarctic (Gerber and Bössneck 2009).

Type species: *Vallonia rosalia* Risso, 1826 (= *Helix pulchella* O.F. Müller, 1774)

Vallonia costohimala Gerber & Bössneck, 2009

Distribution: Endemic to Nepal.

Nepal: Northern districts from Darchula to Panchthar (Gerber and Bössneck 2009).

Vallonia himalaevi Gerber & Bössneck, 2009

Distribution: Endemic to Nepal.

Nepal: Northern districts from Darchula to Panchthar (Gerber and Bössneck 2009).

Vallonia kathrinae Gerber & Bössneck, 2009

Distribution: Endemic to Nepal.

Nepal: Muju and Mustang Districts (Gerber and Bössneck 2009).

Vallonia ladacensis (Nevill, 1878)

Distribution: India: Western Ghats, Jammu and Kashmir; Nepal; Tibet; Tianshan Turkey (Gerber and Bössneck 2009, Raheem et al. 2014).
Nepal: Bajura, Darchula, Humla and Mustang Districts (Gerber and Bössneck 2009).

Vallonia tenuilabris (A. Braun, 1843)

Distribution: Kazakhstan; Tajikistan; NW India: Jammu and Kashmir; Tibet; Siberia; N China; Mongolia to Russia (Gerber and Bössneck 2009).
Nepal: Solukhumbu and Taplejung Districts (Gerber and Bössneck 2009).

Family: Vertiginidae Fitzinger, 1833

Subfamily: Vertigininae Fitzinger, 1833

Genus: *Truncatellina* Lowe, 1852

Distribution: Holarctic (Schileyko 1998).
Type species: *Pupa linearis* Lowe, 1852

Truncatellina sp.

Nepal: Annapurna Conservation Area-Khobang (Kuznetsov and Schileyko 1997).

Subfamily: Gastrocoptinae Pilsbry, 1918

Genus: *Gastrocopta* Wollaston, 1878

Distribution: Almost cosmopolitan extending to all tropical and warm temperate continents but extinct in Europe (Pilsbry 1916–1918).
Type species: *Pupa acarus* Benson, 1856

Gastrocopta huttoniana (Benson, 1849)

Distribution: India: Western Ghats, Himachal Pradesh, Kashmir, Maharashtra (Ramakrishna et al. 2010, Raheem et al. 2014).
Nepal: Annapurna Conservation Area (Kuznetsov and Schileyko 1997).

Superfamily: Enoidea Woodward, 1903

Family: Enidae Woodward, 1903

Subfamily: Pseudonapaeinae Schileyko, 1978

Genus: *Pupinidius* Möllendorff, 1901

Distribution: W China; Nepal (Schileyko 1998, Wu and Zheng 2009).
Type species: *Buliminus pupinidius* Möllendorff, 1901

Pupinidius himalayanus Kuznetsov & Schileyko, 1999

Distribution: Endemic to Nepal.
Nepal: Mustang District, Tukuche to Muktinath trekking route (Kuznetsov and Schileyko 1999).

Pupinidius siniayevi Kuznetsov & Schileyko, 1999

Distribution: Endemic to Nepal.

Nepal: Mustang District-Tukuche to Muktinath trekking route (Kuznetsov and Schileyko 1999).

Pupinidius tukuchensis Kuznetsov & Schileyko, 1997

Distribution: Endemic to Nepal.

Nepal: Mustang District-Tukuche (Kuznetsov and Schileyko 1997).

Genus: *Laevozebrinus* Lindholm, 1925

Distribution: Afghanistan; Iran; mountain regions of C Asia; N Pakistan and adjacent territories of India (Schileyko 1998).

Type species: *Buliminus urgutensis* Kobelt, 1902

Laevozebrinus mustangensis Kuznetsov & Schileyko, 1997

Distribution: Endemic to Nepal.

Nepal: Mustang District-Tukuche to Muktinath trekking route (Kuznetsov and Schileyko 1997).

Laevozebrinus nepalensis Schileyko & Frank, 1994

Distribution: Endemic to Nepal.

Nepal: Annapurna Conservation Area and hills surrounding Kathmandu District (Schileyko and Frank 1994).

Subspecies: *nepalensis* Schileyko & Frank, 1994

Distribution: Mustang District-Khobang, Tukuche, Marpha, Jomsom (Kuznetsov and Schileyko 1997).

Subspecies: *myagdiensis* Kuznetsov & Schileyko, 1997

Distribution: Myagdi District-Sukebagar, Titre, Dana (Kuznetsov and Schileyko 1997).

Genus: *Mirus* Albers, 1850

Distribution: India; Sri Lanka; Myanmar; E Asia; Japan (Kuznetsov and Schileyko 1997, Schileyko 1998, Raheem and Naggs 2006).

Type species: *Bulimus cantorii* Philippi, 1844²⁰

Mirus (?) nilagiricus (L. Pfeiffer, 1846)²¹

Distribution: India: Western Ghats, Tamil Nadu-Nilgiris, Arunachal Pradesh-Dafla Hill, Meghalaya-Khasi Hills; Myanmar (Gude 1914, Ramakrishna et al. 2010, Raheem et al. 2014).

Nepal: Solukhumbu District-Khari Khola (Kuznetsov and Schileyko 1997).

Genus: *Nepalirena* Schileyko & Frank, 1994

Distribution: Endemic to Nepal (Kuznetsov and Schileyko 1997).

Type species: *Bulimus ceratinus* Benson, 1849

Nepalirena ceratina (Benson, 1849)

Distribution: Endemic to Nepal.

Nepal: Kathmandu and Myagdi Districts, Annapurna Conservation Area (Schileyko and Frank 1994, Kuznetsov and Schileyko 1997).

Genus: *Subzebrinus* Westerlund, 1887

Distribution: SE Kazakhstan and adjacent territories of China; India; Japan; Nepal (Gude 1914, Schileyko 1998, Raheem et al. 2010).

Type species: *Buliminus labiellus* Martens, 1881

Subzebrinus rufistriatus (Reeve, 1849)

Distribution: India: Kashmir between Jamuna and Sutlej River, Jhelum Valley (Gude 1914)

New species record for Nepal: Mugu District-Rogumba.

Family: Cerastidae Wenz, 1923

Genus: *Darwininitium* Budha & Mordan, 2012²²

Distribution: Endemic to Nepal.

Type species: *Darwininitium shivalikianum* Budha & Mordan, 2012

Darwininitium shivalikianum Budha & Mordan, 2012

Distribution: Endemic to Nepal.

Nepal: Shiwalik range of C Nepal, Chitwan National Park and Makwanpur District-Taubas, Bhaise (Budha et al. 2012).

Superfamily: Clausilioidea J.E. Gray, 1855

Family: Clausiliidae J.E. Gray, 1855

Subfamily: Phaedusinae A.J. Wagner, 1922

Genus: *Cylindrophaedusa* O. Boettger, 1877²³

Distribution: Pakistan; India; Nepal; Bhutan; Myanmar (Nordsieck 2002).

Type species: *Clausilia cylindrica* L. Pfeiffer, 1846

Subgenus: *Cylindrophaedusa* O. Boettger, 1877

Distribution: India: Punjab, W Bengal (Nordsieck 1973, 2002).

Type species: *Clausilia cylindrica* L. Pfeiffer, 1846

Cylindrophaedusa (Cylindrophaedusa) cylindrica (L. Pfeiffer, 1846)

Distribution: India; Punjab-Muree, W Bengal-Darjeeling (Nordsieck 1973, 2002).
New species record for Nepal: Dadeldhura District.

Subgenus: *Montiphaedusa* Nordsieck, 2002

Distribution: N Pakistan; Nepal; NE India; Bhutan; Myanmar (Nordsieck 2002).
Type species: *Clausilia ioes* Benson, 1852

Cylindrophaedusa (Montiphaedusa) ioes (Benson, 1852)

Distribution: N Pakistan; Nepal; NE India; Bhutan; Myanmar (Nordsieck 2002).

Subspecies: *jiriensis* (Nordsieck, 1973)

Distribution: Endemic to Nepal.
Nepal: Dolakha District-Jiri (Nordsieck 1973).

Cylindrophaedusa (Montiphaedusa) kathmandica (Nordsieck, 1973)

Distribution: Endemic to Nepal.
Nepal: Hills surrounding Kathmandu Valley (Nordsieck 1973).
New distribution records from Nepal: Lalitpur District-Phulchowki Hill, Shivapuri-Nagarjun and Langtang National Parks.

Cylindrophaedusa (Montiphaedusa) martensiana (Nordsieck, 1973)

Distribution: Endemic to Nepal.
Nepal: Lamjung, Myagdi and Mustang Districts (Nordsieck 1973).

Subspecies: *martensiana* (Nordsieck, 1973)

Distribution: Myagdi and Mustang Districts-Dhorpatan, Thakkhola, Lete, Gorepani (Nordsieck 1973).

Subspecies: *dhaulagirica* (Nordsieck, 1973)

Distribution: Lamjung District-Jaljala, Myagdi Khola, Muri (Nordsieck 1973).

Superfamily: Achatinoidea Swainson, 1840

Family: Achatinidae Swainson, 1840

Genus: *Lissachatina* Bequaert, 1950²⁴

Distribution: Originally from E Africa but now globally distributed in tropical to warm temperate areas, i.e. W Africa; N and S America; S and SE Asia; China; Japan; Caribbean countries; Oceania (Tillier et al. 1993, Raut and Barker 2002, EPPO 2013).

Type species: *Achatina fulica* Bowdich, 1822

Lissachatina fulica (Bowdich, 1822)

Distribution: See distribution of *Lissachatina*.

Nepal: Probably introduced into Nepal in the 1930s-40s (Raut 1999). It is now established as a pest in all districts of Tarai and the inner valleys causing significant damage to crops. It has spread into the mid hill districts: Kaski, Baglung, Makwanpur, Chitwan, Myagdi, Tanahun, Dhading, Palpa, Gulmi, Syangja (Budha and Naggs 2008).

New distribution records from Nepal: Dang, Surkhet, Banke, Bardia, Kailali and Kanchanpur Districts.

Family: Ferussaciidae Bourguignat, 1883

Genus: *Cecilioides* Féruccac, 1814²⁵

Distribution: Europe; Africa; S Asia; Philippines; Oceania; American tropics (Thompson 2011).

Type species: *Buccinum acicula* O.F. Müller, 1774

Cecilioides cf. *minuta* Mousson, 1874²⁶

Distribution: Drift debris of the Euphrates (type locality), Sarus River near Adana, SE Asia Minor (Pilsbry and Tryon 1908–1909).

New species record for Nepal: Baitadi District, Far W Nepal.

Family: Subulinidae P. Fischer & Crosse, 1877

Subfamily: Subulininae P. Fischer and Crosse, 1877²⁷Genus: *Allopeas* H.B. Baker, 1935²⁸

Distribution: Tropical, subtropical, and many temperate regions of Africa, S and SE Asia (Schileyko 1999, Thompson 2011).

Type species: *Bulimus gracilis* Hutton, 1834 (= *Allopeas gracile* (Hutton, 1834))

Allopeas clavulinum (Potiez & Michaud, 1838)²⁹

Distribution: Bourbon Island (type locality), other islands of the Indian Ocean; Japan (Pilsbry 1946).

New species records for Nepal: Kathmandu, Kaski and Kailali Districts.

Allopeas gracile (Hutton, 1834)

Distribution: Tropics of both hemispheres, abundant in cultivated districts, perhaps the most widely ranging of all land snails (Pilsbry 1946).

New species records for Nepal: Chitwan and Dhading Districts.

Genus: *Curvella* Chaper, 1885

Distribution: S Africa; India; China; SE Asia (Gude 1914, Schileyko 1999).

Type species: *Curvella sulcata* Chaper, 1885

Curvella sikkimensis Gude, 1914

Distribution: India: W Bengal-Darjeeling, Sikkim (Gude 1914).
 New species record for Nepal: Ilam District-Maipokhari.

Genus: *Paropeas* Pilsbry, 1906

Distribution: Widespread in the tropical Indo-Pacific regions (Naggs 1994).
 Type species: *Bulimus acutissimum* Mousson, 1857

Paropeas achatinaceum (L.Pfeiffer, 1846)

Distribution: Widespread in disturbed habitats in tropical Indo-Pacific regions (Naggs 1994).
 Nepal: Shivapuri-Nagarjun National Park-Nagarjun Forest (Khanal and Budha 2013).
 New distribution record from Nepal: Ramechhap District.

Subfamily: Opeatinae Thiele, 1931

Genus: *Opeas* Albers, 1850³⁰

Distribution: Worldwide in tropical, subtropical and many temperate regions (Schileyko 1999, Thompson 2011).
 Type species: *Helix goodallii* Miller, 1822

Opeas sp.

Nepal: Morang District (Subba and Ghosh 2008).

Subfamily: Glessulinae Godwin-Austen, 1920³¹

Genus: *Bacillum* Theobald, 1870³²

Distribution: NE India: W Bengal-Darjeeling, Sikkim, Assam-North Cachar, Meghalaya-Khasi Hill, Nagaland-Naga Hills (Gude 1914, Ramakrishna et al. 2010).
 Type species: *Achatina cassiaca* Reeve, 1849

Bacillum sp.³³

Nepal: Ilam and Panchthar District (Subba and Ghosh 2008).

Genus: *Glessula* Martens, 1860

Distribution: India; Sri Lanka; Thailand; Malaysia; Vietnam (Gude 1914, Godwin-Austen 1920, Schileyko 2011).
 Nepal: Kathmandu District (Kuznetsov 1996, Schileyko and Kuznetsov 1996).
 Type species: *Achatina ceylanica* L. Pfeiffer, 1845³⁴

Glessula orobia (Benson, 1860)

Distribution: India: W Bengal-Darjeeling (Gude 1914, Ramakrishna et al. 2010).
 New species record for Nepal: Ilam District-Maipokhari.

Glessula subjerdoni Beddome, 1906³⁵

Distribution: S India: Western Ghats, Andhra Pradesh-Golconda Hill, Orissa-Jaypore (Gude 1914, Ramakrishna et al. 2010, Raheem et al. 2014).
Nepal: Kathmandu District-Nagarjun Forest (Kuznetsov 1996).

Genus: *Rishetia* Godwin-Austen, 1920

Distribution: India; Sri Lanka; Nepal; Myanmar; W Bhutan (Godwin-Austen 1920)³⁶.

Type species: *Glessula (Rishetia) longispira* Godwin-Austen, 1920

Rishetia hastula (Benson, 1860)

Distribution: India: W Bengal-Darjeeling (Gude 1914, Godwin-Austen 1920).
New species record for Nepal: Chitwan National Park.

Rishetia tenuispira (Benson, 1836)³⁷

Distribution: India: Western Ghats, W Bengal, Sikkim, Mizoram, Arunachal Pradesh, Maharastra; Myanmar; Bangladesh (Pilsbry and Tryon 1908-1909, Gude 1914, Ramakrishna et al. 2010, Raheem et al. 2014).

Nepal: Shivapuri-Nagarjun National Park-Nagarjun Forest, Balaju (Schileyko and Kuznetsov 1996).

New distribution record from Nepal: Lalitpur District-Phulchowki Hill.

Superfamily: Streptaxoidea J.E. Gray, 1860

Family: Streptaxidae J.E. Gray, 1860

Genus: *Gulella* L. Pfeiffer, 1856

Distribution: Africa; Indo-Pacific (Bruggen 2006, Cole and Herbert 2009, Herbert and Rowson 2011, Rowson et al. 2011).

Type species: *Pupa menkeana* L. Pfeiffer, 1853

Gulella bicolor (Hutton, 1834)³⁸

Distribution: Sri Lanka; throughout India; Myanmar (Blanford and Godwin-Austen 1908); Brazil (Pilsbry 1926, Simone 2013); SE China (Yen 1939); Cuba (Sarasúa 1944, Maceira et al. 2013); Caribbean Islands (Schalie 1948); Philippines; Indonesia (Benthem Jutting 1950); Andaman and Nicobar Islands; Malaysia; Singapore (Benthem Jutting 1961); Kenya (Clench 1964); Venezuela; French Guiana (Tillier 1980); Japan (Azuma 1982); Australia (Stanisic 1981); N America (Dundee and Baerwold 1984); Oman (Mordan 1988); Bel Air area of North Mahé; Seychelles (Naggs 1989); Jamaica (Rosenberg and Muratov 2006); Vietnam (Schileyko 2011).

New species record for Nepal: Chitwan District.

Family: Diapheridae Panha & Naggs, 2010

Subfamily: Enneinae Bourguignat, 1883

Genus: *Sinoennea* Kobelt, 1904

Distribution: Japan; China; Vietnam; Malaysia; Sumatra; India (Gude 1914).

Type species: *Pupa strophoides* Gredler, 1881

Subgenus: *Indoennea* Kobelt, 1904

Distribution: India; Malaysia; Sumatra (Schileyko 2000).

Type species: *Ennea blanfordiana* Godwin-Austen, 1872

Sinoennea (Indoennea) blanfordiana Godwin-Austen, 1872

Distribution: India: Assam-North Cachar (Ramakrishna et al. 2010).

New species record for Nepal: Lalitpur District-Phulchowki Hill.

Subgenus: *Sinoennea* Kobelt, 1904

Distribution: Foothills of Himalaya; S India; China; Malay Peninsula; Sumatra; Japan; S Korea (Schileyko 2000).

Type species: *Pupa strophoides* Gredler, 1881

Sinoennea (Sinoennea) stenopylis (Benson, 1860)

Distribution: NE India: Arunachal Pradesh, Sikkim, Assam, Manipur, Meghalaya, Nagaland (Ramakrishna et al. 2010).

Nepal: Solukhumbu District (Kuznetsov and Schileyko 1997).

Superfamily: Plectopyloidea Möllendorff, 1898

Family: Plectopylidae Möllendorff, 1898

Genus: *Endothyrella* Zilch, 1960³⁹

Distribution: Nepal; NE India: Arunachal Pradesh, Assam, Nagaland, Meghalaya, Manipur, Mizoram, Sikkim (Ramakrishna et al. 2010).

Type species: *Helix pectosoma* Benson, 1836

Endothyrella affinis (Gude, 1897)⁴⁰

Distribution: NE India: Arunachal Pradesh, Assam, Meghalaya-Khasi Hill, Mizoram (Gude 1914, Ramakrishna et al. 2010).

Nepal: Kathmandu District-Swoyambhunath Temple Forest (Kuznetsov and Schileyko 1997).

Endothyrella minor (Godwin-Austen, 1879)

Distribution: India: Manipur, Meghalaya, Nagaland, Sikkim, W Bengal (Gude 1914, Ramakrishna et al. 2010).

New species records for Nepal: Lalitpur District-Phulchowki Hill, Shivapuri-Nagarjun National Park-Chisapani, Baghdwar, Langtang National Park-Golphubhanjyang.

Superfamily: Gastrodontoidea Tryon, 1866

Family: Chronidae Thiele, 1931

Subfamily: Kaliellinae Thiele, 1931

Genus: *Kaliella* W.T. Blanford, 1863

Distribution: Indo-Malayan (Blanford and Godwin-Austen 1908).

Type species: *Helix barrakporensis* L. Pfeiffer, 1853⁴¹

Kaliella barrakporensis (L. Pfeiffer, 1853)

Distribution: India; Sri Lanka; Pakistan; Madagascar; Myanmar; Tropical E Africa and Eastern S Africa (Blanford and Godwin-Austen 1908, Herbert and Kilburn 2004, Verdcourt 2006, Ramakrishna et al. 2010), hot-house alien in Britain (Preece and Naggs 2014).

Nepal: Annapurna Conservation Area (Kuznetsov and Schileyko 1977).

New distribution records from Nepal: Shivapuri-Nagarjun National Park, Lalitpur District-Phulchowki Hill, Kathmandu District-Champadevi Hill, Kirtipur.

Kaliella dikrangensis Godwin-Austen, 1883

Distribution: India: Arunachal Pradesh (Blanford and Godwin-Austen 1908, Ramakrishna et al. 2010).

Nepal: Shivapuri-Nagarjun National Park (Khanal and Budha 2013).

Kaliella fastigiata (Hutton, 1838)

Distribution: India: Himachal Pradesh, Uttarakhand, W Bengal, Arunachal Pradesh, Nagaland; Madagascar; Myanmar (Blanford and Godwin-Austen 1908, Ramakrishna et al. 2010).

Nepal: Kathmandu District-Champadevi Hill (Khanal and Budha 2013).

New distribution record from Nepal: Lalitpur District-Phulchowki Hill.

Kaliella nana (Hutton, 1838)

Distribution: India: Uttarakhand, Himachal Pradesh, W Bengal (Blanford and Godwin-Austen 1908, Ramakrishna et al. 2010).

Nepal: Annapurna Conservation Area (Kuznetsov and Schileyko 1997), Shivapuri-Nagarjun National Park (Khanal and Budha 2013).

New distribution records from Nepal: Lalitpur District-Phulchowki Hill and Kathmandu District-Champadevi Hill.

Kaliella nongsteinensis Godwin-Austen, 1883

Distribution: India: Meghalaya-Khasi Hill (Blanford and Godwin-Austen 1908, Ramakrishna et al. 2010).

Nepal: Solukhumbu District (Kuznetsov and Schileyko 1997).

Family: Euconulidae H.B. Baker, 1928

Subfamily: Euconulinae H.B. Baker, 1928

Genus: *Euconulus* Reinhardt, 1883

Distribution: Holarctic (Roth and Sadeghian 2006).

Type species: *Helix fulva* O.F. Müller, 1774

Euconulus fulvus (O.F. Müller, 1774)⁴²

Distribution: Holarctic (Roth and Sadeghian 2006).

Nepal: Hills surrounding Kathmandu Valley (Kuznetsov and Schileyko 1997).

New distribution records from Nepal: Langtang National Park and Mustang District.

Family: Pristilomatidae Cockerell, 1891

Genus: *Hawaiia* Gude, 1911⁴³

Distribution: N America from Alaska and Maine to Florida and south to Costa Rica, Cuba, Hispaniola, Jamaica, Puerto Rico, and the Virgin Islands; Europe; Japan; Australia (Kerney and Cameron 1979, Rosenberg and Muratov 2006, Sasaki 2008, Thompson 2011).

Type species: *Helix kawaiensis* Reeve, 1854 (= *Helix minuscula* Binney, 1841)

Hawaiia sp.

Nepal: Annapurna Conservation Area (Kuznetsov and Schileyko 1997).

Superfamily: Helicarionoidea Bourguignat, 1877

Family: Helicarionidae Bourguignat, 1877

Subfamily: Durgelinae Godwin-Austen, 1888

Genus: *Durgella* W.T. Blanford, 1863

Distribution: India: Arunachal Pradesh, Assam, Andhra Pradesh, Orissa, Meghalaya, Manipur, Sikkim, W Bengal; Myanmar (Blanford and Godwin-Austen 1908, Ramakrishna et al. 2010).

Type species: *Helix levicula* Benson, 1859

Durgella sp.

New species records for Nepal: Chitwan National Park; Kathmandu and Pokhara Districts.

Genus: *Sitala* H. Adams, 1865

Distribution: India; Sri Lanka; Andaman Islands; SE Asia (Schileyko 2002).

Type species: *Helix infula* Benson, 1848

Sitala rimicola (Benson, 1859)

Distribution: India: Uttarakhand, W Bengal, Sikkim, Assam, Meghalaya, Nagaland (Ramakrishna et al. 2010).

Nepal: Mustang District (Kuznetsov and Schileyko 1997).

New distribution records from Nepal: Dadeldhura, Kathmandu, Rasuwa and Mustang Districts.

Genus: *Cryptaustenia* Cockerell, 1891⁴⁴

Distribution: India; Nepal; Bhutan; Myanmar; Thailand (Schileyko 2003).

Type species: *Vitrina planospira* Benson, 1859 (= *Vitrina succinea* Reeve, 1862)

Cryptaustenia cf. globosa (Godwin-Austen, 1876)

Distribution: India: Arunachal Pradesh (Ramakrishna et al. 2010).

Nepal: Kathmandu District, Annapurna Conservation Area (Kuznetsov and Schileyko 1997).

Cryptaustenia ovata (H.F. Blanford, 1871)

Distribution: India: W Bengal (Blanford and Godwin-Austen 1908).

Nepal: Kathmandu, Panchthar, Taplejung, Morang and Terhathum Districts (Subba and Ghosh 2008), Annapurna Conservation Area (Kuznetsov and Schileyko 1997).

Shivapuri-Nagarjun National Park-Nagarjun Forest (Khanal and Budha 2013).

Genus: *Girasia* J.E. Gray, 1855

Distribution: India: Assam, Arunachal Pradesh, Himachal Pradesh, Meghalaya, Mizoram, Manipur, Nagaland, Sikkim; Myanmar (Blanford and Godwin-Austen 1908, Ramakrishna et al. 2010).

Type species: *Girasia hookeri* J.E. Gray, 1855

Girasia sp.

New species record for Nepal: Langtang National Park.

Family: Ariophantidae Godwin-Austen, 1888

Subfamily: Macrochlamydinae Godwin-Austen, 1888

Genus: *Macrochlamys* Benson in Godwin-Austen, 1883⁴⁵

Distribution: S and SE Asia (Blanford and Godwin-Austen 1908).

Type species: *Macrochlamys indica* Benson in Godwin-Austen, 1883

Macrochlamys indica Benson in Godwin-Austen, 1883

Distribution: India; Andaman Islands; Bangladesh; Sri Lanka (Ramakrishna et al. 2010).

Nepal: Ilam, Sunsari, Dharan, Kathmandu, Lalitpur, Gulmi, Kaski Districts (Subba and Ghosh 2001).

New distribution records from Nepal: Dadeldhura, Baitadi, and Kanchanpur Districts.

Macrochlamys lata Godwin-Austen, 1888

Distribution: India: Meghalaya (Ramakrishna et al. 2010).

Nepal: Annapurna Conservation Area (Kuznetsov and Schileyko 1997).

Macrochlamys longicauda Godwin-Austen, 1883

Distribution: India: Meghalaya (Ramakrishna et al. 2010).

Nepal: Kathmandu District, Annapurna Conservation Area (Kuznetsov and Schileyko 1997).

New distribution records from Nepal: Shivapuri-Nagarjun and Langtang National Parks.

Macrochlamys lubrica (Benson, 1852)

Distribution: India: W Bengal-Darjeeling, Sikkim, Meghalaya (Blanford and Godwin-Austen 1908, Ramakrishna et al. 2010).

Nepal: Mid hills of several districts of E Nepal (Subba and Ghosh 2008).

Macrochlamys nuda (L. Pfeiffer, 1852)

Distribution: NW India: Himachal Pradesh-Simla, Uttarakhand-Kumaon (Blanford and Godwin-Austen 1908).

Nepal: Annapurna Conservation Area (Schileyko and Kuznetsov 1996, 1998b).

Macrochlamys patane (Benson, 1859)

Distribution: NE India: W Bengal-Darjeeling, Sikkim (Blanford and Godwin-Austen 1908).

Nepal: Kathmandu District (Schileyko and Kuznetsov 1996).

Macrochlamys perpaula (Benson, 1859)

Distribution: India: Bihar, Jharkhand, Sikkim, W Bengal-Darjeeling (Ramakrishna et al. 2010).

Nepal: Shivapuri-Nagarjun National Park-Nagarjun Forest (Khanal and Budha 2013).

Macrochlamys sathilaensis Godwin-Austen, 1907

Distribution: NE India: Sikkim-Richila Peak; Bhutan (Blanford and Godwin-Austen 1908).

Nepal: Annapurna Conservation Area, Solukhumbu District (Kuznetsov and Schileyko 1997).

Macrochlamys sequax (Benson, 1859)

Distribution: India: W Bengal-Darjeeling (Blanford and Godwin-Austen 1908).

Nepal: Annapurna Conservation Area (Kuznetsov and Schileyko 1997).

Macrochlamys sequius Godwin-Austen, 1907

Distribution: India: W Bengal-Darjeeling (Blanford and Godwin-Austen 1908).

Nepal: Annapurna Conservation Area (Kuznetsov and Schileyko 1997).

Macrochlamys subiecta (Benson, 1852)

Distribution: India: Jharkhand-Rajmahal Hills, Orrissa-Cuttak (Blanford and Godwin-Austen 1908).

Nepal: Annapurna Conservation Area (Kuznetsov and Schileyko 1997).

New distribution records from Nepal: Widely distributed in W Tarai to the mid hills of C Nepal.

Macrochlamys tugurium (Benson, 1852)⁴⁶

Distribution: India: Manipur, Sikkim, W Bengal-Darjeeling (Ramakrishna et al. 2010).

Nepal: Kathmandu District (Kiauta and Butot 1972).

New distribution record from Nepal: Kaptad National Park.

Genus: *Euaustenia* Cockerell, 1891⁴⁷

Distribution: Afghanistan; Pakistan; NW and NE India: Uttarakhand, Sikkim (Blanford and Godwin-Austen 1908).

Type species: *Vitrina scutella* Benson, 1859 (= *Vitrina monticola* L. Pfeiffer, 1849⁴⁸)

Euaustenia cassida (Hutton, 1838)

Distribution: NW India: Himachal Pradesh, Kashmir, Uttarakhand-Kumaon (Blanford and Godwin-Austen 1908, Ramakrishna et al. 2010).

New species records for Nepal: Baitadi, Darchula, and Dadeldhura Districts.

Euaustenia monticola (L. Pfeiffer, 1849)

Distribution: NW India: Kashmir, Uttarakhand-Nainital (Ramakrishna et al. 2010).

Nepal: Kathmandu District (Schileyko and Frank 1994, Kuznetsov 1996), Annapurna Conservation Area (Kuznetsov and Schileyko 1997).

New distribution records from Nepal: Shivapuri-Nagarjun and Langtang National Parks.

Genus: *Bensonies* H.B. Baker, 1938

Distribution: Afghanistan; Pakistan; India: Uttarakhand, Sikkim (Blanford and Godwin-Austen 1908).

Type species: *Nanina monticola* Hutton, 1838

Bensonies convexa (Reeve, 1852)

Distribution: India: Himachal Pradesh, Uttarakhand (Blanford and Godwin-Austen 1908).

Nepal: Annapurna Conservation Area (Kuznetsov and Schileyko 1997).

New distribution records from Nepal: Lalitpur District-Phulchowki Hill, Kathmandu District- Champadevi Hill, Shivapuri-Nagarjun and Langtang National Parks.

Bensonies jacquemonti (Martens, 1869)

Distribution: Pakistan: Murree; NW India: Himachal Pradesh, Kashmir, Punjab, Uttarakhand (Blanford and Godwin-Austen 1908, Ramakrishna et al. 2010).

New species record for Nepal: Baitadi District.

Bensonies monticola (Hutton, 1838)

Distribution: NW India: Kashmir, Punjab, Uttarakhand (Blanford and Godwin-Austen 1908, Ramakrishna et al. 2010).

New species record for Nepal: Khaptad National Park.

Bensonies nepalensis (W.T. Blanford, 1904)⁴⁹

Distribution: Endemic to Nepal, where it is common in Kathmandu Valley (Blanford 1904, Blanford and Godwin-Austen 1908).

New distribution records from Nepal: Lalitpur, Kavre, Chitwan, Kaski, Gulmi, Syangjha, Parbat, and Myagdi Districts.

Bensonies theobaldiana (Godwin-Austen, 1888)

Distribution: NW India: Himachal Pradesh-Simla, Uttarakhand (Blanford and Godwin-Austen 1908, Ramakrishna et al. 2010).

New species record for Nepal: Khaptad National Park.

Genus: *Himalodiscus* Kuznetsov, 1996

Distribution: Endemic to Nepal.

Nepal: C and W Nepal.

Type species: *Himalodiscus aculeatus* Kuznetsov, 1996

Himalodiscus aculeatus Kuznetsov, 1996⁵⁰

Distribution: Endemic to Nepal.

Nepal: Lalitpur District-Phulchowki Hill (Kuznetsov 1996).

New distribution record from Nepal: Shivapuri-Nagarjun National Park.

Himalodiscus echinatus Schileyko & Kuznetsov, 1998

Distribution: Endemic to Nepal.

Nepal: Annapurna Conservation Area. Only reported from the type locality Lete (Schileyko and Kuznetsov 1998b).

Genus: *Khasiella* Godwin-Austen, 1899

Distribution: E Himalaya from Nepal and India to Myanmar (Blanford and Godwin-Austen 1908).

Type species: *Helix vidua* Hanley & Theobald, 1875⁵¹

Khasiella ornatissima (Benson, 1859)

Distribution: India: W Bengal, Sikkim (Blanford and Godwin-Austen 1908), Uttar Pradesh (Ramakrishna et al. 2010).

Nepal: Lalitpur District-Phulchowki Hill (Kuznetsov 1996).
 New distribution records from Nepal: Chitwan National Park, Chitwan and Nawalparasi Districts.

Khasiella pansa (Benson, 1856)⁵²

Distribution: Myanmar: Ayeyarwady Valley, Sullivan Island, Mergui Archipelago (Blanford and Godwin-Austen 1908).

Nepal: Ilam, Jhapa, Morang, Sunsari, Dharan, Saptari, Udayapur, Kaski, Rupandehi and Kailali Districts (Subba and Ghosh 2001, Subba 2003, Surana et al. 2004).

Genus: *Oxytesta* Zilch, 1956

Distribution: E Himalaya from Nepal and NE India to Myanmar and Laos (Blanford and Godwin-Austen 1908).

Type species: *Helix oxytes* Benson, 1836

Oxytesta blanfordi (Theobald, 1859)

Distribution: India: W Bengal-Darjeeling, Sikkim (Blanford and Godwin-Austen 1908, Ramakrishna et al. 2010).

Nepal: Mustang (Kuznetsov and Schileyko 1997, Schileyko and Kuznetsov 1998b).

New distribution records from Nepal: Rasuwa and Parbat Districts.

Oxytesta cycloplax (Benson, 1852)

Distribution: India: Sikkim (Ramakrishna et al. 2010).

Nepal: Solukhumbu District (Kuznetsov and Schileyko 1997).

New distribution record from Nepal: Sankhuwasabha District.

Oxytesta orobia (Benson, 1848)⁵³

Distribution: India: W Bengal-Darjeeling (Ramakrishna et al. 2010).

Nepal: Hills surrounding Kathmandu Valley (Schileyko and Frank 1994, Kuznetsov and Schileyko 1997).

New distribution records from Nepal: Shivapuri-Nagarjun and Langtang National Parks, Sankhuwasabha District.

Oxytesta sylvicola (W.T. Blanford, 1880)

Distribution: NE India: Assam-Burail range, North Cachar, Nagaland (Blanford and Godwin-Austen 1908, Ramakrishna et al. 2010).

Nepal: Ilam, Morang, Dharan, Udayapur, Kaski, Kathmandu, Lalitpur and Terhathum Districts (Subba and Ghosh 2001).

Genus: *Rotungia* Godwin-Austen, 1918

Distribution: India: Arunachal Pradesh-Abor Hill; Myanmar-Upper Rotung (Ramakrishna et al. 2010).

Type species: *Rotungia williamsoni* Godwin-Austen, 1918

Rotungia williamsoni Godwin-Austen, 1918

Distribution: India: Arunachal Pradesh-Abor Hill (Ramakrishna et al. 2010)
Nepal: Taplejung and Terhathum Districts (Subba and Ghosh 2008).

Genus: *Syama* Blanford & Godwin-Austen, 1908

Distribution: India (Blanford and Godwin-Austen 1908).
Type species: *Nanina (Macrochlamys) prona* Nevill, 1878.

Syama prona (Nevill, 1878)

Distribution: NW India: Himachal Pradesh, Uttarakhand (Blanford and Godwin-Austen 1908, Ramakrishna et al. 2010).
Nepal: Shivapuri-Nagarjun National Park-Nagarjun Forest (Khanal and Budha 2013).

Subspecies: *prona* (Nevill, 1878)

Distribution: Annapurna Conservation Area (Kuznetsov and Schileyko 1997, Schileyko and Kuznetsov 1998b).

Genus: *Rasama* Laidlaw, 1932⁵⁴

Distribution: NE India; W Bhutan (Blanford and Godwin-Austen 1908).
Type species: *Macrochlamys kala* Godwin-Austen, 1883

Rasama kala (Godwin-Austen, 1883)

Distribution: India: Sikkim-Damsang Peak, Dalling Hills; W Bhutan (Blanford and Godwin-Austen 1908).
New species record for Nepal: Ilam District-Maipokhari.

Genus: *Taphrospira* W.T. Blanford, 1905

Distribution: India: Assam; Andaman Islands; Myanmar (Blanford and Godwin-Austen 1908).

Type species: *Helix convallata* Benson, 1856

Taphrospira compluvialis (W.T. Blanford, 1865)

Distribution: India: Assam; Andaman Islands; Myanmar (Blanford and Godwin-Austen 1908).

Nepal: Panchthar, Taplejung and Terhathum Districts (Subba and Ghosh 2008).

Taphrospira convallata (Benson, 1856)

Distribution: Myanmar (Blanford and Godwin-Austen 1908).
New species record for Nepal: Shivapuri-Nagarjun National Park.

Superfamily: Limacoidea Lamarck, 1801

Family: Limacidae Lamarck, 1801

Subfamily: Limicinae Lamarck, 1801

Distribution: W Palearctic region (Wiktor and Bössneck 2004).

Genus: *Limax* Linnaeus, 1758

Distribution: Palearctic region (Wiktor and Bössneck 2004).

Type species: *Limax maximus* Linnaeus, 1758

Limax seticus Wiktor & Bössneck, 2004

Distribution: This is the only *Limax* species recorded from the Himalaya (Wiktor and Bössneck 2004).

Nepal: Endemic to Nepal; probably the slug species with the highest elevation range (up to 5000 m) in the world. This species was reported only from Bajura District.

Genus: *Turcomilax* Simroth, 1901

Distribution: India and Nepal (Wiktor et al. 1999, Bössneck 2006).

Type species: *Gigantomilax (Turcomilax) nanus* Simroth, 1901

Subgenus: *Kasperia* Godwin-Austen, 1914⁵⁵

Distribution: India: Kashmir (Godwin-Austen, 1914).

Type species: *Limax (Kasperia) mayae* Godwin-Austen, 1914 (= *Limax turkestanus* Simroth, 1898)

Turcomilax (Kasperia) oli Wiktor, Naggs & Gupta, 1999⁵⁶

Distribution: India: Kumaun Himalaya (Wiktor et al. 1999).

Nepal: Darchula District (Bössneck 2006).

Family: Agriolimacidae Wagner, 1935

Subfamily: Agriolimicinae Wagner, 1935

Genus: *Deroceras* Rafinesque, 1820

Distribution: Holarctic. From Sahara to NE America and S Asia (Wiktor et al. 2000, Thompson 2011).

Type species: *Limax laevis* O.F. Müller, 1774

Deroceras laeve (O.F. Müller, 1774)

Distribution: Holarctic. From Sahara to NE America. It has been introduced worldwide (Wiktor et al. 2000).

Nepal: Kathmandu, Taplejung and Panchthar Districts (Bössneck 2006).

New distribution record from Nepal: Lalitpur District.

Superfamily: Arionoidea J.E. Gray, 1840

Family: Anadenidae Pilsbry, 1948

Genus: *Anadenus* Heynemann, 1863

Distribution: S China; southern slopes of the Himalaya from Pakistan eastward to Sikkim (Wiktor 2001a).

Type species: *Anadenus giganteus* Heynemann, 1863 [Currently replaced by: *Limax altivagus* Theobald, 1862]⁵⁷

Anadenus altivagus (Theobald, 1862)⁵⁸

Distribution: Southern slopes of the Himalaya from Rawalpindi in the west of N Pakistan through Kashmir and Nepal to Sikkim in NE India (Wiktor 2001a). Nepal: Bajura, Darchula, Humla and Rasuwa Districts (Bössneck 2006).

New distribution records from Nepal: Langtang National Park-Dhunche-Gosainkund-Chisapani trekking route.

Anadenus nepalensis Wiktor, 2001

Distribution: Endemic to Nepal.

Nepal: Hills of Darchula, Dolpa, Humla, Jumla, Lamjung, Kaski, Palpa and Kathmandu Districts (Wiktor 2001a, Bössneck 2006).

Subgenus: *Sagarmathia* Kuzminikh & Schileyko, 2005

Distribution: Endemic to Nepal (Kuzminikh and Schileyko 2005).

Type species: *Anadenus (Sagarmathia) kuznetsovi* Kuzminikh & Schileyko, 2005

Anadenus (Sagarmathia) kuznetsovi Kuzminikh & Schileyko, 2005

Distribution: Endemic to Nepal.

Nepal: Only reported from the type locality, Phuiyan Khola, Solukhumbu District (Kuzminikh and Schileyko 2005).

Family: Philomycidae J.E. Gray, 1847

Genus: *Meghimatium* van Hasselt, 1823

Distribution: Russia; China; Korea; Japan; Borneo; Sumatra; Java; Celebes; Philippines (Wiktor and Jurkowska 2007).

Type species: *Meghimatium striatum* van Hasselt, 1823

Meghimatium cf. pictum (Stoliczka, 1873)⁵⁹

Distribution: China; India (Wiktor et al. 2000).

Nepal: Chitwan National Park (Bössneck 2006).

Superfamily: Helicoidea Rafinesque, 1815

Family: Bradybaenidae Pilsbry, 1934

Subfamily: Bradybaeninae Pilsbry, 1934

Genus: *Bradybaena* Beck, 1837

Distribution: India; Bangladesh; Nepal; China; Myanmar; Thailand; Laos; Vietnam; Cambodia; Indonesia; Malaysia; Singapore (Hoong 1995, Panha 1995–1996, Wu 2002, 2004, Kuznetsov and Schileyko 1997, Ramakrishna et al. 2010, Schileyko 2011); the type species, *B. similaris* is widely introduced in other regions (Carvalho et al. 2008).

Type species: *Helix similaris* Féüssac, 1821

Bradybaena radicicola (Benson, 1848)

Distribution: NW to NE India: Himachal Pradesh, Uttarakhand, Sikkim (Ramakrishna et al. 2010).

Nepal: Annapurna Conservation Area-Kokhethanti, Lete Khola (Kuznetsov and Schileyko 1997, Schileyko and Kuznetsov 1998a, 1998b).

Bradybaena ? thakkholensis Schileyko & Kuznetsov, 1998a⁶⁰

Distribution: Endemic to Nepal.

Nepal: Annapurna Conservation Area. Only known from Thakkhola, the type locality (Schileyko and Kuznetsov 1998a).

Genus: *Plectotropis* Martens, 1860

Distribution: India; China; Japan; Sumatra (Schileyko 2004, Ramakrishna et al. 2010).

Type species: *Helix elegantissima* L. Pfeiffer, 1849

Plectotropis tapeina (Benson, 1836)⁶¹

Distribution: India; Bangladesh; Myanmar (Ramakrishna et al. 2010).

Nepal: Ilam and Panchthar Districts (Subba and Ghosh 2008).

Family: Camaenidae Pilsbry, 1895

Subfamily: Camaeninae Pilsbry, 1895

Genus *Landouria* Godwin-Austen, 1918

Distribution: Sri Lanka; NE India; Nepal; Indonesia; Philippines (Schileyko and Kuznetsov 1998a).

Type species: *Helix buttonii* L. Pfeiffer, 1842⁶²

Landouria aborensis Godwin-Austen, 1918

Distribution: India: Arunachal Pradesh-Abor Hill (Ramakrishna et al. 2010).

Nepal: Dolakha, Lalitpur, Ramechhap and Solukhumbu Districts (Kuznetsov and Schileyko 1997, Schileyko and Kuznetsov 1998a).

Landouria coeni (Preston, 1914)⁶³

Distribution: India: Nagaland (Gude 1914).
Nepal: Solukhumbu District (Schileyko and Kuznetsov 1998a).

Landouria dhaulagirica Schileyko & Kuznetsov, 1998a

Distribution: Endemic to Nepal.
Nepal: Annapurna Conservation Area-Larjung, Kokhethanti, Kalopani (Schileyko and Kuznetsov 1998a).

Landouria buttonii (L. Pfeiffer, 1842)

Distribution: India: Himachal Pradesh, Uttarakhand, W Bengal, Assam, Nagaland (Ramakrishna et al. 2010).
Nepal: Kaski and Myagdi Districts (Kuznetsov and Schileyko 1997, Schileyko and Kuznetsov 1998a).

Landouria rhododendronis Schileyko & Kuznetsov, 1998a

Distribution: Endemic to Nepal.
Nepal: Annapurna Conservation Area-Gorepani, Parbat District (Schileyko and Kuznetsov 1998a).

Landouria savadiensis (Nevill, 1877)

Distribution: Myanmar: Sawady (Nevill, 1877).
Nepal: Shivapuri-Nagarjun National Park-Nagarjun Forest, Tare-Bhir (Schileyko and Kuznetsov 1998a).

Genus: *Ganesella* W.T. Blanford, 1863

Distribution: India; Myanmar; Thailand; Cambodia (Ramakrishna et al. 2010).
Type species: *Helix capitum* Benson, 1848

Ganesella sp.

Nepal: Shivapuri-Nagarjun National Park (Khanal and Budha 2013).
New distribution record from Nepal: Lalitpur District-Phulchowki Hill.

Notes

¹ Eight subgenera of *Cyclophorus* were recognized by Kobelt (1902), one of which, the African *Maizania* Bourguignat, 1889, was elevated to family level Maizaniidae by Tielecke (1940) (see Bouchet and Rocroi 2005: 248). Gude (1921) mentioned only five subgenera viz.: *Glossostylus* Kobelt and Möllendorff, 1897 (S and SE Asia), *Litostylus* Kobelt & Möllendorff, 1897 (S and SE Asia), *Salpingophorus* Kobelt and Möllendorff, 1897 (S and SE Asia), *Cyclophorus* Montfort, 1810 (S and SE Asia) and *Cyclohelix* Mörch, 1852 (Andaman and Nicobar Islands). Wenz

(1939: 458–460) replaced *Salpingophorus* by *Annularia* Schumacher, 1817 and *Cyclohelix* by *Otopoma* Gray, 1850, while Egorov (2006) replaced *Litostylus* by *Kobeltostylus* Egorov, 2006. In this list we follow Gude (1921) but with the adapted names proposed by Wenz (1939) and Egorov (2006). Note that Egorov and Greke (2007) also recognized five subgenera, but they regarded *Salpingophorus* as a junior synonym of *Cyclophorus*, while maintaining *Cricophorus* Kobelt and Möllendorff, 1897 as separate subgenus, next to *Cyclophorus*, *Glossostylus*, *Cyclohelix* and *Kobeltostylus*.

- ² Allozyme data of *Cyclophorus fulgoratus* populations in Thailand suggest that this is a species complex (Prasankok et al. 2009). It remains to be investigated how the Nepalese populations fit into this picture.
- ³ The name *Litostylus* Kobelt & Möllendorff, 1897 is a junior homonym of *Litostylus* Faust, 1893 (= Insecta, Coleoptera, Curculionidae). Egorov (2006) therefore replaced the molluscan name by *Kobeltostylus*.
- ⁴ Wenz (1939) replaced the name *Salpingophorus* Kobelt & Möllendorff, 1897 by *Annularia* Schumacher, 1817, while Egorov and Greke (2007) regarded both these names as junior synonyms of *Cyclophorus* Montfort, 1810 (see note ¹).
- ⁵ *Cyclophorus (A.) aurantiacus* is distributed in SE Asia, so that its presence in Nepal is doubtful and requires confirmation.
- ⁶ Nevill (1878) regarded *Scabrina* W.T. Blanford, 1863 as a subgenus of *Cyclophorus* Montfort, 1810. Kobelt and Möllendorff (1897) raised *Scabrina* to genus rank. *Cyclophorus pinnulifer* Benson, 1857 was fixed as the type species of *Scabrina* by Nevill (1878).
- ⁷ Some authors have erroneously attributed the Alycaeinae to Gray, 1850 (Minato 2005, Tarruella and Domènec 2011). However, according to Bouchet and Rocroi (2005) the correct authorship is ‘W. Blanford, 1864’. Alycaeinae comprises four genera (Tarruella and Domènec 2011) namely *Alycaeus* J.E. Gray, 1850, *Chamalycaeus* Kobelt & Möllendorff, 1897, *Cipangocharax* Shintaro, 1934 and *Dioryx* Benson, 1859. The generic names *Alycaeus* and *Chamalycaeus* have been applied by recent authors (Panha and Burch 2005, Maassen 2006, Dumrongrojwattana and Maassen 2008, Lee et al. 2008, Tarruella and Domènec 2011). Conversely, *Cycloryx* Godwin-Austen, 1914, *Dicharax* Kobelt & Möllendorff, 1900 and *Raptomphalus* Godwin-Austen, 1914 are treated as subgenera of *Chamalycaeus* by Gude (1921). We apply Gude’s (1921) generic categories.
- ⁸ The name of the type species of *Alycaeus* J.E. Gray, 1850, by original designation *Cyclostoma gibbum* Féussac, 1838, is a junior homonym of *Cyclostoma gibbum* Draparnaud, 1805 (Hydrobiidae). Therefore, it has been replaced by *Alycaeus eydouxi* Venmans, 1956.
- ⁹ Some authors attribute authorship of *Chamalycaeus (D.) strangulatus* to Hutton such as Pfeiffer (1846), Nevill (1878), Hanley and Theobald (1878) but without indicating the publication year. Hutton’s name was, however, a manuscript name of no nomenclatural standing. Gude (1921) and Tarruella and Domènec (2011) were correct in assigning authorship to ‘Pfeiffer’.

- ¹⁰ Seven subgenera have been recognized within *Diplommatina* (Kobelt 1902, Kuroda 1928): *Benigoma* Kuroda, 1928, *Diplommatina* Benson, 1849, *Diploptychia* Möllendorff, 1895, *Metadiancta* Möllendorff, 1898, *Moussonia* Semper, 1865, *Pseudopalaina* Möllendorff, 1898, and *Sinica* Möllendorff, 1885.
- ¹¹ *Diplommatina (S.) canarica* is endemic to the Western Ghats (Raheem et al. 2014). Hence, the identification of the Nepalese specimens by Kuznetsov and Schileyko (1997) needs to be verified.
- ¹² In the original description of *Schistoloma funiculatum* Benson (in Hutton and Benson 1838) distinguished this species from the European fossil species *C. mumia* by ‘the rounder and more reflected orange peristome, and by its central position at the base, as well as by the delicate sculpture, and an embossed spiral cord which winds from above the umbilicus to the base, whence the species has received the trivial appellation of ‘Funiculatum’. It is the first known Indian species belonging to pupaeform or subcylindric division of *Cyclostoma*'. Sowerby (1850) changed the spelling ‘*funiculatum*’ to ‘*funiculatum*’ without giving any reason. Gude (1921) considered Benson’s (1838) ‘*funiculatum*’ to be a nomen nudum because of the spelling error (*funiculatum*, laps). Many authors have used ‘*funiculatum*’ (e.g. Gray in Baird 1850, Pfeiffer 1853, Hanley and Theobald 1870, Nevill 1878, Kobelt and Möllendorff 1897, Gude 1921, Ramakrishna et al. 2010). This was, however, an unjustified emendation (ICZN Article 32.2, 32.3) and thus Benson’s name stands. Nepalese specimens differ in possessing a whitish, instead of orange peristome, but the significance of this is unknown.
- ¹³ *Carychium minusculum* Gredler, 1888 was originally described from China, “aus Hupe” (type locality), which is the Chinese Province Hubei (= Hupeh) (See Zilch 1974). The correct publication year is ‘1888’ instead of ‘1887’ as is sometimes mentioned (e.g. Zilch 1974, Nesemann et al. 2007). In Nepal, this species was recorded from two different localities, viz. Kavre District by Nesemann et al. (2007) and Langtang National Park-Syabru by Kuznetsov and Schileyko (1997). PB checked specimens from these two localities (Nesemann’s specimen and an image of Kuznetsov and Schileyko 1997). The shells from these two localities differ by size, shape and sculpture and may be two distinct taxa.
- ¹⁴ *Carychium* shells collected by PB from Phulchowki, Shivapuri-Nagarjun and Langtang National Parks have very fine and strong radial ribs, as well as slender apical whorls. As such they differ from the shells of Nesemann et al. (2007), which are comparatively smooth and smaller. They are therefore, tentatively regarded as separate taxa. The Phulchowki specimen was also compared with images of Schileyko’s *C. minusculum* specimen from Langtang National Park deposited in ZMMU No. Lc-39251 and *C. minusculum* in Zilch (1974: Fig. 13). The peristome along the umbilicus region is more or less straight in the Phulchowki taxon, while it is strongly reflected in Schileyko’s *minusculum*.
- ¹⁵ According to Kennard (1942) the name *Vaginulus alte* was published in 1822, instead of 1821 as is often mentioned in the literature or 1823 as mentioned in Sherborn (1923: 230). The spelling ‘*altae*’ in e.g. Bössneck (2006) and Raheem et al. (2010) is erroneous.

- ¹⁶ The type locality of *Pupilla annandalei* Pilsbry, 1921 was doubtfully recorded as Ava (Myanmar) in the Indian Museum (Nevill 1878). Pilsbry (1920-21) speculated that the holotype may have been collected in Nepal because he associated it with central Asian species. However, confirmed records are restricted to granite mountains between 2,000 and 2,800 m in northern Pakistan (Pokryszko et al. 2009).
- ¹⁷ Pilsbry (1920-21: 204) asserted that *Pupa eurina* Benson, 1864 was collected in Nepal. If correct, Benson's record would be the earliest scientific report of a land snail from Nepal (Budha 2005). Benson (1864: 139) gave the locality in Latin as '*ad Tribeni Ghát fluminis Gogra*', but we have not been able to identify this locality, since 'Tribeni' refers to several localities where two rivers meet and 'ghat' refers either to a place where cremations take place or to sites where people cross a river along a trail by using locally made wooden boats. Godwin-Austen (1899: 260) expanded on Benson's locality information 'the typical specimens were found in the exuviae of the River Gogra at Tribeni Ghat. This river rises in the Tibetan plateau, and these shells may have been brought down thus from far back in the mountain range'. The downstream course of the Karnali river in Nepal is known as Gogra (= Ghaghara) in India immediately after the two branches of Karnali river meet at the Nepal-India border at Katarniya ghat, Uttar Pradesh, India. The confluence of Seti and Karnali river is called Tribeni which is approx. 100 km upstream (north) from the Nepal-India border. There is no clear evidence that William Theobald ever entered Nepal. However, Joseph Hooker, who was among the earliest Europeans to venture into Nepal to investigate its biota, did spend time with Theobald in India (Hooker 1854: 37, 57) and so it is possible that Hooker passed on samples of *Pupa eurina* to Theobald.
- ¹⁸ Hutton and Benson (1838) attributed the authorship of *Helix humilis* to Hutton, but Gude (1914) and Ramakrishna et al. (2010) incorrectly cite 'Benson' while Sherborn (1927: 3062) cited 'Hutton & Benson', 1838 as authors.
- ¹⁹ *Pyramidula kuznetsovi* was misidentified as *P. humilis* by Schileyko and Kuznetsov (1997). Kuznetsov's collections were recently reviewed and Schileyko and Balashov (2012) redescribed the samples as a new species.
- ²⁰ The correct spelling is 'cantorii' and not 'cantori' as some authors mention (e.g. Zilch 1959, Schileyko 1998).
- ²¹ The type locality of *Mirus nilagiricus* (L. Pfeiffer, 1846) is Nilgiris, South India. Although Kuznetsov and Schileyko (1997) reported this species from Nepal, they question whether the Himalayan species belongs to *Mirus*, though without suggesting an alternative generic placement.
- ²² Shortly after the description of *Darwininitium shiwalikianum*, Dr. Somsak Panha communicated that he and Dr. Chirasak Sutcharit (both Chulalongkorn University Bangkok, Thailand) noticed the conchological similarity between this species and *Helix capitulum* Benson, 1848, type species of the camaenid genus *Ganesella* W.T. Blanford, 1863. Further anatomical and DNA studies are needed to verify whether *D. shiwalikianum* and *G. capitulum* are conspecific. Moreover, the family level affiliations of *Darwininitium* and *Ganesella* remain to be assessed since the

Camaenidae may not be monophyletic (e.g. Scott 1996) and the phylogenetic relationships of the Camaenidae are still poorly resolved (Wade et al. 2007). If *D. shiwalikianum* is related or identical to *G. capitulum*, then it does not represent a pseudosigmurethrous orthurethran condition as was originally claimed by Budha et al. (2012). It would also mean that *Darwininitium* Budha & Mordan, 2012 will be a junior synonym of *Ganesella* W.T. Blanford, 1863.

- ²³ Nordsieck (1973) assigned Nepalese Phaedusinae to the genus *Hemiphaedusa* and this was followed by Raheem et al. (2010). Later, Nordsieck (2002) described the new subgenus *Montiphaedusa* Nordsieck, 2002 of the genus *Cylindrophaedusa* and grouped all Himalayan clausiliids in *Montiphaedusa*.
- ²⁴ Although E African *Lissachatina* is distinguished from W African *Achatina* (Bequaert 1950, Mead 1995), both generic names have been applied to this species. This list follows Budha and Naggs (2008) and Raheem et al. (2010, 2014), who used *Lissachatina* at genus level for reasons further explained by Raheem et al. (2014).
- ²⁵ *Ceciliooides* is the name used in the original description but it has been variously spelled by different authors. Hermannsen (1846) emended it to *Caeciliooides*, which was followed by Pilsbry and Tryon (1908–1909) and Gude (1914). *Ceciliooides* has been placed on the official list of generic names (ICZN Opinion 335) and all other spellings are invalid.
- ²⁶ Only a single *Ceciliooides* shell was collected in Nepal (Baitadi District). It measures about 2 mm, has four whorls, and resembles *C. minuta*.
- ²⁷ Many subulinid genera, such as *Opeas*, *Beckianum*, *Leptopeas*, *Lamellaxis* and *Leptinaria* have been confusingly interpreted (Thompson 2011), even if they are conchologically relatively well-differentiated and anatomical data are available for several of them.
- ²⁸ Baker (1935) erected *Allopeas* as a subgenus of *Lamellaxis* Strebler & Pfeiffer, 1882.
- ²⁹ Schileyko and Kuznetsov (1997) identified a Nepalese specimen as *Allopeas mauritianum prestoni* (Sykes, 1898) from Annapurna Conservation Area. Sykes (1898) original combination was *Opeas prestoni*. Naggs and Raheem (2000) placed ‘*prestoni*’ under *Allopeas*. Pilsbry and Tryon (1906) placed *Opeas prestoni* Sykes, 1898 under *O. mauritianum* (Pfeiffer, 1852) as var. *prestoni*. Brodie and Barker (2011) and Bouchet and Closel (1991) also placed ‘*mauritianum*’ under *Opeas*. Some authors assign ‘*prestoni*’ to *Lamellaxis* (e.g. Deisler and Abbott 1984, Nekola 2014). Griffiths and Florens (2006) suggested that *Allopeas mauritianum* is a junior synonym of *A. clavulinum*. The type locality of this species is Mauritius. FN examined the syntype of *mauritianus* and confirmed that it is identical with material identified as *Allopeas clavulinum*. It has been spread by commerce throughout the tropics but its native range is not known (Hanna 1966, Deisler and Abbott 1984).
- ³⁰ As for subulinids in general, *Opeas* species have been confusingly interpreted and have been assigned variously to different genera such as *Allopeas*, *Lamellaxis*, *Paropeas* and *Prosopeas* (although this latter may not even be a subulinid) (Naggs 1994).
- ³¹ We follow Bouchet and Rocroi (2005) and regard Glessuliniae as a subfamily of the Subulinidae.

- ³² The relationships of *Bacillum* are still unclear. Schileyko (1999) placed the genus in the Rishetiinae Schileyko, 1999, together with *Eutomopeas* Pilsbry, 1946, *Tortaxis* Pilsbry, 1906 and *Rishetia* Godwin-Austen, 1920. Based on the half exposed reproductive parts of a specimen labeled as *Bacillum* sp. Godwin-Austen (1920: 7) states ‘The very recent and extended knowledge of the animals of *Bacillum* and *Glessula* shows that the two genera come next to each other....’ (Godwin-Austen (1920: 7). But the same specimen (NHM) from a lot of 3 specimens from Assam, leg. S.L. Hora, Godwin-Austen coll. (Acc. 1830), Reg. 20120113) examined by PB confirmed that it is closer to *Rishetia* than to *Glessula* since it has an elongated flagellum. Because of its truncated columella, and elongately turreted shell, we provisionally retain *Bacillum* in the Glessulinae.
- ³³ Subba and Ghosh (2008) recorded *Bacillum* sp. from E Nepal without a description or figure.
- ³⁴ Although Martens (1860) designated *Cionella gemma* Benson, 1850 as the type species of *Glessula* (e.g. Zilch 1959), the correct type species is *Achatina ceylanica* L. Pfeiffer, 1845 (Gude 1914, Raheem et al. 2014). This is because *A. ceylanica* is the type species (by monotypy) of the genus *Electra* Albers, 1850, which is a junior homonym of *Electra* Lamouroux, 1816 (Ectoprocta). Therefore *Electra* Albers, 1850 was replaced by *Glessula* Martens, 1860 and in such cases ICBN Art. 67.8 rules that the type species of the replaced genus name is automatically also the type species of the new genus name.
- ³⁵ The type locality of *Glessula subjerdoni* is S India: Jaypore and Golconda Hills (Beddome 1906). Specimens from NE India (Darjeeling) were erroneously identified as *G. subjerdoni* by Gude (1914) and were subsequently assigned to *G. crassula* (Reeve, 1850) by Godwin-Austen (1920). Nevertheless, later authors (Kuznetsov 1996, Dey and Mitra 2000, Ramakrishna et al. 2010) have followed Gude (1914). Raheem et al. (2014) consider *G. subjerdoni* to be a ‘nomen dubium’.
- ³⁶ The distribution range of *Rishetia* in this list is based on unpublished anatomical data of specimens from Nepal and Sri Lanka. For example, Dinarzarde Raheem’s unpublished figures of dissected specimen of *Glessula capillacea* (L. Pfeiffer, 1855) from Sri Lanka indicate that it belongs to *Rishetia* because it has an elongated flagellum typical of *Rishetia*.
- ³⁷ Specimens of *Rishetia tenuispira* from Nepal were first described under the genus name *Ranibania* Schileyko & Kuznetsov, 1996. *Ranibania* was subsequently synonymized with *Rishetia* (Schileyko, 1999). However, Schileyko’s (1999) *Rishetia tenuispira* (Benson) from Nepal differs from Benson’s *R. tenuispira* from the type locality, Khasi Hills NE India and is similar to Godwin-Austen’s *Rishetia longispira* Godwin-Austen, 1920. Khanal and Budha (2013) identified specimens of the same locality as Schileyko (1999; Balaju, Raniban, Nepal) as *R. cf. longispira*. Godwin-Austen (1920) gave a very confusing and conflicting account on *longispira* and *tenuispira* (p. 33 same animal characters including reproductive anatomy) but the distribution range of *longispira* was recorded as westward from Bhutan to Sikkim and Darjeeling, whereas *tenuispira* was recorded from the Khasi and Garo Hills (p. 11–12).

- ³⁸ *Gulella bicolor* was originally described as *Pupa bicolor* Hutton, 1834 but Blanford and Godwin-Austen (1908) assigned it to *Ennea* H. Adams & A. Adams, 1855. The species has also been included in the Indo-Chinese streptaxid genus *Sinoennea* Kobelt, 1904. DNA sequence data, however, suggest that *Pupa bicolor* comes within *Gulella* (Rowson et al. 2011).
- ³⁹ The record of *E. plectosoma* (Benson, 1836) from Pegu (=Bamo, Myanmar) (Gude 1914: 81) is probably erroneous (Pál-Gergely et al. unpublished manuscript).
- ⁴⁰ Kuznetsov and Schileyko (1997) recorded *E. affinis* from Swoyambhunath temple forest area, but the material from this area may be a different species (Pál-Gergely et al. unpublished manuscript).
- ⁴¹ Some authors ‘1852’ as the publication year of *Kaliella barrakporensis* (e.g. Godwin-Austen 1882, Blanford and Godwin-Austen 1908). Raheem et al. (2014) pointed out that part 20 p. 156 of the Proceedings of the Zoological Society of London was published in 1854, 1852 being the date when the proceedings were presented at the Society’s meetings (see Duncan 1937). Therefore this publication was preceded by Pfeiffer’s ‘*Helix barrakporensis* 1853 Monographia Heliceorum Viventium 3: 59’. So the correct publication year is 1853, not 1852.
- ⁴² According to Falkner et al. (2002) *Euconulus fulvus* is a species complex.
- ⁴³ The genus *Hawaiia* is assigned to the Vitrinidae by Vaught (1989), to the Zonitidae by Riedel (1980) and to the Pristilomatidae by Anderson (2005). This later placement is followed in this list.
- ⁴⁴ Cockerell (1891, 1893) published the name as ‘*Cryptausteniae*’ (plural), while in 1898 he corrected it to ‘*Cryptaustenia*’ (singular). However, according to Art 11.8 and 33.2.2 of ICZN, the publication date of the corrected name remains ‘1891’.
- ⁴⁵ There is still much nomenclatural and taxonomic confusion with respect to the genus *Macrochlamys* and its type species. This list follows Raheem et al. (2014) in applying the current genus-level interpretation of *Macrochlamys* sensu Godwin-Austen (1883) with *Macrochlamys indica* Benson in Godwin-Austen, 1883 as its type species.
- ⁴⁶ According to Kiauta and Butot (1973) *Macrochlamys tugurium* would be the most common land gastropod of Kathmandu Valley but so far PB has not recorded *M. tugurium* in this area. The most common land gastropod in the Kathmandu Valley is *Bensonies nepalensis*, because of its similar shell shape and size, may have been misidentified as *M. tugurium*.
- ⁴⁷ Cockerell (1891, 1893) published the name as ‘*Euausteniae*’ (plural), while in 1898 he corrected it to ‘*Euaustenia*’ (singular). According to Art. 11.8 and 33.2.2 of the ICZN, the publication date of the corrected name remains ‘1891’.
- ⁴⁸ The publication date of *Vitrina monticola* L. Pfeiffer is ‘1849’ not ‘1848’ as cited by some authors mention (e.g. Blanford and Godwin-Austen 1908, Schileyko 2003, Mitra et al. 2005, Ramakrishna et al. 2010). See Duncan (1937) for the publication date of the *Proceedings of the Zoological Society of London* part 16:107); see also Sherborn (1928).

- ⁴⁹ *Bensonies nepalensis* shows a remarkable shell colour polymorphism that seems to correlate with altitude: at lower altitudes in C Nepal (Chitwan District) the body whorl of shells shows a dark brown band on a chocolate brown or white background. They co-occur with banded shells which are similar to mid hill specimens (PB, unpublished observations).
- ⁵⁰ *Himalodiscus aculeatus* was originally assigned to the Discidae by Kuznetsov (1996) based on conchological features, but based on anatomical data Schileyko and Kuznetsov (1998b) re-assigned it to Ariophantidae.
- ⁵¹ The type species '*Helix vidua*' has been confusingly cited. Zilch (1960) mentions '*Euplecta vidua* W.T. Blanford', but Schileyko (2002) mentions '*Euplecta vidua* Hanley and Theobald, 1875'. Godwin-Austen (1876) mentions '*Euplecta (Rotula) vidua* Blanford', whereas Blanford and Godwin-Austen (1908) list '*K. vidua* Blanford' in the same book under its species description as '*Khasiella vidua* H. & T. (Blf. MSS) (*Helix*)'. We follow Coan and Kabat (2012) in referring the type species to as *Helix vidua* Hanley & Theobald, 1875.
- ⁵² The identification of Nepalese *Khasiella pansa* needs to be verified.
- ⁵³ Schileyko and Frank (1994) and Kuznetsov and Schileyko (1997) reported *Oxytesta orobia* from the neighbourhood of Kathmandu, Nepal. PB checked the syntypes in NHM and specimens available at RBINS and compared these with Nepalese shells and concluded that the Nepalese specimens belong to a different species.
- ⁵⁴ *Sarama* Blanford and Godwin-Austen, 1908 is a junior homonym of *Sarama* Moore, 1887. *Saramina* Wenz, 1947 is a junior synonym of *Rasama* Laidlaw, 1932.
- ⁵⁵ The subgeneric name *Taulimax* Wiktor and Likharev, 1980 is a junior synonym of *Kasperia* Godwin-Austen, 1914 (Wiktor 2001b).
- ⁵⁶ Bössneck (2006) and Raheem et al. (2010) misspelled the genus name as *Turcolimax*.
- ⁵⁷ Although *Anadenus giganteus* Heynemann, 1863 is the type species of *Anadenus* Heynemann, 1863 (Zilch 1959, Wiktor et al. 2000), Wiktor (2001a) proposed to replace it by *Limax altivagus* Theobald, 1862, because he regarded *A. giganteus* Heynemann, 1863 as a 'nomen dubium'. For the time being, we nevertheless maintain *A. giganteus* as the type species as Simroth (1901) did provide anatomical data, including figures of *A. giganteus*.
- ⁵⁸ Three paratypes of *Anadenus nepalensis* from Langtang National Park in fact belong to *A. altivagus* viz. one specimen from 'Chandrabar (= Chandanbari), 3,300 m a.s.l., fir forest' and two specimens from 'Gosainkund, 4,200 m a.s.l.' both collected on 27.09.1981 by A. Kuska (see Wiktor 2001a) (A. Wiktor, pers.comm. 13.10.2009). This was confirmed by their reproductive anatomy (with its typical spines inside the penis) examined by PB.
- ⁵⁹ Wiktor et al. (2000) figured the reproductive anatomy of *Meghimatiumpictum* cf. *pictum* (Stoliczka, 1873) and *M. bilineatum* (Benson, 1842) based on Chinese specimens but found no clear differences and hence were undecided as to whether or not *M. cf. pictum* is a distinct species. The reproductive organs of a specimen from Nepal resemble those of Chinese *M. cf. pictum*.

- ⁶⁰ *Bradybaena (?) thakholensis* was described on the basis of a few juvenile shells by Schileyko and Kuznetsov (1998a), its anatomy is unknown.
- ⁶¹ Subba and Ghosh (2008) misspelled this species name as *Aegista (Placetotropis) tapeina*.
- ⁶² The correct spelling is ‘*buttonii*’ (Raheem et al. 2014; see Pfeiffer 1842: 82), not ‘*buttoni*’ as mentioned by e.g. Zilch (1960) and Schileyko (2004).
- ⁶³ *Landouria coeni* was placed in the subgenus *Plectotropis* of the genus *Aegista* by Ramakrishna et al. (2010). However, these two taxa were treated as distinct genera by Preston (1914), Gude (1914), and Schileyko (2004).

Acknowledgements

PB Budha received a faculty research grant from the University Grants Commission, Sanothimi Bhaktapur, Nepal. This work could not have been undertaken without the support of the Darwin Initiative programme *Developing land snail expertise in South and Southeast Asia*. PB gratefully acknowledges the Royal Belgian Institute of Natural Sciences (RBINS), Brussels, for financial support. We would like to thank the following colleagues for providing specimens and/or advice: Jonathan Ablett (Natural History Museum, London), Dinarzarde Raheem (RBINS), Alexander Sysoev (Zoological Museum of Moscow University, Moscow), Andrzej Wiktor (Museum of Natural History, Wroclaw University, Poland), Rose Sablon (RBINS) and Hasko Nesemann (Aquatic Ecology Centre, Kathmandu University, Nepal). We are grateful to Ramesh Devkota, Srijana Khanal, Sunita Khatiwara and Naresh Kohar for support in the field work. Sanjan Thapa, Hari P. Sharma, Bharat Babu Shrestha, Ripu Kunwar and Janak Khatiwada provided samples from Khaptad National Park, Mugu, Taplejung and Dhading Districts.

References

- Anderson R (2005) An annotated list of the non-marine Mollusca of Britain and Ireland. *Journal of Conchology* 38(6): 607–638.
- Azuma M (1982) Colored illustrations of the land snails of Japan. Hoikusha, Osaka.
- Baird W (1850) Nomenclature of Molluscous Animals and Shells in the collection of the British Museum Part 1. Cyclophoridae. Spottiswoodes and Shaw, London. doi: 10.5962/bhl.title.59531
- Baker HB (1935) Jamaican Land Snails. *The Nautilus* 48(3): 83–88
- Bartsch P (1915) The Philippine land shells of the genus *Schistoloma*. *Proceedings of the United States National Museum* 49(2104): 195–204. doi: 10.5479/si.00963801.49-2104.195
- Beddome RH (1906) Notes on Indian and Ceylonese species of *Glessula*. *Proceedings of the Malacological Society of London* 6: 160–172.
- Benson WH (1864) Characters of *Coilostele*, an undescribed genus of Auriculacea (?), and of species *Helix*, *Pupa*, and *Ancylus*, from India, West Africa, and Ceylon. *Annals and Magazine of Natural History* (3) 13: 136–140.

- Bentham Jutting WSS van (1961) The Malayan Streptaxidae Genera *Huttonella* and *Sinoennea*. Bulletin of the Raffles Museum, Singapore 26: 5–33.
- Bentham Jutting WSS van (1948) Systematic studies on the non-marine Mollusca of the Indo-Australian Archipelago I. Critical Revision of the Javanese Operculate Land-shells of the Families Hydrocenidae, Helicinidae, Cyclophoridae, Pupinidae and Cochlostomatidae. Treubia 19: 539–604.
- Bentham Jutting WSS van (1950) Systematic studies on the non-marine Mollusca of the Indo-Australian Archipelago II. Critical Revision of the Javanese Pulmonate Land-shells of the Families Helicaroniidae, Pleurodontidae, Fruticicolidae and Streptaxidae. Treubia 20: 381–505.
- Bequaert JC (1950) Studies in the Achatininae, a group of African land snails. Bulletin of the Museum of Comparative Zoology, Harvard 105: 1–216.
- Blanford WT (1904) Description of Indian and Burmese land shells referred to the genera *Macrochlamys*, *Bensonia* and *Taphrospira* (gen. nov.), *Microcystina*, *Euplecta* and *Polita*. Proceedings of the Zoological Society of London 2: 441–447.
- Blanford WT, Godwin-Austen HH (1908) The fauna of British India, including Ceylon and Burma. Mollusca, Testacellidae and Zonitidae. London, 311 pp.
- Bössneck U (2006) Ökologie und Verbreitung der Nacktschnecken im Nepal-Himalaya (Molluca: Gastropoda: Pulmonata). In: Hartmann M, Weipert J (Eds) Biodiversität und Naturausstattung im Himalaya. Bandung 2, 39–48.
- Bouchet P, Rocroi JP (2005) Classification and nomenclator of gastropod families. Malacologia 47: 1–397.
- Bouchet P, Cosel B von (1991) Les Mollusques terrestres et fluviatiles des Départements d'Outre-Mer. Muséum national d'Histoire naturelle, Paris, 109 pp.
- Brodie G, Barker GM (2011) Introduced land snails in the Fiji Islands: Are there risks involved? In: Veitch CR, Clout MN, Towns DR (Eds) Island Invasives: eradication and management. IUCN, Gland, Switzerland, 32–36.
- Bruggen AC van (2006) *Gulella johannae* spec. nov. (Gastropoda, Pulmonata, Streptaxidae), a new land snail from the Drakensberg range in Limpopo Province, South Africa, with notes on *G. johannesburgensis* (M. & P.). Zoologische Mededelingen 80(1): 63.
- Budha PB (2005) Nepalese malacology trails behind “Catch up”. Himalayan Journal of Sciences 3(5): 9–10.
- Budha PB, Mordan PB, Naggs F, Backeljau T (2012) *Darwininitium* – a new fully pseudosigmurethrous orthurethran genus from Nepal (Gastropoda, Pulmonata, Cerastidae). ZooKeys 175: 19–26. doi: 10.3897/zookeys.175.2755
- Budha PB, Naggs F (2008) The Giant African Land Snail *Lissachatina fulica* (Bowdich) in Nepal. The Malacologist 50: 19–21.
- Burch JB, Panha S (2002) The land snail genus *Carychium* (Pulmonata: Carychiidae) in Thailand. Walkerana 9(22): 171–175.
- Carvalho CDM, Bessa ECA, D'Ávila S (2008) Life history strategy of *Bradybaena similaris* (Férussac, 1821) (Mollusca: Pulmonata: Bradybaenidae). Molluscan Research 28(3): 171–174.
- Chaijirawong R, Nuamsee K, Dumrongrojwattana P (2008) Shell and radula morphology and reproductive anatomy of the introduced carnivorous snail *Gulella bicolor* (Pulmonata: Streptaxidae) from Chong Buri Province. Kasetsart Journal (Natural Science) 42: 251–255.

- Clench WJ (1964) *Gulella bicolor* (Hutton). The Nautilus 77: 142–143.
- Coan EV, Kabat AR (2012) The malacological works and taxa of Sylvanus Hanley (1819–1899). Malacologia 55(2): 285–359. doi: 10.4002/040.055.0208
- Cockerell TDA (1891) On the geographical distribution of slugs. Proceedings of the Zoological Society of London 59(2): 214–226. doi: 10.1111/j.1096-3642.1891.tb01744.x
- Cockerell TDA (1893) A checklist of the slugs with appendix and notes by Walter E Collinge. The Conchologist 2(7–8): 168–176, 185–232.
- Cole ML, Herbert DG (2009) Description of four new species of *Gulella* Pfeiffer, 1856 from Eastern Cape, South Africa, with additional notes on two poorly known species (Mollusca: Eupulmonata: Streptaxidae). Zoologische Mededelingen 83: 547–564.
- Deisler JE, Abbott RT (1984) Range extensions of some introduced land mollusks in the Bahama Islands, with first reports for four species. The Nautilus 98(1): 12–17.
- Dey A, Mitra SC (2000) Mollusks of the Himalaya. Records of the Zoological Survey of India 98(2): 5–50.
- Dumrongrojwattana P, Maassen WJM (2008) Two new species of *Alycaeus* from Eastern Thailand (Gastropoda, Caenogastropoda, Cyclophoridae). The Thailand Natural History Museum Journal 3(1): 1–4.
- Duncan FM (1937) On the dates of publication of the Society's 'Proceedings', 1859–1926. Proceedings of the Zoological Society of London 107: 71–84.
- Dundee DS, Baerwold RJ (1984) Observations on a micropredator, *Gulella bicolor* (Hutton) (Gastropoda: Pulmonata: Streptaxidae). The Nautilus 98: 63–68.
- Egorov RV (2006) *Kobeltostylus* nom. nov., a replacement name for *Litostylus* Kobelt & Möllendorff, 1897 (non Faust, 1893) (Gastropoda: Littoriniformes: Cyclophoridae) with a revised diagnosis of the subgenus. Club Conchylia Informationen 37: 54–55.
- Egorov RV, Greke K (2007) The genus *Cyclophorus* Montfort, 1810: systematics and nomenclature. Club Conchylia Informationen 38: 57–64.
- EPPO (2013) PQR database. European and Mediterranean Plant Protection Organization, Paris, France. <http://www.eppo.int/DATABASES/pqr/pqr.htm>
- Falkner G, Ripken Th EJ, Falkner M (2002) Mollusques continentaux de France. Liste de référence annotée et bibliographie. Patrimoines naturels 52, 350 pp.
- Gerber J, Bössneck U (2009) The genus *Vallonia* in Nepal (Gastropoda: Pulmonata: Valloniidae). Archiv für Molluskenkunde 138(1): 43–52.
- Gittenberger E, Bank RA (1996) A new start in *Pyramidula* (Gastropoda, Pulmonata: Pyramidulidae). Basteria 60: 71–78.
- Godwin-Austen AA (1876) On the Helicidae collected during the expedition into the Dafla Hills, Assam. The Journal of the Asiatic Society of Bengal 45(2): 311–318.
- Godwin-Austen AA (1882) Land and freshwater Mollusca of India including South Arabia, Baluchistan, Afghanistan, Kashmir, Nepal, Burmah, Pegu, Tenasserim, Malay Peninsula, Ceylon, and other islands of the Indian Ocean. Taylor and Francis, London. Conchologia Indica 1(2): 19–65.
- Godwin-Austen AA (1883) Land and freshwater Mollusca of India including South Arabia, Baluchistan, Afghanistan, Kashmir, Nepal, Burmah, Pegu, Tenasserim, Malay Peninsula,

- Ceylon, and other islands of the Indian Ocean. Taylor and Francis, London. *Conchologia Indica* 1(3): 67–94.
- Godwin-Austen AA (1899) Land and freshwater Mollusca of India including South Arabia, Baluchistan, Afghanistan, Kashmir, Nepal, Burmah, Pegu, Tenasserim, Malay Peninsula, Ceylon, and other islands of the Indian Ocean. Taylor and Francis, London. *Conchologia Indica* 2(9): 87–146.
- Godwin-Austen AA (1914) Land and freshwater Mollusca of India including South Arabia, Baluchistan, Afghanistan, Kashmir, Nepal, Burmah, Pegu, Tenasserim, Malay Peninsula, Ceylon, and other islands of the Indian Ocean. Taylor and Francis, London. *Conchologia Indica* 2(12): 311–442.
- Godwin-Austen AA (1920) Land and freshwater Mollusca of India including South Arabia, Baluchistan, Afghanistan, Kashmir, Nepal, Burmah, Pegu, Tenasserim, Malay Peninsula, Ceylon, and other islands of the Indian Ocean. Taylor and Francis, London. *Conchologia Indica* 3(1): 1–65.
- Gredler PV (1888) Zur Conchylien-Fauna von China. XIII Stück. Jahrbücher der Deutschen-malakzoologischen Gesellschaft 14: 343–373.
- Griffiths OL, Florens VFB (2006) A field guide to the non-marine molluscs of the Mascarene Islands (Mauritius, Rodrigues, and Réunion) and the northern dependencies of Mauritius. Bioculture Press.
- Gude GK (1914) The Fauna of British India, including Ceylon and Burma, Mollusca, Vol II (Trochomorphidae-Janellidae). Today and Tomorrow's Printers and Publishers, New Delhi.
- Gude GK (1921) The Fauna of British India, including Ceylon and Burma, Mollusca, Volume III: Land Operculates (Cyclophoridae, Truncatellidae, Assimineidae, Helicinidae). Today and Tomorrow's Printers and Publishers, New Delhi.
- Hanley S, Theobald W (1870) Illustrations of the land and freshwater shells of British India. *Conchologia Indica*, part 1: 1–10, plates 1–20.
- Hanley S, Theobald W (1875) Illustrations of the land and freshwater shells of British India. *Conchologia Indica*. Part 7: 49–56, plates 121–140.
- Hanna GD (1966) Introduced mollusks of western North America. Occasional papers of the California Academy of Sciences 48, 108 pp.
- Herbert D, Kilburn D (2004) Field Guide to the land snails and slugs of eastern South Africa. Natal Museum, Pietermaritzburg.
- Herbert DG, Rowson B (2011) *Pupa menkeana* Pfeiffer, 1853, type species of the speciose land snail genus *Gulella* Pfeiffer, 1856: correction of longstanding misidentification and designation of neotype (Mollusca: Eupulmonata: Streptaxidae). African Invertebrates 52(2): 1–10. doi: 10.5733/afin.052.0201
- Hermannsen AN (1846) Indicis generum malacozoorum primordia. Fischer Cassel 1: 1–104.
- Hooker JD (1854) Notes of a naturalist in Bengal, the Sikkim and Nepal Himalayas, the Khasia Mountains, &c. Himalayan Journals 1: 1–408.
- Hoong HW (1995) A review of the land snail fauna of Singapore. The Raffles Bulletin of Zoology 43(1): 91–113.
- Hutton T, Benson WH (1838) On the land and freshwater shells of the Western Himalaya. The Journal of the Asiatic Society of Bengal 7: 211–218.

- Kennard AS (1942) The Histoire and Prodrome of Féruccac Part III. Proceedings of the Malacological Society of London 25: 111–118.
- Kerney MP, Cameron RAD (1979) A field guide to the land snails of Britain and north-west Europe. Collins, London.
- Khanal S, Budha PB (2013) Terrestrial gastropod fauna of Nagarjun forest, Shivapuri-Nagarjun National Park, Kathmandu, Nepal. Journal of Institute of Science and Technology, Tribhuvan University 18(1): 113–119.
- Kiauta B, Butot LJM (1973) The chromosomes of *Macrochlamys tugurium* (Benson, 1852) from Nepal (Stylommatophora, Sigmurethra: Ariophantidae). Basteria 37: 145–148.
- Kobelt W (1902) Das Tierreich. Eine Zusammenstellung und Kennzeichnung der rezenten Tierformen. Mollusca: Cyclophoridae, Lief. 16. Berlin.
- Kobelt W, Möllendorff OF von (1897) Catalog der gegenwärtig lebend bekannten Pneumonopomen. Nachrichtsblatt der Deutschen Malakozoologischen Gesellschaft 1(3): 73–192.
- Kongim B, Panha S, Naggs F (2006) Karyotype of land operculate snails of the genus *Cyclophorus* (Prosobranchia: Cyclophoridae). Invertebrate Reproduction & Development 49: 1–8. doi: 10.1080/07924259.2006.9652188
- Kongim B, Sutcharit C, Tongkerd P, Panha S (2013) Karyotypes of the snorkel snail genera *Pterocyclus* and *Rhiostoma* (Prosobranchia: Cyclophoridae). The Raffles Bulletin of Zoology 61(1): 13–20.
- Kuroda T (1928) Catalogue of the shell bearing Mollusca of Amami-Oshima (Oshima, Osumi). Special publication of the Kagoshima-Ken Educational Investigation Committee, Kagoshima, Japan, 126 pp.
- Kuzminykh AA, Schileyko AA (2005) Slugs of the family Anadenidae Pilsbry, 1948, (Gastropoda: Pulmonata) with description of two new subgenera and three new species. Ruthenica 15(2): 109–118.
- Kuznetsov AG (1996) *Himalodiscus aculeatus* Kuznetsov, gen. et sp. nov. (Pulmonata, Endodontidae) from Nepal. Ruthenica 5(2): 163–165.
- Kuznetsov AG, Schileyko AA (1997) New data on Enidae (Gastropoda, Pulmonata) of Nepal. Ruthenica 7(2): 133–140.
- Kuznetsov AG, Schileyko AA (1999) Two new species of the genus *Pupinidius* Moellendorff, 1901 (Enidae, Pulmonata), and the distribution of the genus in Nepal. Ruthenica 9(2): 117–121.
- Lee YC, Leu KY, Wu WL (2008) A molecular phylogenetic investigation of *Cyathopoma* (Prosobranchia: Cyclophoridae) in East Asia. Zoological Studies 47(5): 591–604.
- Maassen WJM (2002) Remarks on Diplommatinidae from Sumatra, Indonesia with descriptions of eleven new species (Gastropoda, Prosobranchia). Basteria 66: 163–182.
- Maassen WJM (2006) Remarks on *Alycaeus* species from South-East Asia, with the descriptions of four new species with keeled shells (Gastropoda, Caenogastropoda, Cyclophoridae). Basteria 70: 133–139.
- Maceira D, Miquel SE, Espinosa J, Virgillito M, Lauranzon B (2013) Moluscos terrestres exóticos y primera cita de la familia Punctidae (Mollusca: Pulmonata: Gastropoda) y de su especie *Paralaoma servilis* para Cuba. Solenodon 11: 95–102.

- Martens E von (1860) Die Heliceen, nach natürlicher Verwandtschaft systematisch geordnet von Joh. Christ. Albers. Zweite Ausgabe nach dem hinterlassenen Manuskript besorgt von Eduard von Martens. Wilhelm Engelmann, Leipzig.
- Mead AR (1995) Anatomical studies reveal new phylogenetic interpretations in *Lissachatina* (Pulmonata: Achatinidae). Journal of Molluscan Studies 61: 257–273. doi: 10.1093/molus/61.2.257
- Minato H (2005) Two new species of *Chamalycaeus* (Gastropoda: Alycaeidae) from Miyazaki Prefecture, Southeastern Kyushu, Japan. Venus 64(1–2): 39–44.
- Mitra SC, Dey A, Ramkrishna (2005) Pictorial handbook: Indian land snails, Zoological Survey of India, 344 pp.
- Mordan PB (1988) Land Molluscs of the Wahiba Sands. Journal of Oman Studies. Special Report no. 3.
- Naggs F (1989) *Gulella bicolor* (Hutton) and its implications for the taxonomy of streptaxids. Journal of Conchology 33: 165–168.
- Naggs F (1994) The reproductive anatomy of *Paropeas achatinaceum* and new concept of *Paropeas* (Pulmonata: Achatinoidea: Subulinidae). Journal of Molluscan Studies 60: 175–191. doi: 10.1093/mollus/60.2.175
- Naggs F (1997) William Benson and the early study of land snails in British India and Ceylon. Archives of Natural History 24(1): 37–88. doi: 10.3366/anh.1997.24.1.37
- Naggs F, Raheem DC (2000) Land snail diversity in Sri Lanka. The Natural History Museum, London, UK, 216 pp.
- Nekola JC (2014) Overview of the North American terrestrial gastropod fauna. American Malacological Bulletin 32(1): 1–35.
- Nesemann H, Sharma S, Sharma G, Khanal SN, Pradhan B, Shah DN, Tachamo RD (2007) Aquatic Invertebrates of the Ganga River basin Vol 1 - Mollusca, Annelida, Crustacea (in part). Published by Hasko Nesemann. ISBN 978-99946-2-674-8
- Nevill G (1877) The list of the Mollusca brought back by Dr J Anderson from Yunnan and Upper Burma with description of new species. The Journal of the Asiatic Society of Bengal 46: 14–41.
- Nevill G (1878) Hand list of Mollusca in the Indian Museum Calcutta, Part I. Gastropoda, Pulmonata and Prosobranchia-Neurobranchia. Calcutta, 338 pp. doi: 10.5962/bhl.title.23978
- Nordsieck H (1973) Zur Anatomie und Systematik der Clausiliens, XII, Phaedusinae, I: Phae dusenaus Nepal und ihre systematische Stellung innerhalb der Unterfamilie. Archiv für Molluskenkunde 103: 63–85.
- Nordsieck H (2002) Annotated checklist of the South East Asian Phaedusinae with the description of new taxa (Gastropoda, Pulmonata, Clausiliidae). Basteria 66: 85–100.
- Panha S (1995–1996) A checklist and classification of the terrestrial pulmonate snails of Thailand. Walkerana 8(19): 31–40.
- Panha S, Burch JB (2005) An introduction to the microsnails of Thailand. Malacological Review 37/38: 1–155.
- Patterson CM (1975) *Quickia aldabraensis* (Mollusca, Gastropoda: Pulmonata, Succineidae), a new species of land snail from Aldabra atoll, western Indian Ocean. Bulletin of the British Museum (Natural History) Zoology 28: 177–186.

- Pfeiffer L (1842) *Symbolae ad historiam heliceorum. Sectio altera.* Fischer, Cassel.
- Pfeiffer L (1846) Nachtrag zur Revision der Gattung *Cyclostoma*. Zeitschrift für Malakozoologie 3: 86.
- Pfeiffer L (1853) Die gedeckelten Lungenschnecken (Helicinacea et Cyclostomacea) in Abbildungen nach der Natur. Systematisches Conchylien-Cabinet 1(19): 8–400.
- Pilsbry HA (1916–1918) Pupillidae (Gastrocoptinae). Manual of Conchology, Second Series: Pulmonata 24. Academy of Natural Sciences of Philadelphia, 1–380.
- Pilsbry HA (1920–1921) Pupillidae (Vertiginidae and Pupillinae). Academy of Natural Sciences of Philadelphia. Manual of Conchology, Second Series: Pulmonata 26: 1–254.
- Pilsbry HA (1926) The land mollusks of the Republic of Panama and the Canal zone. Proceedings of the Academy of Natural Sciences of Philadelphia 78: 57–126.
- Pilsbry HA (1946) Notes on the anatomy of Australian and Galapagos Bulimulidae (Mollusca, Pulmonata). Notulae Naturae of the Academy of Natural Sciences of Philadelphia 168: 1–4.
- Pilsbry HA, Tryon GW (1906) Achatinidae: Stenogyrinae and Coeliaxinae. Academy of Natural Sciences of Philadelphia. Manual of Conchology, Second Series: Pulmonata 18: 1–357.
- Pilsbry HA, Tryon GW (1908–1909) Structural and Systematic with illustrations of the species: *Caecilioides*, *Glessula* and *Partulidae*. Academy of Natural Sciences of Philadelphia. Manual of Conchology, Second Series: Pulmonata 20(77–78): 1–154.
- Pokryszko BM, Auffenberg K, Hlaváč JC, Naggs F (2009) Pupilloidea of Pakistan (Gastropoda: Pulmonata): Truncatellininae, Vertigininae, Gastrocoptinae, Pupillinae (in part). Annales Zoologici (Warszawa) 59(4): 423–458. doi: 10.3161/000345409X484847
- Prasankok P, Sutcharit C, Tongkerd P, Panha S (2009) Biochemical assessment of the taxonomic diversity of the operculate land snail, *Cyclophorus fulgoratus* (Gastropoda: Cyclophoridae) from Thailand. Biochemical Systematics and Ecology 36: 900–906. doi: 10.1016/j.bse.2008.11.005
- Preece RC, Naggs F (2014) *Kaliella barrakporensis* (Pfeiffer), a new hot-house alien in Britain. Journal of Conchology 41(6): 781–782.
- Preston HB (1914) Characters of new land and freshwater shells from the Naga Hills, Assam. Proceedings of the Malacological Society of London 11: 19–24.
- Raheem DC, Budha PB, Naggs F, Preece RC (2010) An illustrated guide to the terrestrial snails of Nepal. Natural History Museum, London.
- Raheem DC, Taylor H, Ablett J, Preece RC, Aravind NA, Naggs F (2014) A systematic revision of the land snails of the Western Ghats of India. Tropical Natural History, Supplement 4. Chulalongkorn University, Bangkok, Thailand, 1–285.
- Raheem DC, Naggs F (2006) An illustrated guide to the land snails of Sri Lankan natural forest and cultivated habitats. Natural History Museum, London.
- Ramakrishna, Mitra SC, Dey A (2010) Annotated checklist of Indian land molluscs. Zoological Survey of India, Kolkata.
- Raut SK (1999) The giant African land snail *Achatina fulica* Bowdich in Nepal and Bhutan. Journal of the Bombay Natural History Society 96(1): 173.
- Raut SK, Barker GM (2002) *Achatina fulica* Bowdich and other Achatinidae as pests in tropical agriculture. In: Barker GM (Ed.) Molluscs as crop pests. CAB International, 55–114.
- Riedel A (1980) Genera zonitidarum. Dr. W. Backhuys, Publisher, Rotterdam.

- Rosenberg G, Muratov IV (2006) Status report on the terrestrial Mollusca of Jamaica. Proceedings of the Academy of Natural Sciences, Philadelphia 155: 117–161. doi: 10.1635/i0097-3157-155-1-117.1
- Roth B, Sadeghian PS (2006) Checklist of the land snails and slugs of California, Second Edition. Contributions in Science, Santa Barbara Museum of Natural History 3: 1–82.
- Rowson B, Tattersfield P, Symondson WOD (2011) Phylogeny and biogeography of tropical carnivorous land-snails (Pulmonata: Streptaxoidea) with particular reference to East Africa and Indian Ocean. *Zoologica Scripta* 40(1): 85–98. doi: 10.1111/j.1463-6409.2010.00456.x
- Sarasúa PH (1944) Los Moluscos de los Jardines Cubanos. *Revista de la Sociedad Malacológica “Carlos de la Torre”* 2(2): 59–63.
- Sasaki T (2008) Micromolluscs in Japan: taxonomic composition, habitats, and future topics. *Zoosymposia* 1: 147–232. doi: 10.11646/zosymposia.1.1.12
- Schalie H van der (1948) The land- and freshwater mollusks of Puerto Rico. Miscellaneous Publications, Museum of Zoology, University of Michigan No. 70. University of Michigan Press, 134 pp.
- Schileyko AA (1998) Treatise on recent terrestrial pulmonate molluscs Part 2. Gastrocoptidae, Hypselostomatidae, Vertiginidae, Truncatellinidae, Pachnodidae, Enidae, Sagdidae. Ruthenica, Supplement 2: 129–256.
- Schileyko AA (1999) Treatise on recent terrestrial pulmonate molluscs Part 4. Draparnau- diidae, Caryodidae, Macrocyclidae, Acavidae, Clavatoridae, Dorcasiidae, Sculptariidae, Corillidae, Plectopylidae, Megalobulimidae, Strophocheilidae, Cerionidae, Achatinidae, Subulinidae, Glessulidae, Micractaeonidae, Ferrussaciidae. Ruthenica, Supplement 2: 437–556.
- Schileyko AA (2000) Treatise on recent terrestrial pulmonate molluscs Part 6. Rhytididae, Chlamydephoridae, Systrophiidae, Haplotremaidae, Streptaxidae, Spiraxidae, Oleacinidae, Testacellidae. Ruthenica, Supplement 2: 731–880.
- Schileyko AA (2002) Treatise on recent terrestrial pulmonate molluscs Part 9. Helicarionidae, Gymnarionidae, Rhysotinidae, Ariophantidae. Ruthenica, Supplement 2: 1167–1302.
- Schileyko AA (2003) Treatise on recent terrestrial pulmonate molluscs Part 11. Trigono- chlamydidae, Papillodermidae, Vitrinidae, Limacidae, Bielziidae, Agriolimacidae, Boettgerilidae, Camaenidae. Ruthenica, Supplement 2: 1467–1626.
- Schileyko AA (2004) Treatise on recent terrestrial pulmonate molluscs Part 12. Bradybaenidae, Monadeniidae, Xanthonychidae, Epiphramphoridae, Helminthoglyptidae, Elobidae, Humboldtianidae, Sphincterochilidae, Cochlicellidae. Ruthenica, Supplement 2: 1627–1763.
- Schileyko AA (2007) Treatise on recent terrestrial pulmonate molluscs Part 15. Oopeltidae, Anadenidae, Arionidae, Philomycidae, Succineidae, Athoracophoridae. Ruthenica, Supplement 2: 2049–2210.
- Schileyko AA (2011) Checklist of land pulmonate molluscs of Vietnam (Gastropoda: Stylom- matophora). Ruthenica 21(1): 1–68.
- Schileyko AA, Balashov IA (2012) *Pyramidula kuznetsovi* sp. nov. a new species of land molluscs from Nepal (Pulmonata, Pyramidulidae). Ruthenica 22(1): 41–45.

- Schileyko AA, Frank C (1994) Some terrestrial Mollusca of the Nepalesian fauna. Archiv für Molluskenkunde 123(1/6): 127–136.
- Schileyko AA, Kuznetsov AG (1996) A new genus of the Subulinidae (Pulmonata) from Nepal. Ruthenica 5(2): 158–160.
- Schileyko AA, Kuznetsov AG (1998a) Land snails of the genus *Landouria* Godwin-Austen, 1918 and some other Bradybaenidae of Nepal (Gastropoda, Pulmonata). Ruthenica 8(1): 43–54.
- Schileyko AA, Kuznetsov AG (1998b) On the taxonomic position of the genus *Himalodiscus* Kuznetsov, 1996 with a description of a new species. Ruthenica 8(1): 85–88.
- Scott B (1996) Phylogenetic relationships of the Camaenidae (Pulmonata: Stylommatophora: Helicoidea). Journal of Molluscan Studies 62: 65–73. doi: 10.1093/mollus/62.1.65
- Sherborn CD (1923) Index Animalium Part II. Index Aff.-Anus, 129–384. http://www.sil.si.edu/digitalcollections/indexanimalium/volumes/pagedisplay.cfm?filename=SIL34_02_02_0104&recordid=74260
- Sherborn CD (1927) Index Animalium Part XII. Index haani-implicatus, 2881–3136. http://www.sil.si.edu/digitalcollections/indexanimalium/volumes/pagedisplay.cfm?filename=SIL34_02_12_0184&recordid=207160
- Sherborn CD (1928) Index Animalium Part XVI. Index melanophysa-munchissoni, 3971–4194. http://www.sil.si.edu/digitalcollections/indexanimalium/volumes/pagedisplay.cfm?filename=SIL34_02_16_0188&recordid=262638
- Simone LRL (2013) Anatomy of predator snail *Huttonella bicolor*, an invasive species in Amazon rainforest, Brazil (Pulmonata, Streptaxidae). Papéis Avulsos de Zoologia 53(3): 47–58.
- Simroth H (1901) [1902] Die Nacktschneckenfauna des Russischen Reiches. Kaiserlichen Akademie der Wissenschaften, St. Petersburg. doi: 10.5962/bhl.title.46383
- Sowerby GB (1850) Descriptions of some additional species of the genus *Cyclostoma*. Thesaurus Conchyliorum 1(1): 157–168.
- Stanisic J (1981) The Carnivorous Land Snail *Gulella (Huttonella) bicolor* (Hutton, 1834) in Australia (Pulmonata: Streptaxidae). Journal of the Malacological Society of Australia, 5(1–2): 84–86. doi: 10.1080/00852988.1981.10673939
- Stanisic J (1998) Systellommatophora. In: Beesly PL, Ross GJB, Wells A (Eds) Mollusca: The Southern Synthesis. Fauna of Australia. Volume 5. CSIRO Publishing, Part B, 1060–1067.
- Subba BR (2003) Molluscan checklist of Ghodaghodi Tal area, Kailali District. Our Nature 1: 1–2.
- Subba BR, Ghosh TK (2001) Terrestrial mollusks from Nepal. Journal of the Bombay Natural History Society 98(1): 58–61.
- Subba BR, Ghosh TK (2008) Report on some terrestrial molluscs from different regions of Nepal. Journal of Natural History Museum, Tribhuvan University 23: 78–81.
- Surana R, Subba BR, Limbu KP (2004) Report of mollusks from Chimdi Lake (Birju Tal), Sunsari District eastern Nepal. Our Nature 2: 45–46.
- Sutcharit C, Naggs F, Wade CM, Fontanilla I, Panha S (2010) The new family Diapheridae, a new species of *Diaphera* Albers from Thailand, and the position of the Diapheridae within a molecular phylogeny of the Streptaxoidea (Pulmonata: Stylommatophora). Zoological Journal of the Linnean Society 160: 1–16.

- Sykes ER (1898) List of the species of *Cataulus* found in Ceylon with descriptions of some new land-shells from that Island. Proceedings of the Malacological Society of London 3: 65–74.
- Sysoev A, Schileyko A (2009) Land snails and slugs of Russia and adjacent countries. Pensoft Series Faunistica No 87. Sofia, Bulgaria, 312 pp., 142 plates.
- Tarruela A, Domènech JL (2011) Listado taxonómico ilustrado de la familia Cyclophoridae J.E. Gray, 1847 (Mollusca: Gastropoda): Parte 2. La subfamilia Alycaeinae J.E. Gray, 1850. Spira 4(1): 71–76.
- Thompson FG (2011) An annotated checklist and bibliography of the land and freshwater snails of Mexico and Central America. Bulletin of the Florida Museum of Natural History 50(1): 1–299.
- Tielecke H (1940) Anatomie, Phylogenie und Tiergeografie der Cyclophoridaen. Berlin.
- Tillier S (1980) Gastéropodes terrestres et fluviatiles de Guyane Francaise. Mémoires du Muséum national d'histoire naturelle. Paris, Sér. A, Zoologie 118: 1–19.
- Tillier S, Jackson GVH, MacFarlane R (1993) Giant African snail, 2nd edition. Pest Advisory Leaflet - South Pacific Commission, No. 6: 4.
- Tumpeesuwan S, Panha S (2008) First record of the genus *Schistoloma* Kobelt, 1902 (Prosobranchia: Pupinidae) in Thailand. The Natural History Journal of Chulalongkorn University 8(1): 65–67.
- Vaught KC (1989) A classification of the living Mollusca. American Malacologists, Melbourne, Florida, 195 pp.
- Venmans LAWC (1956) Notes on *Alycaeus*. Proceedings of the Malacological Society of London 32(8): 81–87.
- Verdcourt B (2006) A revised list of the non-marine Mollusca of East Africa (Kenya, Uganda and Tanzania, excluding Lake Malawi). Published by the author, Maidenhead, England, 75 pp.
- Wade CM, Hudelot C, Davison A, Naggs F, Mordan PB (2007) Molecular phylogeny of the helicoid land snails (Pulmonata: Stylommatophora: Helicoidea), with special emphasis on the Camaenidae. Journal of Molluscan Studies 73(4): 411–415. doi: 10.1093/mollus/eym030
- Webster NB, Van Dooren TJ, Schilthuizen M (2012) Phylogenetic reconstruction and shell evolution of the Diplommatinidae (Gastropoda: Caenogastropoda). Molecular Phylogenetics and Evolution, 63(3): 625–638. doi: 10.1016/j.ympev.2012.02.004
- Wenz W (1939) Allgemeiner Teil und Prosobranchia. In: Schindewolf OH (Ed.) Handbuch der Paläozoologie, Band 6, Gastropoda. Berlin, 481–720.
- Wiktor A (2001a) A review of Anadenidae (Gastropoda: Pulmonata) with a description of a new species. Folia Malacologica 9(1): 3–26. doi: 10.12657/folmal.009.002
- Wiktor A (2001b) Taxonomic position of *Kasperia* Godwin-Austen, 1914 (Gastropoda: Pulmonata: Limacidae). Folia Malacologica 9(1): 37–38. doi: 10.12657/folmal.009.004
- Wiktor A, Bössneck U (2004) *Limax (Limax) seticus* n. sp. from high mountains in Nepal (Gastropoda: Pulmonata: Limacidae). Folia Malacologica 12(4): 183–187. doi: 10.12657/folmal.012.017
- Wiktor A, Chen De-Niu, Wu M (2000) Stylommatophoran slugs of China (Gastropoda: Pulmonata) *Prodromus*. Folia Malacologica 8(1): 3–35. doi: 10.12657/folmal.008.002

- Wiktor A, Jurkowska J (2007) The collection of terrestrial slugs (Gastropoda: Pulmonata) at the Museum of Natural History, Wroclaw University (Poland). *Folia Malacologica* 15(2): 83–93.
- Wiktor A, Naggs F, Gupta PK (1999) *Turcomilax (Taulimax) oli* sp. n. from the Kumaun Himalaya, India (Gastropoda: Pulmonata: Limacidae). *Malakologische Abhandlungen* 19: 225–231.
- Wu M (2002) Study on Bradybaenid land snails in NW Sichuan (Gastropoda: Pulmonata: Stylommatophora). *Zoological Research* 23(6): 504–513.
- Wu M (2004) Preliminary phylogenetic study of Bradybaenidae (Gastropoda: Stylommatophora: Helicoidea). *Malacologia* 46(1): 79–125.
- Wu M, Zheng W (2009) A review of Chinese *Pupinidius* Moellendorff (Gastropoda, Stylommatophora: Enidae), with the description of new species. *Zootaxa* 2053: 1–31.
- Yen TC (1939) Die chinesischen Land- und Süßwasser-Gastropoden des Natur-Museums Senckenberg. *Abhandlungen der Senckenbergischen Naturforschenden Gesellschaft* 444: 1–233.
- Zilch A (1953) Die Typen und Typoide des Natur-Museums Senckenberg, 9: Mollusca, Cyclophoridae, Diplommatininae. *Archiv für Molluskenkunde* 82: 1–47.
- Zilch A (1956) Die Typen und Typoide des Natur-Museums Senckenberg, 17: Mollusca, Cyclophoridae, Cyclophorinae-Cyclophoreae (3). *Archiv für Molluskenkunde* 85: 33–54.
- Zilch A (1959) Euthyneura. Handbuch der Paläozoologie Band 6, Gastropoda 1–2: 1–400.
- Zilch A (1960) Euthyneura. Handbuch der Paläozoologie Band 6, Gastropoda 3–4: 401–835.
- Zilch A (1974) Vinzenz Gredler und die Erforschung der Weichtiere Chinas durch Franziskaner aus Tirol. *Archiv für Molluskenkunde* 104(4–6): 171–228.