

Revision of the genus *Epimesoplecia* Zhang, 2007 (Diptera, Nematocera, Protopleciidae) with five new species

Xiuqin Lin¹, Chungkun Shih¹, Dong Ren¹

¹ Key Lab of Insect Evolution and Environmental Change, College of Life Sciences, Capital Normal University, 105 Xisanhuanbeilu, Haidian District, Beijing 100048, China

Corresponding author: Dong Ren (rendong@mail.cnu.edu.cn)

Academic editor: V. Blagoderov | Received 18 December 2014 | Accepted 13 March 2015 | Published 30 March 2015

<http://zoobank.org/3DE583E5-6795-4A77-B70F-7B085B0A91F0>

Citation: Lin X, Shih C, Ren D (2015) Revision of the genus *Epimesoplecia* Zhang, 2007 (Diptera, Nematocera, Protopleciidae) with five new species. ZooKeys 492: 123–143. doi: 10.3897/zookeys.492.6852

Abstract

The genus *Epimesoplecia* Zhang, 2007 of Protopleciidae is revised based on five new species, *E. plethora* sp. n., *E. prosoneura* sp. n., *E. stana* sp. n., *E. macrostrena* sp. n., and *E. ambloneura* sp. n., described and illustrated from the Jiulongshan Formation of China. These new species, with clearly preserved characters of (1) compound eyes connected in males; (2) antennae, filiform or moniliform, with 16 segments; (3) r-m reaching the middle of the wing; (4) R_{4+5} ending very close to wing apex; (5) ratio of bRs/dRs ranging from 1.6 to 10.5; (6) M_2 more than 3 times as long as dM_{1+2} ; (7) legs thin and long, femur slender, almost equal to tibia; (8) tibial spurs minute; and (9) male genitalia (previously unknown), enable us to emend the diagnosis of *Epimesoplecia* Zhang, 2007. In addition, all described species of *Epimesoplecia* are characterized, their features summarized, and a key to *Epimesoplecia* species is given.

Keywords

Insects, fossil, taxonomy, Jiulongshan Formation, late Middle Jurassic

Introduction

Protopleciidae Rohdendorf, 1946, reported from the Jurassic, is a paraphyletic stem group to the Bibionidae (Blagoderov et al. 2002; Grimaldi and Engel 2005). Rohdendorf (1946) erected the Protopleciidae with three genera *Protoplecia* Handlirsch, 1906, *Mesoplecia* Rohdendorf, 1938 and *Mesopleciella* Rohdendorf, 1946 (Evenhuis 1994). Kovalev (1987) transferred 14 species in *Rhaetofungivora* Rohdendorf, 1964 of Pleciofungivoridae to Protopleciidae, but later, some of those were assigned to several different genera (Blagoderov 1996). The earliest record of the Protopleciidae is *Macropeza liasina* Geinitz, 1884 from the Early Jurassic in upper Liassic of Germany. Ansorge (1996) provided an updated description for *Protoplecia liasina* (Geinitz, 1884) and reported *Protoplecia klafackii* from the upper Liassic of Germany. Ansorge (1996) considered the affiliation of *Mesoplecia* and *Mesopleciella* with the Protopleciidae questionable based on the clearly shorter Sc, and suggested that *Archipleciomima* Rohdendorf, 1962 is the stem group of Pleciofungivoridae and Pleciomimidae due to long Rs stem. On the other hand, Lin (1976), Hong (1984), and Hong and Wang (1990) documented several genera and species in China, but many of them have been subsequently removed from this family (Blagoderov 1996). Recently, from the Jiulongshan Formation of China, Zhang (2007) described *Epimesoplecia* with two species, emended the diagnosis of *Mesoplecia* Rohdendorf, 1938, added two species to the genus, and excluded *Paraoligus exilis* Lin, 1976 and *Mesoplecia xinboensis* Hong, 1984 from this family, but stated that an alternative placement could not be suggested. Hao and Ren (2009) described three species of *Mesoplecia*. Lin et al. (2014) described two species as members of *Mesoplecia*, while transferring *M. antiqua* Hao & Ren, 2009 to Mesosciophilidae, because R_{2+3} of *M. antiqua* Hao & Ren, 2009 reaching R_1 forming a cell r, instead of reaching the anterior margin as all other protopleciids. After documented corrections and transfers, there are 33 species in seven genera described in Protopleciidae to date (Lin et al. 2014).

Herein, based on fourteen specimens collected from the Jiulongshan Formation in Daohugou Village, Ningcheng County, Inner Mongolia, China, five new species are described in *Epimesoplecia*, *E. plethora* sp. n., *E. prosoneura* sp. n., *E. stana* sp. n., *E. macrostrena* sp. n., and *E. ambloneura* sp. n. with ten specimens. One of the remaining four specimens is identified as a new material for *E. elenae* Zhang, 2007, while the other three cannot be assigned to species owing to lack of preserved diagnostic characters. These five new species are assigned to *Epimesoplecia* by a combination of the following characters: (1) antenna long, at least twice the head length; (2) wing narrow and long; (3) Sc elongate, nearly half of wing length; (4) R_{2+3} long, more than two-thirds of R_{4+5} length.

The Jiulongshan Formation of Inner Mongolia in China is very rich in fossil insects' record (Shi et al. 2011; Ren et al. 2010; Liu et al. 2012; Wang et al. 2013 and Li et al. 2013). Because of new calibrations for the Jurassic System, this deposit should be now considered as latest Middle Jurassic (late Callovian) in age (Walker

et al. 2013). The paleoenvironment reconstructed for that time was a volcanic region with mountains, streams and lakes under a humid and warm climate (Ren et al. 2002; Gao and Ren 2006).

Material and methods

All the type materials were collected from the Jiulongshan Formation (Fig. 1A) of Daohugou Village in Ningcheng County of Inner Mongolia, China (Fig. 1B) (after Ren et al. 2002). The specimens are housed in the Key Laboratory of Insect Evolution and Environmental Changes, College of Life Sciences, Capital Normal University, Beijing, China (CNU; Dong Ren, Curator). The specimens were examined and photographed using a Leica MZ12.5 dissecting microscope with a Leica DFC 500 digital camera and illustrated with the aid of camera lucida attached to the microscope. The line drawings were drawn by Adobe Photoshop CS5. The wing venation nomenclature used in this paper is based on the interpretations and system proposed by Shcherbakov et al. (1995) and Wootton and Ennos (1989).

Systematic Paleontology

Order Diptera Linnaeus, 1758

Suborder Nematocera Latreille, 1825

Family Protopleciidae Rohdendorf, 1946

Genus *Epimesoplecia* Zhang, 2007

Type species. *Epimesoplecia shcherbakovi* Zhang, 2007

Species included. Type species, *E. elenae* Zhang, 2007, *E. plethora* sp. n., *E. prosononeura* sp. n., *E. stana* sp. n., *E. macrostrena* sp. n. and *E. ambloneura* sp. n.

Revised diagnosis. Compound eyes connected in males. Antennae filiform or moniliform, with 16 segments, at least twice of head length or slightly less than twice of head length; wings narrow and long; Sc elongate, at or near the same level of r-m; bRs at least 4 times as long as r-m; R_{2+3} long, more than two-thirds of R_{4+5} , R_{2+3} slightly sigmoidly curved or straight, reaching anterior margin distad of the apex of R_1 ; r-m reaching the middle of the wing; R_{4+5} ending very close to wing apex; M_{1+2} furcated distinctly proximad or distad of R_{2+3} ; M_2 more than 3 times as long as dM_{1+2} ; bM_{3+4} longer or slightly shorter than m-cu; pterostigma absent; bM_{1+2} longer or shorter than dM_{1+2} . Legs thin and long, femur slender, almost equal to tibia; tibial spurs minute. Male genitalia: abdomen cylindrical; genitalia complex, narrower than the 8th segment, with gonocoxites rounded; gonostylus elongated, shorter than gonocoxites. Female genitalia: the 8th segment smaller than preceding segments, genitalia simple, with 2-segmented cerci, the basal segment of cerci longer than the terminal one.




Figure 1. A Measured stratigraphic section at the Jiulongshan Formation of northeastern China **B** Map showing the fossil locality (after Ren et al. 2002).

***Epimesoplecia plethora* sp. n.**

<http://zoobank.org/86178CAD-809C-4637-B64A-33B6BFBA6030>

Figs 2–3

Etymology. The epithet of *plethora* is derived from the Greek word “plethore”, meaning “fullness”, emphasizing the body covered with dense pubescence. The specific epithet is a noun in apposition.

Diagnosis. Compound eyes crescent. Antennae moniliform. Sc very close to the level of r-m; fork of Rs distad of fork of M_{1+2} ; Rs distad of crossvein r-m; bRs less than 2 times (1.6–1.8) as long as dRs, the latter about 3 times (2.6–3) as long as r-m; R_{2+3} , sigmoidly curved, distinctly shorter than bRs and dRs combined; bM_{1+2} shorter than dM_{1+2} ; bM_{3+4} shorter than m-cu; cell bp as wide as cell ba terminally.

Material. Holotype: Female. NO. CNU-DIP-NN2013202, a well-preserved almost complete body with left haltere, both wings and part of legs (Fig. 2A). Paratype: NO. CNU-DIP-NN2013209p/c, part and counterpart, lateral view, only right wing and legs preserved, head and abdomen incomplete (Fig. 3A, B).

Horizon and locality. All specimens were collected from the Jiulongshan Formation, late Middle Jurassic age (Late Callovian) from Daohugou Village, Ningcheng County, Inner Mongolia Autonomous Region in China.

Description. Head (Fig. 2D, F): Ovipiform. Compound eyes crescent in females. Antennae: scape and pedicel thick and stout, 1st flagellomere slender, the remaining ones becoming thinner toward apex.

Thorax (Figs 2A, 3A, B): Scutum convex; scutellum clearly projecting; haltere depressed.

Legs (Figs 2A, 3A, B): Forelegs comparatively thin and slender, femur slightly thicker than tibia, covered with dense setae as preserved. The 1st tarsomere 2 times as long as the 2nd tarsomere; the 3rd to 5th tarsi gradually thinned, claws small. Mid legs: femur long and slender, almost equal to tibia, tibial spurs minute, claws well-preserved. Hind legs: femur more than four-fifths of tibia; distinctly longer than forelegs and mid legs, tibia less than 2 times as long as femur; the 1st tarsomere more than 2 times as long as the 2nd tarsomere; with two pretarsal claws.

Wings (Figs 2A, C and 3A): Wing long and narrow, 2.6–2.8 times as long as width (length 8.4–9.0 mm, width 3.0–3.4 mm); Sc terminating at the middle of the anterior margin, the costal field narrow; bRs 4–5 times as long as r-m; R_{2+3} slightly sigmoidly curved; Rs arising from one-fourth of wing length, furcating distal level of fork of M_{1+2} ; stem of Rs longer than stem R, the former longer than R_{2+3} ; R_{4+5} weakly curved upward medially, ending just below apex of wing; both R_{4+5} and M_1 subparallel; crossvein m-cu as long as r-m; CuA strongly curved, distad of M forking, ending at posterior margin of wing; A_1 not preserved.

Female genitalia (Fig. 2E, G): The 8th segment slightly smaller than preceding segments, genitalia simple, with 2-segmented cerci, the basal segment of cerci thicker and longer than the terminal one.




Figure 2. *Epimesoplecia plethora* sp. n. (CNU-DIP-NN2013202). Holotype. **A** Photograph of habitus; Line drawings of **B** Habitus **C** Left wing; Photographs of **D** Details of head (under alcohol) **E** Details of female genitalia (under alcohol); Line drawings of **F** Head **G** Female genitalia. Scale bars = 1 mm.

Dimensions (in mm). [Measurements for the paratype CNU-DIP-NN2013209p/c in brackets, if different]. Holotype: female. CNU-DIP-NN2013202, Body length 1.0 [5 as preserved], maximal width of body 2.2 [2.4]. Head length 0.6, width 0.8. Fore-legs: femur 1.7 as preserved [2.2]; tibia 1.4 as preserved [3.4]. Mid legs: femur 3 [2.8], tibia 3.5 [3]. Hind legs: femur 3.6 [4], tibia 4.4 [5]. Wing: length 9.0 [8.2], width 3.4 [3.0], R_{2+3} 2.4, bRs 2.2 [2.1], dRs 1.2, R_{4+5} 3 [3.4].

Remarks. The new species is differentiated from *E. shcherbakovi* Zhang, 2007 by the following features: bRs less than 2 times (1.6–1.7) as long as dRs (vs. bRs 4.5 times as long as dRs); Rs bifurcation distad to fork of M_{1+2} (vs. Rs bifurcation at the same level of fork of M_{1+2}); dm_{1+2} longer than bm_{1+2} (vs. dm_{1+2} shorter than bm_{1+2}). The




Figure 3. *Epimesoplecia plethora* sp. n. (CNU-DIP-NN2013209p/c). Paratype. **A, B** Photographs of part and counterpart **C, D** Line drawings of part and counterpart. Scale bars = 1 mm.

new species differs from *E. elenae* Zhang, 2007 in having antennae moniliform (vs. fili-form); bRs short, less than 2 times (1.6–1.7) as long as dRs (vs. bRs long, 2.5 times as long as dRs); bM_{3+4} clearly shorter than m-cu (vs. bM_{3+4} as long as m-cu). Comparisons with other species are listed in Table 1.

***Epimesoplecia prosoneura* sp. n.**

<http://zoobank.org/260A2E9B-8331-483A-9BBB-E67D58EB0B32>

Figs 4–5

Etymology. The epithet of *prosoneura* is derived from the Greek preposition “pro”, meaning “before”, and Greek word “neura”, meaning “string or sinew”, referring to proximal position of the fork of Rs. The specific epithet is a noun in apposition.

Diagnosis. Compound eyes crescent. Antennae moniliform. Sc exceeding the level of r-m or very close to r-m; fork of Rs proximad of fork of M_{1+2} ; R_{2+3} very close to crossvein r-m; R_{2+3} , almost straight, distinctly longer than bRs and dRs combined; bRs about 10 times (9.4–10.5) as long as dRs, the latter as long as r-m; bM_{1+2} shorter than dM_{1+2} ; bM_{3+4} shorter than m-cu; cell bp wider than cell ba terminally.

Table 1. Summary of data for all species of *Epimesoplecia* Zhang, 2007.

Species	Specimen numbers	H/P	Sex	BL	WL	L/W	bRs/dRs	dRs/r-m	bM ₁₊₂ /dM ₁₊₂	M ₇ /dM ₁₊₂	bM ₃₊₄ /m-cu	bRs/r-m	Rs vs. M ₁₊₂
<i>E. plethora</i> sp. n.	CNU-DIP-NN2013202	H	♀	10.6	9	2.6	1.6	2.6	0.6	4.2	shorter	4	DS
	CNU-DIP-NN2013209 p/c	P	NA	5 (ic)	8.4	2.8	1.75	3	0.8	6	NA	5	DS
<i>E. prosoneura</i> sp. n.	CNU-DIP-NN2013207 p/c	H	♂	8.3	8	3.1	10.5	0.3	0.4	4.2	shorter	4.8	PX
	CNU-DIP-NN2013214	A	♀	8.2	8.9	3.3	9.4	0.6	0.8	5.2	shorter	5.8	PX
<i>E. stana</i> sp. n.	CNU-DIP-NN2013201 p/c	H	♀	10.2	11.2	3.3	2.5	2	2.3	8	shorter	5.4	DS
	DHG200384	H	NA	7 (ic)	9.7	2.7	4.5	0.9	1.5	5	longer	4.6	SL
<i>E. elenae</i> Zhang, 2007	DHG200385	H	♀	10.5	10	2.4	2.5	1.8	0.6	5	shorter	4	DS
	CNU-DIP-NN2013213 p/c	N	♀	11.2	7.6	2.5	2.7	1.7	0.5	5.6	shorter	4.6	DS
<i>E. macrostrena</i> sp. n.	CNU-DIP-NN2013211	H	NA	9 (ic)	7.4 (ic)	>3	5	1.1	0.7	≈3	NA	5.6	PX
	CNU-DIP-NN2013212	P	♀	10.3	7.1	2.5	4.7	1.1	0.7	3	shorter	4.6	PX
	CNU-DIP-NN2013206 p/c	P	NA	7 (ic)	8	3.5	4.2	1.2	0.7	6.2	longer	4.4	PX
<i>E. ambloneura</i> sp. n.	CNU-DIP-NN2013215	H	♀	12.7	9.1	2.5	7	0.7	2.8	11	shorter	5	SL
	CNU-DIP-NN2013208	P	NA	5 (ic)	7.4	2.3	6	0.7	1.3	8	shorter	5	SL

Notes: Abbreviations: 1. H = Holotype, P = Paratype, A = Allotype, N = New material; 2. BL = Body length (mm); 3. WL = Wing length (mm); 4. L/W = The ratio of wing length and width; 5. ic = incomplete; 6. p/c = Part and counterpart; 7. DS = Fork of Rs distad of fork of M₁₊₂; 8. PX = Fork of Rs proximad of fork of M₁₊₂; 9. NA = Not Available; 10. SL = Rs and M₁₊₂ at the same level.




Figure 4. *Epimesoplecia prosoneura* sp. n. (CNU-DIP-NN2013207p/c). Holotype. Photographs of **A, B** Part and counterpart **C** Details of head (under alcohol) **D** Details of male genitalia (under alcohol); Line drawings of **E** Counterpart **F** Head **G** Male genitalia **H** Wing. Scale bars = 1 mm, gc—gonocoxite; gs—gonostylus.

Material. Holotype: male, NO. CNU-DIP-NN2013207p/c, part and counterpart, an almost complete specimen with well-preserved antennae, both wings and part of legs (Fig. 4A, B). Allotype (paratype): female. NO. CNU-DIP-NN2013214, in dorsal view, a specimen with well-preserved wings and body (Fig. 5A, B).

Horizon and locality. All specimens were collected from the Jiulongshan Formation, late Middle Jurassic age (Late Callovian) from Daohugou Village, Ningcheng County, Inner Mongolia Autonomous Region in China.

Description. Head (Figs 4C and 5C): Ovipositor. Compound eyes protrusive in males. Antennae long, scape and pedicel thick and stout, the 1st flagellomere slender and long, the remaining ones becoming thinner toward apex.

Thorax (Figs 4A, B and 5A): Prothorax barely visible; scutum of mesothorax broad and oval, convex obviously; scutellum of metathorax projecting, semicircle; haltere depressed.




Figure 5. *Epimesoplecia prosoneura* sp. n. (CNU-DIP-NN2013214). Allotype. **A** Photograph of habitus **B** Line drawing of habitus; Photographs of **C** Details of head (under alcohol); **D** Details of female genitalia (under alcohol). Scale bars = 1 mm.

Legs (Figs 4A, B and 5A): Forelegs relatively slender; femur slender and long, covered with dense setae, slightly shorter than tibia; tarsi not preserved. Mid legs similar to forelegs, femur long and slender, tibia thinner than femur as preserved; Hind legs: femur slightly expanded, tibia slightly longer than femur as preserved.

Wings (Figs 4H and 5B): Wing long and narrow (length: 8–8.9 mm, width: 2.6–2.7 mm), apex of wings covering the abdominal terminalia. Costal field long and thin, Sc reaching C at the middle of anterior margin; Rs arising from basal one-fourth of wing length, furcating distad to fork of M_{1+2} ; bRs about 5 times (4.8–5.7) as long as r-m; crossvein m-cu slightly longer than r-m; CuA slightly curved, ending at posterior margin distad of mid wing; vein A_1 nearly straight, reaching posterior margin.

Male genitalia (Fig. 4G): Abdomen cylindrical; genitalia complex, slightly narrower than the 8th segment, with gonocoxites robust and rounded; gonostylus cylindrical and elongated, shorter than gonocoxites.

Female genitalia (Fig. 5D): The 8th segment slightly smaller than preceding segments, genitalia simple, with 2-segmented cerci, the basal segment of cerci longer than the terminal one.

Dimensions of holotype (in mm). [Measurements for the paratype CNU-DIP-NN2013214 in brackets, if different]. Holotype: male, CNU-DIP-NN2013207p/c, Body length 9.3 [8.2], maximal width of body 1.6 [1.4]. Antennae length: 1.8 (segments 1–16) [1.4 (segments 1–15)]. Foreleg: femur 1.6 as preserved; tibia 2.5 as preserved. Mid leg: femur 1.9 as preserved; tibia 2.7 as preserved. Hind leg: femur 2.2

as preserved [2 as preserved], tibia 3.5 as preserved [1.3 as preserved]. Wing: length 8 [8.9], width 2.6 [2.7]; R_{2+3} 3 [3.7]; bRs 1.9 [2.3]; dRs 0.1 [0.3]; R_{4+5} 3.6 [4].

Remarks. The new species is similar to *E. shcherbakovi* Zhang, 2007 but differs from the latter in having bRs about 10 times (9.4–10.5) as long as dRs (vs. 4.5 times); Rs bifurcation proximad of fork of M_{1+2} (vs. Rs bifurcation at the same level of fork of M_{1+2}); dM_{1+2} longer than bM_{1+2} (vs. dM_{1+2} shorter than bM_{1+2}); R_{2+3} , very close to the position crossvein r-m, distinctly longer than bRs and dRs combined (vs. R_{2+3} , distad of the position crossvein r-m, clearly shorter than bRs and dRs combined). The new species differs from *E. plethora* sp. n. in having bRs about 10 times (9.4–10.5) as long as dRs (vs. less than 2 times); bM_{3+4} clearly shorter than m-cu (vs. bM_{3+4} shorter than m-cu); Rs bifurcation proximad of fork of M_{1+2} (vs. Rs bifurcation distad of fork of M_{1+2}); dRs distinctly shorter than r-m (vs. dRs about 3 times as long as r-m); R_{2+3} , very close to the position crossvein r-m, distinctly longer than bRs and dRs combined (vs. R_{2+3} , sigmoidly curved, distinctly shorter than bRs and dRs combined). Comparisons with other species are listed in Table 1.

***Epimesoplecia stana* sp. n.**

<http://zoobank.org/323B8EC8-B13C-4360-BCFF-82E436529BB0>

Fig. 6

Etymology. The epithet is an arbitrary combination of letters used as a nun in apposition.

Diagnosis. Antennae moniliform. Sc very close to the level of r-m; fork of Rs distad of fork of M_{1+2} ; R_{2+3} distad of crossvein r-m; R_{2+3} , straight, distinctly less than bRs and dRs combined; bRs 2.5 times as long as dRs, the latter 2 times as long as r-m; bM_{1+2} significantly longer than dM_{1+2} (2.3 times); bM_{3+4} slightly longer than m-cu; cell bp narrower than cell ba terminally.

Material. Holotype: female, NO. CNU-DIP-NN-2013201p/c, in lateral view, a well-preserved body with partial antennae, almost complete wings and legs.

Horizon and locality. The specimen was collected from the Jiulongshan Formation, late Middle Jurassic age (Late Callovian) from Daohugou Village, Ningcheng County, Inner Mongolia Autonomous Region in China.

Description. Head (Fig. 6C): small, in lateral view; Antennae with segments 1–11 visible as preserved, moniliform. Maxillary palpi barely visible.

Thorax (Fig. 6A, B): Prothorax barely visible; scutum of mesothorax broad and oval, convex obviously; scutellum of metathorax projecting, semicircular; haltere depressed.

Legs (Fig. 6A, B): Forelegs relatively slender, femur slightly expanded, slightly shorter than tibia; tarsi not preserved. Mid legs similar to forelegs, femur thicker than tibia; tibial spurs minute; tarsi not preserved. Hind legs: femur slightly expanded, slightly shorter than tibia; tibia slender and long as preserved.

Wings (Fig. 6F): Wing long and narrow, more than 3 times as long as width (length: 11.2 mm, width: 3.4 mm as preserved); apex of wings covering the abdominal terminalia. Costal field long and thin, Sc reaching C very close to the position of r-m;




Figure 6. *Epimesoplecia stana* sp. n. (CNU-DIP-NN2013201p/c). Holotype. Photographs of **A**, **B** Part and counterpart; **C** Details of head (under alcohol) **D** Details of female genitalia (under alcohol); Line drawings of **E** Part **F** Left wing of counterpart **G** Partial antenna. Scale bars = 1 mm.

Rs arising from basal one-fourth of wing length, furcating distad to level of fork of M_{1+2} . Stem of Rs longer than stem of R, the former longer than R_{2+3} , bRs 5 times as long as r-m; crossvein r-m longer than m-cu; CuA slightly curved, ending at posterior margin; vein A_1 short, slightly longer than half of cell bp.

Female genitalia (Fig. 6D): The 8th segment slightly smaller than preceding segments, genitalia simple, cerci segments not visible.

Dimensions (in mm). Holotype: female, CNU-DIP-NN2013201p/c, Body length 10.2, maximal width of body 2.2. Antennae 1.1 (segments 1–11). Foreleg: femur 2.5; tibia 2.7. Mid leg: femur 3; tibia 3.4. Hind leg: femur 3.7, tibia 3.4. Wing: length 11.2, width 3.4; R_{2+3} 3; bRs 2.5; dRs 1.1; R_{4+5} 3.7.

Remarks. The new species having ratio of bRs/dRs of 2.5 is similar to *E. elenae* Zhang, 2007, but is distinguished from the latter by having Rs bifurcation

significantly distad of fork of M_{1+2} (vs. Rs bifurcation slightly distad of fork of M_{1+2}); bM_{1+2} distinctly longer than dM_{1+2} (vs. bM_{1+2} clearly shorter than dM_{1+2}); dM_{1+2} clearly shorter than r-m (vs. dM_{1+2} as long as r-m); dM_{1+2} short, almost one-eighth of M_2 (vs. dM_{1+2} long, one-fifth of M_2). Comparisons with other species are listed in Table 1.

***Epimesoplecia macrostrena* sp. n.**

<http://zoobank.org/8A937A8A-0349-45A6-B5FB-D5287856A37F>

Figs 7–8

Etymology. The epithet of *macrostrena* is derived from the Greek prefix “macro-”, meaning “large”, and Greek word “strenos”, meaning “insolence or excess of strength”, referring to the large wings and strong body of this species. The specific epithet is a noun in apposition.

Diagnosis. Antennae filiform. Sc very close to the level of r-m; fork of Rs proximal of fork of M_{1+2} ; R_{2+3} distad of crossvein r-m; R_{2+3} , straight, slightly shorter than bRs and dRs combined; bRs about 5 times (4.2–5) as long as dRs, the latter almost equal to r-m; bM_{1+2} shorter than dM_{1+2} (0.7 times); bM_{3+4} shorter than m-cu (barely longer than m-cu); cell bp slightly wider than cell ba terminally.

Materials. Holotype: sex unknown. NO. CNU-DIP-NN-2013211, in lateral view, a well-preserved specimen with partial antennae, wings and body as preserved (Fig. 7). Paratypes: CNU-DIP-NN-2013206p/c, sex unknown, a well-preserved specimen with almost complete wings and body (Fig. 8A, B), NO. CNU-DIP-NN-2013212, female, in ventral view, a well-preserved specimen with complete wings and body as preserved (Fig. 8E).

Horizon and locality. All specimens were collected from the Jiulongshan Formation, late Middle Jurassic age (Late Callovian) from Daohugou Village, Ningcheng County, Inner Mongolia Autonomous Region in China.

Description. Head (Figs 7C, D and 8C): Ovipositor and very small in lateral view; antennae filiform, segments 1–9 well-preserved, scape and pedicel slightly compressed; flagellar segments slender and long, becoming thinner and shorter toward apex (moniliform in CNU-DIP-NN-2013206p/c in Fig. 8C); maxillary palpi segments not visible.

Thorax (Figs 7A, B and 8E): Scutum well-developed, arched convex; scutellum depressed; haltere clearly depressed.

Legs (Figs 7A, B and 8E): Forelegs: femur thicker than tibia; almost four-fifths of tibia; the 1st tarsomere longer than half of tibia, the 2nd–5th greatly thinned than the former. Mid legs: femur slightly thicker than tibia; tibial spurs minute; the 1st tarsomere slightly longer than the 1st tarsomere of forelegs. Hind legs: femur expanded almost equal to tibia; tibial spurs minute; tarsi not preserved.

Wings (Figs 7E, 8D and F): Wing long and narrow (length: 7.1–8 mm, width: 2.3–3.2 mm), apex of wings not reaching the abdominal terminalia. Costal field long and thin, apex of Sc gradually tapering to the end; Rs arising almost from basal one-




Figure 7. *Epimesoplecia macrostrena* sp. n. (CNU-DIP-NN2013211). Holotype. **A** Photograph of habitus **B** Line drawing of habitus **C** Photograph of details of head (under alcohol); Line drawings of **D** Head **E** Wing. Scale bars = 1 mm.

fifth of wing length, furcating proximad of fork of M_{1+2} bRs 4.4–5.6 times as long as r-m; crossvein m-cu slightly shorter than r-m; CuA arched near anal margin; vein A_1 nearly straight, reaching posterior margin.

Female genitalia (Fig. 8F): In lateral view, genitalia simple, with 2-segmented cerci.

Dimensions (in mm). [Measurements for the holotype in brackets]. Body length 7 (as preserved)–11.3 [10.3 as preserved]; antennae 0.9 (as preserved)–1.4 (segments 1–16) [0.9 segments 1–9]. Forelegs: femur 2.1–2.5 [1.8 as preserved]; tibia 2.9–3.1 [2.7 as preserved]. Mid leg: femur 2.4–3.8 as preserved [3.8 as preserved]; tibia 2.7–3.6 as preserved [3.6 as preserved]. Hind leg: femur 3.4–4.7 [4.7], tibia 3.8–4.8 [4.8]. Wing: length 7.1–8 [7.4 as preserved], width 2.3–3.3 [3.3]; R_{2+3} 2.4–3.2 [3.2]; bRs 1.9–2.9 [2.9]; dRs 0.5–0.8 [0.6]; R_{4+5} 3–3.7 [3.7].

Remarks. The new species is similar to *E. shcherbakovi* Zhang, 2007, but is differentiated from the latter by having Rs bifurcation proximad of fork of M_{1+2} (vs. Rs bifurcation at the same level of fork of M_{1+2}); b M_{1+2} shorter than d M_{1+2} (vs. b M_{1+2} longer than d M_{1+2}); dRs as long as r-m (vs. dRs clearly shorter than r-m); d M_{1+2} long, almost one-third of M_2 (vs. d M_{1+2} short, significantly less than one-third of M_2). Comparisons with other species are listed in Table 1.




Figure 8. *Epimesoplecia macrostrena* sp. n. (CNU-DIP-NN2013206p/c). Paratype. Photographs of **A, B** Part and counterpart **C** Details of head; Line drawing of **D** Counterpart; (CNU-DIP-NN2013212) Paratype **E** Photograph of habitus **F** Line drawing of habitus. Scale bars = 1 mm.

***Epimesoplecia ambloneura* sp. n.**

<http://zoobank.org/268EE32F-E347-45C1-8FF7-F3DC03805D09>

Figs 9–10

Etymology. The epithet of *ambloneura* is derived from the Greek prefix “ambl-”, meaning “obtuse”, and Greek word “neura”, meaning “string or sinew”, referring to the blunt caudal vein of this species. The specific epithet is a noun in apposition.




Figure 9. *Epimesoplecia ambloneura* sp. n. (CNU-DIP-NN2013215). Holotype. **A** Photograph of habitus **B** Line drawing of habitus; Photographs of **C** Details of head (under alcohol) **D** Details of female genitalia (under alcohol). Scale bars = 1 mm.

Diagnosis. Antennae filiform. Sc very close to the level of r-m; fork of Rs at the same level of fork of M_{1+2} ; R_{2+3} proximad of crossvein r-m; R_{2+3} , straight, distinctly longer than bRs and dRs combined; bRs about 6–7 times as long as dRs, the latter clearly shorter than r-m; bM_{1+2} significantly longer than dM_{1+2} (1.3–2.8 times); M_2 8–11 times as long as dM_{1+2} ; bM_{3+4} shorter than m-cu; cell bp slightly wider than cell ba terminally.

Materials. Holotype: male. NO. CNU-DIP-NN-2013215, in ventral view, a well-preserved specimen with partial antennae, complete wings and body (Fig. 9A). Paratype: sex unknown. NO. CNU-DIP-NN-2013208, in dorsal view, a specimen with only right wing well-preserved, but fragments of body as preserved (Fig. 10).

Horizon and locality. All specimens were collected from the Jiulongshan Formation, late Middle Jurassic age (Late Callovian) from Daohugou Village, Ningcheng County, Inner Mongolia Autonomous Region in China.

Description. Head (Fig. 9C): Ovoid and very small in lateral view; antennae long, with 16 segments, scape and pedicel slightly compressed; flagellar segments slender and long, becoming thinner and shorter toward apex; maxillary palpi segments barely visible.

Thorax (Figs 9A and 10A): Scutum well-developed, arched convex; scutellum depressed; haltere not visible.




Figure 10. *Epimesoplecia ambloneura* sp. n. (CNU-DIP-NN2013208). Paratype. **A** Photograph of habitus **B** Line drawing of habitus. Scale bars = 1 mm.

Legs (Fig. 9A): Forelegs: femur slightly thicker than tibia; almost equal to tibia; tarsi not preserved. Mid legs: femur thicker than tibia; tibial spurs minute; tarsi not preserved. Hind legs: femur expanded, almost equal to tibia; tibial spurs minute; the 1st tarsomere longer than half of tibia; tarsi 2nd–5th not preserved.

Wings (Figs 9B and 10B): Wing long and narrow (length: 7.4–9.1 mm, width: 3.2–3.6 mm), apex of wings not reaching the abdominal terminalia. Costal field long and thin, apex of Sc gradually tapering to the end; Rs arising almost from basal one-fifth of wing length, furcating at the same level of fork of M_{1+2} ; bRs 5 times as long as r-m; CuA slightly oblique; vein A_1 nearly straight, not reaching posterior margin, exceeding the level of m-cu.

Female genitalia (Fig. 9D): Genitalia with 2-segmented cerci.

Dimensions of holotype (in mm). [Measurements for the paratype, CNU-DIP-NN2013208 in brackets]. Holotype: male. NO. CNU-DIP-NN-2013215, body length 12.7 as preserved [5 as preserved]; antennae 1.5 as preserved. Forelegs: femur 3.5 as preserved; tibia 3.9. Mid leg: femur 3.9; tibia 4.2. Hind leg: femur 4.2, tibia 4.8. Wing: length 9.1 [7.4], width 3.6 [3.2]; R_{2+3} 3.5 [2.9]; bRs 2.5 [1.9]; dRs 0.35 [0.3]; R_{4+5} 4 [3.3].

Remarks. The new species is similar to *E. shcherbakovi* Zhang, 2007, but differs from the latter in having bRs about 6–7 times as long as dRs (vs. 4.5 times); R_{2+3} clearly longer than bRs and dRs combined (vs. R_{2+3} significantly shorter than bRs and dRs combined); Rs bifurcation proximad of r-m (vs. Rs distad of r-m); M_2 8–11 times as long as dM_{1+2} (vs. 5 times); bM_{3+4} shorter than m-cu (vs. bM_{3+4} longer than m-cu). Comparisons with other species are listed in the Table 1.

Key to the species of *Epimesoplecia* Zhang, 2007

- | | | |
|---|--|------------------------------|
| 1 | Fork of Rs proximad fork of M_{1+2} | 2 |
| – | Fork of Rs distad or at same level fork of M_{1+2} | 3 |
| 2 | Fork of Rs proximad of r-m; bRs about 10 times as long as dRs | |
| – | <i>E. prosoneura</i> sp. n. | |
| – | Fork of Rs distad of r-m; bRs significantly less than 10 times as long as dRs... | |
| – | <i>E. macrostrena</i> sp. n. | |
| 3 | Fork of Rs at the same level of M_{1+2} ; dRs shorter than r-m..... | 4 |
| – | Fork of Rs distad of M_{1+2} ; dRs longer than r-m..... | 5 |
| 4 | R_{2+3} longer than Rs; bM_{1+2} clearly shorter than m-cu .. | <i>E. ambloneura</i> sp. n. |
| – | R_{2+3} distinctly shorter than Rs; bM_{1+2} longer than m-cu | |
| – | <i>E. shcherbakovi</i> Zhang, 2007 | |
| 5 | bM_{1+2} longer than dM_{1+2} | <i>E. stana</i> sp. n. |
| – | bM_{1+2} distinctly shorter than dM_{1+2} | 6 |
| 6 | Antennae moniliform; bRs significantly less than 2.5 times as long as dRs.... | |
| – | <i>E. plethora</i> sp. n. | |
| – | Antennae filiform; bRs 2.5 times as long as dRs..... | <i>E. elenae</i> Zhang, 2007 |




Figure 11. *Epimesoplecia elenae* Zhang, 2007. (CNU-DIP-NN2013213). New material. Photographs of **A, B** Part and counterpart **C** Details of female genitalia (under alcohol); Line drawings of **D** Counterpart **E** Wing. Scale bars = 1 mm.

Discussion

The generic diagnosis of *Epimesoplecia* Zhang, 2007 is revised based on eleven well-preserved new specimens, among which ten are used to describe the afore-mentioned five new species. One is identified as a new material for *E. elenae* Zhang, 2007 (Fig. 11). In total, seven species with 13 specimens have been described in *Epimesoplecia* so far, all from the Jiulongshan Formation of China (Table 1).

It is interesting to note that only one of the 13 specimens reported so far is male, i.e. the holotype of *E. prosoneura* sp. n. (CNU-DIP-NN2013207 p/c) with well-preserved male genitalia. We also describe a female paratype of *E. prosoneura* sp. n. (CNU-DIP-NN2013214) with well-preserved female genitalia. Since both specimens have similar body size, wing length and venational characters, the sexual dimorphism of this species seems to be not significant.

The measurements of body length, wing length and other important characters of wings are summarised in Table 1. The data and information suggest that the following characters are stable within a species, but differ among different species: (1) fork of Rs vs. fork of M_{1+2} ; (2) ratio range of bRs and dRs; (3) dRs longer or shorter than r-m; (4) bM_{1+2} longer or shorter than m-cu; and (5) antennae moniliform or filiform. Based on these taxonomic characters, a key to the species of *Epimesoplecia* Zhang, 2007 is provided.

Acknowledgements

We sincerely appreciate comments and suggestion from Dr. Vladimir Blagoderov and two anonymous reviewers in improving this manuscript. We are grateful to Dr. Gao Taiping, Li Longfeng and Peng Yuanyuan for their constructive and critical comments on the manuscript. This research was supported by the National Basic Research Program of China (973 Program) (No. 2012CB821906), the National Natural Science Foundation of China (Nos. 31230065 and 41272006), Great Wall Scholar and KEY project of Beijing Municipal Commission of Education (KZ201310028033) and Program for Changjiang Scholars and Innovative Research Team in University (IRT13081).

References

- Ansorge J (1996) Insekten aus dem oberen Lias von Grimmen (Vorpommern/Norddeutschland). *Neue Paläontologische Abhandlungen* 2: 1–132.
- Blagoderov VA (1996) Revision of the Dipteran family Protopleciidae (Insecta, Diptera) from the Early Jurassic Soguty locality, Kyrgyzstan. *Paleontologicheskii Zhurnal*, 85–90.
- Blagoderov VA, Lukashevich ED, Mostovski MB (2002) Order Diptera Linné, 1758. The true flies (= Muscida Laicharting, 1781). In: Rasnitsyn AP, Quicke DLJ (Eds) *History of insects*. Kluwer Academic Publisher, Dordrecht, 227–240.
- Evenhuis NL (1994) *Catalogue of the Fossil Flies of the world (Insecta: Diptera)*. Backhuys Publishers, Leiden, 600 pp.
- Gao KQ, Ren D (2006) Radiometric Dating of Ignimbrite from Inner Mongolia Provides no Indication of a Post-Middle Jurassic Age for the Daohugou Beds. *Acta Geologica Sinica* 80: 42–45. [English Edition] doi: 10.1111/j.1755-6724.2006.tb00793.x
- Geinitz FE (1884) Über die Fauna des Dobbertiner Lias. *Zeitschrift der Deutschen Geologischen Gesellschaft* 36: 566–583.
- Grimaldi D, Engel MS (2005) *Evolution of the Insects*. Cambridge University Press, New York, 755 pp.
- Handlirsch A (1906) Die fossilen Insekten und die Phylogenie der rezenten Formen. Ein Handbuch für Paläontologen und Zoologen. Engelmann, Leipzig, 800 pp.
- Hao JY, Ren D (2009) Middle Jurassic Protopleciidae from Daohugou, Inner Mongolia, China. *Acta Zootaxonomica Sinica* 34(3): 554–559. [In Chinese, English summary]
- Hong YC (1984) Tracheata, Insecta. In: *The Tianjin Institute of Geology and Mineral Resources (Ed.) Palaeontological Atlas of North China II. Mesozoic Volume*. Geological Publishing House, Beijing, 128–185. [In Chinese]
- Hong YC, Wang WL (1990) Fossil insects from the Laiyang basin, Shandong province. Geological Publishing House, Beijing, 189 pp. [In Chinese, English summary]
- Kovalev VG (1987) Mesozoic Mycetophiloid Dipterans of the Family Plecofungivoridae. *Paleontologicheskii Zhurnal* 2: 69–82. [In Russian]
- Latreille PA (1825) *Familles naturelles du règne animal, exposé succinctement et dans un ordre analytique avec l'indication de leurs genres*. J. B. Baillière, Paris, 570 pp.

- Li S, Shih CK, Wang C, Pang H, Ren D (2013) Forever love: The hitherto earliest record of copulating insects from the Middle Jurassic of China. *PLoS ONE* 8(11): e78188. doi: 10.1371/journal.pone.0078188
- Lin QB (1976) The Jurassic fossil insects from western Liaoning. *Acta Palaeontologica Sinica* 15(1): 97–116. [In Chinese, English summary]
- Lin XQ, Shih CK, Ren D (2014) Two new species of *Mesoplecia* (Insecta: Diptera: Protopleciiidae) from the late Middle Jurassic of China. *Zootaxa* 3838(5): 545–556. doi: 10.11646/zootaxa.3838.5.3
- Linnaeus C (1758) *Tomus I. Systema naturae per regna tria naturae, secundum classes, ordines, genera, species, cum characteribus, differentiis, synonymis, locis. Editio decima, reformata.* Laurentii Salvii, Holmiae, 824 pp.
- Liu LX, Shih CK, Ren D (2012) Two new species of Ptychopteridae and Trichoceridae from the Middle Jurassic of northeastern China (Insecta: Diptera: Nematocera). *Zootaxa* 3501: 55–62.
- Ren D, Gao KQ, Guo ZG, Ji SA, Tan JJ, Song Z (2002) Stratigraphic division of the Jurassic in the Daohugou Area, Ningcheng, Inner Mongolia. *Geological Bulletin of China* 21: 584–591. [In Chinese, English abstract]
- Ren D, Shih CK, Gao TP, Yao YZ, Zhao YY (2010) *Silent stories-insect fossil treasures from dinosaur era of the northeastern China.* Science Press, Beijing, 324 pp.
- Rohdendorf BB (1938) Mesozoic Diptera from Karatau. I Brachycera and some Nematocera. *Trudy paleontologicheskogo instituta akademii nauk SSSR, Moscow* 7(3): 29–67, text fig. 1–21.
- Rohdendorf BB (1946) The evolution of the Wing and the Phytogeny of Oligoneura (Diptera, Nematocera). *Trudy Paleontologicheskogo Instituta Akademii Nauk SSSR Moscow, Leningrad*, 1–108. [In Russian, with English summary]
- Rohdendorf BB (1964) Historical development of the Diptera. *Trudy Paleontologicheskogo Instituta. Akademii Nauk SSSR* 100: 1–311. [In Russian]
- Shi CF, Yang Q, Ren D (2011) Two new fossil lacewing species from the Middle Jurassic of Inner Mongolia, China (Neuroptera: Grammolegiidae). *Acta Geologica Sinica* 85(2): 482–489. [English Edition] doi: 10.1111/j.1755-6724.2011.00416.x
- Shcherbakov DE, Lukashevich ED, Blagoderov VA, (1995) Triassic Diptera and initial radiation of the order. *International Journal of Dipterological Research* 6(2): 75–115.
- Walker JD, Geissman JW, Bowring SA, Babcock LE (2013) The Geological Society of America Geologic Time Scale. *Geological Society of America Bulletin* 125: 259–272. doi: 10.1130/B30712.1
- Wang Q, Shih CK, Ren D (2013) The Earliest Case of Extreme Sexual Display with Exaggerated Male Organs by Two Middle Jurassic Mecopterans. *PLoS ONE* 8(8): e71378. doi: 10.1371/journal.pone.0071378
- Wootton RJ, Ennos AR (1989) The implications of function on the origin and homologies of the dipterous wing. *Systematic Entomology* 14: 507–520. doi: 10.1111/j.1365-3113.1989.tb00300.x
- Zhang JF (2007) New Mesozoic Protopleciiidae (Insecta: Diptera: Nematocera) from China. *Cretaceous Research*, 289–296. doi: 10.1016/j.cretres.2006.05.009