

Trioxys liui Chou & Chou, 1993 (Hymenoptera, Braconidae, Aphidiinae): an invasive aphid parasitoid attacking invasive *Takecallis* species (Hemiptera, Aphididae) in the Iberian Peninsula

Ehsan Rakhshani¹, Jose Michelena Saval², Nicolas Pérez Hidalgo³,
Xavier Pons⁴, Nickolas G. Kavallieratos⁵, Petr Starý⁶

1 Department of Plant Protection, College of Agriculture, University of Zabol, Zabol, 98613–35856, IR Iran
2 Cavanilles Institute of Biodiversity and Evolutionary Biology (ICBiBE), Universitat de València, 46980, València, Spain
3 Instituto de Biología Integrativa de Sistemas (I2SysBio), Universidad de València-CSIC, 46071, València, Spain
4 Universitat de Lleida, Department of Crop and Forest Sciences, Agrotecnio Centre, Rovira Roure 191, 25198, Lleida, Spain
5 Laboratory of Agricultural Zoology and Entomology, Department of Crop Science, Agricultural University of Athens, 75 Iera Odos str., 11855, Athens, Attica, Greece
6 Laboratory of Aphidology, Institute of Entomology, Biology Center AVCR, Branišovská 31, 370 05, České Budějovice, Czech Republic

Corresponding author: Ehsan Rakhshani (rakhshani@uoz.ac.ir)

Academic editor: K. van Achterberg | Received 22 February 2020 | Accepted 26 April 2020 | Published 30 June 2020

<http://zoobank.org/07C2FC85-4ECC-4D87-AC4C-716624D9F99D>

Citation: Rakhshani E, Saval JM, Pérez Hidalgo N, Pons X, Kavallieratos NG, Starý P (2020) *Trioxys liui* Chou & Chou, 1993 (Hymenoptera, Braconidae, Aphidiinae): an invasive aphid parasitoid attacking invasive *Takecallis* species (Hemiptera, Aphididae) in the Iberian Peninsula. ZooKeys 944: 99–114. <https://doi.org/10.3897/zookeys.944.51395>

Abstract

Biological invasion of aphids and other insects has been increased due to long distance commercial transportation of plant material. The bamboo-aphid-parasitoid association is strictly specific and even though it does not develop interactions with the local environment it should be listed as part of the fauna of southwestern Europe. On-going research regarding aphids and their aphidiine parasitoids in Spain has yielded a new association of *Trioxys liui* Chou & Chou, 1993 with an undescribed species of *Takecallis* aphids on bamboo, *Phyllostachys* spp. Here we present the first association of *T. liui* with aphids of the genus *Takecallis* that attack bamboos. *Trioxys liui* is known as a parasitoid of *Cranaphis formosana* (Takahashi, 1924) and *Phyllaphoides bambusicola* Takahashi, 1921 on bamboos in China and Russia. The accidental introduction of this parasitoid species to southwestern Europe has been probably realized through transportation of contaminated bamboo plant material. In the current study, a new host association is recorded for *T. liui*. Its potential to invade other bamboo-associated aphids and the significance of the tritrophic bamboo-aphid-parasitoid interactions in the new environments are also discussed.

Keywords

Bamboo, invasive species, new association, parasitoid

Introduction

Bamboo is a common name that encompasses at least 1250 species and 75 plant genera (Scurlock et al. 2000) within the family Poaceae (subfamily Bambusoideae). Although bamboos are distributed mostly in the tropics, they also naturally occur in the subtropical and temperate zones of all continents except Europe (Ohrnberger 2002). China (Qiu et al. 1992) and India (Shanmughavel and Francis 1996) are regions where the highest number of bamboo species are growing, mainly as natural stands. On the other hand, species of the genera *Phyllostachys* and *Pleioblastus* are ornamental plants that have been introduced into Europe for commercial purposes (Schilberszky 1911). Furthermore, there is an increasing interest in the use of bamboos as energy plants (Scurlock et al. 2000; Wright 2006; Potters et al. 2009) and for other industrial applications (van der Lugt et al. 2006; Lipp-Symonowicz et al. 2011). The well known group of aphids that belongs to genus *Takecallis* Mastumura is associated with bamboos. These aphids have been considered as invasive organisms in different parts of the world where bamboos are purposefully or accidentally introduced (Stroyan 1964; Cof-felt and Schultz 1990; Limonta 1990; Lazzari et al. 1999; Trejo-Loyo et al. 2004; Valenzuela et al. 2010). Three invasive *Takecallis* species, i.e., *Takecallis arundinariae* (E-sig, 1917), *Takecallis taiwanus* (Takahashi, 1926) and *Takecallis arundicolens* (Clarke, 1903) (Calaphidinae, Panaphidini) have already been recorded in several European countries (see Rakhshani et al. 2017). It should be noted that the three aforementioned aphid species occur in Spain (Nieto Nafría and Mier Durante 1998; Suay-Cano and González-Funes 1998; Pons and Lumbierres 2004).

A new parasitoid species, *Trioxys remaudierei* Starý & Rakhshani, 2017 (Hymenoptera, Braconidae, Aphidiinae), has been recently described and associated with two bamboo aphids, *T. arundinariae* and *T. taiwanus* in France and Spain (Rakhshani et al. 2017). This evidence led to an exhaustive investigation for parasitoids of bamboo aphids in southwestern Europe that resulted in detection of a *Trioxys* species emerging from *Takecallis* aphids, which infest bamboo groves in Spain. Surprisingly, this parasitoid species was not conspecific with *T. remaudierei*. Here we determine and illustrate a new aphid parasitoid association from bamboos in Spain. An annotated world-review of *Takecallis* aphids attacking bamboos is also provided.

Material and methods

Research on bamboo aphids was carried out in the east and northeast of the Iberian Peninsula: Valencia, Barcelona and Lleida (Fig. 1). In Valencia, samples were collected



Figure 1. General scheme of the sampling localities in the Iberian Peninsula **A** patch of *Phyllostachys aurea* in the Arboretum of Lleida **B** street garden with rows of bamboo (*Phyllostachys* sp.) in Barcelona.

from the Botanical Garden of the University of Valencia during spring 2017 (March to June). All species of bamboos that are growing in the Botanical Garden were sampled: *Bambusa ventricosa*, *Dendrocalamus giganteus*, *Phyllostachys nigra*, *Phyllostachys viridis*, *Phyllostachys aurea*, *Pleioblastus linearis*, *Pleioblastus pumilus*, *Pleioblastus pygmaeus* and *Shibataea kumasasa*. The aphid colonies were inspected in the field and the laboratory so as to be checked for any occurrence of parasitoids. At the beginning of April (10.iv.2017), several mummies of *Takecallis* spp. were recorded on mixed colonies of the three species of *Takecallis* present in the Iberian Peninsula (*T. arundicolens*, *T. arundinariae* and *T. taiwanus*). Samples from Barcelona originated from two locations: 1) a street garden (41°24'24.15"N, 2°11'49.78"E, 5 m a.s.l.), where a 100 m × 2 m row of bamboo (*Phyllostachys* nr. *aurea*) are planted; 2) the gardens of the Royal Palace of Pedralbes, where different irregular size patches of bamboo (mainly *Phyllostachys* nr. *aurea*), of a whole area of about 300 m² are planted (41°23'16.05"N, 2°07'03.53"E, 99 m a.s.l.). Both locations are separated by a 7 km straight line. Visual inspection determined the presence of some common aphid species on bamboo (*Melanaphis bambusae* (Fullaway, 1910) (Aphidinae, Aphidini); *T. taiwanus* and *T. arundinariae*), but the plants from the first location were mostly infested by *Takecallis* species. Mummies and live aphids within isolated or mixed colonies of *T. taiwanus* and *T. arundinariae* were collected in May and June 2018. In Lleida, samples from *Indocalamus tessellatus* and *P. aurea* were collected from the Arboretum and Botanical Garden Pius Font i Quer (41°37'29.96"N, 0°36'11.70"E, 182 m a.s.l.). *Phyllostachys aurea* was the dominant bamboo species there, occupying most of the bamboo plantation area (700 m²), but there were also small patches of other bamboos species such as *Phyllostachys aureosulcata*, *P. nigra*, *Pleioblastus fortunei*, *Pseudosasa usawae*, *S. kumasasa*, *I. tessellatus* and *Fargesia scabrida*. Aphids from the genus *Takecallis* were found on *P. aurea* and *I. tessellatus*.

Aphid colonies, containing both alive individuals and mummies, were sampled with small pieces of bamboo plants that were gently cut with scissors. Samples were transferred to the laboratory, where they were maintained at room temperature. Some adult aphids were preserved in a solution containing two parts of 90% ethanol to one

part of 75% lactic acid (Eastop and van Emden 1972) for identification. Then, they were compared with the keyed material according to Nieto Nafria and Mier Durante (1998) and Blackman and Eastop (2019). Aphid nomenclature and classification follows Favret (2019). The emerged parasitoids were captured with an aspirator and directly dropped into 70% ethanol. The external morphology of parasitoids was studied using a Nikon Eclips E200 stereomicroscope (Nikon Corp., Japan). The parasitoid specimens were identified according to Chen and Shi (2001) and Davidian (2005). Illustrations were traced on the digital photographs captured from the slides with a Canon EOS 700D (Canon Inc., Japan) in Adobe Illustrator CS5 and were processed in Photoshop CS5 (Adobe systems Inc., San Jose, USA).

A series of voucher specimens of the emerged aphid parasitoids was sorted and preserved in absolute ethanol, kept in a refrigerator for DNA extraction. Total DNA was extracted separately from two individuals (a male and a female) following the HotSHOT method (Truett et al. 2000) using 60 µl of both alkaline lysis and neutralizing reagents. A 710 bp fragment of the 5' region of the mitochondrial gene coding the cytochrome c oxidase subunit 1 (COI) was sequenced, using the primer pair LCO1490 and HCO2198 as described by Folmer et al. (1994). PCR and sequencing procedures are outlined in Pérez Hidalgo et al. (2012). After removing sequences corresponding to primers used in the PCR reaction, the sequences obtained from each sample consisted of 658 nucleotides. DNA sequences of both male (a) and female (b) parasitoids were deposited in the GenBank database under the accession numbers MT324250 and MT324249, respectively. The existing mtCOI sequences in NCBI for *Trioxys* and *Binodoxys* spp. were retrieved and aligned in MEGA X (Kumar et al. 2018) with the integrated ClustalW using default parameters. The evolutionary history was inferred using the Neighbor-Joining method (Saitou and Nei 1987) with pairwise deletion of missing sites and Kimura-2-Parameter (K2P) distances (Kimura 1980). The optimal tree with the sum of branch length = 1.48087269 is shown. The percentage of replicate trees in which the associated taxa clustered together in the bootstrap test (1000 replicates) are shown next to the branches. *Ephedrus persicae* Froggatt, 1904 (Hymenoptera, Braconidae, Aphidiinae – KY213710.1) and *Praon volucre* (Haliday, 1833) (Hymenoptera, Braconidae, Aphidiinae – KJ698515.1) were used as outgroups for the molecular analysis. The tree is drawn to scale, with branch lengths in the same units as those of the evolutionary distances used to infer the phylogenetic tree. The analysis involved 34 nucleotide sequences. Codon positions included were 1st+2nd+3rd+Noncoding. All ambiguous positions were removed for each sequence pair (pairwise deletion option). There were a total of 519 positions in the final dataset.

The material is deposited in the Entomology Collection of the University of Valencia of the Cavanilles Institute of Biodiversity and Evolutionary Biology, in the Laboratory of Entomology of the Department of Crop and Forest Sciences of the University of Lleida, in the collection of P. Stary (Academy of Sciences of the Czech Republic), in the Laboratory of Agricultural Zoology and Entomology of the Agricultural University of Athens and in the collection of Department of Plant Protection, University of Zabol.

Results

Four aphid species are present in the Botanical Garden of the University of Valencia living in mixed colonies only on *P. viridis*, *P. aurea*, and *P. linearis*. Three of them, which are not attended by ants, belong to the genus *Takecallis* (*T. arundicolens*, *T. arundinariae*, *Takecallis* sp. (probably a new species)). However, the fourth of these species, *M. bambusae*, was strongly attended by the ant *Lasius grandis* Forel, 1909 (Hymenoptera, Formicidae). Out of all parasitoid individuals emerged from 20 mummies of *Takecallis* sp. collected from Valencia we obtained six males and six females of an aphidiine whose morphological characters clearly matched those of *T. liui*. Samples collected from Barcelona and Lleida led to the emergence of additional 11 female and 13 male specimens of *Trioxys liui* Chou & Chou (Hymenoptera, Braconidae, Aphidiinae) originating from *T. taiwanus* and *T. arundinariae* which are reported below. Although *T. liui* has been originally figured (Chou and Chou 1993) and later keyed by Chen and Shi (2001) and Davidian (2005, 2007), additional illustrations (Figs 2, 3) are provided to increase the taxonomical evidence for comparison with other parasitoid species of bamboo aphids.

Trioxys liui Chou & Chou, 1993

Figures 2, 3

Material examined. 6♀ 6♂, SPAIN: Valencia, Botanical Garden, 39°28'36.6"N, 0°23'09.1"W, 17 m a.s.l., collected from 15.iv.2018 to 6.vi.2018, ex *Takecallis* sp. on *Phyllostachys aurea*, J.M. Michelena leg.; 2♀, Barcelona, public garden, 41°24'24.25"N, 2°11'49.78"E, 2 m a.s.l., 06.v.2018, ex *Takecallis taiwana* (Takahashi) on *Phyllostachys* sp., X. Pons leg. [Sample B-1052]; 2♀ 4♂, Barcelona, public garden, 41°24'24.15"N, 2°11'49.78"E, 2 m a.s.l., 18.v.2018, ex *Takecallis taiwana* (Takahashi) on *Phyllostachys* sp., X. Pons leg. [Sample B-1054]; 2♀ 4♂, same collecting data as for preceding, 06.vi.2018, ex *Takecallis arundinariae* (Essig) on *Phyllostachys* sp., X. Pons leg. [Sample B-1064]; 3♀, same collecting data as for preceding, ex *Takecallis taiwana* (Takahashi) on *Phyllostachys* sp., X. Pons leg. [Sample B-1065]; 2♀ 1♂, same collecting data as for preceding, captured on *Phyllostachys* sp., X. Pons leg. [Samples B-1064 + B-1065]; 1♂, Barcelona, public park, 41°23'16.05"N, 2°07'03.53"E, 71 m a.s.l., 15.vi.2018, captured on *Phyllostachys* sp., X. Pons leg. [Sample B-1070]; 1♂, Lleida Arboretum, 41°37'29.96"N, 0°36'11.70"E, 181 m a.s.l., 24.v.2018, captured on *Indocalamus tessellatus*, X. Pons leg. [Sample L-1059].

Morphological diagnosis. Female – Body length: 1.4–1.6 mm, forewing length 1.5–1.6 mm. **Chlypeus** (Fig. 2A) narrow with 6 long setae on dorsal surface. Maxillary palp with 4 palpomeres, labial palp with 2 palpomeres (Fig. 2B). **Antenna** (Fig. 2C) filiform, with 11 antennomeres, covered mainly with semierect setae, slightly shorter than the diameter of segments. Flagellomere 1 (F1) 1.15–1.22× as long as F2 and 3.60–3.75× as long as maximally wide. F1 and F2 with 1–2 and 1–3 longitudinal




Figure 2. *Trioxys liui* – female **A** head, frontal view **B** maxillary and labial palps **C** antenna **D** mesonotum and scutellum, dorsal view **E** forewing **F** propodeum **G** petiole, dorsal view **H** genitalia, lateral aspect.

placodes, respectively. **Mesonotum** (Fig. 2D) smooth with notaulices hardly visible at anterior part, sparsely setose. **Forewing** (Fig. 2E) stigma elongate, triangular with slightly convex outline, $2.90\text{--}3.10\times$ as long as wide and $3.00\text{--}3.20\times$ as long as R1. Vein r&RS extended beyond R1. Wing margin with very long fringes. **Propodeum** (Fig. 2F) with well developed carinae, irregularly branched at anterior and lateral parts. Central



Figure 3. *Trioxys liui* – general habitus **A** female **B** male.

areola partially divided by irregular internal carinae. Upper and lower parts of propodeum with 8 and 2 long erected setae, respectively. **Petiole** (Fig. 2G) short, 1.55–1.80× as long as wide at spiracles with a pair of long setae near each prominent spiracular tubercle and a single seta at postero-dorsal area. **Ovipositor sheath** (Fig. 2H) stout, with smooth dorsal outline, deeply concaved ventrally in anterior edge, sharply expanded into a ventral projection bearing a pair of long setae. Prongs distinctly separated at base, almost stright, progressively constricted and upcurved at apex. Ventral side of prongs bear 4–5 prependicular setae, with single claw-shape seta and pair of short setae at apex.

Color (Fig. 3A). Head and mesosoma dark brown to black, gaster brown. Antenna brown, mouth parts, pedicel, F1, legs and petiole yellowish brown. Wings infumated. Apex of ovipositor sheath dark brown.

Male (Fig. 3B) – Antennae with 13 antennomeres, body length 1.4–1.5 mm.

Molecular data. The DNA sequences of the mtCOI gene were obtained from a single specimen of both male and female of *T. liui*, with no intraspecific genetic distance (0.0%). The interspecific genetic distances of mtCOI within *Trioxys* species ranged from 10.9% to 14.7%. The uncorrected pairwise genetic distances (p-distance) of mtCOI, separated *T. liui* from all other *Trioxys* species, with at least 10% of the sequence divergence (Table 1). After the reconstruction of the Neighbor-Joining tree, *T. liui* stands basal to the branch that includes *Trioxys pallidus* (Haliday, 1833) and *Trioxys complanatus* Quilis, 1931 (Fig. 4). No further analysis was possible because almost all deposited sequences originated from specimens that were identified only at generic level.

Discussion

Trioxys liui is an aphidiine parasitoid of two bamboo aphids in China: *Cranaphis formosana* (Takahashi, 1924) (Liu 1975) and *Phyllaphoides bambusicola* Takahashi, 1921 (Calaphidinae, Panaphidini) on *Phyllostachys makinoi* (Chou and Chou 1993). So far, *T. arundinariae* in Spain and *T. taiwanus* in France, both attacking *Phyllostachys* spp.,




Figure 4. Neighbor-Joining tree based on the partial mtCOI sequences from *Trioxys* and *Binodoxys* spp., including *Trioxys liui*, with *Praon volucre* and *Ephedrus persicae* as outgroups (NCBI accession no). Numbers next to nodes are the bootstrap values.

Table 1. Uncorrected pairwise genetic distances between *Trioxys liui* and other *Trioxys*/*Binodoxys* species based on mtCOI sequences resulted from MEGA X.

Species	Accession No.	1	2	3	4	5	6	7	8	9	10
1 <i>T. liui</i>	MT324249										
2 <i>T. pallidus</i>	KM973271.1	0.139									
3 <i>T. complanatus</i>	KJ848479.1	0.147	0.052								
4 <i>Trioxys</i> sp.	KR411291.1	0.112	0.137	0.134							
5 <i>Trioxys</i> sp.	KR420424.1	0.109	0.134	0.134	0.005						
6 <i>T. auctus</i>	KY887993.1	0.133	0.145	0.136	0.141	0.138					
7 <i>T. parauctus</i>	MK080164.1	0.163	0.128	0.153	0.136	0.136	0.125				
8 <i>T. sunnysidensis</i>	JN288965.1	0.125	0.137	0.136	0.128	0.128	0.083	0.086			
9 <i>B. brevicornis</i>	MK080162.1	0.144	0.153	0.158	0.133	0.133	0.135	0.133	0.122		
10 <i>B. acalephae</i>	MK080161.1	0.122	0.128	0.128	0.131	0.128	0.104	0.104	0.101	0.101	
11 <i>B. angelicae</i>	MK080159.1	0.128	0.143	0.151	0.137	0.134	0.127	0.130	0.081	0.112	0.109

have been reported as the only hosts of the newly described aphidiine parasitoid, *T. remaudierei* in Western Europe (Rakhshani et al. 2017). Our findings add *T. liui* as a new member of the parasitoid fauna of Spain that parasitizes even a new species of *Takecallis* (Valencia). This evidence contributes both to the increase in the number of exotic parasitoid species on bamboo aphids and the potential of interspecific relations of parasitoids. The new association, *T. liui*/*Takecallis* spp. (Valencia, Barcelona and Lleida) is also a new assemblage apart from *T. remaudierei*/*Takecallis* spp. (Paris-France and Lleida-Spain). Therefore, our results clearly document the invasion and/or subsequent adaptation of the southeastern Asian endemic species into Western Europe. The lack of information on *T. liui*, with the exception of the knowledge about its original region, China, has indicated a direct interaction of southeastern Asia with Spain through bamboo plant material, that accidentally included aphid and parasitoid contamination. Furthermore, secondary exchanges of shipments between the gardeners are also considered possible. However, similar evidence from insects in other areas has not been documented yet. It is noteworthy to emphasize the obvious morphological differences between the two parasitoids of *Takecallis* in Europe, which might reveal new bamboo-aphid-parasitoid associations. *Trioxys remaudierei* has long ventral prongs fused over two-thirds of their length. However, prongs of *T. liui* are short and completely separated. Among European species, *Trioxys betulae* Marshall, 1896 has also partially fused prongs. However, this parasitoid has a different host range since it includes aphid species that do not belong to genus *Takecallis* (Rakhshani et al. 2017). Despite the complexity of the taxonomy of the genus *Trioxys*, little attention has been devoted to using molecular data for identification and resolving questions on their classification (mainly unpublished). None of these efforts include the parasitoids of *Takecallis* aphids.

Generally, bamboos grow and spread fast (Cao et al. 2011), a trait that can crowd out native plant species. The allelopathic secretions of various bamboo species can also prohibit the growth of the nearby plant species (Chou and Yang 1982; Chou 1999; Rawat et al. 2017). However, ornamental bamboos should be protected against invasive herbivorous insects that could be distributed along with their host plant material in botanical gardens in Europe. With few exceptions (Potenza 2005), there is no record

Table 2. *Takecallis* species associated with various bamboos and their distribution.

Aphid species	Distribution	Host plant	References
<i>Takecallis affinis</i> Ghosh, 1986	India	<i>Bambusa</i> sp. <i>Chimonobambusa jaunsarensis</i>	Ghosh 1986
<i>Takecallis alba</i> Lee, 2018	South Korea	<i>Pseudosasa</i> sp. <i>Sasa</i> spp.	Lee and Lee 2018
<i>Takecallis arundicola</i> (Clarke, 1903)	China, Eastern Russia, Europe (Spain, France, Netherlands, Serbia), Japan, Korea, North America, Taiwan	<i>Arundinaria japonica</i> <i>Bambusa</i> sp. <i>Phyllostachys</i> sp. <i>Pleioblastus chino</i> <i>Pseudosasa japonica</i> <i>Sasa nipponica</i> <i>Sasa palmate</i> <i>Sasa paniculata</i> <i>Sasa senaenensis</i> <i>Sasaella ramosa</i>	Leclant 1966; Higuchi 1972; Lampel and Meier 2003; Qiao and Zhang 2004; Pons and Lumbierres 2004; Blackman and Eastop 2019; Piron 2009; Lee and Lee 2018; Petrović-Obradović et al. 2018
<i>Takecallis arundinariae</i> (Essig, 1917)	Australia, Central America (Mexico), China, Eastern Russia, Europe (Greece, Hungary, Italy, Netherlands, Portugal, Spain, Switzerland, United Kingdom), India, Japan, Korea, North America, South America (Brazil), Taiwan	<i>Arundinaria graminea</i> <i>Arundinaria japonica</i> <i>Bambusa bambos</i> <i>Bambusa rigida</i> <i>Bambusa stenostach</i> <i>Bambusa textilis</i> <i>Dendrocalamus asper</i> <i>Phyllostachys aurea</i> <i>Phyllostachys bambusoides</i> <i>Phyllostachys castillonis</i> <i>Phyllostachys dulcis</i> <i>Phyllostachys edulis</i> <i>Phyllostachys iridescens</i> <i>Phyllostachys mannii</i> <i>Phyllostachys puberula</i> <i>Phyllostachys viridiglaucescens</i> <i>Pseudosasa japonica</i> <i>Sasa nipponica</i> <i>Sasa palmate</i> <i>Sasa senaenensis</i> <i>Sinoarundinaria nitakayamensis</i> <i>Sinobambusa tootsik</i>	Higuchi 1968; Higuchi 1972; Raychaudhuri 1973; Ghosh 1980; Ghosh and Quednau 1990; Coffelt and Schultz 1990; Giacalone and Lampel 1996; Aguiar and Ilharco 1997; Nieto Nafría and Mier Durante 1998; Suay-Cano and González-Funes 1998; Lazzari et al. 1999; Qiao and Zhang 2004 Trejo-Loyo et al. 2004; Pons and Lumbierres 2004; Cœur d'acier et al. 2010; Blackman and Eastop 2019; Barbagallo and Ortu 2009; Piron 2009; Valenzuela et al. 2010; Basky and Neményi 2014; Lee and Lee 2018
<i>Takecallis asumentus</i> Qiao & Zhang, 2004	China	<i>Bambusa</i> sp.	Qiao and Zhang 2004
<i>Takecallis himalayensis</i> Chakrabarti, 1988	India	<i>Arundinaria jounsarensis</i> <i>Bambusa</i> sp.	Chakrabarti 1988; Ghosh and Quednau 1990
<i>Takecallis sasae</i> (Matsumura, 1917)	Japan	<i>Phyllostachys</i> sp. <i>Pleioblastus</i> sp. <i>Sasa nipponica</i> <i>Sasa paniculata</i>	Higuchi 1968, 1972
<i>Takecallis taiwanus</i> (Takahashi, 1926)	Central America (Mexico), China, Europe (Hungary, Croatia, Netherlands, Spain, United Kingdom, Russia), Georgia, Japan, Korea, New Zealand, North America, South Africa, South America (Argentina, Brazil, Chile), Taiwan	<i>Arundinaria anceps</i> <i>Arundinaria gigantea</i> <i>Bambusa stenostach</i> <i>Dendrocalamus asper</i> <i>Phyllostachys arcana</i> <i>Phyllostachys aurea</i> <i>Phyllostachys bambusoides</i> <i>Phyllostachys castukkinis</i> <i>Phyllostachys dulcis</i> <i>Phyllostachys nigra</i> <i>Phyllostachys sulphurea</i> <i>Phyllostachys viridiglaucescens</i> <i>Pleioblastus amarus</i> <i>Pleioblastus variegatus</i> <i>Sasa</i> spp. <i>Shibataea kumasasa</i>	Stroyan 1964; Higuchi 1968; Higuchi 1972; Giacalone and Lampel 1996; Nieto Nafría and Mier Durante 1998; Suay-Cano and González-Funes 1998; Foureaux and Kato 1999; Lazzari et al. 1999; Gonzales et al. 2000; Peronti and Sousa-Silva 2002; Qiao and Zhang 2004; Pons and Lumbierres 2004; Trejo-Loyo et al. 2004; Ortego et al. 2004; Blakman and Eastop 2019; Ripka 2008; Simala et al. 2008; Maslyakov and Izhevsky 2011; Lee and Lee 2018

of aphid contamination hazards on bamboos, but their associations should be considered and listed in the Iberian Peninsula. The above mentioned case was the infestation of bamboo hedges by *T. taiwanus*, which were successfully controlled by the application of chemical pesticides.

Aphids of the genus *Takecallis* and their parasitoids are strictly associated with bamboo while any other interactions with the environmental local fauna have not been determined yet (Rakhshani et al. 2017). Therefore, in terms of the specific associations of *Takecallis* spp. and their parasitoids on bamboos, no hazardous effects are expected on the environment that is reserved for their plantation (i.e., arboretums, parks, public and private gardens). Although genus *Takecallis* is represented by seven valid species worldwide (Table 2), a vast number of aphid genera are recorded in association with bamboos and other plant species (e.g., Anacardiaceae, Cyperaceae, Poaceae, Rosaceae, Styracaceae) (Blackman and Eastop 1994, 2006). Apart from aphids, a rich fauna of insects is also associated with bamboos in their area of origin (Revathi and Remadevi 2011).

Conclusion

The increasing attention of bamboos from both ornamental and industrial aspects will evidently lead to the invasion of more alien species into Europe and other parts of the world. A concrete knowledge on the status of those insects in southeastern Asia needs to be elaborated, which is the current bamboo source to Europe. Phytosanitary authorities should very carefully examine all imported bamboo material at any entry point of Europe to intercept alien insect species that could be a potential threat to the local plantations and entomofauna.

Acknowledgments

The research was supported by the Grant No. UOZ-GR-9618-6, University of Zabol and Grant No. CGL2015-68188-P, funded by “Ministerio de Economía, Industria y Competitividad” of Spain (MIMECO). Thanks are expressed to Xue-Xin Chen (Zhejiang University, Fuzhou), Isabel Mateu and Jaime Güemes (Jardí Botànic Universitat de València), Javad Karimi (Ferdowsi University of Mashad, Iran) and Behnam Motamedinia (Canadian National Collection of Insects) for their technical assistance.

References

- Aguiar AMF, Ilharco FA (1997) New records of aphids (Homoptera, Aphidoidea) from Madeira Island. *Boletín de Sanidad Vegetal Plagas* 23: 565–570.
- Barbagallo S, Ortu S (2009) Contribution to the knowledge of the Sardinian aphid fauna (Hemiptera, Aphididae). *Bollettino di Zoologia Agraria e di Bachicoltura* 41: 181–206. [In Italian]

- Basky Z, Neményi A (2014) *Takecallis arundinariae* (Essig, 1917) new record for the Hungarian aphid fauna on *Phyllostachys iridescens* (CY Yao and SY Chen) bamboo species. *Acta Phytopathologica et Entomologica Hungarica* 49: 281–287. <https://doi.org/10.1556/APhyt.49.2014.2.15>
- Blackman RL, Eastop VE (1994) *Aphids on the World's Trees. An Identification and Information Guide*. CAB International, Wallingford, 1024 pp.
- Blackman RL, Eastop VE (2006) *Aphids on the World's Herbaceous Plants and Shrubs*. John Wiley and Sons Ltd., Chichester, 1456 pp.
- Blackman RL, Eastop VE (2019) *Aphids on the world plants. An online identification and information guide*. <http://www.aphidsonworldsplants.info/> [Date accessed: 1 July 2019]
- Cao ZH, Zhou GM, Wong MH (2011) Special issue on bamboo and climate change in China. *The Botanical Review* 77: 188–189. <https://doi.org/10.1007/s12229-011-9064-1>
- Chakrabarti S (1988) Revision of Drepanosiphinae (Homoptera, Aphidoidea) from the Indian Subregion. *Oriental Insects* 22: 1–86. <https://doi.org/10.1080/00305316.1988.11835482>
- Chen J, Shi Q (2001) *Systematic Studies on Aphidiidae of China (Hymenoptera, Aphidiidae)*. Fujian Science and Technology Press, Fuzhou, 234 pp. [In Chinese]
- Chou CH (1999) Roles of allelopathy in plant biodiversity and sustainable agriculture. *Critical Reviews in Plant Sciences* 18: 609–636. <https://doi.org/10.1080/07352689991309414>
- Chou CH, Yang CM (1982) Allelopathic research of subtropical vegetation in Taiwan II. Comparative exclusion of understory by *Phyllostachys edulis* and *Cryptomeria japonica*. *Journal of Chemical Ecology* 8: 1489–1507. <https://doi.org/10.1007/BF00989105>
- Chou LY, Chou KC (1993) A new species of *Trioxys* (Hymenoptera, Braconidae) from Taiwan. *Chinese Journal of Entomology* 13: 375–378.
- Clarke WT (1903) A list of California Aphididae. *The Canadian Entomologist* 35(9): 247–254. <https://doi.org/10.4039/Ent35247-9>
- Cœur d'acier A, Pérez Hidalgo N, Petrović-Obradović O (2010) Aphids (Hemiptera, Aphididae) Chapter 9.2. *BioRisk* 4: 435–474. <https://doi.org/10.3897/biorisk.4.57>
- Coffelt MA, Schultz PB (1990) Seasonal abundance and population dynamics of a bamboo aphid, *Takecallis arundinaria* (Homoptera, Aphididae). *Journal of Entomological Science* 25: 526–534. <https://doi.org/10.18474/0749-8004-25.4.526>
- Davidian EM (2005) A review of species of the subgenus *Trioxys* s.str., genus *Trioxys* Haliday (Hymenoptera, Aphidiidae) of Russia and adjacent countries. *Entomologicheskoe Obozrenie* 84: 579–609. [In Russian]
- Davidian EM (2007) Aphidiidae. In: Lelej AS (Ed.) *Key to the Insects of Russian Far East (Vol. IV)*. Neuropteroidea, Mecoptera, Hymenoptera, Dalnauka, Vladivostok, 192–254. [In Russian]
- Eastop VE, van Emden HF (1972) The insect material. In: van Emden HF (Ed.) *Aphid Technology*, Academic Press, London, 45 pp.
- Essig EO (1917) *Aphididae of California*. University of California publications 1: 301–346.
- Favret C (2019) *Aphid Species File*. Ver 5.0. <http://aphid.speciesfile.org/> [1 July 2019]
- Folmer O, Black M, Hoeh W, Lutz R, Vrijenhoek R (1994) DNA primers for amplification of mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates. *Molecular Marine Biology and Biotechnology* 3: 294–299.

- Forel A (1909) Fourmis d'Espagne récoltées par M. O. Vogt et Mme Cécile Vogt, Docteurs en médecine. *Annales de la Société Entomologique de Belgique* 53: 103–106. <https://doi.org/10.5962/bhl.part.21869>
- Foureaux LV, Kato CM (1999) First record of *Takecallis taiwanus* (Takahashi) (Homoptera, Aphididae) in Brazil. *Anais da Sociedade Entomológica do Brasil* 28: 183–184. [In Portuguese] <https://doi.org/10.1590/S0301-80591999000100021>
- Froggatt WW (1904) Experimental work with the peach aphid (*Aphis persicae-niger*, Sm.). *Agricultural Gazette of New South Wales* 15: 603–612.
- Fullaway DT (1910) Report of the entomologist: general insect notes. *Annual Report of the Hawaiian Agricultural Experiment Station 1909*: 17–46.
- Ghosh AK (1980) Floral assemblage and faunal diversity in Aphidoidea (Homoptera, Aphididae) in Eastern India. *Bulletin of the Zoological Survey of India* 2: 171–176.
- Ghosh AK, Quednau FW (1990) Homoptera Aphidoidea. Subfamily Drepanosiphinae. The Fauna of India and adjacent countries. *Zoological Survey of India, Technical Monograph* 5: 229–240.
- Ghosh LK (1986) A conspectus of Aphididae (Homoptera) of Himachal Pradesh in Northwest Himalaya, India. *Zoological Survey of India, Technical Monograph* 16: 1–282.
- Giacalone I, Lampel G (1996) Pucerons (Homoptera, Aphidina) de la région insubrique tessinoise d'origine subméditerranéenne, méditerranéenne, est-européenne, asiatique et américaine. *Mitteilungen der Schweizerischen Entomologischen Gesellschaft* 69: 229–260.
- Gonzales, WL, Fuentes-Contreras E, Niemeyer HM (2000) Record of a new aphid introduced into Chile: *Takecallis taiwanus* (Takahashi) (Hemiptera, Aphididae, Drepanosiphinae). *Revista Chilena de Entomología* 26: 53–55.
- Haliday AH (1833) An essay on the classification of the parasitic Hymenoptera of Britain, which correspond with the Ichneumonones minuti of Linnaeus. *Entomological Magazine* 1: 480–491.
- Higuchi H (1968) A revision of the genus *Takecallis* Matsumura (Homoptera, Aphididae). *Insecta Matsumurana* 31: 25–33.
- Higuchi H (1972) A taxonomic study of the subfamily Callipterinae in Japan (Homoptera, Aphididae). *Insecta Matsumurana* 35: 19–126.
- Kimura M (1980) A simple method for estimating evolutionary rate of base substitutions through comparative studies of nucleotide sequences. *Journal of Molecular Evolution* 16: 111–120. <https://doi.org/10.1007/BF01731581>
- Kumar S, Stecher G, Li M, Knyaz C, Tamura K (2018) MEGA X: Molecular Evolutionary Genetics Analysis across computing platforms. *Molecular Biology and Evolution* 35: 1547–1549. <https://doi.org/10.1093/molbev/msy096>
- Lampel G, Meier W (2003) *Fauna Helvetica* 8. Hemiptera: Sternorrhyncha. part 1: Non-Aphididae. *Centre Suisse de Cartographie de la Faune, Schweizerische Entomologische Gesellschaft, Switzerland, Neuchâtel*, 312 pp.
- Lazzari SMN, Zonta de Carvalho RC, Cardoso JT (1999) *Takecallis arundinariae* (Essig) (Aphididae, Drepanosiphinae, Phyllaphidini): first record in Brazil and comparison to *Takecallis taiwanus* (Takahashi). *Revista Brasileira de Zoologia* 16: 865–870. <https://doi.org/10.1590/S0101-81751999000300023>

- Leclant F (1966) Contribution à l'étude des Aphidoidea du Languedoc meridional. Annales de la Société d'Horticulture et d'Histoire Naturelle de l'Hérault 106: 119–134.
- Lee Y, Lee S (2018) A review of the genus *Takecallis* Mastumura in Korea with the description of a new species (Hemiptera, Aphididae). ZooKeys 748: 131–149. <https://doi.org/10.3897/zookeys.748.23140>
- Limonta L (1990) Callaphididae (Aphidoidea) new to Italy. Bollettino di Zoologia Agraria e di Bachicoltura 22: 93–99.
- Lipp-Symonowicz B, Sztajnowski S, Wojciechowska D (2011) New commercial fibres called 'bamboo fibres' – Their structure and properties. Fibres & Textiles in Eastern Europe 19: 18–23.
- Liu CS (1975) Aphid parasites and their propagation in Taiwan. Journal of Agricultural Research of China 24: 62–84. [In Chinese]
- Maslyakov VY, Izhevsky SS (2011) Alien Phytophagous Insect Invasions in the European Part of Russia. IGRAS, Moscow, 272 pp. [In Russian]
- Matsumura S (1917) A list of the Aphididae of Japan, with description of new species and genera. The Journal of the College of Agriculture, Tohoku Imperial University, Sapporo, Japan 7(6): 351–414.
- Nieto Nafría JM, Mier Durante MP (1998) Fauna Ibérica (Vol. 11). Hemiptera, Aphididae I. Museo Nacional de Ciencias Naturales and Consejo Superior de Investigaciones Científicas. Madrid, 428 pp.
- Ohrnberger D (2002) Bamboos of the World: Annotated Nomenclature and Literature of the Species and Higher and Lower Taxa. Elsevier Science B.V., Amsterdam, 596 pp.
- Ortego J, Difabio ME, Mier Durante MP (2004) New records and actualized check-list of aphids (Hemiptera, Aphididae) from Argentina. Revista de la Sociedad Entomológica Argentina 63: 19–30.
- Pérez Hidalgo N, Martínez-Torres D, Collantes-Alegre JM, Villalobos Muller W, Nieto Nafría JM (2012) A new species of *Rhopalosiphum* (Hemiptera, Aphididae) on *Chusquea tomentosa* (Poaceae, Bambusoideae) from Costa Rica. ZooKeys 166: 59–73. <https://doi.org/10.3897/zookeys.166.2387>
- Peronti AL, Sousa-Silva CR (2002) Aphids (Hemiptera, Aphidoidea) of ornamental plants from Sao Carlos, Sao Paulo state, Brazil. Revista de Biologia Tropical 50: 137–144.
- Petrović-Obradović O, Radonjić A, Jovičić I, Petrović A, Kocić K, Tomanović Ž (2018) Alien species of aphids (Hemiptera, Aphididae) found in Serbia, new to the Balkan Peninsula. Phytoparasitica 46: 653–660. <https://doi.org/10.1007/s12600-018-0693-3>
- Piron PG (2009) New aphid (Aphidoidea) records for the Netherlands (1984–2005). Mitteilungen des Internationalen Entomologischen Vereins 33: 119–126.
- Pons X, Lumbierres B (2004) Aphids on ornamental shrubs and trees in an urban area of the Catalan coast: bases for an IPM programme. In: Simon JC, Dedryver CA, Rispe C, Hullé M (Eds) Aphids in a New Millennium, Institut National de la Recherche Agronomique, Paris, 359–364.
- Potenza MR (2005) Chemical control of the aphid *Takecallis taiwanus* (Takahashi) (Homoptera, Aphididae) in ornamental Bamboo (*Bambusa gracilis horti*). Arquivos do Instituto Biológico 72: 495–497.

- Potters G, Brems A, Valcke R, Dewil R, D'Haese L, Samson R, Gielis J (2009) Energy crops in Western Europe: Is bamboo an acceptable alternative. 8th World Bamboo Congress (Vol. 3). World Bamboo Organization, Bangkok, 22–24.
- Qiao GX, Zhang GX (2004) Review of the genus *Takecallis* Matsumura (Homoptera, Aphididae, Myzocallidinae) from China and description of one new species. *The Raffles Bulletin of Zoology* 52: 373–378.
- Qiu GX, Shen YK, Li DY, Wang ZW, Hudang QM, Yang DD, Gao AX (1992) Bamboo in subtropical eastern China. In: Long SP, Jones MB, Roberts MJ (Eds) *Primary Productivity of Grass Ecosystems of the Tropics and Sub-tropics*, Chapman and Hall, London, 159–188.
- Quilis Perez M (1931) *Especies nuevas de Aphidiidae espanoles* (Hym., Brac.). *Revista Espanola de Entomologia* 7: 25–84.
- Rakhshani E, Pons X, Lumbierres B, Havelka J, Pérez Hidalgo N, Tomanovič Ž, Starý P (2017) A new parasitoid (Hymenoptera, Braconidae, Aphidiinae) of the invasive bamboo aphids *Takecallis* spp. (Hemiptera, Aphididae) from western Europe. *Journal of Natural History* 51: 1237–1248. <https://doi.org/10.1080/00222933.2017.1327622>
- Rawat P, Narkhede SS, Rane AD, Mhaiske VM, Dalvi VV (2017) Allelopathic effect of leaf leachates of solid bamboo *Dendrocalamus stocksii* (Munro.) on growth and yield of *Eleusine coracana* L. (Gaertn.). *Indian Journal of Agroforestry* 19: 79–82.
- Raychaudhuri DN (1973) *Taxonomy of the Aphids of the Eastern Himalayas*. PL 480 Project Technical Report, University of Calcutta, Calcutta, 93 pp.
- Revathi TG, Remadevi OK (2011) Review on beneficial insects and report on ants occurring on cultivated bamboos in India. *Current Biotica* 5: 107–120.
- Ripka G (2008) Checklist of the Aphidoidea and Phylloxeroidea of Hungary (Hemiptera, Sternorrhyncha). *Folia Entomologica Hungarica* 69: 19–157.
- Saitou N, Nei M (1987) The Neighbor-Joining method: a new method for reconstructing phylogenetic trees. *Molecular Biology and Evolution* 4: 406–425. <https://doi.org/10.1093/oxfordjournals.molbev.a040454>
- Scurlock JMO, Dayton DC, Hames B (2000) Bamboo: An overlooked biomass resource? *Biomass and Bioenergy* 19: 229–244. [https://doi.org/10.1016/S0961-9534\(00\)00038-6](https://doi.org/10.1016/S0961-9534(00)00038-6)
- Shanmughavel P, Francis K (1996) Above ground biomass production and nutrient distribution in growing bamboo (*Bambusa bambos* (L.) Voss). *Biomass and Bioenergy* 10: 383–391. [https://doi.org/10.1016/0961-9534\(95\)00124-7](https://doi.org/10.1016/0961-9534(95)00124-7)
- Schilberszky K (1911) A bambusznádról. (Bamboo). *Termés Közlön* 43: 161–178. [In Hungarian]
- Simala M, Seljak G, Milek TM (2008) Leaf aphid *Takecallis taiwanus* Takahashi, 1926 (Hemiptera, Aphididae, Myzocallidinae) as a serious pest of golden bamboo (*Phyllostachys aurea* A. and C. Revière) in Croatia. *Glasiló Biljne Zastite* 8: 29–32.
- Stroyan HLG (1964) Notes on hitherto unrecorded or overlooked British aphid species. *Transactions of the Royal Entomological Society of London* 116: 29–72. <https://doi.org/10.1111/j.1365-2311.1964.tb00824.x>
- Suay-Cano VA, González-Funes P (1998) Estudio de los pulgones (Hemiptera, Aphididae) de la provincia de Valencia. III: Subfamilias Eriosomatinae, Anoeciinae, Thelaxinae, Myzocallidinae, Drepanosiphinae, Chaitophorinae, Lachninae y Pterocommatinae. *Boletín de la Asociación Española de Entomología* 22: 113–123.

- Takahashi R (1921) Aphididae of Formosa, Part I, Report of the Department of Agriculture Government Research Institute Formosa 20: 1–97.
- Takahashi R (1924) Aphididae of Formosa, Part III. Report of the Department of Agriculture Government Research Institute Formosa 10: 1–121.
- Takahashi R (1926) The aphids of *Myzocallis* infesting the bamboo. Proceedings of the Entomological Society of Washington 28(7): 159–162.
- Trejo-Loyo A, Peñaa-Martínez R, Villegas-Jiménez N (2004) Afdofauna (Hemiptera, Aphididae) de Cuernavaca, Morelos, Mexico. Folia Entomológica Mexicana 43: 191–202.
- Truett AA, Walker JA, Warman ML, Truett GE, Heeger P, Mynatt RL (2000) Preparation of PCR-quality mouse genomic DNA with Hot Sodium Hydroxide and Tris (HotSHOT). BioTechniques 29: 52–53. <https://doi.org/10.2144/00291bm09>
- Valenzuela I, Boulton A, Malipatil MB (2010) First record of *Takecallis arundinariae* (Essig) (Hemiptera, Aphididae) from Australia. General & Applied Entomology 39: 23–25.
- van der Lugt P, van den Dobbelssteen AAJF, Janssen JJA (2006) An environmental, economic and practical assessment of bamboo as a building material for supporting structures. Construction and Building Materials 20: 648–656. <https://doi.org/10.1016/j.conbuildmat.2005.02.023>
- Wright L (2006) Worldwide commercial development of bioenergy with a focus on energy crop-based projects. Biomass and Bioenergy 30: 706–714. <https://doi.org/10.1016/j.biombioe.2005.08.008>