

First record of the family Colinauropodidae (Myriapoda, Pauropoda) in China, with the description of three new species

Yun Bu¹

¹ Natural History Research Center, Shanghai Natural History Museum, Shanghai Science & Technology Museum, Shanghai 200041, China

Corresponding author: Yun Bu (buy@sstm.org.cn)

Academic editor: Pavel Stoev | Received 28 April 2020 | Accepted 30 May 2020 | Published 8 July 2020

<http://zoobank.org/7E8524DC-A5E1-4F8A-BE1F-089CDFEE7F8>

Citation: Bu Y (2020) First record of the family Colinauropodidae (Myriapoda, Pauropoda) in China, with the description of three new species. ZooKeys 947: 53–70. <https://doi.org/10.3897/zookeys.947.53723>

Abstract

The pauropod family Colinauropodidae Scheller, 1985 is recorded from China for the first time. Three new species of the genus *Colinauropus* Remy, 1956 are described: *Colinauropus chinensis* **sp. nov.** and *C. chongzhoui* **sp. nov.** from Jiangsu Province, and *C. foliosus* **sp. nov.** from Sichuan Province. They can be easily separated from similar species by the number and the shape of sclerotized plates on the tergites, setae on the body and the anal plate. A key for all species of the genus is provided.

Keywords

anal plate, bothriotricha, pauropod, sclerotized plate, taxonomy

Introduction

The family Colinauropodidae Scheller, 1985 includes the single genus *Colinauropus* Remy, 1956 and contains three species in the world: *Colinauropus regis* Remy, 1956 from Réunion and Mauritius (Remy 1956, 1959), *C. schelleri* Hagino, 1991 from Japan (Hagino 1991, 2005), and *C. haginoi* Scheller, 2009 from Philippines (Scheller, 2009). Their most charming character lies in the tergites which split into several distinctly sclerotized plates of irregular shape (Scheller 2011).

In the original description, the genus *Colinauropus* was considered to be affiliated with species of the family Brachypauropodidae Silvestri, 1902 according to the fragmented tergites and the shape of anal plate (Remy 1956). Its taxonomic position was reconsidered and placed in the family Pauropodidae Lubbock 1867, under the new subfamily Colinauropodinae, which was supposed to be closely related to the subfamily Scleropauropodinae (Scheller 1985). In the latest classification system, the subfamily Colinauropodinae was upgraded to family Colinauropodidae (Scheller 2009, 2011).

The purposes of this study are 1) to record the occurrence of family Colinauropodidae Scheller, 1985 in China for the first time; 2) to describe three new species of the genus *Colinauropus* Remy, 1956 from China; 3) to give a key to the species of the genus.

Materials and methods

All pauropods were collected using a Tullgren's funnel. The specimens were sorted under a stereomicroscope and preserved in 80% alcohol. They were mounted on slides using Hoyer's solution and dried in an oven at 50 °C. Observations were performed under a phase contrast microscope (Leica DM 2500). Photos were taken using a digital camera (Leica DMC 4500). Line drawings were made using a drawing tube. All specimens were deposited in the collection maintained by the Shanghai Natural History Museum.

Abbreviations used in the descriptions follow Qian et al. (2018). Absolute lengths of all other body parts are given in mm and μm . Otherwise, the text refers to relative lengths. For the description of the new species, measurements and indices of paratypes are given in brackets.

Results

Taxonomy

Family Colinauropodidae Scheller, 1985

Genus *Colinauropus* Remy, 1956

Type species. *Colinauropus regis* Remy, 1956.

Diagnosis. Body fusiform; head and pygidium free; tergites divided into sclerotized coarse plates, partly of irregular shape; stalk of antennal globulus g shorter than globulus itself; adults with first and last pair of legs 5-segmented, remaining pairs 6-segmented; pygidial sternum with two pairs of setae $b_1 + b_2$ (Scheller 2011).

Distribution. Ethiopian, Palearctic, and Oriental regions.

***Colinauropus chinensis* sp. nov.**

<http://zoobank.org/DFA53888-8023-4745-B84C-82E51BCB5E57>

Figures 1–3

Material examined. *Holotype*, female adult with 9 pairs of legs (slide no. JS-WX-PA2017033), China, Jiangsu Province, Wuxi City, Daji Mountain, extracted from soil samples in bamboo forest, elev. 5 m, 31°32'N, 120°12'E, 9-X-2017, coll. Y. Bu. *Paratypes*, 2 female adults with 9 pairs of legs (slides no. JS-WX-PA2017031, JS-WX-PA2017032), same data as holotype; 1 female adult with 9 pairs of legs (slide no. JS-WX-PA2018006), same locality as holotype, 9-X-2018, coll. Y. Bu.

Diagnosis. *Colinauropus chinensis* sp. nov. is characterized by the cylindrical, annulate setae on head, antennae and tergites; tergite I without distinct sclerotized plates; tergite II with 2 large and 4 small sclerotized plates; tergites III–V each with 4 large and 4 small plates; tergite VI with 2 large plates; seta *st* on tergum of pygidium cylindrical; bothriotrichum T_3 with thicker axis and dense tufted pubescence distally.

Description. Adult body length (0.88–) 0.96 (–0.98) mm ($N = 4$); body white-yellow in alcohol, sclerotized plates on tergites brown (Fig. 2A).

Head (Figs 1A, 2D). Dorsal setae cylindrical, annulate, first and second rows shorter than posterior rows. Relative lengths of setae, 1st row: $a_1 = 10$, $a_2 = 8$ (–9); 2nd row: $a_1 = 13$ (–14), $a_2 = 7$ (–9), $a_3 = 7$; 3rd row: $a_1 = (18–) 20$, $a_2 = (23–) 24$; 4th row: $a_1 = 16$ (–17), $a_2 = 16$ (–17), $a_3 = 22$ (–25), $a_4 = 14$ (–16); lateral group setae $l_1 = 21$ (–26), $l_2 = 26$ (–31), $l_3 = 29$ (–35); the ratio $a_1/a_1 - a_1$ in 1st row 0.7 (–0.9), 2nd row 0.5, 3rd row 1.2 and 4th row 0.7 (–0.8). Temporal organs oval in dorsal view, their length 0.8 of their shortest distance apart. Pistil present. Head cuticle faintly granular.

Antennae (Figs 1E, 2B, C). Antennal segments 1–3 with 2, 2, 3 short, cylindrical, annulate setae respectively, and 1 rudimentary setae present on segment 3. Antennal segment 4 with 4 cylindrical setae; relative lengths of setae: $p = 10$, $p' = 6$, $p'' = 5$, $r = 5$; tergal seta p (1.3–) 1.4 times as long as tergal branch t ; the latter cylindrical, 1.7 (–1.8) times as long as its greatest diameter and 0.8 of sternal branch s , which itself is 1.6 times as long as its greatest diameter. Seta q cylindrical, annulate, 0.9 of s . Relative lengths of flagella (base segments included) and base segments: $F_1 = 100$, $bs_1 = 8$ (–11); $F_2 = (41–) 49$, $bs_2 = (5–) 6$; $F_3 = (84–) 92$, $bs_3 = 9$ (–10). F_1 (6.6–) 7.2 times as long as t , F_2 and F_3 (2.3–) 2.7 and (4.8–) 5.1 times as long as s respectively. Distal calyces spherical; apex of flagella fusiform, with a short lateral flap. Globulus g 1.7 times as long as wide; about 12 bracts, capsule spherical; width of g (0.5–) 0.6 of the greatest diameter of t . Antennal cuticle granulated.

Trunk. Setae on collum segment cylindrical, annulate; sublateral setae length (20–) 22 μ m, (1.9–) 2.0 times as long as submedian setae; sternite process triangular, furcate and granulated; appendages barrel shaped (Fig. 2E). Tergite I with 4+4 short, cylindrical setae (14–15 μ m), posteriorly with two patches of thickened cuticles but not form distinct sclerotized plates (Fig. 2F); Tergite II with 6+6 setae (9–20 μ m), 4 small anterior and 2 large posterior sclerotized plates (Figs 1B, 2G); Tergites III–V each with 6+6 setae (9–21 μ m), 4 large and 4 small sclerotized plates (Fig. 2H–J); Ter-




Figure 1. *Colinauropus chinensis* sp. nov. **A** head, dorsal view, right side **B** tergite II, right side **C** T_5 **D** T_3 **E** right antenna, tergal view **F** tergum of pygidium **G** sternum of pygidium and anal plate. Scale bars: 20 μm.




Figure 2. *Colinauropus chinensis* sp. nov. **A** habitus, tergal view **B** F_1 and F_3 of right antenna **C** globulus and F_2 of right antenna **D** head, dorsal view **E** collum segment, sternal view **F** tergite I **G** tergite II **H** tergite III **I** tergite IV **J** tergite V **K** tergite VI and tergum of pygidium **L** sternum of pygidium and anal plate. Scale bars: 100 μm (**A**); 20 μm (**B**–**L**).

gite VI with 4+2 setae and 2 large plates (Fig. 2K), posterior setae 10 (–11) μm long, their mutual distance 20 (–23) μm (Fig. 2K). Sclerotized plates with dense, brown granules, diameter 1.5–3.2 μm (Fig. 1B). Other areas of cuticle on tergites with pale and fine granules.

Bothriotricha. Relative lengths: $T_1 = 100$, $T_2 = (110\text{--}) 117$, $T_3 = (122\text{--}) 128$, $T_4 = 133(140)$, $T_5 = (167\text{--}) 178$. T_1 , T_2 , T_4 and T_5 long, with short erect and oblique pubescence on axis (Fig. 1C). T_3 with thicker axis and dense tufted pubescence distally (Fig. 1D).

Legs. First and last pair of legs 5-segmented, others 6-segmented (Fig. 3A–C). Setae on coxa and trochanter of legs 1–8 cylindrical, annulate (Fig. 3A, C), length 13 (–14) μm and 18 (–20) μm respectively. Setae on coxa of leg 9 cylindrical, annulate, length (15–) 17 μm (Fig. 3B, D). Setae on trochanter of leg 9 furcate, with subcylindrical, annulate, blunt branches, shorter one about (0.6–) 0.7 of longer one (Fig. 3B, D). Tarsi 1–8 with short, annulate distal seta (6 μm) only (Fig. 3A, C). Tarsus of leg 9 tapering, 35 μm in length, 3.2 (–3.5) times as long as its greatest diameter (Fig. 3B), proximal seta slender, pointed, striate, 10 (–13) μm in length; distal seta cylindrical, annulate, 6 (–7) μm in length, about 0.2 of the tarsal length. Cuticle of tarsus pubescent.

Pygidium. Tergum. Posterior margin waved. Relative lengths of setae: $a_1 = 10$, $a_2 = 13$, $a_3 = 15$, $st = 10$ (–12). Setae distinctly differentiated, a_1 short, clavate, pubescent; a_2 and a_3 slender and pubescent (Figs 1F, 2K); st thick and pubescent (Figs 1G, 2K). Distance a_1 – a_1 as same long as a_1 ; distance a_1 – a_2 2.0 (–2.5) times as long as a_2 – a_3 ; distance st – st (1.5–) 1.6 times as long as st and 1.6 (–1.8) times as long as distance a_1 – a_1 .

Sternum (Figs 1G, 2L). Posterior margin with a deep indentation between b_1 . Relative lengths of setae ($a_1=10$): $b_1 = 33(35)$, $b_2 = 13$ (–15). Seta b_1 cylindrical, thick and annulate; b_2 slender and short, pubescent. Distance b_1 – b_1 (0.7–) 0.8 of length of b_1 ; distance b_1 – b_2 (0.7–) 0.9 of b_2 .

Anal plate linguiform, glabrous, 2.0 times longer than broad, lateral margins concave in anterior part, posterior margin with three small lobes; two pairs of appendages present: inner one tiny and conical; outer one cylindrical and longer, (0.4–) 0.5 of the length of plate and with short pubescence (Figs 1G, 2L).

Etymology. The species is named after China where the type specimens were collected.

Distribution. China (Jiangsu). Only known from the type locality.

Remarks. *Colinauropus chinensis* sp. nov. is most similar to *C. haginoi* Scheller, 2009 from Philippines in the similar shape of the anal plate and absence of sclerotized plates on tergite I. They can be easily distinguished by the number of sclerotized plates on tergites II and VI (6 and 2 in *C. chinensis* sp. nov., vs. 8 and 4 in *C. haginoi*), length of setae on collum segment (sublateral setae 1.9–2.0 times as long as submedian setae in *C. chinensis* sp. nov. vs. 3.2 times in *C. haginoi*), and the shape of T_3 (subcylindrical, not clavate in *C. chinensis* sp. nov. vs. proximal half distinctly clavate in *C. haginoi*).




Figure 3. *Colinauropus chinensis* sp. nov. **A** leg 1 **B** leg 9 **C** leg 4 **D** coxa and trochanter of leg 9. Scale bars: 20 μ m.

***Colinauropus chongzhoui* sp. nov.**

<http://zoobank.org/9DB912C3-DB79-4AC1-8649-AAAC2985E274>

Figures 4–6

Material examined. *Holotype*, female adult with 9 pairs of legs (slide no. JS-WX-PA2018007), China, Jiangsu Province, Wuxi City, Daji Mountain, extracted from

soil samples in bamboo forest, elev. 5 m, 31°32'N, 120°12'E, 8-X-2018, coll. Y. Bu. Non-type specimens, 1 juvenile with 8 pairs of legs (slides no. JS-WX-PA2017034), 2 juveniles with 6 pairs of legs (slides no. JS-WX-PA2018008, JS-WX-PA2018009), same data as holotype.

Diagnosis. *Colinauropus chongzhoui* sp. nov. is characterized by the slender, annulate-striate setae on head, antennae and tergites; tergite I with 1 large sclerotized plate; tergite II with 6 small and 2 large sclerotized plates; tergites III–V each with 4 large and 4 small plates; tergite VI with 2 large plates; seta *st* on tergum of pygidium clavate; bothriotrichum T_3 brush-shaped, with branched pubescence distally.

Description. Adult body length 0.97 mm ($N = 1$); body white-yellow in alcohol, sclerotized plates on tergites brown (Fig. 5A).

Head (Figs 4A, 5C). Dorsal setae short, cylindrical, annulate-striate, except seta a_3 of second row which is slender and tapering. Relative lengths of setae, 1st row: $a_1 = 10$, $a_2 = 10$; 2nd row: $a_1 = 8$, $a_2 = 14$, $a_3 = 14$; 3rd row: $a_1 = 9$, $a_2 = 10$; 4th row: $a_1 = 12$, $a_2 = 14$, $a_3 = 21$, $a_4 = 12$; lateral group setae $l_1 = 23$, $l_2 = 21$, $l_3 = 19$; the ratio $a_1/a_1 - a_1$ in 1st row 1.5, 2nd row 0.6, 3rd row 0.9 and 4th row 0.7. Temporal organs oval in dorsal view, their length 1.1 times as long as their shortest distance apart. Pistil present. Head cuticle with dense granules.

Antennae (Figs 4E, 5B). Antennal segments 1–3 with 2, 2, 3 short, cylindrical, annulate setae respectively, and 1 rudimentary seta present on segment 3. Antennal segment 4 with 4 cylindrical setae and rudimentary seta *u*; relative lengths of setae: $p = 10$, $p' = 6$, $p'' = 6$, $r = 6$, $u = 1$; tergal seta *p* 1.1 times as long as tergal branch *t*; the latter cylindrical, 2.2 times as long as its greatest diameter and 0.9 of sternal branch *s*, which itself is 1.9 times as long as its greatest diameter. Seta *q* cylindrical, annulate, 1.3 times as long as *s*. Relative lengths of flagella (base segments included) and base segments: $F_1 = 100$, $bs_1 = 10$; $F_2 = 52$, $bs_2 = 5$; $F_3 = 88$, $bs_3 = 9$. F_1 6.4 times as long as *t*, F_2 and F_3 2.9 and 4.9 times as long as *s* respectively. Distal calyces spherical; apex of flagella fusiform, with a short lateral flap. Globulus *g* 1.7 times as long as wide; about 12 bracts, capsule spherical; width of *g* 0.5 of the greatest diameter of *t*. Antennal cuticle densely granulated.

Trunk. Setae on collum segment cylindrical, annulate; sublateral setae length 23 μ m, 2.1 times as long as submedian setae; sternite process triangular, furcate and granulated; appendages cylindrical and tapering (Fig. 5D). Tergite I with 4+4 cylindrical setae (12–13 μ m) and 1 large sclerotized plate (Fig. 5E) (2 plates in juveniles, Fig. 5F); Tergite II with 6+6 setae (12–23 μ m), 6 small anterior and 2 large posterior sclerotized plates (Figs 4B, 5G); Tergites III–V each with 6+6 setae (6–27 μ m), 4 large and 4 small sclerotized plates (Fig. 5H–J); Tergite VI with 4+2 setae and 2 large plates (Fig. 5K), posterior setae 23 μ m long, their mutual distance 18 μ m (Fig. 5K). Sclerotized plates with dense, brown granules, diameter 2–4 μ m, and each granule with one short straight apical hair (Fig. 4B). Other areas of cuticle on tergites with coarse granules.




Figure 4. *Colinauropus chongzhoui* sp. nov. **A** head, dorsal view, right side **B** tergite II, right side **C** T_5 **D** T_3 **E** right antenna, tergal view **F** tergum of pygidium **G** sternum of pygidium and anal plate. Scale bars: 20 μm.




Figure 5. *Colinauropus chongzhoui* sp. nov. **A** habitus, tergal view **B** left antenna, sternal view **C** head, dorsal view **D** collum segment, sternal view **E** tergite I of adult **F** tergite I of juvenile **G** tergite II **H** tergite III **I** tergite IV **J** tergite V **K** tergite VI. Scale bars: 100 μm (**A**); 20 μm (**B–K**).

***Bothriotricha*.** Relative lengths: $T_1 = 100$, $T_2 = 113$, $T_3 = 86$, $T_4 = 118$, $T_5 = 167$. T_1 , T_2 , T_4 and T_5 thin, long, with short erect or oblique pubescence on axis (Fig. 4C). T_3 brush-shaped, with thicker axis and branched pubescence in distal 2/3 (Figs 4D, 6E).

***Legs*.** First and last pair of legs 5-segmented, others 6-segmented. Setae on coxa and trochanter of legs 1–8 cylindrical, annulate, length 13–15 μm and 16–20 μm respectively (Fig. 6D). Setae on coxa of leg 9 cylindrical, annulate, length 12 μm (Fig. 6C). Seta




Figure 6. *Colinauropus chongzhoui* sp. nov. **A** sternum of pygidium and anal plate **B** tarsus of leg 9 **C** coxa and trochanter of leg 9 **D** coxa and trochanter of leg 1 **E** T_3 . Scale bars: 20 μm .

on trochanter of leg 9 furcate, with two subcylindrical, annulate, blunt branches, shorter one about 0.5 of longer one (Fig. 6C). Tarsi 1–8 with short annulate distal seta (8 μm) only. Tarsus of leg 9 tapering, 40 μm in length, 3.6 times as long as its greatest diameter (Fig. 6B), proximal seta slender, pointed, striate, 11 μm in length; distal one cylindrical, annulate, 9 μm in length, about 0.2 of the tarsal length. Cuticle of tarsus pubescent.

Pygidium. Tergum. Posterior margin blunt triangular. Relative lengths of setae: $a_1 = 10$, $a_2 = 12$, $a_3 = 16$, $st = 5$. Setae distinctly differentiated, a_1 cylindrical, pubescent; a_2 and a_3 slender, pubescent (Fig. 4F); st short, clavate, pubescent (Figs 4G, 6A). Distance $a_1 - a_1$ 0.7 of length of a_1 ; distance $a_1 - a_2$ 1.6 times as long as $a_2 - a_3$; distance $st - st$ 2.0 times as long as st and 1.5 times as long as distance $a_1 - a_1$.

Sternum (Figs 4G, 6A). Posterior margin with one lower indentation between b_1 . Relative lengths of setae ($a_1 = 10$): $b_1 = 28$, $b_2 = 12$. Seta b_1 cylindrical, thick, annulate; b_2 slender, short, pubescent. Distance $b_1 - b_1$ 0.8 of length of b_1 ; distance $b_1 - b_2$ 0.9 of b_2 .

Anal plate linguiform, 1.7 times longer than broad; a pair of clavate appendage inserted posteriorly, 0.4 of the length of plate, and with short pubescence (Figs 4G, 6A).

Etymology. The species is dedicated to the honor of the late Professor Chongzhou Zhang (1930–2014) who was an eminent zoologist from Institute of Zoology, Chinese Academy of Sciences, for his great contribution to the knowledge of Myriapoda of China (Stoev et al. 2014).

Distribution. China (Jiangsu). Only known from the type locality.

Remarks. *Colinauropus chongzhoui* sp. nov. is similar to *Colinauropus regis* Remy, 1956 in the shape of anal plate. They can be easily distinguished by the number of sclerotized plates on tergite I (1 large plate in *C. chongzhoui* sp. nov. vs. 2 in *C. regis*) and tergite II (8 in *C. chongzhoui* sp. nov., vs. 6 in *C. regis*), shape of setae on tergites (slender and striate in *C. chongzhoui* sp. nov. vs. clavate and pubescent in *C. regis*), and the shape of seta a_1 on pygidium (tapering in *C. chongzhoui* sp. nov. vs. clavate in *C. regis*).

***Colinauropus foliosus* sp. nov.**

<http://zoobank.org/867E5192-CE51-4339-947B-7C17D668C86E>

Figures 7–9

Material examined. *Holotype*, female adult with 9 pairs of legs (slide no. SC-PA2017002), China, Sichuan Province, Ganzi Tibetan Autonomous Region, Kangding City, Yala town, 30°06'N, 101°57'E, elev. 3100 m, soil samples from mixed forest, 11-VIII-2017, coll. C.W. Huang. *Paratypes*, 1 male adult with 9 pairs of legs (slides no. SC-PA2017001) and 1 female adult with 9 pairs of legs (slide no. SC-PA2017003), same data as holotype.

Diagnosis. *Colinauropus foliosus* sp. nov. is characterized by the leaf-shaped pubescent setae on head and tergites; tergite I with one large sclerotized plate; tergites II–IV each with 4 large and 4 small plates; tergite V with 4 large and 2 small middle sclerotized plates; tergite VI with 2 large plates; granules on plates ovoid, each inserted with one fine hair; seta st on tergum of pygidium clavate; bothriotrichum T_3 with thick axis and dense tufted pubescence distally.

Description. Adult body length 1.28 (–1.32) mm ($N = 3$); body white-yellow in alcohol, sclerotized plates on tergites brown (Fig. 8A).

Head (Figs 7A, 8D). Dorsal setae distinctly differentiated, on first and second rows cylindrical to tapering; on third and fourth rows leaf-shaped and with long pubescence; seta a_3 of second row slender and tapering. Relative lengths of setae, 1st row: $a_1 = 10$, $a_2 = 10$ (–12); 2nd row: $a_1 = 10$ (–11), $a_2 = (11–) 12$, $a_3 = 12$ (–13); 3rd row: $a_1 = (18–) 20$, $a_2 = 18$ (–20); 4th row: $a_1 = 13$ (–16), $a_2 = (15–) 17$, $a_3 = 20$ (–23), $a_4 = 16$ (–17); lateral group setae $l_1 = 18$ (–24), $l_2 = 18$ (–23), $l_3 = 25$ (–32); the ratio $a_1/a_1 - a_1$ in 1st row (1.6–) 1.7, 2nd row (0.7–) 0.8, 3rd row 1.0 (–1.1) and 4th row 0.7 (–0.8). Temporal organs oval in dorsal view, their length (0.8–) 0.9 of their shortest distance apart. Pistil present. Head cuticle with coarse granules.

Antennae (Figs 7C, 8B, C). Antennal segments 1–3 with 2, 2, 3 short cylindrical pubescent setae respectively, and 1 rudimentary seta present on segment 3. Antennal segment 4 with 4 tapering setae and a short, rudimentary u ; relative lengths of setae: $p = 10$, $p' = 7$ (–8), $p'' = (6–) 7$, $r = 5$ (–6), $u = 1$; tergal seta p (0.9 of –) 1.0 times as long as tergal branch t ; the latter cylindrical, 1.8 (–2.0) times as long as its greatest diameter and 0.7 (–0.9) of sternal branch s , which itself about 2.0 times as long as its greatest diameter. Seta q cylindrical, annulate, 0.8 (–1.0 times as long as) of s . Relative lengths of flagella (base segments included) and base segments: $F_1 = 100$, $bs_1 = 8$ (–11); $F_2 = (35–) 42$, $bs_2 = (4–) 5$; $F_3 = (78–) 93$, $bs_3 = 7$ (–9). F_1 (5.8–) 8.6 times as long as t , F_2




Figure 7. *Colinauropus foliosus* sp. nov. **A** head, dorsal view, left side **B** tergite II, right side **C** right antenna, tergal view **D** T_3 **E** T_5 **F** tergum of pygidium **G** sternum of pygidium and anal plate. Scale bars: 20 μm.

$a_{nd F3}$ 2.0 (–2.1) and 4.4 (–4.8) times as long as s respectively. Distal calyces spherical; apex of flagella fusiform, on F_1 and F_3 with a short lateral flap. Globulus g 1.7 times as long as wide; about 12 bracts, capsule spherical; width of g (0.4–) 0.6 of the greatest diameter of t . Antennal cuticle granulated.




Figure 8. *Colinauropus foliosus* sp. nov. **A** habitus, tergal view **B** terminal part of F_1 **C** sternal branch of left antenna, show F_2 , F_3 and globulus **D** head, dorsal view **E** collum segment, sternal view **F** tergite I **G** tergite II **H** tergite III **I** tergite IV **J** tergite V **K** tergite VI and tergum of pygidium **L** sternum of pygidium and anal plate. Scale bars: 100 μm (**A**); 20 μm (**B–K**).

Trunk. Setae on collum segment cylindrical, annulate; sublateral setae length 22 (–34) μm , (1.4–) 1.7 times as long as submedian setae; sternite process triangular, furcate and granulated; appendages tapering (Fig. 8E). Tergite I with 4+4 leaf-shaped setae (22–27 μm) and 1 large sclerotized plate (Fig. 8F); Tergites II–IV each with 6+6 setae (18–40 μm), 4 large and 4 small sclerotized plates (Figs 7B, 8G–I); Tergite V with 6+6 slender setae (20–40 μm), 4 large and 2 small sclerotized plates, posterior plates square (Figs 8J, 9A); Tergite VI with 4+2 setae and 2 large plates (Fig. 8K), posterior setae 35 μm long, their mutual distance 24 (–26) μm (Fig. 8K). Sockets of some setae on tergites and bothriotricha with distinct thickened cuticle surrounded (Figs 7B, D, E, 8F–J). Sclerotized plates with ovoid, brown granules, diameter 1.5–5.0 μm and each with one long curved hair (Figs 7B, 9A). Cuticle granulated or pubescent.

Male genital papillae (Fig. 9E) glabrous, subuliform, 1.5 times as long as greatest diameter; seta 0.5 of the length of papilla. Seta on coxa of leg 2 in male with two adjacent setae (only 1 thick setae in female, 20–25 μm), both cylindrical and annulate, one thick and short, 17 μm in length, another slender and longer, 20 μm in length (Fig. 9E).

Bothriotricha. Relative lengths: $T_1 = 100$, $T_2 = (110\text{--}) 100$, $T_3 = (95\text{--}) 105$, $T_4 = 114$ (–120), $T_5 = (115\text{--}) 120$. T_1 , T_2 , T_4 and T_5 thin, long, with short erect pubescence on axes (Fig. 7E). T_3 cylindrical, with thicker axis and dense tufted pubescence in distal 2/3 part (Fig. 7D).

Legs. First and last pair of legs 5-segmented, others 6-segmented (Fig. 9C, D). Setae on coxa and trochanter of legs 1–8 cylindrical, annulate (Fig. 9B), length 23 (–26) μm and 23 (–27) μm respectively. Seta on coxa of leg 9 cylindrical, annulate, length 22 (–25) μm (Fig. 9D). Seta on trochanter of leg 9 furcate, with subcylindrical, annulate, blunt branches, shorter branch about 0.5 of longer one (Fig. 9D). Tarsi 1–8 with short, annulate distal seta (6–8 μm) only (Fig. 9C). Tarsus of leg 9 tapering, 48 (–55) μm in length, 3.7 (–4.2) times as long as its greatest diameter (Fig. 9F), proximal seta slender, pointed, pubescent, 12 (–15) μm in length; distal one cylindrical, annulate, 6 (–8) μm in length, about 0.1 of the tarsal length. Cuticle of tarsus pubescent.

Pygidium. Tergum. Posterior margin straight. Relative lengths of setae: $a_1 = 10$, $a_2 = (8\text{--}) 9$, $a_3 = (12\text{--}) 16$, $st = (7\text{--}) 9$. Setae a_1 cylindrical, pubescent; a_2 and a_3 slender and pubescent (Figs 7F, 8K); st clavate, pubescent (Fig. 7G). Distance $a_1\text{--}a_1$ (0.6–) 0.8 of a_1 ; distance $a_1\text{--}a_2$ 2.0 (–2.5) times as long as $a_2\text{--}a_3$; distance $st\text{--}st$ (1.5–) 1.6 times as long as st and (1.7–) 2.0 times as long as distance $a_1\text{--}a_1$.

Sternum (Figs 7G, 8L). Posterior margin straight between b_1 . Relative lengths of setae ($a_1=10$): $b_1 = (25\text{--}) 29$, $b_2 = 14$ (–15). Seta b_1 cylindrical, thick, annulate; b_2 tapering, short, pubescent. Distance $b_1\text{--}b_1$ (0.7–) 0.8 of length of b_1 ; distance $b_1\text{--}b_2$ (0.6–) 0.7 of b_2 .

Anal plate round, glabrous, 1.5 times longer than broad, lateral margins bulged in middle part, posterior part divided into two round, pubescent branches, two tiny lobes present at inner side (Figs 7G, 8L).

Etymology. The species name “*foliosus*” from the Latin “*foliose*”, leaf-shaped, referring to the leaf-shaped setae on head and tergites.

Distribution. China (Sichuan). Only known from the type locality.




Figure 9. *Colinauropus foliosus* sp. nov. **A** posterior plate on tergite V, left side **B** coxa and trochanter of leg 1 **C** leg 8 **D** leg 9 **E** male genital papillae and coxa of leg 2 **F** tarsus of leg 9. Scale bars: 20 μ m.

Remarks. *Colinauropus foliosus* sp. nov. differs from all other congeners by having 6 sclerotized plates on tergite V, and the posterior two square-shaped, compared with 4 or 8 irregular plates in congeners. It is similar to *C. regis* Remy, 1956 in the leaf-shaped setae on tergites, but they can be easily distinguished by the number of sclerotized

plates on tergite I and II (1 and 8 in *C. foliosus* sp. nov. vs. 2 and 6 in *C. regis*), shape of the setae in the posterior two rows of the head (leaf-shaped in *C. foliosus* sp. nov. vs. cylindrical in *C. regis*), and the shape of anal plate (with two round posterior branches in *C. foliosus* sp. nov. vs. with two clavate appendages in *C. regis*).

Key to the species of the genus *Colinauropus* Remy, 1956

- 1 Tergite I without sclerotized plates, at most with two posterior thickened patches.....2
- Tergite I with distinct sclerotized plates.....3
- 2 Tergite II with 4 large and 4 small sclerotized plates, tergite VI with 4 plates..... ***C. haginoui* Scheller, 2009 (Philippines)**
- Tergite II with 2 large and 4 small sclerotized plates, tergite VI with 2 plates..... ***C. chinensis* sp. nov. (China)**
- 3 Tergite I with 1 large sclerotized plate4
- Tergite I with 2 sclerotized plates5
- 4 Setae on head and tergite cylindrical, tergite II with 2 large and 6 small sclerotized plates..... ***C. chongzhoui* sp. nov. (China)**
- Setae on head and tergite leaf-shaped, tergite II with 4 large and 4 small sclerotized plates..... ***C. foliosus* sp. nov. (China)**
- 5 Tergites II and V with 6 and 8 sclerotized plates respectively, anal plate rounded with 2 posterior appendages, setae on tergites II–V clavate..... ***C. regis* Remy, 1956 (Réunion, Mauritius)**
- Tergites II and V with 8 and 4 sclerotized plates respectively, anal plate indented mediodistally without appendages, setae on tergites II–V cylindrical...
..... ***C. schelleri* Hagino, 1991 (Japan)**

Discussion

The genus *Colinauropus* Remy, 1956 is well defined by the presence of sclerotized plates on the tergites. The number of plates on tergites I, II, V and VI, which varies from 1 to 8, are good characters for species identification, while tergites III and IV always have 8 plates in all species. The shape and arrangement of the plates are also taxonomically informative for species definition. On tergite I, the plates are absent or at most with small patches of thickened cuticle posteriorly (*C. haginoui*, *C. chinensis* sp. nov.), with 1 complete large plate (*C. chongzhoui* sp. nov., *C. foliosus* sp. nov.) or with 2 axially separated plates (*C. regis*, *C. schelleri*). On tergite II, 4 small anterior plus 2 large posterior plates are present in *C. regis* and *C. chinensis* sp. nov., 6 small anterior plus 2 large posterior plates are present in *C. chongzhoui* sp. nov., while there are 4 small plus 4 large plates in the remaining three species. On tergite V, the number of plates can be 4 in *C. schelleri*, 4 large plus 2 small middle plates in *C. foliosus* sp. nov., and 4 large plus 4 small plates in others. On tergite VI, all species have 2 large plates, except *C. haginoui* which has 4 plates. The shapes of plates are usually ovoid, round, sub-triangular, or irregular, while the two pos-

terior large plates on tergite V are nearly square-shaped in *C. foliosus* sp. nov. Variation of plates within a species has *never been reported in former studies but is observed here in C. chongzhoui* sp. nov., which exhibits 2 plates on tergite I in juveniles vs. 1 complete plate in adults. Thus, caution is advised when describing species of this genus, which should be based on fully mature specimens. As an additional taxonomic character, the bothriotrichum T_3 is also well differentiated and nicely separates species. The most informative characters are the shape and appendages of the anal plates. The six known species of the genus *Colinauropus* Remy, 1956 can be distinguished by the key provided above.

Acknowledgements

I thank Dr Osami Nakamura (Japan) and Prof. Claude Tautel (France) for their help in sending some reprints of references, Dr Chang-Yuan Qian also shared several references, Mr Chengwang Huang collected the specimens from Sichuan. I also cordially thank Dr Nikolaus Szucsich (Austria) and another anonymous referee for their valuable comments during the review of the manuscript. This research is supported by the National Natural Science Foundation of China (no. 31772509) and the Research Foundation of Shanghai Science and Technology Museum.

References

- Hagino Y (1991) New species of the family Pauropodidae (Pauropoda) from Central Japan. *The Canadian Entomologist* 123(5): 1009–1045. <https://doi.org/10.4039/Ent1231009-5>
- Hagino Y (2005) Contribution to the knowledge of the Japanese pauropod fauna I: A cumulative list of identified pauropod specimens from Japan during 1985–2003. *Natural History Research* 8(2): 15–51.
- Qian CY, Bu Y, Dong Y, Luan YX (2018) Study on the Pauropoda from Tibet, China. Part I. The genera *Decapauropus* and *Hemipauropus* (Myriapoda). *ZooKeys* 754: 33–46. <https://doi.org/10.3897/zookeys.754.24210>
- Remy PA (1956) Un nouveau Pauropode de l'île de la Réunion: *Colinauropus regis* n. g., n. sp. *Bulletin du Muséum national d'Histoire naturelle* 28(1): 119–123.
- Remy PA (1959) Pauropodes de l'île Maurice. *The Mauritius Institute Bulletin* 5(5): 149–194.
- Scheller U (1985) On the classification of the family Brachypauropodidae (Myriapoda; Pauropoda). *Bijdragen tot de Dierkunde* 55(1): 202–208.
- Scheller U (2009) Records of Pauropoda (Pauropodidae, Brachypauropodidae, Eurypauropodidae) from Indonesia and the Philippines with descriptions of a new genus and 26 new species. *International Journal of Myriapodology* 2: 69–148. <https://doi.org/10.1163/187525409X12577705044548>
- Scheller U (2011) Pauropoda. In: Minelli A (Ed.) *Treatise on Zoology-Anatomy, Taxonomy, Biology: the Myriapoda* (Vol. 1). Brill, Leiden, 467–508. https://doi.org/10.1163/9789004188266_022
- Stoev P, Li SQ, Meng K (2014) In memoriam Chong-zhou Zhang [1930–2014]. *Bulletin du Centre International de Myriapodologie* 47: 25–29.