

# Endonura Cassagnau in Iran, with a key to species of the genus (Collembola, Neanuridae, Neanurinae)

Adrian Smolis<sup>1</sup>, Morteza Kahrarian<sup>2</sup>, Agata Piwnik<sup>1</sup>, Dariusz Skarżyński<sup>1</sup>

<sup>1</sup> Institute of Environmental Biology, University of Wrocław, Przybyszewskiego 63-77, 51-148 Wrocław, Poland

<sup>2</sup> Department of Plant breeding and Agronomy, College of Agriculture, Kermanshah Branch, Islamic Azad University, Kermanshah, Iran

Corresponding author: Dariusz Skarżyński ([dariusz.skarzynski@uni.wroc.pl](mailto:dariusz.skarzynski@uni.wroc.pl))

---

Academic editor: L. Deharveng | Received 8 April 2015 | Accepted 26 November 2015 | Published 14 January 2016

---

<http://zoobank.org/A031D5CB-3874-40F2-8D7D-206D4437B99E>

---

**Citation:** Smolis A, Kahrarian M, Piwnik A, Skarżyński D (2016) *Endonura* Cassagnau in Iran, with a key to species of the genus (Collembola, Neanuridae, Neanurinae). ZooKeys 553: 53–71. doi: 10.3897/zookeys.553.6009

---

## Abstract

Three new species of *Endonura* are described from Iran. *Endonura dichaeta* sp. n. can be recognized by an ogival labrum, head without chaetae O and E, chaeta D connected with tubercle Cl, tubercle Dl with five chaetae on head, absence of tubercles Di on thorax I and tubercle (Di+Di) of thorax V with 2+2 chaetae. *Endonura ceratolabralis* sp. n. is characterized by large body size, reduction of labral chaetotaxy, ogival labrum, head without chaeta O and fusion of tubercles Di and De on first thoracic segment. *Endonura persica* sp. n. is distinguished from its congeners by a nonogival labrum, absence of chaeta O, tubercles Dl and (L+So) with five and eight chaetae respectively and claw with inner tooth. The key to all species of the genus is given.

## Keywords

Springtails, taxonomy, new species

## Introduction

*Endonura* was established by Cassagnau (1979) as one of four subgenera within the genus *Neanura* MacGillivray, 1893. Later, Deharveng (1982) raised it to the generic level. At present, *Endonura* is one of the largest (37 valid species) and most accurately studied genera within the subfamily Neanurinae (Dallai 1983, Deharveng 1979, 1982, Fanciulli and Dallai 2008, Pomorski and Skarżyński 2000, Pozo and Simón 1982, Smolis and Kaprus' 2003, 2009, Smolis 2006, Smolis et al. 2007, 2011). It is mostly a Palaearctic genus and only one species, *E. reticulata* (Axelson, 1905), is known from the Nearctic (Alaska, Smolis et al. 2011). According to a recent definition (Smolis 2008), *Endonura* is characterized by the following characters: 0–2 ocelli, reduced mouth parts with a thin mandible and a styliform maxilla, separate tubercles Di and De on the head, the non-cross-type of chaetotaxy on the head and three or two tubercles on abdomen V. The highest species diversity is observed in Europe (32 from among the 37 known species). However, this may be a false picture because many areas of the Palaearctic have been poorly studied by collembologists. Undoubtedly, one of such regions is Central Asia, but in this case the situation is rapidly and positively changing (Arbea and Kahrarian 2015, Kahrarian 2014, Kahrarian et al. 2013, Mayvan et al. 2015, Shayanmehr et al. 2013, Smolis et al. 2012). In the present paper, three new non-European *Endonura* from the western part of Iran are described. An updated key to all species of the genus is included.

## Terminology

Terminology for the description follows that given in Deharveng (1983), Deharveng and Weiner (1984), Smolis and Deharveng (2006) and Smolis (2008).

## Abbreviations used:

General morphology: abd. – abdomen, ant. – antenna, AOIII – sensory organ of antennal segment III, Cx – coxa, Fe – femur, Scx2 – subcoxa 2, T – tibiotarsus, th. – thorax, Tr – trochanter, VT – ventral tube.

Groups of chaetae: Ag – antegenital, An – chaetae of anal lobes, ap – apical, ca – centroapical, cm – centromedial, cp – centroposterior, d – dorsal, Fu – furcal, vc – ventrocentral, Ve or ve – ventroexternal, Vea – ventroexternoanterior, Vem – ventroexternomedial, Vep – ventroexternoposterior, Vel – ventroexternolateral, Vec – ventroexternocentral, Vei – ventroexternointernal, Vi or vi – ventriinternal, Vl – ventrolateral.

Tubercles: Af – antenno-frontal, Cl – clypeal, De – dorsoexternal, Di – dorsointernal, Dl – dorsolateral, L – lateral, Oc – ocular, So – subocular.

Types of chaetae: Ml – long macrochaeta, Mc – short macrochaeta, Mcc – very short macrochaeta, me – mesochaeta, mi – microchaeta, ms – s-microchaeta or microsensillum, S or s – chaeta s, bs – border s-chaeta on ant. IV, miA – microchaetae on

ant. IV, iv – ordinary chaetae on ventral ant. IV, or – organite of ant. IV, brs – border s–chaeta on ant. IV, i – ordinary chaeta on ant. IV, mou – cylindrical s–chaetae on ant. IV („soies mousses”), x – labial papilla x, L’ – ordinary lateral chaeta on abd. V, B4, B5 – ordinary chaetae on tibiotarsi.

## Materials and methods

The specimens were cleared in Nesbitt's fluid, subsequently mounted on slides in Swan's medium and observed using a phase contrast microscope Nikon E600. Photographs were made using a camera Nikon D5100 mounted on a microscope mentioned above. Photographs were stacked using Helicon Focus 6.2.2. and prepared for publication using Adobe Photoshop CS3. Material is deposited in the Department of Invertebrate Biology, Evolution and Conservation, Institute of Environmental Biology, University of Wrocław, Poland.

## Taxonomy

### *Endonura dichaeta* sp. n.

<http://zoobank.org/4CBE64B2-069B-4254-AF20-43D26E6CFE10>


Figs 1–4, Table 1

**Type material.** Holotype: adult female on slide, Iran, Osmanevand area, near Golestan village (N33°55', E47°06', 1241 m a.s.l.), litter in oak forest, 13.XII.2013, leg. M. Kahrarian. Paratypes: female, two males and two juveniles on slides, same data as holotype.

**Other material.** Two females and male on slide, Iran, Osmanevand area, near Chelkooshk village (N34°03', E47°12', 1516 m a.s.l.), litter in oak forest, 31.I.2014, leg. M. Kahrarian; three juveniles on slide, Iran, Paveh county, near Shabankereh village (N34°52.978', E46°30.760', 1632 m a.s.l.), litter in oak forest, 20.I.2014, leg. M. Kahrarian; two females and juvenile, Iran, Kermanshah county, near Chahar zebra-e-oliya village (N34°13', E46°40', 1592 m a.s.l.), litter in oak forest, 24.I.2014, leg. M. Kahrarian.

**Etymology.** The species name refers to rare feature within the genus - only two chaetae Di on each side of tubercle (Di+Di) of abdomen V.

**Diagnosis.** Habitus typical of the genus *Endonura*. Dorsal tubercles present and well developed, except tubercles Di on th. I. 2+2 unpigmented eyes. Buccal cone long, labrum ogival. Head with chaetae A, B, C, D, F and G. Chaetae O and E absent. Tubercles Cl and Af separate. Tuberclle Cl with chaetae D. Tubercles Dl and (L+So) on head with five and eight chaetae respectively. Tubercles De on th. II and III with three and four chaetae respectively. Tubercles L on abd. III and IV with three and six chaetae respectively. Abd. IV and V with eight and three tubercles respectively. Claw without inner tooth. Tibiotarsi with chaetae B4 and B5 short.



**Figures 1–4.** *Endonura dichaeta* sp. n.: **1** head (holotype), dorsal and lateral chaetotaxy **2** ventral sclerification of labrum **3** dorsal chaetotaxy of thorax **4** dorsal chaetotaxy of abdomen III–VI. Arrows indicate the position of eyes.

**Description.** Habitus typical of the genus. Body length (without antennae): 0.75–1.55 mm (holotype 1.30 mm). Colour of the body white. 2+2 medium unpigmented eyes (Fig. 1).

Types of dorsal ordinary chaetae. Macrochaetae M<sub>1</sub> relatively long, strongly thickened, almost cylindrical, arc-like or straight, narrowly sheathed, feebly serrated, apical-

ly pointed (Figs 1, 3–4); macrochaetae Mc and Mcc thickened, straight and pointed; mesochaetae and microchaetae short, thin, feebly serrated and pointed.

Head. Labrum ogival, with ventral sclerifications as in Fig. 2. Labrum chaetotaxy 2/2, 4. Labium with four basal, three distal and four lateral chaetae, papillae x absent. Maxilla styliform, mandible thin tridentate. Chaetotaxy of antennae as in Table 1c. Apical vesicle distinct, trilobed. S–chaetae of ant.IV long and moderately thickened. Chaetotaxy of head as in Table 1a, b, and Fig. 1. Chaeta D connected with tubercle Cl. Tubercl Af on head longer than tubercles Oc. Tubercl Dl with five chaetae, chaeta Dl3 absent, chaeta Dl6 as minute microchaeta and hard to detect (Fig. 1). Tubercl (L+So) with eight chaetae, chaetae So2 and L3 absent, chaeta So6 as Mc (Fig. 1). Elementary tubercles BE and CD present. Chaeta A shorter than B.

Thorax, abdomen, legs. Body s–chaeta thin and smooth, shorter than nearby macrochaetae (Figs 3, 4). Chaetotaxy of th. and abd. as in Table 1d and in Figs 3, 4. Tubercl Di on th.I not differentiated. Chaetae De3 on th. III and abd. I–III as Mcc. Chaetae De2 on th. II–III and De3 on th. III connected with tubercle De. Chaetae De3 on abd. I–III connected with tubercle De (Fig. 4). The line of chaetae De1–chaeta

**Table 1a.** Chaetotaxy of *Endonura dichaeta* sp. n.: Cephalic chaetotaxy–dorsal side.

Tubercl	Number of chaetae	Types of chaetae	Names of chaetae
Cl	6	Ml Mc	F D, G
Af	6	Ml Mc	A B, C
Oc	3	Ml Mc	Ocm, Ocp Oca
Di	2	Ml Mc	Di1 Di2
De	2	Ml Mc	De1 De2
Dl	5	Ml Mc Mcc mi	Dl1, Dl5 Dl4 Dl2 Dl6
(L+So)	8	Ml Mc me	L1, L4, So1 L2, So6 So3–5

**Table 1b.** Chaetotaxy of *Endonura dichaeta* sp. n.: Cephalic chaetotaxy–ventral side.

Group	Number of chaetae
Vi	6
Vea	3
Vem	3
Vep	4
labium	11, 0x

**Table 1c.** Chaetotaxy of *Endonura dichaeta* sp. n.: Chaetotaxy of antennae.

Segment, Group	Number of chaetae	Segment, Group	Number of chaetae adult
I	7	IV ap	or, 8 S, i, 12 mou, 6 brs, 2 iv
II	11		
III ve	5 sensilla AO III		8 bs, 5 miA
	5		2 bs, 3 miA
vc	4		3 bs, 1 miA
vi	4		8 miA, 1 brs
d	5		

**Table 1d.** Chaetotaxy of *Endonura dichaeta* sp. n.: Postcephalic chaetotaxy.

	Di	Terga De	Dl	L	Scx2	Cx	Legs Tr	Fe	T
th. I	1	2	1	-	0	3	6	13	19
th. II	3	2+s	3+s+ms	3	2	7	6	12	19
th. III	3	3+s	3+s	3	2	8	6	11	18
							Sterna		
abd. I	2	3+s	2	3	VT: 4				
abd. II	2	3+s	2	3	Ve: 4–5	Ve1 -	present		
abd. III	2	3+s	2	3	Vel: 4–5			Fu: 5–6me	2–4mi
abd. IV	2	2+s	3	6	Vel: 4	Vec: 2	Wei: 2	Vl: 4	
abd. V	(2+2)		5+s		Ag: 2			Vl: 1	L: 1
abd. VI		7			Ve: 13–14			An: 2mi	

s not perpendicular to the dorsomedian line on abd I–III. Furca rudimentary with 2–4 microchaetae. Tuberles Di on abd. V fused, with chaetae Di2 as Mc or Mcc, chaetae Di3 absent (Fig. 4). Chaetae L' and Vl on abd. V present. IV abd. with 2+2 chaetae Ag. No cryptopyggy. Chaetotaxy of legs as in Table 1d.

**Remarks.** In general appearance (shape of dorsal chaetae, chaetotaxy of central area of head and dorsal side of thorax and abdomen, complete absence of pigmentation and absence of cryptopyggy), *E. dichaeta* sp. n. strongly resembles *E. tartaginensis* Deharveng, 1980 described from Corsica. Nevertheless, both taxa differ in some essential characters, important from taxonomic point of view: presence/absence of chaetae E on head (*dichaeta* sp. n. absent, *tartaginensis* present), number of chaetae Dl on head (*dichaeta* sp. n. five, *tartaginensis* six), number of chaetae (L+So) (*dichaeta* sp. n. eight, *tartaginensis* nine), presence/absence of elementary tubercle EE on head (*dichaeta* sp. n. absent, *tartaginensis* present), presence/absence of tubercles Di on the first thoracic segment (*dichaeta* sp. n. absent, *tartaginensis* present), number of chaetae Di on abd. V (*dichaeta* sp. n. 2+2, *tartaginensis* 3+3) and presence/absence of tooth on claw (*dichaeta* sp. n. absent, *tartaginensis* present). In addition, the new species is characterized by only 2+2 antegenital chaetae (*tartaginensis* 3+3) and ogival labrum (unknown in *tartaginensis*), characters rarely observed within the genus.

***Endonura ceratolabralis* sp. n.**

<http://zoobank.org/FC09DDF3-EB60-416D-B31C-A290A4E812D5>

Figs 5–9, Table 2

**Type material.** Holotype: adult female on slide, Iran, Osmanevand area, near Markhor village ( $N33^{\circ}53'$ ,  $E47^{\circ}05'$ , 1389 m a.s.l.), litter in oak forest, 13.XII.2013, leg. M. Kahrarian. Paratypes: 3 females on slide, same data as holotype.

**Other material.** Three females on slide, Iran, Osmanevand area, near Ghadermarz village ( $N34^{\circ}01.030'$ ,  $E47^{\circ}12.415'$ , 1682 m a.s.l.), litter in oak forest, 31.I.2014, leg. M. Kahrarian.

**Etymology.** The species name refers to sharp labral apex which looks like a horn (“cera” in latin).


**Diagnosis.** Habitus typical of the genus *Endonura*. Dorsal tubercles present and well developed. 2+2 eyes darkly pigmented. Buccal cone long. Head with chaetae A, B, C, D, E, F and G. Chaeta O absent. Tuberles Cl and Af separate. Tuberles Dl and (L+So) on head with six and nine chaetae respectively. Tuberles Di and De on th. I fused. Tuberles De on th. II and III with three and four chaetae respectively. Tuberles L on abd. III and IV with three and 6–7 chaetae respectively. Abd. IV and V with eight and three tubercles respectively. Claw without inner tooth. Tibiotarsi with chaetae B4 and B5 short.

**Description.** Habitus typical of the genus. Body length (without antennae): 2.25–2.55 mm (holotype: 2.55 mm). Colour of the body bluish grey. 2+2 medium dark-pigmented eyes (Fig. 5).

Types of dorsal ordinary chaetae. Macrochaetae Ml thickened, relatively long, arc-like or straight, narrowly sheathed, feebly serrated, apically pointed or rarely rounded (Figs 5, 7–9); macrochaetae Mc and Mcc thickened, straight, pointed or apically rounded; mesochaetae and microchaetae short, thin and pointed.

Head. Buccal cone very long. Labrum ogival, with ventral sclerifications as in Fig. 6. Labrum chaetotaxy 0/2, 2. Labium with four basal, three distal and four lateral chaetae, papillae x absent. Maxilla styliform, mandible thin with two basal and two apical teeth. Chaetotaxy of antennae as in Table 2c. Apical vesicle distinct, trilobed. S–chaetae of ant.IV of medium length and moderately thickened. Chaetotaxy of head as in Table 2a, b, and Fig. 5. Tuberles Cl and Af separate. Tuberle Af on head longer than tuberles Oc. Chaeta O absent. Chaeta D free. Tuberle Dl with six chaetae, chaeta Dl3 present. Tuberle (L+So) with nine chaetae, chaeta So2 absent and chaeta So3 as Mc (Fig. 5). Elementary tubercles BE and CD present. Chaeta A shorter than B.

Thorax, abdomen, legs. Body s–chaetae thin and smooth, shorter than nearby macrochaetae (Figs 7–9). Chaetotaxy of th. and abd. as in Table 2d and in Figs 7–9. Tuberles Di on th.I differentiated and fused with De (Fig. 7). Dorsal side of th. and abd. without free chaetae De. The line of chaetae De1–chaeta s perpendicular to the dorsomedian line on abd I–III. Furca rudimentary with two or without microchaetae. Tuberles Di on abd. V fused, with chaetae Di2 and Di3 as Mc (Fig. 9). Chaetae L' and VI on abd. V present. No cryptopygy. Chaetotaxy of legs as in Table 2d.



**Figures 5–9.** *Endonura ceratolabralis* sp. n.: **5** head (holotype), dorsal and lateral chaetotaxy **6** ventral sclerification of labrum **7** dorsal chaetotaxy of thorax **8** dorsal chaetotaxy of abd. II **9** dorsal chaetotaxy of abdomen IV–VI. Arrows indicate the position of eyes.

**Remarks.** Because of the very characteristic long and pointed labrum, *E. ceratolabralis* sp. n. seems to be most similar to *E. cretensis* (Ellis, 1976) (Crete) and *E. gracilirostris* Smolis et al. 2007 (Crimea). Nevertheless, the new species can be easily distinguished from these two taxa by the following combination of characters: maximum length of the body without antennae (*ceratolabralis* sp. n. 2.55 mm; *gracilirostris*

**Table 2a.** Chaetotaxy of *Endonura ceratolabralis* sp. n.: Cephalic chaetotaxy—dorsal side.

Tubercle	Number of chaetae	Types of chaetae	Names of chaetae
Cl	4	Ml	F
		Mc	G
Af	10	Ml	A
		Mc	B, C, D, E
Oc	3	Ml	Ocm, Ocp
		me	Oca
Di	2	Ml	Di1
		Mc	Di2
De	2	Ml	De1
		Mc	De2
Dl	6	Ml	Dl1, Dl5
		Mc	Dl2, Dl3, Dl4
		mi	Dl6
(L+So)	9	Ml	L1, L4, So1
		Mc	L2, L3, So3
		me	So4–6

**Table 2b.** Chaetotaxy of *Endonura ceratolabralis* sp. n.: Cephalic chaetotaxy—ventral side.

Group	Number of chaetae
Vi	6
Vea	3–4
Vem	3
Vep	4
Labium	11, 0x

**Table 2c.** Chaetotaxy of *Endonura ceratolabralis* sp. n.: Chaetotaxy of antennae.

Segment, Group	Number of chaetae	Segment, Group	Number of chaetae adult
I	7	IV ap	or, 8 S, i, 12 mou, 6 brs, 2 iv
II	12–14		
III	5 sensilla AO III		8 bs, 5 miA
ve	5		2 bs, 3 miA
vc	4	ca	3 bs, 1 miA
vi	4	cm	8 miA, 1 brs
d	5	cp	

1.45 mm; *cretensis* 0.8 mm), labral formula (*ceratolabralis* sp. n. 0/2, 2; *gracilirostris* 0/2, 4; *cretensis* 2/2, 4), presence/absence of chaeta O on head (*ceratolabralis* sp. n. absent, in others present), number of chaetae Dl on head (*cretensis* three, in others six), number of chaetae (L+So) on head (*cretensis* seven, in others nine), fusion/separation of tubercles Di and de on th. I (*gracilirostris* separate, in others fused), number of ordinary chaetae De on th. III (*cretensis* two, others three), presence/absence of free chaetae on thorax (*ceratolabralis* sp. n. absent, in others present) and number of chaetae Di on abd. V (*cretensis* 1–2, others three).

**Table 2d.** Chaetotaxy of *Endonura ceratolabralis* sp. n.: Postcephalic chaetotaxwy.

	<b>Di</b>	<b>Terga De</b>	<b>Dl</b>	<b>L</b>	<b>Scx2</b>	<b>Cx</b>	<b>Legs Tr</b>	<b>Fe</b>	<b>T</b>
th. I	1	2	1	-	0	3	6	13	19
th. II	3	2+s	3+s+ms	3	2	7	6	12	19
th. III	3	3+s	3+s	3	2	8	6	11	18
						Sterna			
abd. I	2	3+s	2	3	VT: 4				
abd. II	2	3+s	2	3	Ve: 5–6	Ve1 -	Present		
abd. III	2	3+s	2	3	Vel: 4–5			Fu: 3–6me	0–2mi
abd. IV	2	2+s	3	6–7	Vel: 4	Vec: 2	Vei: 2	Vl: 4	
abd. V	(3+3)		5+s		Ag: 3			Vl: 1	L: 1
abd. VI		7			Ve: 13–14			An: 2mi	

***Endonura persica* sp. n.**

<http://zoobank.org/9CFE5947-62CC-4A3E-ABF7-5B84EA69A21A>

Figs 10–13, Table 3

**Type material.** Holotype: adult female on slide, Iran, Kermanshah area, near Ghaleh shahin village ( $N34^{\circ}25.590'$ ,  $E05^{\circ}12.415'$ , 566 m a.s.l.), litter in willow shrubs, 7.IV.2014, leg. M. Kahrarian. Paratypes: two females, four males and four juveniles on slides, same data as holotype.


**Other material.** Female on slide, Iran, Kermanshah Province, Halashi County, near Sarfiroozabad village ( $N34^{\circ}02'$ ,  $E47^{\circ}10'$ , 1624 m a.s.l.), litter in oak forest, 15.II.2014, leg. M. Kahrarian; female and male on slide, Iran, Osmanevand area, near Sarjoob village ( $N33^{\circ}56'$ ,  $E47^{\circ}08'$ , 1240 m a.s.l.), litter in oak forest, 13.XII.2013, leg. M. Kahrarian.

**Etymology.** The species name refers to the historic name of Iran, Persia.

**Diagnosis.** Habitus typical of the genus *Endonura*. Dorsal tubercles present and generally well developed, only tubercles Di on th. I weakly differentiated. 2+2 large dark-pigmented eyes. Buccal cone rather short. Head with chaetae A, B, C, D, E, F and G. Chaeta O absent. Tubercles Cl and Af separate. Tubercles Dl and (L+So) on head with five and eight chaetae respectively. Tubercles De on th. II and III with three and four chaetae respectively. Tubercles L on abd. III and IV with four and 6–7 chaetae respectively. Abd. IV and V with eight and three tubercles respectively. Claw with inner tooth. Tibiotarsi with chaetae B4 and B5 long.

**Description.** Habitus typical of the genus. Body length (without antennae): 0.75–1.90 mm (holotype: 1.10 mm). Colour of the body bluish grey. 2+2 large dark pigmented eyes (Fig. 10).

Types of dorsal ordinary chaetae. Macrochaetae Ml thickened, relatively long, arc-like or straight, narrowly sheathed, feebly serrated, apically rounded or rarely pointed (Figs 10, 12); macrochaetae Mc and Mcc thickened, straight and not pointed; mesochaetae and microchaetae short, thin and pointed.



**Figures 10–13.** *Endonura persica* sp. n.: 10 head and th. I, dorsal and lateral chaetotaxy 11 ventral sclerification of labrum 12 dorsal chaetotaxy of abdomen III–VI (holotype) 13 tibiotarsi and claw of leg III. Arrows indicate the position of eyes.

Head. Buccal cone short. Labrum rounded, with ventral sclerifications as in Fig. 11. Labrum chaetotaxy 4/2, 4. Labium with four basal, three distal and four lateral chaetae, papillae x absent. Maxilla styliform, mandible thin with two basal and two apical teeth. Chaetotaxy of antennae as in Table 3c. Apical vesicle distinct trilobed. S–chaetae of ant.IV of medium length and moderately thickened. Chaetotaxy of head as in Table 3a, b, and Fig. 10. Chaetae D and E free. Tuberles Cl and Af separate.

**Table 3a.** Chaetotaxy of *Endonura persica* sp. n.: Cephalic chaetotaxy—dorsal side.

Tubercl	Number of chaetae	Types of chaetae	Names of chaetae
Cl	4	Ml Mc	F G
Af	10	Ml Mc Mc or Mcc	B A, C, E D
Oc	3	Ml Mc mi	Ocm Ocp Oca
Di	2	Ml Mcc	Di1 Di2
De	2	Ml Mcc	De1 De2
Dl	5	Ml Mc or Mcc	Dl1, Dl5 Dl2, Dl4, Dl6
(L+So)	8	Ml Mc me or mi	L1, L4, So1 L2 So3–6

**Table 3b.** Chaetotaxy of *Endonura persica* sp. n.: Cephalic chaetotaxy—ventral side.

Group	Number of chaetae
Vi	6
Vea	3–4
Vem	3
Vep	4
labium	11, 0x

**Table 3c.** Chaetotaxy of *Endonura persica* sp. n.: Chaetotaxy of antennae.

Segment, Group	Number of chaetae	Segment, Group	Number of chaetae adult
I	7		
II	12–14		
III	5 sensilla AO III		
ve	5	IV ap	or, 8 S, i, 12 mou, 6 brs, 2 iv
vc	4		8 bs, 5 miA
vi	4	ca	2 bs, 3 miA
d	5	cm	3 bs, 1 miA
		cp	8 miA, 1 brs

Tubercl Af on head longer than tubercles Oc. Tubercl Dl with five chaetae, chaeta Dl3 absent. Tubercl (L+So) with eight chaetae, chaetae So2 and L3 absent (Fig. 10). Elementary tubercle BE absent. Chaeta A shorter than B.

Thorax, abdomen, legs. Body s—chaetae fine and smooth, distinctly shorter than nearby macrochaetae (Fig. 12). Chaetotaxy of th. and abd. as in Table 3d and in Figs

**Table 3d.** Chaetotaxy of *Endonura persica* sp. n.: Postcephalic chaetotaxy.

	Di	Terga De	Dl	L	Scx2	Cx	Legs Tr	Fe	T
th. I	1	2	1	-	0	3	6	13	19
th. II	3	2+s	3+s+ms	3	2	7	6	12	19
th. III	3	3+s	3+s	3	2	8	6	11	18
						Sterna			
abd. I	2	3+s	2	3	VT: 4				
abd. II	2	3+s	2	3	Ve: 5–6	Ve1 -	present		
abd. III	2	3+s	2	4	Vel: 5			Fu:5–10me	0 mi
abd. IV	2	2+s	3	6–7	Vel: 4	Vec: 2	Vei: 2	Vl: 4	
abd. V	(3+3)		8+s		Ag: 3			Vl: 1	L: 1
abd. VI		7			Ve: 13–14			An: 2mi	

10, 12. Tuberles Di on th.I differentiated or not. Chaetae De2 on th. II–III and De3 on th. III free. Chaetae De3 on abd. I–III free (Fig. 12). The line of chaetae De1–chaeta s parallel to the dorsomedian line on abd. I–III. Furca rudimentary without microchaetae. Tuberles Di on abd. V fused, with chaetae Di2 as Mcc and Di3 as mi (Fig. 12). Chaetae VI on abd. V present. Cryptopygy slightly developed. Chaetotaxy of legs as in Table 3d. Tibiotarsi with rather long chaetae B4 and B5. Claw with inner tooth (Fig. 13).

**Remarks.** In general appearance and presence of inner tooth on claw, characters rarely observed within the genus, *E. persica* sp. n. strongly resembles to *E. dentifera* Smolis et al. 2007 (described from Crimea). However, the new species can be reliably separated from Crimean species with the following characters: number of chaetae Dl on head (*persica* sp. n. five, *dentifera* six), number of chaetae (L+So) on head (*persica* sp. n. eight, *dentifera* ten), presence/absence of tubercles Di on the first thoracic segment (*persica* sp. n. present, *dentifera* absent) and number of chaetae L of abd. IV (*persica* sp. n. 6–7 chaetae, *dentifera* 8–9).

### Key to the genus *Endonura*

In 1982, Deharveng, in his PhD thesis, elevated *Endonura* to the generic level and prepared a key to the genus that comprised 23 species. Nowadays, including the taxa described herein, the genus contains 40 members and is the second largest of the tribe Neanurini, after *Deutonura* Cassagnau, 1979. Moreover, after the publication of Deharveng's paper (date), a few species were redescribed and one taxon was synonymised

(Smolis and Kaprus' 2003, Smolis 2008, Smolis et al. 2007, 2011). Considering these facts, the preparation of an updated key to all species of the genus seemed to be highly recommended.

1	Head with fusion of tubercles Af and Cl .....	2
—	Head with separation of tubercles Af and Cl .....	7
2.	Chaeta O on head present .....	3
	Chaeta O on head absent.....	4
3	Tubercles Di on th. I present and fused with De, tubercle (Di+Dl+L) on abd. V with nine chaetae ..... <i>E. poinsotae</i> Deharveng, 1980 (France, Corsica)	
—	Tubercles Di on th. I absent, tubercle (Di+Dl+L) on abd. V with seven chaetae..... <i>E. ichnusae</i> Dallai, 1983 (Italy, Sardinia)	
4	Tubercles De on abd. I–III with four chaetae.....	5
—	Tubercles De on abd. I–III with three chaetae..... .... <i>E. granulata</i> (Cassagnau & Delamare Deboutteville, 1955) (Lebanon)	
5	Tubercles Di and De on th. I fused, cryptopygy strongly developed .....	
	..... <i>E. gladiirostra</i> Smolis & Kaprus', 2003 (Israel)	
—	Tubercles Di and De on th. I separate, cryptopygy absent or weakly developed .....	6
6	Chaeta E on head present, Tuberle Dl on head with four chaetae .....	
	..... <i>E. tyrrhenica</i> Dallai, 1983 (Italy, Sardinia)	
—	Chaeta E on head absent, Tuberle Dl on head with six chaetae.....	
	..... <i>E. pejai</i> Deharveng, 1980 (France, Corsica)	
7	Tuberle Af on head equal or shorter than tubercles Oc .....	8
—	Tuberle Af on head longer than tubercles Oc.....	9
8	Labrum with ventral sclerifications ogival and without prelabral chaetae.....	
	..... <i>E. gracilirostris</i> Smolis et al., 2007 (Crimea, Moldova)	
—	Labrum with ventral sclerifications nonogival and with prelabral chaetae.....	
	..... <i>E. taurica</i> (Stach, 1951) (Crimea)	
9	Chaeta O on head present .....	10
—	Chaeta O on head absent.....	26
10	Eyes completely absent.....	11
—	Eyes present .....	12
11	Tubercles Di on th. I present... <i>E. arbascensis</i> Deharveng, 1979 (France, Spain)	
—	Tubercles Di on th. I absent ... <i>E. caeca</i> (Gisin, 1963) (Bosnia and Herzegovina)	
12	Anterior eye present and located outside tubercle Oc .....	
	..... <i>E. asiatica</i> Smolis et al., 2011 (Kyrgyzstan)	
—	Anterior eye present or absent, if present located within tubercle Oc .....	13
13	Anterior eye present .....	14
—	anterior eye absent ..... <i>E. immaculata</i> Deharveng, 1980 (France, Corsica)	
14	Claw with inner tooth, tibiotarsi with long chaetae B4 and B5 .....	15
—	Claw without tooth, tibiotarsi with short chaetae B4 and B5 .....	16

15	Tubercle Dl on head with three chaetae, tubercles Di on th. II–III with two chaetae.....	<i>E. tetrophthalma</i> (Stach, 1929) (Hungary)
	Tubercle Dl on head with five chaetae, tubercles Di on th. II–III with three chaetae.....	<i>E. lusatica</i> (Dunger, 1966) (Germany, Poland, Ukraine)
16	Chaeta E on head absent.....	17
–	Chaeta E on head present .....	18
17	Tubercle Cl on head with chaetae D, elementary tubercle DF present .....	
	..... <i>E. colorata</i> (Gama, 1964) (Portugal)	
–	Tubercle Cl on head without chaetae D, elementary tubercle DF absent.....	
	..... <i>E. cantabrica</i> (Deharveng, 1979) (Spain)	
18	Tubercle Dl on head with six chaetae .....	19
–	Tubercle Dl on head with less number of chaetae.....	25
19	Tubercles Di on head present.....	20
–	Tubercles Di on head absent .....	
	..... <i>E. dalensi</i> Deharveng, 1979 (Andorra, France, Spain, Italy)	
20	Body white .....	21
–	Body blue or bluish–grey .....	22
21.	Tubercle (L+So) on head with nine chaetae, macrochaetae thin and pointed ..	
	..... <i>E. deharvengi</i> Cassagnau & Péja, 1979 (Greece)	
–	Tubercle (L+So) on head with eight chaetae, macrochaetae thickened and blunt.....	
	..... <i>E. levantica</i> Smolis et al., 2011 (Israel)	
22	Tubercle De on th. III with two ordinary chaetae .....	
	..... <i>E. gladiolifer</i> (Cassagnau, 1954) (Algeria, Spain)	
–	Tubercle De on th. III with three ordinary chaetae.....	23
23	Tubercle Cl on head with chaetae D, furca rudimentary with microchaetae....	
	..... <i>E. alavensis</i> Pozo & Simon, 1982 (Spain)	
–	Tubercle Cl on head without chaetae D, furca rudimentary without microchaetae.....	24
24	Tubercle (L+So) on head with nine chaetae, free chaeta L on abd. IV present...	
	..... <i>E. quadriseta</i> Cassagnau & Péja, 1979 (Greece, Turkey, Crimea)	
–	Tubercle (L+So) on head with ten chaetae, free chaeta L on abd. IV absent.....	
	..... <i>E. reticulata</i> (Axelson, 1905) (Finland; Russia; Sweden; United States, Alaska)	
25	Tubercle Dl on head with four chaetae, tubercles Di and De on th. I separate .....	
	..... <i>E. occidentalis</i> (Deharveng, 1979) (Spain)	
–	Tubercle Dl on head with three chaetae, tubercles Di and De on th. I fused.....	
	..... <i>E. cretensis</i> (Ellis, 1976) (Greece, Israel)	
26	Cryptopygy strong and complete, tubercles of abd. VI invisible in dorsal view .....	
	..... <i>E. ludoviciae</i> (Denis, 1948) (France, Corsica)	
–	Cryptopygy absent or weak, tubercles of abd. VI well or partially visible in dorsal view .....	27
27	Body bluish–grey .....	28
–	Body white .....	30

28	Claw with inner tooth, labrum chaetotaxy 4/2, 4.....	29
–	Claw without inner tooth, labrum chaetotaxy 0/2, 2 .....	
	..... <i>E. ceratolabralis</i> sp. n. (Iran)	
29	Tubercle Dl on head with five chaetae, tubercles Di on th. I present .....	
	..... <i>E. persica</i> sp. n. (Iran)	
–	Tubercle Dl on head with six chaetae, tubercles Di on th. I absent.....	
	..... <i>E. dentifera</i> Smolis et al., 2007 (Crimea)	
30	Chaeta C on head absent .....	31
–	Chaeta C on head present.....	33
31	Macrochaetae Di1 on abd. V distinctly thickened and club-like.....	
	..... <i>E. baculifer</i> Deharveng, 1979 (Portugal)	
–	Macrochaetae Di1 on abd. V slightly thickened and cylindrical .....	32
32	Eyes present, tubercles Di on th. I present.....	
	..... <i>E. transcaucasica</i> (Stach, 1951) (Georgia)	
–	Eyes absent, tubercles Di on th. I absent.... <i>E. carpatica</i> Smolis, 2006 (Poland)	
33	Tubercle Cl on head with chaetae D, elementary tubercle DF present .....	34
–	Tubercle Cl on head without chaetae D, elementary tubercle DF absent... <i>E. tartaginensis</i> Deharveng, 1980 (France, Corsica)	35
34	Chaeta E on head present, tubercle Dl on head with six chaetae .....	
–	..... <i>E. dichaeta</i> sp. n. (Iran)	
35	Chaeta E on head present .....	
	..... <i>E. urotuberculata</i> Pomorski & Skarżyński, 2000 (Bulgaria)	
–	Chaeta E on head absent.....	36
36	Chaeta L4 on head free, eyes absent or present unpigmented.....	37
–	Chaeta L4 within tubercle (L+So), eyes present and pigmented .....	39
37	Abd. V with two tubercles .....	
	..... <i>E. incolorata</i> (Stach, 1951) (Poland, Ukraine, Romania)	
–	Abd. V with three tubercles.....	38
38	Abd. IV with eight tubercles, macrochaetae Ml relatively short .....	
	..... <i>E. tatricola</i> (Stach, 1951) (Poland, Slovakia)	
–	Abd. IV with five tubercles, macrochaetae Ml long .....	
	..... <i>E. dudichi</i> (Loksa, 1967) (Hungary, Poland, Slovakia)	
39	Tubercle Dl on head with six chaetae, tubercle L on abd. III with three chaetae.....	
	..... <i>E. centaurea</i> Cassagnau & Péja, 1979 (Greece)	
–	Tubercle Dl on head with five chaetae, tubercle L on abd. III with four chaetae..... <i>E. saleri</i> Facciulli & Dallai, 2008 (Italy)	

## Discussion

Considering the data presented here and those obtained from the literature (Mayvan et al. 2015, Shayanmehr et al. 2013, Smolis et al. 2012), Neanurinae fauna of Iran

comprises ten species and seven genera: *Bilobella aurantiaca* (Caroli, 1912), *Cryptonura persica* Smolis et al., 2012, *C. maxima* Smolis et al., 2012, *Deutonura decolorata* (Gama & Gisin, 1964) (Gisin 1964), *Endonura ceratolabralis* sp. n., *E. dichaeta* sp. n., *E. persica*, sp. n., *Neanura muscorum* (Templeton, 1835), *Persanura hyrcanica* Mayvan et al. 2015, *Thaumanura echinata* (Kos, 1940). It should be noted, however, that until now only the western part of Iran has been roughly studied. Although future research may change the present picture of the subfamily diversity in the studied country and region, some preliminary conclusions can be drawn. The first is related to the higher systematic pattern and composition of Neanurinae of Iran. This fauna consists almost exclusively of members of the tribe Neanurini, the most diverse and dominant among Neanurinae in the western Palaearctic. To date, none of the Lobellini and Paranurini genera have been found in Iran, although they are numerous and widely distributed in south, south-east and east Asia. The second conclusion seems to be more expected, *Endonura* species from Iran resemble those known from south-east Europe. It suggests their close affinity and the historical connection between these faunas. The third conclusion sheds light on the distribution and the history of this genus. Most *Endonura* species were recorded from Mediterranean and temperate zones of Europe, where they live predominantly in forests. It is worth saying that the greatest diversity of the genus through the continent is more or less correlated to the areas of land that have never been subjected to glaciations. Till now, the occurrence of only a few species is documented outside Europe, especially in the Middle East (Smolis and Kaprus 2003, 2009; Smolis et al. 2011). The recent and present discoveries of *Endonura* species in Kyrgyzstan (Smolis et al. 2011) and Iran significantly expand the list of species and also our knowledge on the genus. Undoubtedly, diverse forest habitats of the coastal and montane regions of Iran and adjacent countries hide a rich fauna of Neanurinae. We therefore hope that a more comprehensive study in the future will allow us to present a better picture of the distribution of *Endonura* in Iran and the near East.

## References

- Arbea JI, Kahrarian M (2015) The genus *Folsomides* Stach (Collembola, Isotomidae) in Kerman-shah Province (W Iran) with the description of two new species. Zootaxa 3925: 281–290. doi: 10.11646/zootaxa.3925.2.9
- Axelson WM (1905) Einige neue Collembolen aus Finland. Zoologisher Anzeiger 28: 788–794.
- Caroli E (1912) Contribuzioni alla conoscenza dei Collemboli italiani. I. La tribu degli Achorutini CB (1906). Archivio zoologico italiano 6: 349–374.
- Cassagnau P (1954) Sur un rudiment de furca chez les Neanurinae et sur quelques espèces de ce groupe. Extrait Du Bulletin de la Société D'Histoire Naturelle de Toulouse 89: 27–34.
- Cassagnau P (1979) Les Collemboles Neanuridae des Pays Dinaro-Balkaniques: leur intérêt phylogénétique et biogéographique. Biologia Gallo-Hellenica 8: 185–203.
- Cassagnau P (1989) Les Collemboles Neanurinae; éléments pour une synthèse phylogénétique et biogéographique. In: Dallai R (Ed.) 3<sup>rd</sup> International Seminar on Apterygota, Siena, 171–182.

- Cassagnau P, Delamare Deboutteville C (1955) Mission Henri Coiffait (1951). 3. Collemboles. Archives de Zoologie Expérimentale et Générale 91: 365–395.
- Cassagnau P, Péja N (1979) Diagnoses préliminaires de quelques Neanuridae de Grèce et d'Albanie. Biologia Gallo-Hellenica 8: 205–222.
- Dallai R (1983) Interesse biogeografico dei Neanuridi (Collembola) della Sardegna e delle isole dell'Arcipelago Toscano. Lavori della Societa Italiana di Biogeografia (N.S.) 8: 417–465.
- Deharveng L (1979) Contribution à la connaissance des Collemboles Neanurinae de France et de la Peninsule Iberique. Travaux du Laboratoire d'Écobiologie des Arthropodes Édaphiques, Toulouse 1: 1–61.
- Deharveng L (1980) Contribution à la connaissance systématique, écologique et biogéographique des collemboles Neanurinae de Corse. Revue d'écologie et de biologie du sol 17(3): 419–435.
- Deharveng L (1982) Contribution à l'étude des Collemboles Neanurinae: évolution, spéciation, polymorphisme somatique et chromosomique des formes européennes. PhD thesis, University of Paul Sabatier, Toulouse, France.
- Deharveng L (1983) Morphologie évolutive des Collemboles Neanurinae en particulier de la lignée Neanurinae. Travaux de Laboratoire d'Écobiologie des Arthropodes Édaphiques, Toulouse 4: 1–63.
- Deharveng L, Weiner W (1984) Collemboles de Corée du Nord III–Morulinae et Neanurinae. Travaux de Laboratoire d'Écobiologie des Arthropodes Édaphiques, Toulouse 4: 1–61.
- Denis JR (1948) Sur la faune Française des Aptérygotes (XXIV<sup>e</sup> note). Étude du matériel récolté en Corse par P. et L. Rémy en 1942. Annales des Sciences Naturelles Zoologie 11<sup>e</sup> série, 9: 1–12.
- Dunger W (1966) Zur Kenntnis von Neanura tetrophthalmus (Stach) (Collembola: Apterygota). Abhandlungen und Berichte des Naturkundemuseums Görlitz 41(4): 1–11.
- Ellis WN (1976) Autumn fauna of *Collembola* from Central Crete. Tijdschrift voor Entomologie 119: 221–236.
- Fanciulli PP, Dallai R (2008) Three new species of Collembola from north-east Italy. Zootaxa 1701: 15–28.
- Gama MM (1964) Colêmbolos de Portugal Continental. Mémorias e Estudos do Museu Zoológico da Universidade de Coimbra 292: 1–252.
- Gisin H (1963) Sieben neue Arten von Collembolen aus Bosnien und Wiederbeschreibung von *Onychiurus serratuberculatus* Stach. Extrait du Godišnjak Biološkog Instituta Univerziteta u Sarajevu 14: 3–13.
- Gisin H (1964) Collemboles d'Europe. VI. Revue suisse de Zoologie 74(20): 383–400. doi: 10.5962/bhl.part.75615
- Kahrarian M (2014) New records of Poduromorpha for the Iranian springtail fauna (Collembola). Natura Somogyensis 25: 21–26.
- Kahrarian M, Vafei-Shooshtari R, Skarżyński D, Konikiewicz M, Soleymannezhadyan E, Shayan Mehr M, Shams B (2013) A new species and new records of the genus *Hypogastrura* Bourlet, 1839 (Collembola, Hypogastruridae) from Iran. Zootaxa 3709: 89–94. doi: 10.11646/zootaxa.3709.1.4
- Kos F (1940) Terricole Collembolen aus Slovenien. Glasnik, Bulletin de la Société Scientifique de Skoplje 22: 136–168.

- Loksa I (1967) Vier neue Höhlencollembolen aus Ungarn (Biospeologica Hungarica, XIII). Opuscula Zoologica Budapest VI (2): 289–296.
- Mayvan MM, Shayanmehr M, Smolis A, Skarżyński D (2015) *Persanura hyrcanica*, a new genus and species of Neanurinae (Collembola: Neanuridae) from Iran, with a key to genera of the tribe Neanurini. Zootaxa 3918: 552–558. doi: 10.11646/zootaxa.3918.4.4
- Pomorski RJ, Skarżyński D (2000) A new species of *Endonura* Cassagnau, 1979 from Bulgaria with remarks on *Endonura centaurea* (Cassagnau & Peja, 1979) (Collembola, Neanuridae). Zoosystema 22: 117–120.
- Pozo J, Simón JC (1982) Colémbolos del País Vasco I. *Neanura (Endonura) occidentalis alawensis* nov. ssp. y *Neanura (Deutonura) plena* ssp. *plena* Stach, 1951. Eos 57: 221–229.
- Shayanmehr M, Yahyapour E, Kahrarian M, Lafooraki E (2013) An introduction to Iranian Collembola (Hexapoda): an update to the species list. Zookeys 335: 69–83. doi: 10.3897/zookeys.335.5491
- Smolis A (2006) *Endonura carpatica*, a new species from Poland (Collembola, Neanuridae, Neanurinae). Genus 17: 471–476.
- Smolis A (2008) Redescription of four Polish *Endonura* Cassagnau, 1979 (Collembola, Neanuridae, Neanurinae), with a nomenclature of the ventral chaetae of antennae. Zootaxa 1858: 9–36.
- Smolis A, Deharveng L (2006) *Vitronura mascula*, a new species of Neanurinae (Collembola: Neanuridae) from northern Vietnam, with a key to the species of the genus. Revue suisse de Zoologie 113: 263–268. doi: 10.5962/bhl.part.80349
- Smolis A, Deharveng L, Kaprus' IJ (2011) Studies on the non-European *Endonura* Cassagnau, 1979 (Collembola, Neanuridae, Neanurinae). Zootaxa 3004: 45–56.
- Smolis A, Falahati A, Skarżyński D (2012) The genus *Cryptonura* Cassagnau, 1979 (Collembola, Neanuridae, Neanurinae) in Iran. Zootaxa 3530: 51–58.
- Smolis A, Kaprus' IJ (2003) A new species of the genus *Endonura* Cassagnau, 1979 from Israel (Collembola: Neanuridae). Genus 14: 325–329.
- Smolis A, Kaprus' IJ (2009) Redescription of *Endonura cretensis* (Ellis, 1976) (Collembola: Neanuridae: Neanurinae) with a new record from Israel. Annales Zoologici 59: 1–6. doi: 10.3161/000345409X432538
- Smolis A, Skarżyński D, Pomorski RJ, Kaprus' IJ (2007) Redescription of *Endonura taurica* (Stach, 1951) and *E. quadriseta* Cassagnau & Peja, 1979, and description of two new species of the genus *Endonura* Cassagnau, 1979 (Collembola: Neanuridae: Neanurinae) from the Crimea (Ukraine). Zootaxa 1442: 19–35.
- Stach J (1929) Verzeichnis der Apterygogenea Ungarns. Annales Musei Nationalis Hungarici, Budapest, 26: 282–283.
- Stach J (1951) The Apterygotan fauna of Poland in relation to the world-fauna of this group of insects. Family: Bilobidae. Polska Akademia Umiejętności, Acta monographica Musei Historiae Naturalis, Kraków, 97 pp.
- Templeton R (1835) Thysanura Hibernicae, or descriptions of such species of spring-tailed insects (Podura and Lepisma, Lin.) as have been observed in Ireland. Transactions of the Entomological Society of London 1(2): 89–98. doi: 10.1111/j.1365-2311.1838.tb00147.x