

A new species, a new combination, and a new record of *Crossotarsus* Chapuis, 1865 (Coleoptera, Curculionidae, Platypodinae) from China

Shengchang Lai¹, Ling Zhang², You Li³, Jianguo Wang²

1 College of Forestry, Nanjing Forestry University, Nanjing, Jiangsu 210037, China **2** College of Agricultural Sciences, Jiangxi Agricultural University, Nanchang, Jiangxi 330045, China **3** School of Forest Resources and Conservation, University of Florida, Gainesville, FL, 32611, USA

Corresponding author: Jianguo Wang (jgwang@jxau.edu.cn)

Academic editor: M. Alonso-Zarazaga | Received 22 November 2020 | Accepted 22 February 2021 | Published 6 April 2021

<http://zoobank.org/3279FAE9-E002-4142-930F-96DF49B9E959>

Citation: Lai S, Zhang L, Li Y, Wang J (2021) A new species, a new combination, and a new record of *Crossotarsus* Chapuis, 1865 (Coleoptera, Curculionidae, Platypodinae) from China. ZooKeys 1028: 69–83. <https://doi.org/10.3897/zookeys.1028.61018>

Abstract

This study describes a new species, *Crossotarsus beaveri* Lai & Wang, **sp. nov.**, designates a new combination, *C. brevis* (Browne, 1975, **comb. nov.** from *Platypus* Herbst, 1793), and notes a new record, *C. emorus* Beeson, 1937, from China. Genetic data from four genes indicate that the new species and *C. brevis* form a clade clustered with other *Crossotarsus* species. Molecular phylogeny and morphological characters support their taxonomic placement.

Keywords

Ambrosia beetle, Fujian, Jiangxi, molecular phylogeny, pinhole borer, taxonomy

Introduction

The genus *Crossotarsus* Chapuis 1865 was erected for 29 species of pinhole borers. *Crossotarsus wallacei* (Thomson, 1857) was designated as the type species of the genus (Hopkins 1914). Wood (1993) revised the genera of Platypodidae and placed *Crossotarsus* in the subfamily Platypodinae, tribe Platypodini. *Crossotarsus* is distinguished from

other Platypodine genera primarily by the following combination of characters (Browne 1961; Wood 1993; Beaver and Sanguansub 2015): 1. Labial palps two-segmented, with basal segments fused in the midline; 2. Sexually dimorphic protibiae, the outer face of the protibia transversely carinate in the male and finely granulate in the female; 3. Pronotum without specialized mycangial pores in either sex; 4. Femoral grooves angulate at the anterior extremity and gently rounded behind; 5. Metacoxa strongly projecting with a deep vertical posterior face. But Wood's (1993) generalisation that the female pronotum of *Crossotarsus* species has numerous mycangial pores is incorrect (Beaver 2004).

The catalog of Wood and Bright (1992) includes 118 species of *Crossotarsus*. As a result of taxonomic changes since that time, 116 species are currently recognised. Most species of *Crossotarsus* occur in the Oriental region, extending from India across Southeast Asia and Indonesia to Australia and the Pacific islands, and northward to Taiwan and Japan (Wood 1993). *Crossotarsus externedentatus* (Fairmaire 1849) is also widespread in the Afrotropical forests.

The Platypodinae have been almost entirely neglected in China. Only a few papers include original records of *Crossotarsus* from the country. Yin and Huang (1987) recorded three species: *C. coniferae* Stebbing, 1906, *C. squamulatus* Chapuis, 1865, and *C. wallacei* from Yunnan. Yin et al. (2002) added two species: *C. externedentatus* and *C. terminatus* Chapuis, 1865 from Hainan island, and Zhang et al. (2008) cited 13 species from China. After taxonomic changes (Beaver 2004; 2005; 2016; Bright 2014), the following 13 species are currently known from China: *C. coniferae* (Yunnan, Sichuan, Xizang); *C. emancipatus* Murayama, 1934 (Taiwan); *C. externedentatus* (Hainan, Taiwan); *C. flavomaculatus* Strohmeier, 1912 (Taiwan); *C. formosanus* Strohmeier, 1912 (Taiwan); *C. niponicus* Blandford, 1894 (Taiwan); *C. piceus* Chapuis, 1865 (Taiwan); *C. saltatorinus* (Schedl, 1954) (Fujian); *C. sauteri* (Strohmeier, 1913) (Taiwan); *C. simplex* Murayama, 1925 (Taiwan); *C. squamulatus* (Yunnan); *C. terminatus* (Hainan, Yunnan, Xizang); *C. wallacei* (Hainan, Taiwan).

In this study, we describe a new species of *Crossotarsus* from China, give a new distribution record, transfer a previously described species to the genus, and provide molecular data of Chinese species for molecular phylogenetic analyses.

Materials and methods

Abbreviations used for collections

BMNH The Natural History Museum, London, United Kingdom.

JXAU College of Agricultural Sciences, Jiangxi Agricultural University, Nanchang, China.

KIZCAS Kunming Institute of Zoology, Chinese Academy of Sciences, Kunming, China.

NIAES National Institute of Agro-Environmental Sciences (ITLJ), Tsukuba, Ibaraki, Japan.

- NMNS** National Museum of Natural Science, Taichung, Taiwan.
NZMC National Zoological Museum of China, Institute of Zoology, Chinese Academy of Science, Beijing, China.
RAB Private collection of Roger A. Beaver, Chiang Mai, Thailand.
RIFID Research Institute of Forest Insect Diversity, Namyangju, South Korea.
SYU Museum of Biology, Sun Yat-sen University, Guangzhou, China.
USNM National Museum of Natural History, Washington D.C., USA.
ZIN Zoological Institute. Russian Academy of Sciences, St. Petersburg, Russia.

Adults of the new species were collected by log dissection. The samples were immediately preserved in tubes containing 99.9% ethyl alcohol, which were stored at -20°C for DNA extraction and examination. Specimens were examined using an Olympus SZX160 stereoscopic zoom microscope. Photographs were taken with a KEYENCE VHX-6000 Digital Microscope System. All photos were further adjusted and assembled with Adobe Photoshop CS6. Body length was measured between the anterior margin of the pronotum and the elytral apex (head not included).

Genomic DNA was extracted from the adult head. The total genomic DNA was extracted from each individual using the Ezup Column Animal Genomic DNA Purification Kit (Sangon Biotech Co. Ltd). Amplification of four gene fragments (COI, EF-1 α , CAD, 28S) was made by PCR, using primers (Table 1) and cycling conditions previously described (Jordal et al. 2011). The PCR products were sent to Sangon Biotech Co. Ltd (Shanghai, China) for sequencing, and the sequences were analyzed using the software DNASTar. Additional information on *Crossotarsus* material was collected by the authors in China or downloaded from NCBI (The National Center for Biotechnology Information) (Table 2). Concatenated DNA sequence data from Jordal (2013) were analysed in MrBayes v. 3.2.6 (Ronquist et al. 2012). Partitions and models were estimated by PartitionFinder 2 (Lanfear et al. 2017) and ModelFinder (Kalyana-moorthy et al. 2017) respectively in PhyloSuite (Zhang et al. 2020), GTR+G+I were selected for each partition. 10 million generations were run, with 25% of the generations as burn-in. PSRF close to 1.0 and standard deviation of split frequencies below 0.01 were accepted.

Table 1. Gene fragments targeted for PCR and the primers used. Sequencing primers were identical to those used in PCR.

Gene	Primer name	Annealing	Primer sequence	Reference
COI	S1718	46	5'-GGAGGATTTGGAAATTGATTAGTTCC-3'	Jordal et al. 2011
	A2411		5'-GCTAATCATCTAAAAACTTTAATTCCWGTWG-3'	
28S	S3690	55	5'-GAGAGTTMAASAGTACGTGAAAC-3'	Jordal et al. 2011
	A4394		5'-TCGGAAGGAACCAGCTACTA-3'	
EF-1 α	S149	52	5'-ATCGAGAAGTTCGAGAAGGAGGCYCARGAAATGGG-3'	Jordal et al. 2011
	A1043		5'-GTATATCCATTGGAATTTGACCCNGGRTGRTT-3'	
CAD	CAD for4	50	5'-TGGAARGARTBGARTACGARGTGGTYCG-3'	Jordal et al. 2011
	CAD rev1mod		5'-GCCATYRCYTCBCCYACRCTYTTTCAT-3'	

Table 2. Material used for phylogenetic analyses, including their GenBank accession numbers.

No.	Taxon	Country	CAD	COI	EF-1 α	28S	Reference
1	<i>C. beaveri</i> sp. nov.	China: Jiangxi	LC616080	LC613149	–	LC613157	This study
2	<i>C. brevis</i> (Browne, 1975)	China: Yunnan	LC616086	LC613154	LC616520	LC613163	This study
3	<i>C. chalcographus</i> Schedl, 1972	Papua New Guinea	KR261163	KR261313	–	–	Jordal 2015
4	<i>C. emorsus</i> Beeson, 1937	China: Yunnan	LC616087	LC613155	–	LC613164	This study
5	<i>C. externedentatus</i> (Fairmaire, 1849)	China: Yunnan	LC616083	LC613152	LC616518	LC613160	This study
6	<i>C. externedentatus</i> (Fairmaire, 1849)	Tanzania	KR261162	KR261312	–	KR261216	Jordal 2015
7	<i>C. externedentatus</i> (Fairmaire, 1849)	Madagascar	KR261166	KR261316	KR261275	KR261218	Jordal 2015
8	<i>C. fractus</i> Sampson, 1912	Papua New Guinea	KR261165	KR261315	KR261274	–	Jordal 2015
9	<i>C. minusculus</i> Chapuis, 1865	Papua New Guinea	HQ883809	HQ883669	HQ883739	HQ883579	Jordal 2015
10	<i>C. niponicus</i> Blandford, 1894	China: Sichuan	–	LC613156	–	LC613165	This study
11	<i>C. nitescens</i> Schedl, 1979	Australia	KR261161	KR261311	KR261272	–	Jordal 2015
12	<i>C. sauteri</i> (Strohmeyer, 1913)	China: Jiangxi	LC616081	LC613150	LC616516	LC613158	This study
13	<i>C. squamulatus</i> Chapuis, 1865	China: Yunnan	LC616084	LC613153	–	LC613161	This study
14	<i>C. terminatus</i> Chapuis, 1865	China: Jiangxi	LC616082	LC613151	LC616517	LC613159	This study
15	<i>C. wallacei</i> (Thomson, 1857)	China: Yunnan	LC616085	–	LC616519	LC613162	This study
16	<i>P. contaminatus</i> (Blandford, 1894)	China: Jiangxi	LC387560	LC383433	LC387562	LC386151	Lai et al. 2019

Results

Crossotarsus beaveri Lai & Wang, sp. nov.

<http://zoobank.org/B8D65F2C-90C7-4B5B-84D0-AA714D42A565>

Figures 1, 2

Type material. Holotype. Male, CHINA: Jiangxi Province, Ganzhou City, Longnan County, Jiulianshan national nature reserve of Jiangxi, Hualu Village, 24°37'19"N, 114°29'57"E, 2.VII.2020, log dissection, host *Paulownia fortunei*, Shengchang Lai leg. (deposited in NZMC IOZ(E)225775).

Allotype. Female, same data as holotype (deposited in NZMC IOZ(E)225776).

Paratypes. 6 males, 6 females, same data as holotype, but host *Phoebe zhennan* and *Liquidambar formosana* (5 males, 5 females JXAU; 1 male, 1 female NZMC); 11 male, 6 females, as holotype except: Xunwu County, Xiangshan Town, Congkeng Village, 24°54'20"N, 115°52'44"E, ca 650m, 15.IX.2017, log dissection, host *Castanopsis fargesii* and *Vernicia montana*, Shengchang Lai leg. (10 males, 5 females JXAU; 1 male, 1 female RAB); 6 males, 6 females, as holotype except: Xunwu County, Liuche Town, Luanluozhang, 24°40'41"N, 115°44'9"E, ca 640 m, 22.VIII.2017, log dissection, host *Castanopsis carlesii*, Shengchang Lai leg. (5 males, 5 females JXAU; 1 male, 1 female RAB); 38 males, 38 females, China: Fujian Province, Zhangzhou City, Yunxiao County, Xiahe Town, Qigaoqi Village, 24°1'31"N, 117°10'36"E, 8.VII.2019, log dissection, host *C. carlesii*, Ling Zhang leg. (2 males, 2 females BMNH; 2 males, 2 females KIZCAS [KIZ0121459–0121462]; 2 males, 2 females NIAES; 2 males, 2 females NMNS; 2 males, 2 females RAB; 2 males, 2 fe-




Figure 1. Male of *Crossotarsus beaveri* sp. nov. **A** dorsal view **B** head **C** lateral view **D** declivity. Scale bars: 0.5 mm.

males RIFID; 2 males, 2 females SYU; 2 males, 2 females USNM; 2 males, 2 females ZIN; 20 males, 20 females JXAU).

Description. Male. 3.58–4.01 mm long (mean = 3.78; $n = 20$); 2.75–2.95 times as long as wide. Head and pronotum dark brown, disc of elytra reddish brown becoming dark brown, declivity of elytra nearly black.

Head. Frons flat, slightly shining, with irregular large punctures; finely, sparsely punctured above the epistoma, bearing bristly, erect, long setae, weakly concave, smooth around short median line, upper part of frons with scattered, coarse punctures, the punctures with moderate, semierect, dorsally directed setae. Antennal scape clavate with scattered, forwardly directed hairs on apical half; club oval, flattened, evenly covered with short setae. Labial palps two-segmented, with basal segments fused along the midline.

Pronotum. About 1.2 times longer than wide, shining, no mycangial pores, the lateral femoral grooves angulate anteriorly, pronotum widest in front of the grooves, with finely, scattered, irregular punctures, a few semierect backwardly pointed hairs close to anterior margin, median line extending about 1/4 from base.

Scutellum. Depressed below level of elytra, with a median longitudinal groove between lateral carinae.

Elytra. About 2.0 times as long as wide, about 1.4 times as long as pronotum. Surface of disc smooth, shining, striae distinctly impressed for almost their entire length, except striae 6 and 7, other striae with circular, distinct, shallow punctures, the bases of striae 1 and 2, striae 3 and 4, respectively, conjoint, more impressed; interstriae slightly raised on disc, interstriae 1, 3 and 5 distinctly raised and conjoint at base,

interstriae 8 and 9 fused at apex of disc, forming ventral, rounded angle; cylindrical declivity obliquely truncate, acutely margined all around except at sutural apex, strongly concave, forming a cup-like structure, surface shining, with 4 rows of longitudinal granules bearing erect, long, golden setae, a row of sparse, medially directed, erect golden setae at the inner margin of declivity, elytral apex broadly emarginate, the main emargination approximately U-shaped, about as wide as deep, extending about 1/3 of the height of the declivity, at its inner end a much smaller, V-shaped second emargination (Fig. 1A, D).

Protibia. Five transverse carinae at tibial apex, transverse rugae at base.

Abdomen. Abdominal ventrites 1–4 moderately finely punctured, with irregular rows of erect, short hairs at both sides posteriorly, ventrite 5 strongly concave at middle, with dense, large, circular punctures.

Female. 3.64–4.42 mm long (mean = 3.96 mm; $n = 20$); 2.79–2.93 times as long as wide. Head and pronotum brown, disc of elytra reddish brown becoming dark brown to apex.

Head. Similar to male, but frons flatter, very shining, smooth, with shallow, small punctures; finely, sparsely punctured above the epistoma, bearing bristly, erect, long setae; very shallowly concave in median line, upper part of frons with scattered, shallow, small punctures, the punctures with moderate, semierect, dorsally directed setae.

Pronotum. Similar to male.

Elytra. About 1.8 times as long as wide, about 1.5 times as long as pronotum, sides subparallel. Similar to male, but disc of elytra shining, with dense, longitudinal, semierect, backwardly pointed hairs at apex and declivity, striae weakly impressed, interstriae smoother, declivity vertical, a few irregularly granules, sparsely hairy.

Protibia. Three transverse carinae at tibial apex, fine, confused granules at base.

Abdomen. Surface of abdominal ventrites smooth, rounded, sparsely hairy, ventrite 5 without concavity, punctures shallow.

Etymology. The species is named for Roger A. Beaver to honor his contributions to the study of platypodines and scolytines.

Host plants. *Castanopsis carlesii* (Hemsl.) Hayata, *C. fargesii* Franch. (Fagaceae), *Liquidambar formosana* Hance (Altingiaceae), *Phoebe zhennan* S.K.Lee & F.N.Wei (Lauraceae), *Paulownia fortunei* (Seem.) Hemsl. (Paulowniaceae), *Vernicia montana* Lour. (Euphorbiaceae).

Distribution. China (Jiangxi, Fujian).

Diagnosis. The species is placed in *Crossotarsus* because it possesses a combination of characters similar to that cited in the introduction.

Crossotarsus beaveri is very similar to *Crossotarsus brevis* (Browne, 1975) (new combination, see below) and *Crossotarsus platypoides* (Browne, 1955). They can be easily distinguished from other *Crossotarsus* species by the male elytral apex truncate with a large, circular, concave declivity. The elytral apex of male of *C. beaveri* and that of *C. brevis* possess a deep, acutely margined declivity, with a broad, almost circular, apical emargination.




Figure 2. Female of *Crossotarsus beaveri* sp. nov. **A** dorsal view **B** head **C** lateral view **D** declivity. Scale bars: 0.5 mm.

Key to the species of *Crossotarsus* with a circular, truncate elytral declivity

- 1 Male elytral apex truncate, with a circular, shallow, concave, bluntly margined declivity; sutural apex of declivity slightly dehiscent without apical emargination. Female smaller and stouter, 2.60–2.70 mm long, 2.70–2.75 times as long as wide ***C. platypoides* Browne**
- Male elytral apex truncate, with a circular, deep, concave, acutely margined declivity, with a broad, almost circular, apical emargination. Female larger and more elongate, 3.00–3.90 mm long, 2.79–3.44 times as long as wide **2**
- 2 Male striae weakly impressed on disc of elytra (Fig. 1A); declivity gradually, obliquely truncate, its face shining, cylindrical, apex rounded with a double sutural emargination, borders of inner emargination weakly elevated, outer emargination forming pointed angles; surface of declivity with 4 longitudinal rows of granules, bearing erect, long golden setae (Fig. 1D). Female frons flat, more shining, smoother, very shallowly concave in median line; dense, shallow, small punctures bearing semierect hairs on upper part; almost flat above the epistoma below median line (Fig. 2B); striae weakly impressed on disc of elytra (Fig. 2A). 3.64–3.90 mm long ***C. beaveri* sp. nov.**
- Male striae moderately impressed on disc of elytra (Fig. 3A); declivity abruptly, vertically truncate, its face subnitid, cylindrical, apex rounded with a double sutural emargination, borders of inner emargination distinctly elevated and dilated, outer emargination forming obtuse angles; surface of declivity with

sparse, obscure granules, bearing erect, long golden setae (Fig. 3D). Female frons slightly shining, reticulate, very distinctly concave, smooth around median line; dense, deep, large punctures bearing semierect hairs on upper part; weakly, irregularly impressed above the epistoma below median line (Fig. 4B); striae moderately impressed on disc of elytra (Fig. 4A). 2.96–3.44 mm long..

..... ***C. brevis* (Browne)**

***Crossotarsus brevis* (Browne, 1975), comb. nov.**

Figures 3, 4

Platypus brevis Browne: Beaver and Browne 1975: 306.

Dinoplatypus brevis Browne: Beaver 1998:184.

Material examined. 7 males, 5 females (JXAU); 1 male, 1 female (RAB): CHINA: Yunnan Province, Xishuangbanna Dai Autonomous Prefecture, Jinghong City, Damanmi Village, 22°02'50"N, 100°48'27"E, ca 580 m, 20.I.2018, log dissection, host unknown, Shengchang Lai leg.

Taxonomy. The specimens in the RAB were identified by comparison to a paratype *C. brevis*, which is also in the RAB. Browne put this species in *Platypus* Herbst, noting that the apical emargination of the elytra was rather similar to that of *Platypus caliculus* Chapuis 1865 (Beaver and Browne 1975). In fact, *C. brevis* has the typical characters of *Crossotarsus*: labial palps two-segmented, with basal segments fused in the midline, whereas *Platypus* has the labial palps three-segmented, with separate basal segments. Beaver (1998) transferred the species from *Platypus* to *Dinoplatypus* Wood following Wood's (1993) attempt to split the genus *Platypus*. Wood diagnosed *Dinoplatypus* largely on the basis of the circular, truncate, elytral declivity of the male, with the sutural apex emarginate. However, this is an adaptive character of the declivity which has evolved independently more than once in the Platypodinae, as it has in the Scolytinae (Hulcr et al. 2015). Molecular phylogenetic study also shows that the few morphological characters used by Wood (1993) to erect several groups of Neotropical and Indo-Malayan/Australasian species in Platypodini to new genera are not sufficiently diagnosable for all those groups (Jordal 2015).

Browne (1961) and Beaver and Sanguansub (2015) suggested that the adult generic characters of primary value in *Crossotarsus* included the structure of the labial and maxillary palps, the form of the pronotum, the sexual dimorphism of the protibia, and various modifications of the abdominal sternites in the male. Based on the two-segmented labial palps, the lateral pronotal emarginations angulate anteriorly, the pronotum without mycangial pores, and the sexual dimorphism of the protibiae, *Platypus brevis* belongs in the genus *Crossotarsus*, and is here transferred to that genus.

Distribution. Thailand (Beaver and Liu 2013). New to China (Yunnan).

Host. *Castanopsis* sp. (Fagaceae) (Beaver and Liu 2013).




Figure 3. Male of *Crossotarsus brevis* (Browne) **A** dorsal view **B** head **C** lateral view **D** declivity. Scale bars: 0.5 mm.




Figure 4. Female of *Crossotarsus brevis* (Browne) **A** dorsal view **B** head **C** lateral view **D** declivity. Scale bars: 0.5 mm.

New record

Crossotarsus emorsus Beeson, 1937

Figures 5, 6

Crossotarsus emorsus Beeson, 1937: 87.

Material examined. 4 males, 1 female (JXAU) CHINA: Yunnan Province, Xishuangbanna Dai Autonomous Prefecture, Jinghong City, Nabanhe River Watershed National Nature Reserve, Guomenshan, ca 1030 m, 22°14'46"N, 100°36'10"E, 27.I.2018, log dissection, host *Dalbergia assamica*, Shengchang Lai leg.; 1 male, 1 female (RAB); 1 male (JXAU) China: Yunnan Province, Xishuangbanna Dai Autonomous Prefecture, Jinghong City, Damanmi Village, ca 580 m, 22°02'50"N, 100°48'27"E, 20.I.2018, log dissection, host *Cassia siamea*, Shengchang Lai leg.

Diagnosis. *C. emorsus* is similar to *C. terminatus* but can be distinguished using the characters given in Table 3.

Distribution. Myanmar, Thailand, Laos (Beaver and Liu 2013; Beaver 2016). New to China (Yunnan).

Host. The species is recorded from trees in the families Lecythidaceae, Fabaceae, Sterculiaceae and Verbenaceae (Beeson 1937), and is presumably polyphagous (Beaver 2016). Host plants recorded here are: *Senna siamea* (Lam.) H.S.Irwin & Barneby and *Dalbergia assamica* Benth. (Fabaceae).

Molecular data. The phylogenetic tree for analyzing the evolutionary relationships of 13 taxa including the ingroups (*Crossotarsus* species) and the outgroups (*P. contaminatus*) was constructed based on four genes (Fig. 7). The BI tree shows the new species (*C. beaveri*) and the new combination (*C. brevis*) forming a clade, with high node support. These group with Schedl's (1972a) '*Crossotarsi coleoprati*' (*C. fractus* Sampson, 1912, *C. squamulatus* and *C. terminatus*) and cluster with all remaining *Crossotarsus* species. It confirms that the taxonomic changes and the relationship of *C. brevis* and *C. brevis* are correct. It also indicates that *C. emorsus*, *C. fractus*, *C. squamulatus* and *C. terminatus* should be considered distinct species (as by Beaver and Liu 2013), and not considered synonyms or subspecies (Schedl 1972a).

Discussion

Crossotarsus beaveri is clearly related to *C. brevis*. They are the sister lineage to the *Crossotarsi coleoprati* group, not the genus *Dinoplatypus*. This is a good example that declivity in males usually is an adaptive character and not of generic significance. We consider that the morphologically diagnosable characters of the genus *Crossotarsus* should refer to the summary by Browne (1961) and Beaver and Sanguansub (2015, 2020), as introduction.




Figure 5. Male of *Crossotarsus emorsus* Beeson **A** dorsal view **B** head **C** lateral view **D** declivity. Scale bars: 0.5 mm.




Figure 6. Female of *Crossotarsus emorsus* Beeson **A** dorsal view **B** head **C** lateral view **D** declivity. Scale bars: 0.5 mm.

References

- Beaver RA (1998) New synonymy, new combinations and taxonomic notes on Scolytidae and Platypodidae (Insecta: Coleoptera). *Annalen des Naturhistorischen Museums in Wien. Serie B für Botanik und Zoologie* 100: 179–192.
- Beaver RA (2004) The genus *Crossotarsinulus* Schedl (Coleoptera: Platypodidae). *Entomologist's Monthly Magazine* 140: 243–245.
- Beaver RA (2005) New synonymy in Taiwanese ambrosia beetles (Coleoptera: Curculionidae: Platypodinae). *Plant Protection Bulletin* 47: 195–200.
- Beaver RA (2016) The platypodine ambrosia beetles of Laos (Coleoptera: Curculionidae: Platypodinae). *Entomologica Basiliensia et Collectionis Frey* 35: 487–504.
- Beaver RA, Browne FG (1975) The Scolytidae and Platypodidae (Coleoptera) of Thailand: a checklist with biological and zoogeographical notes. *Oriental Insects* 9(3): 283–211. <https://doi.org/10.1080/00305316.1975.10434499>
- Beaver RA, Liu LY (2013) A synopsis of the pin-hole borers of Thailand (Coleoptera: Curculionidae: Platypodinae). *Zootaxa* 3646(4): 447–486. <https://doi.org/10.11646/zootaxa.3646.4.7>
- Beaver RA, Sanguansub S (2015) A review of the genus *Carchesiopygus* Schedl (Coleoptera: Curculionidae: Platypodinae), with keys to species. *Zootaxa* 3931(1): 401–412. <https://doi.org/10.11646/zootaxa.3931.3.4>
- Beaver RA, Sanguansub S (2020) New synonymy and taxonomic changes in Australian and oriental pin-hole borers (Coleoptera: Curculionidae, Platypodinae). *Entomologist's Monthly Magazine* 156(2): 79–86. <https://doi.org/10.31184/M00138908.1562.4024>
- Beaver RA, Shih HT (2003) Checklist of Platypodidae (Coleoptera: Curculionoidea) from Taiwan. *Plant Protection Bulletin* 44: 75–90.
- Beeson CFC (1937) New *Crossotarsus* (Platypodidae, Col.). *The Indian Forest Records* 3: 47–103.
- Bright DE (2014) A catalog of Scolytidae and Platypodidae (Coleoptera), Supplement 3 (2000–2010), with notes on subfamily and tribal reclassifications. *Insecta Mundi* 0356: 1–336.
- Blandford WFH (1894) The rhynchophorous Coleoptera of Japan. Part III. Scolytidae. *Transactions of the Entomological Society of London*, 53–141.
- Browne FG (1961) Taxonomic notes on Platypodidae (Coleoptera). *Annals and Magazine of Natural History, Series 13*, 4(47): 641–656. <https://doi.org/10.1080/00222936108651189>
- Chapuis F (1865) *Monographie des Platypides*. H. Dessain, Liège, 344 pp.
- Fairmaire L (1849) *Essai sur les coléoptères de la Polynésie*. *Revue et Magasin de Zoologie Pure et Appliquée, Série 2*, 2: 1–102.
- Goto H (2009) Taxonomic history of Japanese bark and ambrosia beetles with a check list of them. *Journal of Japanese Forest Society* 91: 479–485. <https://doi.org/10.4005/jjfs.91.479>
- Hopkins AD (1914) List of generic names and their type-species in the coleopterous superfamily Scolytoidea. *Proceedings of the United States National Museum* 48(2066): 115–136. <https://doi.org/10.5479/si.00963801.2066.115>
- Hulcr J, Atkinson TH, Cognato AI, Jordal BH, Mckenna DD (2015) Morphology, taxonomy, and phylogenetics of bark beetles. In: Vega FE, Hofstetter RW (Eds) *Bark Beetles. Biology and Ecology of Native and Invasive Species*. Academic Press, London, 41–84. <https://doi.org/10.1016/B978-0-12-417156-5.00002-2>

- Jordal BH (2013) Deep phylogenetic divergence between *Scolytoplatypus* and *Remansus*, a new genus of Scolytoplatypodini from Madagascar (Coleoptera, Curculionidae, Scolytinae). *ZooKeys* 352: 9–33. <https://doi.org/10.3897/zookeys.352.6212>
- Jordal BH (2015) Molecular phylogeny and biogeography of the weevil subfamily Platypodinae reveals evolutionarily conserved range patterns. *Molecular Phylogenetics and Evolution* 92: 294–307. <https://doi.org/10.1016/j.ympev.2015.05.028>
- Jordal BH, Sequeira AS, Cognato AI (2011) The age and phylogeny of wood boring weevils and the origin of subsociality. *Molecular Phylogenetics and Evolution* 59: 708–724. <https://doi.org/10.1016/j.ympev.2011.03.016>
- Kalyaanamoorthy S, Minh BQ, Wong TK, von Haeseler A, Jermini LS (2017) ModelFinder: fast model selection for accurate phylogenetic estimates. *Nature Methods* 14(6): 587–589. <https://doi.org/10.1038/nmeth.4285>
- Lai SC, Liao JX, Dai XH, Wang YX, Wang JG (2019) Identification of hawthorn trunk borer (*Platypus contaminatus*), an important insect pest on hawthorn. *Plant Quarantine* 33(1): 48–51. [in Chinese with English summary]
- Lanfear R, Frandsen PB, Wright AM, Senfeld T, Calcott B (2017) PartitionFinder 2: new methods for selecting partitioned models of evolution for molecular and morphological phylogenetic analyses. *Molecular Biology and Evolution* 34(3): 772–773. <https://doi.org/10.1093/molbev/msw260>
- Murayama J (1925) Supplementary notes on the Platypodidae of Formosa. *Journal of the College of Agriculture, Hokkaido Imperial University* 15: 229–235.
- Murayama J (1934) Supplementary notes on the platypodidae of Formosa IV. *Journal of the Faculty of Agriculture, Hokkaido Imperial University* 35(3): 133–149.
- Ronquist F, Teslenko M, van der Mark P, Ayres DL, Darling A, Höhna S, Larget B, Liu L, Suchard MA, Huelsenbeck JP (2012) MrBayes 3.2: efficient Bayesian phylogenetic inference and model choice across a large model space. *Systematic Biology* 61: 539–542. <https://doi.org/10.1093/sysbio/sys029>
- Sampson FW (1912) Some new species of Ipidae and Platypodidae in the British Museum. *Annals and Magazine of Natural History, Series 8, 10*: 245–250. <https://doi.org/10.1080/00222931208693228>
- Schedl KE (1954) Fauna Indomalayensis IV. *Philippine Journal of Science* 83: 137–159.
- Schedl KE (1972a) Monographie der Familie Platypodidae Coleoptera. W. Junk, Den Haag, 322 pp.
- Schedl KE (1972b) Schedl K E. Scolytidae and Platypodidae from the Papuan subregion and Australia. 279. Contribution to the morphology and taxonomy of the Scolytidae. *Papua New Guinea Agricultural Journal* 23(3/4): 61–72.
- Schedl KE (1979) Bark and timber beetles from Australia. 326. Contribution to the morphology and taxonomy of the Scolytoidea. *Entomologische Arbeiten aus dem Museum G. Frey* 28: 157–163.
- Stebbing EP (1906) Departmental notes on insects that affect forestry. No. 3. Office of Superintendent of Government Printing, Calcutta, 335–469.
- Strohmeyer H (1912) H. Sauter's Formosa-ausbeute, Ipidae und Platypodidae (Col.). *Entomologische Mitteilungen* 1: 38–42. <https://doi.org/10.5962/bhl.part.25902>

- Strohmeyer H (1913) Neue Platypodiden. Entomologische Blätter 9: 161–165.
- Thomson J (1857) Description de trois espèces de coléoptères. Archives Entomologiques 1: 341–344.
- Wood SL (1993) Revision of the genera of Platypodidae (Coleoptera). Great Basin Naturalist 53(3): 259–281. <https://doi.org/10.5962/bhl.part.16605>
- Wood SL, Bright DE (1992) A catalog of Scolytidae and Platypodidae (Coleoptera), Part 2: taxonomic index. Great Basin Naturalist Memoirs 13: 1–1553.
- Yin HF, Huang FS, (1987) Platypodidae. In: Huang FS, Zheng LY (Eds) Forest Insects of Yunnan. Yunnan Science and Technology Press, Kunming, 854–858. [in Chinese]
- Yin HF, Huang FS, Zeng R, Li H (2002) Coleoptera: Platypodidae. In: Huang FS (Eds) Forest Insects of Hainan. Science Press, Beijing, 472–473. [in Chinese]
- Zhang D, Gao F, Jakovlić I, Zou H, Zhang J, Li WX, Wang GT (2020) PhyloSuite: an integrated and scalable desktop platform for streamlined molecular sequence data management and evolutionary phylogenetics studies. Molecular Ecology Resources 20(1): 348–355. <https://doi.org/10.1111/1755-0998.13096>
- Zhang Y, Du YZ, Zhu HB, Gu J, Zhang YZ (2008) Species record of Chinese *Crossotarsus*. Entomological Journal of East China 17(3): 205–212. [in Chinese with English summary]