

Contributions to the knowledge of oribatid mites of Indonesia. 2. The genus *Pergalumna* (Galumnidae) with description of a new species and key to known species in the Oriental region (Acari, Oribatida)

Sergey G. Ermilov¹, Dorothee Sandmann², Bernhard Klarner²,
Rahaju Widystuti³, Stefan Scheu²

1 Tyumen State University, Tyumen, Russia **2** Georg August University Göttingen, J.F. Blumenbach Institute of Zoology and Anthropology, Göttingen, Germany **3** Institut Pertanian Bogor, Bogor, Indonesia

Corresponding author: Sergey G. Ermilov (ermilovacari@yandex.ru)

Academic editor: V. Pesic | Received 31 August 2015 | Accepted 15 September 2015 | Published 26 October 2015

<http://zoobank.org/6C38D7A7-63BC-4314-8E4C-5BD45EBCC89A>

Citation: Ermilov SG, Sandmann D, Klarner B, Widystuti R, Scheu S (2015) Contributions to the knowledge of oribatid mites of Indonesia. 2. The genus *Pergalumna* (Galumnidae) with description of a new species and key to known species in the Oriental region (Acari, Oribatida). ZooKeys 529: 87–103. doi: 10.3897/zookeys.529.6421

Abstract

A new species of oribatid mite of the genus *Pergalumna* (Oribatida, Galumnidae) is described from litter and soil materials in Sumatra, Indonesia. *Pergalumna paraindistincta* sp. n. is morphologically most similar to *P. indistincta* Ermilov & Anichkin, 2011, *P. pertrichosa* Mahunka, 1995 and *P. sura* Balogh, 1997; however, the new species differs from *P. indistincta* by the smaller body size, presence of long adanal setae *ad*₁, and large, single median pore in females and males; from *P. pertrichosa* by the smaller body size, presence of three pairs of notogastral porose areas, elongated *A1* and minute anal setae; from *P. sura* by the presence of strong adanal setae *ad*₁, large, single median pore in females and males, and shorter bothridial setae. Furthermore, *Pergalumna hawaiiensis hawaiiensis* (Jacot, 1934) and *P. panayensis* Ermilov & Corpuz-Raros, 2015 are recorded for the first time in the Indonesian fauna. An identification key to the known species of *Pergalumna* in the Oriental region is given.

Keywords

Oribatid mites, *Pergalumna*, new species, new record, key, Indonesia, Oriental region

Introduction

This work is a part of a continuing study on the Indonesian fauna of oribatid mites, and it includes the data on the genus *Pergalumna* Grandjean, 1936 (Oribatida, Galumnidae). During taxonomic identification, four species were identified, including one new to science. The primary goal of the paper is to present data on the specific localities, notes on new records and overall known distributions of registered taxa and to describe the new species.

Pergalumna is a genus that was proposed by Grandjean (1936) with *Oribata nervosa* Berlese, 1914 as type species. Based on an updated generic diagnosis (Ermilov et al. 2013b), it comprises more than 140 species (Subías 2004, updated 2015; Ermilov and Bayartogtokh 2015; Ermilov and Corpuz-Raros 2015) having collectively a cosmopolitan distribution (Subías 2004, updated 2015). The identification keys to selected species were given by Shaldybina (1975), Balogh and Balogh (1990, 2002), Weigmann (2006), and Ermilov et al. (2014b). The secondary goal of the paper is to provide an identification key to known species of *Pergalumna* in the Oriental region.

Materials and methods

Exact collection locality and habitat are given in the respective “Material examined” section for each species.

Specimens were mounted in lactic acid on temporary cavity slides for measurement and illustration. The body length was measured in lateral view, from the tip of the rostrum to the posterior edge of the ventral plate. Notogastral width refers to the maximum width in dorsal aspect. Lengths of body setae were measured in lateral aspect. All body measurements are presented in micrometers. Formulas for leg setation are given in parentheses according to the sequence trochanter–femur–genu–tibia–tarsus (famulus included). Formulas for leg solenidia are given in square brackets according to the sequence genu–tibia–tarsus.

General terminology used in this paper follows that of Grandjean (summarized by Norton and Behan-Pelletier 2009).

Drawings were made with a camera lucida using a Carl Zeiss transmission light microscope “Axioskop-2 Plus”.

Description

Pergalumna paraindistincta sp. n.

<http://zoobank.org/F1F876B8-867F-4EA9-851D-FB4186C05342>

Figs 1–9

Diagnosis. Body size: 415–481 × 298–365. Rostral, lamellar and interlamellar setae well developed, barbed. Bothridial setae long, setiform, ciliate. Anterior notogastral




Figure 1. *Pergalumna paraindistincta* sp. n., adult: dorsal view. Scale bar 100 μ m.

margin not developed. Three pairs of elongate oval porose areas on notogaster, *Aa* transversally oriented, located between *la* and *lm*, *A1* longitudinally oriented. Median pore single, large. Adanal setae *ad*₁ of medium size, straight, heavily barbed. Postanal porose area absent.

Description. Measurements. Body length: 431 (holotype: male), 415–481 (10 paratypes: three females and seven males); notogaster width: 298 (holotype), 298–365 (10 paratypes). Without sexual dimorphism.

Integument. Body color brown. Body surface microgranulate, visible under high magnification, $\times 1000$ (diameter of granules less than 1).

Prodorsum (Figs 1, 3, 5). Rostrum broadly rounded. Lamellar (*L*) and sublamellar (*S*) lines distinct, parallel, curving backwards. Rostral (*ro*, 41–49), lamellar (*le*, 69–77) and interlamellar (*in*, 86–90) setae setiform, barbed. Bothridial setae (*bs*, 114–127)




Figure 2. *Pergalumna paraindistincta* sp. n., adult: ventral view (gnathosoma and legs not shown). Scale bar 100 μm .

setiform, densely ciliate in medio-distal parts. Exobothridial setae and their alveoli absent. Porose areas *Ad* narrowly elongate oval, transversally oriented ($16–20 \times 4$).

Notogaster (Figs 1, 3, 4). Anterior notogastral margin not developed. Dorsophragmata (*D*) of medium size, elongated longitudinally. Notogastral setae represented by 10 pairs of alveoli. Three pairs of porose areas: *Aa* clearly bordered, elongate oval, transversally oriented ($28–32 \times 8–12$), *A1* distinctly bordered only in anterior part, elongate oval to slightly elongate triangular, longitudinally oriented ($57–68 \times 12–16$), *A3* without clear borders, elongate oval ($24–28 \times 8–12$). Areas *Aa* located between setal alveoli

la and *lm*, equal distanced from them. Median pore present in males and females, comparatively large (diameter 10–16). All lyrifissures (*ia*, *im*, *ip*, *ih*, *ips*) distinct, *im* located antero-laterally to *A1*. Opisthonotal gland openings (*gla*) located laterally to *A1*.

Gnathosoma (Fig. 6). Morphology of subcapitulum, palps and chelicerae typical for *Pergalumna* (see Engelbrecht 1972; Ermilov and Anichkin 2011a, b). Subcapitulum size: 102–106 × 102–106. Subcapitular setae setiform, slightly barbed, *m* (14–16) shorter than *a* and *b* (both pairs 18–20); *a* thickest, *m* thinnest. Two pairs of adoral setae (*or₁*, *or₂*, 12–14) setiform, hook-like distally, barbed. Palps (90–94) with typical setation: 0–2–1–3–9(+ω). Axillary sacculi (*sac*) distinct. Chelicerae (164) with two setiform, barbed setae; *cha* (41) longer than *chb* (24). Trägårdh's organ long, tapered.


Epimeral and lateral podosomal regions (Fig. 2). Anterior tectum of epimere I smooth. Apodemes 1, 2, sejugal and 3 well visible. Six pairs of setae, setal formula: 1–0–2–3. Setae thin, slightly barbed, *1a*, *3c* and *4c* (24) longer than *3b* (10–14) and *4a* and *4b* (4); *1a* thicker than others. Pedotecta II (Pd II) scale-like in lateral view, rounded distally in ventral view. Discidia (*dis*) sharply triangular. Circumpedal carinae (*cp*) slightly developed.

Anogenital region (Figs 2, 4, 7, 8). Six pairs of genital (*g₁*, *g₂*, 12; *g₂–g₃*, 8), one pair of aggenital (*ag*, 8), two pairs of anal (*an₁*, *an₂*, 8) and two pairs of adanal setae (*ad₂*, *ad₃*, 8) thin, indistinctly barbed. One pair of adanal setae (*ad₁*, 24–36) thickened, straight, heavily barbed, however, in three paratypes one seta of the *ad₁* pair is short, as *ad₂* and *ad₃* in morphology. Adanal setae *ad₃* inserted laterally to adanal lyrifissures (*iad*). Genital plates with two or three setae on anterior edge of each plate. Postanal porose area absent.


Legs (Fig. 9). Morphology of leg segments, setae and solenidia typical for *Pergalumna* (see Engelbrecht 1972; Ermilov and Anichkin 2011a, b). Claws smooth. Formulas of leg setation and solenidia: I (1–4–3–4–20) [1–2–2], II (1–4–3–4–15) [1–1–2], III (1–2–1–3–15) [1–1–0], IV (1–2–2–3–12) [0–1–0]; homology of setae and solenidia indicated in Table 1. Solenidion φ of tibiae IV inserted dorsally in posterior part of segments.

Material examined. Holotype (male): Indonesia, Sumatra, Harapan landscape, secondary rainforest, research site HF1, 02°09'09.9"S, 103°21'43.2"E, 76 m a.s.l., from forest floor litter material. Six paratypes (two females and four males): Indonesia, Sumatra, Harapan landscape, rubber plantation, research site HR2, 01°52'44.5"S, 103°16'28.4"E, 59 m a.s.l., from forest floor litter material. Four paratypes (one female and three males): Sumatra, Indonesia, Harapan landscape, jungle rubber agro-forest, research site HJ1, 01°55'40.0"S, 103°15'33.8"E, 51 m a.s.l., from forest floor litter material. All specimens were collected by Bernhard Klärner (15.XI.2013) and identified and collected to morphospecies level by Dorothee Sandmann.

Type deposition. The holotype is deposited in LIPI (Indonesian Institute of Science) Cibinong, Indonesia; three paratypes are in the collection of the Senckenberg Museum, Görlitz, Germany; seven paratypes are in the collection of the Tyumen State University Museum of Zoology, Tyumen, Russia.



Figures 3–4. *Pergalumna paraindistincta* sp. n., adult: **3** anterior part of body, lateral view (gnathosoma and leg I not shown) **4** posterior view. Scale bar 100 μ m.



Figures 5–9. *Pergalumna paraindistincta* sp. n., adult: 5 bothridial seta 6 anterior part of left half of subcapitulum, ventral view 7 genital plate, right 8 anal plate, left, and adanal setae 9 tibia of leg IV, right, antiaxial view. Scale bar 20 μ m.

Table I. Leg setation and solenidia of adult *Pergalumna paraindistincta* sp. n.

| Leg | Tr | Fe | Ge | Ti | Ta |
|-----|------|----------------|-------------------|----------------------------------|--|
| I | v' | $d, (l), bv''$ | $(l), v', \sigma$ | $(l), (v), \varphi_1, \varphi_2$ | $(ft), (tc), (it), (p), (u), (a), s, (pv), v', (pl), l'', \varepsilon, \omega_1, \omega_2$ |
| II | v' | $d, (l), bv''$ | $(l), v', \sigma$ | $(l), (v), \varphi$ | $(ft), (tc), (it), (p), (u), (a), s, (pv), \omega_1, \omega_2$ |
| III | v' | d, ev' | l', σ | $l', (v), \varphi$ | $(ft), (tc), (it), (p), (u), (a), s, (pv)$ |
| IV | v' | d, ev' | d, l' | $l', (v), \varphi$ | $ft'', (tc), (p), (u), (a), s, (pv)$ |

Note: Roman letters refer to normal setae, Greek letters to solenidia (except ε = famulus). Single prime ($'$) marks setae on the anterior and double prime ($''$) setae on the posterior side of a given leg segment. Parentheses refer to a pair of setae. Tr – trochanter, Fe – femur, Ge – genu, Ti – Tibia, Ta – tarsus.

Etymology. The specific name *paraindistincta* refers to the morphological similarity of the new species to *Pergalumna indistincta* Ermilov & Anichkin, 2011.

Remarks. *Pergalumna paraindistincta* sp. n. is morphologically most similar to *P. indistincta* Ermilov & Anichkin, 2011 from Vietnam (see Ermilov and Anichkin 2011b) and *P. sura* Balogh, 1997 from the Neotropical region (see Balogh 1997; Ermilov et al. 2014a) in having rounded rostrum, well-developed prodorsal setae, setiform and ciliate of bothridial setae, three pairs of notogastral porose areas, transversally oriented *Aa* and strongly elongated, longitudinally oriented *A1*, and the absence of anterior notogastral margin as well as postanal porose area. However, the new species differs from both by the presence of strong adanal setae *ad*₁ (vs. minute in *P. indistincta* and *P. sura*) and large, single median pore in females and males (vs. median pore absent in males and represented by several foveae in females in *P. indistincta*, and absent in females and males in *P. sura*). Additionally, the new species is smaller than *P. indistincta* (415–481 × 298–365 vs. 547–614 × 381–415) and has shorter bothridial setae than *P. sura*.

Furthermore, *P. paraindistincta* sp. n. is morphologically similar to *P. pertrichosa* Mahunka, 1995 from Borneo (see Mahunka 1995) in having a rounded rostrum, well developed prodorsal setae, setiform and ciliate bothridial setae, strong adanal setae *ad*₁, a single median pore, and the absence of an anterior notogastral margin. However, the new species differs from the latter by the smaller body size (415–481 × 298–365 vs. 550–608 × 413–454 in *P. pertrichosa*), presence of three pairs of notogastral porose areas with elongated *A1* (vs. four pairs of porose areas with *A1* rounded in *P. pertrichosa*) and minute anal setae (vs. well developed in *P. pertrichosa*).

Records

Pergalumna hawaiiensis hawaiiensis (Jacot, 1934) (see Jacot 1934a). Distribution: Pacific Islands. New record for Indonesia.

Material examined. One specimen: Indonesia, Sumatra, Harapan landscape, Jungle rubber agroforest, research site HJ1, 01°55'40.0"S, 103°15'33.8"E, 51 m a.s.l., from upper soil layer (0–5 cm), 15.XI.2013 (B. Klarner). Three specimens: Indonesia, Sumatra, Bukit Duabelas landscape, rubber plantation, research site BR2, 02°05'06.8"S, 102°47'20.7"E, 95 m a.s.l., from upper soil layer (0–5 cm), 15.XI.2013 (B. Klarner). One specimen: Indonesia, Sumatra, Bukit Duabelas landscape, oil palm plantation, research site BO3, 02°04'15.2"S, 102°47'30.6"E, 71 m a.s.l., from upper soil layer (0–5 cm), 15.XI.2013 (B. Klarner).

Pergalumna panayensis Ermilov & Corpuz-Raros, 2015 (see Ermilov and Corpuz-Raros 2015). Distribution: Philippines. New record for Indonesia.

Material examined. One specimen: Indonesia, Sumatra, Harapan landscape, rubber plantation, research site HR2, 01°52'44.5"S, 103°16'28.4"E, 59 m a.s.l., from forest

floor litter material, 15.XI.2013 (B. Klärner). One specimen: same data, but in upper soil layer (0–5 cm). One specimen: Indonesia, Sumatra, Bukit Duabelas landscape, jungle rubber agroforest, research site BJ4, 02°00'57.3"S, 102°45'12.3"E, 60 m a.s.l., from upper soil layer (0–5 cm), 15.XI.2013 (B. Klärner).

Pergalumna pterinervis (Canestrini, 1898) (see Mahunka 1992). Distribution: Oriental region. New record for Indonesia.

Material examined. One specimen: Indonesia, Sumatra, Harapan landscape, jungle rubber agroforest, research site HJ2, 01°49'31.9"S, 103°17'39.2"E, 84 m a.s.l., from forest floor litter material, 15.XI.2013 (B. Klärner). One specimen: Indonesia, Sumatra, Harapan landscape, jungle rubber, research site HJ4, 01°47'07.3"S, 103°16'36.9"E, 57 m a.s.l., from upper soil layer (0–5 cm), 15.XI.2013 (B. Klärner). One specimen: Indonesia, Sumatra, Bukit Duabelas landscape, secondary rainforest, research site BF2, 01° 58'55.1"S, 102°45'02.7"E, 77 m a.s.l., from upper soil layer (0–5 cm), 15.11.2013 (B. Klärner). Three specimens: Indonesia, Sumatra, Harapan landscape, jungle rubber agroforest, research site HJ2, 01°49'31.9"S, 03°17'39.2"E, 84 m a.s.l., from forest floor litter material, 15.XI.2013 (B. Klärner).

Key to known species of *Pergalumna* in the Oriental region

At present, 45 species/subspecies of *Pergalumna* are known in the Oriental region (Subías 2004, updated 2015; including present and personal data of the first author).

Pergalumna heroica (Willmann, 1931) from Java (see Willmann 1931), *P. medialis* (Sellnick, 1925) from Sumatra (see Sellnick 1925) and *P. obsessa* Subías, 2004 from Taiwan (see Tseng 1984 as *Galumna pallida* Tseng, 1984) are excluded from the key because these species have been poorly described.

Pergalumna curva curva (Ewing, 1907) from the Holarctic and Oriental regions (see Ewing 1907; Jacot 1934b), *P. curva ventralis* (Willmann, 1931) from the Holarctic, Neotropical and Oriental regions and Polynesia (see Willmann 1931; Jacot 1934b; Hammer 1958, 1961, 1972), *P. obvia obvia* (Berlese, 1914) from the Ethiopian, Neotropical, Oriental and Palaearctic regions, and Hawaii and U.S.A. (see Weigmann 2006; Ermilov et al. 2013c), *P. operata* Tseng, 1984 from Taiwan (see Tseng 1984) and *P. pyramidalis* (Tseng, 1984) from Taiwan (see Tseng 1984) were not included because systematic placement of these species is not clear. We consider *P. obvia obvia* as a representative of the genus *Galumna* Heyden, 1826 (see Ermilov et al. 2013c). *Pergalumna curva curva*, *P. curva ventralis* and *P. operata* Tseng, 1984 have distinct notogastral setae, which are not traits for *Pergalumna* (see generic diagnosis in Ermilov et al. 2013b); these species are poorly described and redescribed; however, based on available data they should be considered as representatives of the genus *Allogalumna* Grandjean, 1936 or *Trichogalumna* Balogh, 1960 (depending on presence or absence of lamellar lines). *Pergalumna pyrami-*

dalis has 14 pairs of notogastral setal alveoli, developed lamellae directed to insertions of lamellar setae and pteromorphs without setae (Tseng 1984); most likely this species is a representative of the subgenus *Neoribates* (*Neoribates*) Berlese, 1914 (Oripodoidea, Parakalummidae), and it is morphologically most similar to *N. (N.) paratuberculatus* Ermilov, Shtanchaeva & Subías, 2014 from Vietnam (see Ermilov et al. 2014d) and *N. (N.) tuberculatus* Willmann, 1956 from “Czechoslovakia” (see Willmann 1956) in having tubercles on pteromorphs and some other characters.

- | | | |
|---|---|----|
| 1 | Anterior margin of notogaster of specific structure, tuberculate | 2 |
| — | Anterior margin of notogaster simple, smooth or not developed..... | 3 |
| 2 | Genital plates with several striae; notogastral porose areas of medium size, larger than diameter of bothridia; body size: 451–490 × 328–366..... | |
| | <i>P. margaritata</i> Mahunka, 1989 (Mahunka 1989). Distribution: Vietnam | |
| — | Genital plates with one pair of striae; notogastral porose areas small, similar to diameter of bothridia; body size: 402–447 × 281–315..... <i>P. pseudomargaritata</i> Mahunka, 1994 (see Mahunka 1994). Distribution: Thailand | |
| 3 | Anterior margin of notogaster distinctly developed, complete..... | 4 |
| — | Anterior margin of notogaster not developed..... | 12 |
| 4 | Rostrum pointed | 5 |
| — | Rostrum rounded | 7 |
| 5 | Four pairs of notogastral porose areas; <i>Aa</i> elongate triangular, transversally oriented; lateral parts of pteromorphs with strong ridges forming slightly visible reticulate pattern; body size: 517–670 × 397–525..... | |
| | <i>P. altera</i> (Oudemans, 1915) (see Aoki 1961 as <i>P. harunaensis</i> Aoki 1961, 1975; Engelbrecht 1972; Weigmann 2006). Distribution: Semicosmopolitan | |
| — | Three pairs of notogastral porose areas; <i>Aa</i> rounded; pteromorphs without strong ridges and reticulate pattern | 6 |
| 6 | Interlamellar setae long; posterior part of notogaster without furrows; body size: 664–830 × 498–630..... <i>P. yurtaevi</i> Ermilov & Anichkin, 2011 (see Ermilov and Anichkin 2011a; Ermilov et al. 2012a). Distribution: Vietnam | |
| — | Interlamellar setae represented by alveoli; posterior part of notogaster with two parallel, longitudinal furrows; body size: 664–830 × 498–630 | |
| | <i>P. asetosa</i> Ermilov, Shtanchaeva, Kalúz & Subías, 2013 (see Ermilov et al. 2013a). Distribution: India | |
| 7 | Bothridial setae setiform; body size: 520–676 × 502 | |
| | <i>P. foveolata</i> Hammer, 1973 (see Hammer 1973; Bayartogtokh and Chatterjee 2010). Distribution: Australian, Neotropical and Oriental region | |
| — | Bothridial setae with developed head | 8 |
| 8 | Interlamellar setae minute; body surface foveolate; body size: 222–235 × 177–190 | |
| | <i>P. annulata</i> Mahunka, 1995 (see Mahunka 1995). Distribution: Borneo | |
| — | Interlamellar setae long; body surface not foveolate..... | 9 |

- 9 Three pairs of notogastral porose areas; Aa rounded; body length: 820..... ***P. corniculata* (Berlese, 1905)** (see Berlese 1905; Mahunka 1992). Distribution: Java 10
- Four pairs of notogastral porose areas; Aa elongated, transversally oriented ... ***P. taprobanica* Balogh, 1988** (see Balogh 1988). Distribution: Oriental region 11
- 10 Notogastral porose areas Aa triangular; median pore present; body size: 623×533 ***P. andhraense* Raju, Appalanaidu & Rao, 1981** (see Raju et al. 1981). Distribution: India 12
- Bothridial setae fusiform, with well-developed head rounded distally; postanal porose area present; body size: 672×528 ***P. paraelongata* Ermilov & Anichkin, 2012** (see Ermilov et al. 2012b). Distribution: Vietnam 13
- 12 Rostrum trapezoid; anal setae comparatively long, longer than width of anal plate; body size: $1278\text{--}1311 \times 976\text{--}1045$ ***P. paraclericata* Ermilov, Chatterjee, Das & Bordoloi, 2014** (see Ermilov et al. 2014c). Distribution: India 14
- Rostrum pointed 14
- Rostrum rounded 18
- 14 Four pairs of notogastral porose areas; Aa located nearer to setal alveoli la than lm ; body size: $730\text{--}780 \times 564\text{--}597$ ***P. cattienica* Ermilov & Anichkin, 2011** (see Ermilov and Anichkin 2011a). Distribution: Vietnam 15
- Three pairs of notogastral porose areas; Aa located nearer to setal alveoli lm or distanced equal from la and lm 15
- 15 Interlamellar setae represented by alveoli; anterior part of prodorsum with two longitudinal ridges; notogastral porose areas Aa located nearer to setal alveoli lm than la ; body size: $1162\text{--}1278 \times 898\text{--}1012$ ***P. minipora* Ermilov, Chatterjee, Das & Bordoloi, 2014** (see Ermilov et al. 2014c). Distribution: India 16
- Interlamellar setae of medium size or long; prodorsum without ridges; notogastral porose areas Aa distanced equal from la and lm 16
- 16 Notogastral porose areas $A1$ elongated, longitudinally oriented; body surface foveolate; genital plates not striate; body size: $365\text{--}415 \times 265\text{--}332$ ***P. paratsurusakii* Ermilov, Shtanchayeva, Kalúz & Subías, 2013** (see Ermilov et al. 2013a). Distribution: India 17
- Notogastral porose areas $A1$ rounded; body surface not foveolate; genital plates striate 17
- 17 Adanal setae ad_1 and ad_2 comparatively long, not shorter than width of anal plate; median pore absent; interlamellar setae longer than bothridial setae; body size: $597\text{--}680 \times 431\text{--}498$ ***P. paracattienica* Ermilov, Chatterjee, Das & Bordoloi, 2014** (see Ermilov et al. 2014c). Distribution: India

- Adanal setae ad_1 and ad_2 minute; median pore present; interlamellar setae shorter than bothridial setae; body size: $498\text{--}531 \times 381\text{--}398$ ***P. mabunkai* Ermilov, Shtanchaeva, Kalúz & Subías, 2013** (see Ermilov et al. 2013a). Distribution: India
18 Four pairs of notogastral porose areas 19
- Three pairs of notogastral porose areas 22
- 19 Interlamellar setae represented by alveoli; notogastral porose areas Aa located anteriorly to setal alveoli la ; body length: 730 ***P. corolevuensis* Hammer, 1973** (see Hammer 1973). Distribution: Fiji and India
— Interlamellar setae of medium size or long; notogastral porose areas Aa located between setal alveoli la and lm 20
- 20 Notogastral porose areas $A1$ located antero-medially to $A2$; interlamellar setae of medium size; body size: $745\text{--}842 \times 567\text{--}640$
..... ***P. hauseri* Mahunka, 1995** (see Mahunka 1995). Distribution: Borneo
— Notogastral porose areas $A1$ located anteriorly to $A2$; interlamellar setae long 21
- 21 Adanal setae ad_1 and ad_2 similar in length; median pore absent; body size: $510\text{--}630 \times 410\text{--}481$
..... ***P. pterinervis* (Canestrini, 1898)** (see Canestrini 1898; Berlese 1905, 1914; Mahunka 1992; including our data). Distribution: Oriental region
— Adanal setae ad_1 longer than ad_2 ; median pore present; body size: $550\text{--}608 \times 413\text{--}454$
..... ***P. pertrichosa* Mahunka, 1995** (see Mahunka 1995). Distribution: Borneo
22 Notogastral porose areas Aa located nearer to setal alveoli la than lm ; bothridial setae clavate 23
- Notogastral porose areas Aa located nearer to setal alveoli lm than la or distanced equal from them; bothridial setae setiform or with slightly dilated, elongate head 24
- 23 Interlamellar setae minute, shorter than diameter of bothridia; body surface not foveolate; median pore represented by several foveae; body size: $262\text{--}282 \times 192\text{--}209$ ***P. pseudosejugalis* Ermilov & Anichkin, 2012** (see Ermilov and Anichkin 2012). Distribution: Vietnam
— Interlamellar setae short, but longer than diameter of bothridia; body surface foveolate; median pore absent; body size: $246\text{--}275 \times 186\text{--}212$
... ***P. crassipora* Mahunka, 1995** (see Mahunka 1995). Distribution: Borneo
24 Notogastral porose areas Aa located nearer to setal alveoli lm than la 25
- Notogastral porose areas Aa distanced equal from la and lm 30
- 25 Notogastral porose areas minute, smaller than diameter of bothridia; body size: $527\text{--}612 \times 390\text{--}428$ ***P. imadatei* Aoki & Hu, 1993** (see Aoki and Hu 1993). Distribution: Oriental region
— Notogastral porose areas well developed, larger than diameter of bothridia ... 26

| | | | |
|----|--|--|----|
| 26 | Body surface slightly striate; median pore represented by several foveae; body size: 610–715 × 475–545..... | <i>Pergalumna hawaiiensis hawaiiensis</i> (Jacot, 1934) (see Jacot 1934a; including our data) | 99 |
| – | Body surface not striate; median pore single or absent | 27 | |
| 27 | Interlamellar setae minute, shorter than diameter of bothridia; body length: 720..... | <i>P. bimaculata</i> | |
| – | Hammer, 1973 (see Hammer 1973). Distribution: Polynesia and Philippines | | |
| – | Interlamellar setae of medium size, longer than diameter of bothridia..... | 28 | |
| 28 | Median pore present, large; body length: 720 | <i>P. remota</i> | |
| – | (Hammer, 1968) (see Hammer 1968). Distribution: New Zealand and India | | |
| – | Median pore absent | 29 | |
| 29 | Bothridial setae densely ciliate; body size: 451–490 × 300–334 | <i>P. kotschyti</i> Mahunka, 1989 (see Mahunka 1989). Distribution: Vietnam | |
| – | Bothridial setae smooth; body size: 398–453 × 275–340 | 30 | |
| – | <i>P. indivisa</i> Mahunka, 1995 (see Mahunka 1995). Distribution: Borneo | | |
| 30 | Bothridial setae with slightly dilated, elongated head | 31 | |
| – | Bothridial setae setiform | 33 | |
| 31 | Body surface heavily tuberculate; body size: 385–425 × 285–331 | <i>P. granulata</i> Balogh & Mahunka, 1967 (see Balogh and Mahunka 1967). Distribution: Vietnam and Japan | |
| – | Body surface not tuberculate..... | 32 | |
| 32 | Body surface heavily granulate; body size: 302–356 × 237–262 | <i>P. punctulata</i> Balogh & Mahunka, 1967 (see Balogh and Mahunka 1967). Distribution: Vietnam | |
| – | Body surface smooth; body size: 437–465 × 310–324 | 34 | |
| 33 | <i>P. intermedia intermedia</i> Aoki, 1963 (see Aoki 1963, 1966). Distribution: Palaearctic and Oriental regions | | |
| – | Notogastral porose areas <i>A1</i> elongated, longitudinally oriented..... | 34 | |
| – | Notogastral porose areas <i>A1</i> rounded to oval | 36 | |
| 34 | Adanal setae <i>ad</i> ₁ long, not shorter than width of anal plate; median pore single; body size: 415–481 × 298–365 | <i>P. paraindistincta</i> sp. n. Distribution: Indonesia | |
| – | Adanal setae <i>ad</i> ₁ minute; median pore absent or represented by several foveae.. | 35 | |
| 35 | Postanal porose area absent; median more present in females; body size: 547–614 × 381–415 | <i>P. indistincta</i> Ermilov & Anichkin, 2011 (see Ermilov and Anichkin 2011b). Distribution: Vietnam | |
| – | Postanal porose area present; median more absent in females; body size: 576 × 426 | <i>P. magnipora capensis</i> Engelbrecht, 1972 (see Engelbrecht 1972). Distribution: Southern Africa and India | |
| 36 | Body surface striate and short ridges; posterior part of notogaster with longitudinal concavity; body size: 408–485 × 298–352 | <i>P. menglunensis</i> Aoki & Hu, 1993 (see Aoki and Hu 1993). Distribution: southern China | |

- Body not striate and without short ridges; posterior part of notogaster without concavity 37
- 37 Adanal setae *ad*₁ and *ad*₂ comparatively long, not shorter than width of anal plate; setae *c* developed on pteromorphs; body size: 514–597 × 365–431 *P. minituberculata* Ermilov & Martens, 2014 (see Ermilov and Martens 2014). Distribution: Nepal
- Adanal setae *ad*₁ and *ad*₂ shorter than width of anal plate; setae *c* represented by alveoli on pteromorphs 38
- 38 Interlamellar setae represented by alveoli; median pore present; body size: 863–1145 × 639–970 *P. panayensis* Ermilov & Corpuz-Raros, 2015 (see Ermilov and Corpuz-Raros 2015)
- Interlamellar setae of medium size or long; median pore absent 39
- 39 Bothridial setae densely ciliate 40
- Bothridial setae densely smooth 41
- 40 Notogastral porose areas amorphous, without distinct borders; genital plates not striate; body size: 332–377 × 245–276
... *P. amorphia* Mahunka, 2008 (see Mahunka 2008). Distribution: Thailand
- Notogastral porose areas with distinct borders; genital plates striate; body size: 390–435 × 282–315 *P. intermedia retroversa* Aoki & Hu, 1993 (see Aoki and Hu 1993). Distribution: southern China
- 41 Interlamellar setae comparatively short, about 1/3 as long as their mutual distance; genital plates smooth; body size: 742–845 × 589–653 *P. magnipora capillaris* Aoki, 1961 (see Aoki 1961). Distribution: Palaearctic and Oriental regions
- Interlamellar setae of medium size, about 1/2 as long as their mutual distance; genital plates striate; body size: 822–840 × 618–650 *P. magnipora xishuangbanna* Aoki & Hu, 1993 (see Aoki and Hu 1993). Distribution: southern China

Acknowledgements

We cordially thank Prof. Dr. Badamdorj Bayartogtokh (National University of Mongolia, Ulaanbaatar, Mongolia) and Prof. Dr. Gerd Weigmann (Free University of Berlin, Institute of Zoology, Berlin, Germany) for their valuable comments, Kristina Richter (Georg August University Göttingen, Göttingen, Germany) for assistance in building up the Indonesian oribatid mite morphospecies collection, the State Ministry of Research and Technology of Indonesia (RISTEK) for the research permit and the Indonesian Institute of Sciences (LIPI) and Ministry of Forestry (PHKA) for the collection permit, the village heads, local site owners, PT REKI and Bukit Duabelas National Park for granting access to their properties and the many colleagues and helpers for support in the field.

Financial support was provided by the German Research Foundation (DFG) in the framework of the collaborative German – Indonesian research project CRC990 (EFForTS). The taxonomic study on Galumnoidea was supported by the Russian Foundation for Basic Research (project: 15-04-02706 A).

References

- Aoki J (1961) On six new oribatid mites from Japan. *Sanitary Zool* 12(4): 233–238.
- Aoki J (1963) Einige neue oribatiden aus dem kaiserlichen palastgarten Japans. *Ann Zool Japon* 36(4): 218–224.
- Aoki J (1966) The large-winged mites of Japan (Acari: Cryptostigmata). *Bul Nat Mus Nat Sci, Tokyo* 9(3): 257–275.
- Aoki J (1975) Oribatid mites from Korea (Acari). II. *Acta Zool Acad Sci Hung* 21(3–4): 233–239.
- Aoki J, Hu S (1993) Oribatid mites from tropical forests of Yunnan Province in China. II. Families Galumnidae and Galumnellidae. *Zool Sci* 10(5): 835–848.
- Balogh J, Balogh P (1990) Oribatid mites of the Neotropical region. II. Budapest, Akadémiai Kiadó press, 333 pp.
- Balogh J, Balogh P (2002) Identification keys to the oribatid mites of the Extra-Holarctic regions. Vol. 1. Miskolc, Well-Press Publ. Limited, 453 pp.
- Balogh J, Mahunka S (1967) New oribatids (Acari) from Vietnam. *Acta Zool Acad Sci Hung* 13(1–2): 39–74.
- Balogh P (1988) Oribatid mites (Acari) from Sri Lanka. *Acta Zool Hung* 34(2–3): 171–189.
- Balogh P (1997) New species of oribatids (Acari) from the Neotropical region. *Opusc Zool Budapest* 29–30: 21–30.
- Bayartogtokh B, Chatterjee T (2010) Oribatid mites from marine littoral and freshwater habitats in India with remarks on world species of *Thalassozetes* (Acari: Oribatida). *Zool Stud* 49(6): 839–854.
- Berlese A (1905) Acari nuovi. *Manipulus IV* (Acari di Giava). *Redia* 2: 154–176.
- Berlese A (1914) Acari nuovi. *Manipulus IX*. *Redia* 10: 113–150.
- Canestrini G (1898) Nuovi Acaroidei della N. Guinea (Seconda serie). *Természetrajzi Füzetek* 21: 193–197.
- Engelbrecht CM (1972) Galumnids from South Africa (Galumnidae, Oribatei). *Acarologia* 14(1): 109–140.
- Ermilov SG, Alvarado-Rodríguez O, Retana-Salazar AP (2014a) Contribution to the knowledge of Costa Rican oribatid mite fauna, with supplementary descriptions of *Pergalumna silvatica* and *P. sura* (Acari: Oribatida Galumnidae). *Syst Appl Acarol* 19(2): 216–222. doi: 10.11118/saa.19.2.12
- Ermilov SG, Alvarado-Rodríguez O, Retana-Salazar AP (2014b) Two new species of *Pergalumna* (Acari, Oribatida, Galumnidae) from Costa Rica, including a key to all species of the genus from the Neotropical region. *ZooKeys* 435: 7–23. doi: 10.3897/zookeys.435.8213
- Ermilov SG, Anichkin AE (2011a) New oribatid mites of the genera *Pergalumna* and *Galumnella* (Acari, Oribatida, Galumnoidea) from Vietnam. *Acarina* 19(2): 242–251.
- Ermilov SG, Anichkin AE (2011b) The galumnoid fauna (Acari: Oribatida) of Cat Tien National Park (southern Vietnam) with description of two new species. *Int J Acarol* 37 (Suppl 1): 85–94. doi: 10.1080/01647954.2010.539982
- Ermilov SG, Anichkin AE (2012) Two new species of oribatid mites (Acari: Oribatida) from Bu Gia Map National Park (Vietnam). *Zoosyst Ros* 21(1): 18–27.

- Ermilov SG, Anichkin AE, Wu D (2012a) Oribatid mites from Bu Gia Map National Park (Southern Vietnam), with description of a new species of *Dolicheremaeus* (Tetracondylidae) (Acari: Oribatida). *Genus* 23(4): 591–601.
- Ermilov SG, Bayartogtokh B (2015) Systematic placement of some taxa in the family Galumnidae (Acari, Oribatida). *Zootaxa* 3964(4): 489–493. doi: 10.11646/zootaxa.3964.4.8
- Ermilov SG, Chatterjee T, Das MK, Bordoloi S (2014c) Three new species of oribatid mites of the genus *Pergalumna* (Acari: Oribatida: Galumnidae) from India. *Biologia* 69(4): 489–497. doi: 10.2478/s11756-014-0332-5
- Ermilov SG, Corpuz-Raros LA (2015) New species of galumnid oribatid mites of the genera *Mirogalumna* and *Pergalumna* from the Philippines (Acari, Oribatida, Galumnidae). *Syst Appl Acarol* 20(5): 556–566. doi: 10.11158/saa.20.5.9
- Ermilov SG, Martens J (2014) Two new species of oribatid mites of the genera *Pergalumna* and *Carinogalumna* (Acari, Oribatida, Galumnidae) from Nepal. *Syst Appl Acarol* 19(4): 462–470. doi: 10.11158/saa.19.4.9
- Ermilov SG, Niedbała W, Anichkin AE (2012b) Oribatid mites of Dong Nai Biosphere Reserve (=Cat Tien National Park) of Southern Vietnam, with description of a new species of *Pergalumna* (Acari, Oribatida, Galumnidae). *Acarina* 20(1): 20–28.
- Ermilov SG, Shtanchaeva UYa, Kalúz S, Subías LS (2013a) Three new species of the genus *Pergalumna* (Acari: Oribatida: Galumnidae) from India. *Zootaxa* 3682(3): 412–420. doi: 10.11646/zootaxa.3682.3.2
- Ermilov SG, Shtanchaeva UYa, Subías LS (2014d) New species of oribatid mites (Acari: Oribatida) of the genera *Suctobelbella* (Suctobelbidae) and *Neoribates* (Parakalummidae) from Vietnam. *Biologia* 69(11): 1593–1600. doi: 10.2478/s11756-014-0457-6
- Ermilov SG, Starý J, Sandmann D, Marian F, Maraun M (2013b) New taxa and new records of oribatid mites of the family Galumnidae (Acari: Oribatida) from Ecuador. *Zootaxa* 3700(2): 259–270. doi: 10.11646/zootaxa.3700.2.4
- Ermilov SG, Weigmann G, Tolstikov AV (2013c) Morphology of adult and juvenile instars of *Galumna obvia* (Acari, Oribatida, Galumnidae), with discussion of its taxonomic status. *ZooKeys* 357: 11–28. doi: 10.3897/zookeys.357.6404
- Ewing HE (1907) New Oribatidae. *Phyche* 14: 111–115. doi: 10.1155/1907/296057
- Grandjean F (1936) Les Oribates de Jean Frédéric Hermann et de son pere. *Ann Soc Ent France* 105: 27–110.
- Hammer M (1958) Investigations on the oribatid fauna of the Andes Mountains. I. The Argentine and Bolivia. *Det Kong Dansk Vidensk Selsk Biol Skr* 10(1): 1–129.
- Hammer M (1961) Investigations on the oribatid fauna of the Andes Mountains. II. Peru. *Det Kong Dansk Vidensk Selsk Biol Skr* 13(1): 1–157.
- Hammer M (1968) Investigations on the Oribatid fauna of New Zealand. Part III. *Det Kong Dansk Vidensk Selsk Biol Skr* 16(2): 1–96.
- Hammer M (1972) Investigations on the oribatid fauna of Tahiti, and some oribatids found on the Atoll Rangiroa. *Det Kong Dansk Vidensk Selsk Biol Skr* 19(3): 1–66.
- Hammer M (1973) Oribatids from Tongatapu and Eua, the Tonga Islands, and from Upolu, Western Samoa. *Det Kong Dansk Vidensk Selsk Biol Skr* 20(3): 1–70.

- Jacot AP (1934a) Some Hawaiian Oribatoidea (Acarina). Bernice P Bishop Mus Bul, Honolulu 121: 1–99.
- Jacot AP (1934b) The Galumnas (Oribatoidea—Acarina) of the Northeastern United States. J New York Ent Soc 42(1): 87–125.
- Mahunka S (1989) A survey of the Oribatid fauna (Acari) of Vietnam, III. Folia Ent Hung 50: 47–59.
- Mahunka S (1992) “Pelops” and “Oribates” species in the Berlese-collection (Acari). Acta Zool Hung 38(3–4): 213–260.
- Mahunka S (1994) Two new Galumnid species (Acari: Oribatida) from Thailand. Acta Zool Acad Sci Hung 40(4): 351–357.
- Mahunka S (1995) Oribatids from Sabah, East Malaysia (Acari: Oribatida, Parakalummoidea, n. stat. and Galumnoidea). Trop Zool 8: 269–308. doi: 10.1080/03946975.1995.10539286
- Mahunka S (2008) A new genus and some other data of oribatids from Thailand (Acari: Oribatida). Acta Zool Acad Sci Hung 54(2): 125–150.
- Norton RA, Behan-Pelletier VM (2009) Chapter 15. Oribatida. In: Krantz GW, Walter DE (Eds) A Manual of Acarology. Texas Tech Univ Press, Lubbock, 430–564.
- Raju VV, Appalainaidu V, Rao KH (1981) *Pergalumna andhraense* sp. n. (Galumnidae: Oribatei) from Visakhapatnam, Andhra Pradesh. Geobios 8: 219–221.
- Sellnick M (1925) Fauna sumatrensis. Suppl Ent 11: 79–89.
- Shaldybina ES (1975) The family Galumnidae. In: Ghilyarov MS (Ed.) Key to Soil Inhabiting Mites. Sarcoptiformes. Nauka Press, Moscow, 347–363.
- Subías LS (2004) Listado sistemático, sinonímico y biogeográfico de los ácaros oribátidos (Acariformes: Oribatida) del mundo (excepto fósiles). Graellsia 60 (número extraordinario): 3–305. Online version accessed in March 2015, 587 pp.
- Tseng Y (1984) Taxonomical study of oribatid mites from Taiwan (Acarina: Astigmata) (II). Chin J Ent 4: 27–74.
- Weigmann G (2006) Hornmilben (Oribatida). Die Tierwelt Deutschlands. Teil 76. Goecke and Evers, Keltern, 520 pp.
- Willmann C (1931) Oribatei (Acari), gesammelt von der deutschen limnologischen sundae-expedition. Arch Hydrobiol, Suppl 9 “Trop Binneng 2: 240–305.
- Willmann C (1956) Milben aus naturschutzgebient auf dem Spiegitzer (Glatzer) Schneeberg. Českoslov parasitol 3: 211–273.