

An annotated checklist of the eukaryotic parasites of humans, exclusive of fungi and algae

Blaine A. Mathison¹, Sarah G. H. Sapp²

1 *Institute for Clinical and Experimental Pathology, ARUP Laboratories, Salt Lake City, UT, USA* **2** *Parasitic Diseases Branch, Division of Parasitic Diseases and Malaria, Centers for Disease Control and Prevention, Atlanta, GA, USA*

Corresponding author: Blaine A. Mathison (blaine.mathison@aruplab.com)

Academic editor: Matthew Wayland | Received 15 April 2021 | Accepted 20 August 2021 | Published 9 November 2021

<http://zoobank.org/1561AF88-B389-4F1E-9F39-3890381271EE>

Citation: Mathison BA, Sapp SGH (2021) An annotated checklist of the eukaryotic parasites of humans, exclusive of fungi and algae. ZooKeys 1069: 1–313. <https://doi.org/10.3897/zookeys.1069.67403>

Abstract

The classification of “parasites” in the medical field is a challenging notion, a group which historically has included all eukaryotes exclusive of fungi that invade and derive resources from the human host. Since antiquity, humans have been identifying and documenting parasitic infections, and this collective catalog of parasitic agents has expanded considerably with technology. As our understanding of species boundaries and the use of molecular tools has evolved, so has our concept of the taxonomy of human parasites. Consequently, new species have been recognized while others have been relegated to synonyms. On the other hand, the decline of expertise in classical parasitology and limited curricula have led to a loss of awareness of many rarely encountered species. Here, we provide a comprehensive checklist of all reported eukaryotic organisms (excluding fungi and allied taxa) parasitizing humans resulting in 274 genus-group taxa and 848 species-group taxa. For each species, or genus where indicated, a concise summary of geographic distribution, natural hosts, route of transmission and site within human host, and vectored pathogens are presented. Ubiquitous, human-adapted species as well as very rare, incidental zoonotic organisms are discussed in this annotated checklist. We also provide a list of 79 excluded genera and species that have been previously reported as human parasites but are not believed to be true human parasites or represent misidentifications or taxonomic changes.

Keywords

Acanthocephalans, arthropods, cestodes, leeches, nematodes, parasitology, protozoa, trematodes

Table of contents

Introduction.....	2
Materials and methods	6
Checklist of eukaryotic human parasites.....	8
Amoebozoa Lühe, 1913, amend Cavalier-Smith, 1998.....	8
Tubulinea Smirnov et al., 2005	8
Echinamoebida Cavalier-Smith in Cavalier-Smith et al. 2004	8
Elardia Kang et al., 2017.....	9
Evosea Kang et al., 2017	9
Archamoeba Cavalier-Smith, 1993 sensu Cavalier-Smith et al., 2004.....	9
Discosea Cavalier-Smith et al., 2004	13
Flabellinia Smirnov et al., 2005.....	13
Centramoebia Cavalier-Smith et al., 2016.....	13
SAR (Stremanopiles-Alveolata-Rhizaria) Burki et al., 2008, emend. Adl et al., 2012...16	
Stramenopiles Patterson 1989, emend. Adl et al., 2005.....	16
Bigyra Cavalier-Smith, 1998, emend. Cavalier-Smith et al., 2006.....	16
Alveolata Cavalier-Smith, 1991	18
Colpodellida Cavalier-Smith, 1993, amend. Adl et al. 2005, 2019	18
Apicomplexa Levine, 1970, emend. Adl et al. 2005.....	19
Ciliophora Doflein, 1901.....	32
Excavates Cavalier-Smith, 2002, emend. Simpson, 2003.....	33
Metamonada Grassé, 1952.....	33
Fornicata Simpson, 2003	33
Parabasalia Honigberg, 1973.....	34
Discoba Simpson in Hampl et al., 2009.....	36
Heterolobosea Page & Blanton, 1985	36
Euglenozoa Cavalier-Smith, 1981, emend. Simpson, 1997	38
Opisthokonta Cavalier-Smith, 1987, emend. Adl et al., 2005	47
Holozoa Lang et al., 2002	47
Metazoa Haeckel, 1874, emend. Adl et al. 2005	47
Excluded species.....	246
Acknowledgements.....	254
References	255

Introduction

The online Oxford English dictionary defines a parasite as: ‘an organism that lives in or on an organism of another species (its host) and benefits by deriving nutrients at the other’s expense’ (LEXICO 2020). From a strictly biological perspective, this definition can cover a wide variety of organisms, including, but not limited to, bacteria, viruses, fungi, helminths (worms), protozoans, cnidarians, tardigrades, rotifers, mollusks, and arthropods. However, in the historical concept of the medical community, the term

‘parasite’ refers to animals and other eukaryotic organisms with animal-like affinities, and is usually limited to the protozoans, helminths, and arthropods.

Historically, human parasites have been classified within the broad Linnaean kingdom of ‘Animals’ and divided into taxa with formal categorical names (phylum, class, order, family, etc.). While traditional Linnaean taxa are still commonly used for helminths and arthropods, it is becoming more common to use clade-based systems devoid of formal designations, especially with the taxa collectively called protozoans. Much of this work comes from a better understanding of the evolutionary relationships of organisms based on both morphological and molecular analyses (Adl and Mathison 2019).

Currently, eukaryotes are classified into two domains, Amorphea and Diaphoretickes, as well as additional large clades that do not fit into either of those domains. Within domains are large clades commonly referred to as supergroups. Eukaryotic parasitic organisms fall into five supergroups within these domains: SAR, Archaeplastida, Excavates, Amoebozoa, and Opisthokonta (Adl et al. 2019; Adl and Mathison 2019). For the purpose of this work, the focus is on the traditional human parasitic protozoans (Amoebozoa, SAR, Excavates) and helminths, annelids, and arthropods (all in Opisthokonta). Two groups of organisms that were historically considered protozoans, but are now considered fungi and are not included here, are *Pneumocystis jirovecii* Frenkel, 1999 (syn. *P. carinii* f. sp. *hominis*) and the microsporidia. The genus *Prototheca* is another opportunistically parasitic eukaryote, but is currently aligned with the algae in the clade Archaeplastida and is also not covered here (Lass-Flörl and Mayr 2007). For detailed descriptions on the higher classifications of eukaryotes, the readers are referred to the work by Adl and colleagues (Adl et al. 2019).

The Amoebozoa include many of the human parasites historically referred to as amebae, including the intestinal amebae *Entamoeba*, *Endolimax*, and *Iodamoeba*, and several of the ‘free-living’ amebae such as *Balamuthia* and *Acanthamoeba* (Adl and Mathison 2019). These organisms feed by phagocytosis and move by pseudopods. They lack cilia and most have an environmentally hardy stage (cyst) for dispersal and infection of a new host (Adl et al. 2019; Adl and Mathison 2019). The free-living amebae are not adapted for true parasitism and are only opportunistic parasites when conditions allow for colonization of the human host. Clinically, *Acanthamoeba* species are usually characterized by their genotypes rather than traditional Linnaean binominal nomenclature (Booton et al. 2005).

The SAR (which commonly refers to Stramenopiles-Alveolates-Rhizaria) includes parasitic coccidians (e.g., *Toxoplasma*, *Cyclospora*, *Cystoisospora*), gregarines (*Cryptosporidium*), piroplasmids (*Babesia*), haemosporidians (*Plasmodium*), ciliophores (*Balantioides*, formerly *Balantidium*), and the stremenopile *Blastocystis* (Adl and Mathison 2019).

Within the SAR, the coccidians, gregarines, piroplasmids, and hemsporidians are obligate parasites and part of the clade Apicomplexa that are characterized by a structure called an apical complex that is used for penetration of the host cell. Motility is usually by gliding or body flexion and feeding is by pinocytosis (ingestion by budding of small vesicles from the cell membrane). Apicomplexans may have one or two hosts and reproduce by sexual reproduction, the end result of which is an oocyst that when mature contains sporozoites that are the infectious stage for a new host (Adl et al. 2019;

Adl and Mathison 2019). The Haemospororida (e.g., *Plasmodium*) and Piroplasmorida (e.g., *Babesia*) are vectored by sanguivorous arthropods. Humans are intermediate hosts for *Plasmodium*, which are vectored by mosquitoes that serve as the definitive hosts. *Babesia* are zoonotic in humans and in nature typically cycle between a tick vector (definitive host) and a mammalian intermediate host (Adl and Mathison 2019).

The only human parasitic ‘ciliate’ is *Balantioides* (formerly *Balantidium*, syn. *Neobalantidium*). Movement is by numerous cilia and feeding is by phagocytosis. It forms an environmentally-hardy cyst that is the infectious stage for a new host (Adl et al. 2019; Adl and Mathison 2019).

Blastocystis is an enigmatic organism that has historically been variably classified among coccidians, fungi, and algae. The life cycle is still not completely understood, but it has been proposed that it reproduces by binary fission of vacuolar forms within the host and infects a new host via environmentally hardy cysts. Both vacuolar forms and cysts can be detected in stool specimens, and the variability in morphology of the former has created numerous descriptive terms. It has also been suggested it has a third form, an amoeboid form, that is responsible for pathogenesis in the host, but this hypothesis is not yet broadly accepted (Tan 2008). As the molecular epidemiology is becoming better understood, it is becoming less common to recognize *Blastocystis* by traditional Linnaean binominal nomenclature and more common to base identification on their genetic subtypes (Stensvold et al. 2007; Tan 2008).

The Excavates includes the traditional ‘flagellates’, such as *Giardia*, *Trypanosoma*, *Dientamoeba*, *Chilomastix*, *Leishmania*, and *Trichomonas*, as well as amoeboid flagellates, such as *Naegleria* (Adl and Mathison 2019). All excavates move by cilia (historically referred to as flagella, which are not analogous to the bacterial flagellum) at some point in their life cycles. Most feed by pinocytosis (although the amoeboid form of *Naegleria* and its relatives feed by phagocytosis). The life cycle patterns vary greatly among the excavates. Intestinal flagellates typically multiply by binary fission in the lumen of the intestine of their hosts and have an environmentally hardy cyst stage that is infectious for a new host. The trichomonads lack a cyst stage, and *Dientamoeba* is unusual in that it lacks external cilia and moves by pseudopods and feeds by phagocytosis, much like the Amoebozoa. The ‘hematoflagellates’ (*Leishmania*, *Trypanosoma*, *Crithidia*, *Endotrypanum*) have two host life cycles and are transmitted to the human host via an arthropod vector. Amoeboflagellates, such as *Naegleria*, *Paravahlkampfia*, *Tetramitus*, *Vahlkampfia*, *Allovahlkampfia* are not true parasites and only infect the human host when opportunities allow for infection and colonization (Adl et al. 2019; Adl and Mathison 2019).

The Opisthokonta includes the arthropods, leeches, and helminths, all within the clade Metazoa (Adl and Mathison 2019). The term ‘helminth’ is not a taxonomic unit in any classification scheme, but instead a basket term for parasitic worms of humans, which is divided into four broad categories: cestodes (tapeworms), trematodes (flukes), nematodes (roundworms), and acanthocephalans (thorny-headed worms) (Adl and Mathison 2019). Leeches (Annelida), while not historically considered helminths are blood-feeding animals that will opportunistically feed on humans. Within the Metazoa are the Spiralia (acanthocephalans, annelids, cestodes, trematodes) and Ecdysozoa (arthropods and nematodes).

The cestodes are obligate parasites that usually reside in the small intestine of their definitive host. Most have multiple-host life cycles, and the definitive host becomes infected after ingestion of a larval stage encysted in the tissues of an intermediate host. Adults are long, ribbon-like organisms that have three main body regions: scolex (for attachment to the intestinal mucosa of the definitive host), neck (the base of the strobila), and the strobila (the main body, which is made of up individual segments called proglottids). Historically, cestodes that parasitize humans were divided into two categories: Pseudophyllidea (*Dibothriocephalus*, *Adenocephalus*, *Diphyllobothrium*), and Cyclophyllidea (*Taenia*, *Dipylidium*, *Hymenolepis*, several others). However, Pseudophyllidea has been abandoned in favor of two broad groups, Bothriocephalidea and Diphyllobothriidea, the latter of which contains human parasites (Kuchta et al. 2008). From the perspective of species that parasitize humans, Diphyllobothriidea are characterized by adult worms that attach to the intestine of the definitive host by means of bothria (grooves in the scolex). Eggs have an operculum and are unembryonated when shed in feces, and the infectious stage for the definitive host is the plerocercoid larva (e.g., sparganum). The Cyclophyllidea however attach to the intestine of their definitive hosts by means of four suckers and (often) a rostellum, that may be armed with hooklets or not. Eggs lack an operculum, are shed embryonated, and contain a hooked oncosphere that is immediately infectious for an intermediate host. The infectious larva is highly variable and terminology is often taxon-dependent (cysticercus, hydatid cyst, tetrahyridium, etc.) (Chervy 2002; Adl and Mathison 2019).

The trematodes that infect humans are all within Digenea and are also obligate parasites with multi-host life cycles. All parasitic species of humans have a snail as a first intermediate host, and many digeneans have second intermediate hosts or paratenic hosts that may comprise a variety of animals; in some species, infective stages are encysted in the environment, including on the surface of plants. Adults are usually hermaphroditic, apart from the schistosomes which are dioecious but live *in copula*. Adults have two muscular suckers, an oral sucker for attachment and feeding and a ventral sucker (acetabulum) for attachment. Infection of the human host is either by direct penetration of the cercaria larval stage (*Schistosoma*) or ingestion of the metacercaria larval stage in an intermediate animal host or on contaminated plants (Adl and Mathison 2019).

The acanthocephalans are zoonotic parasites in humans. While superficially similar to nematodes, they are more closely related to the rotifers, or possibly nested within Rotifera, proper (Garcia-Varela et al. 2000). Adults are large, pseudocoelomate animals that attach to the intestinal mucosa of the definitive host by means of a heavily armed proboscis. Human infection usually occurs from the ingestion of an infected intermediate or paratenic host (Mathison et al. 2016). The leeches (Annelida: Hirudinea) include blood-feeding members that will opportunistically feed on humans. Most attacks on humans usually come from swimming, fishing, and wading in fresh water.

The nematodes are a large and diverse group of organisms, most of which are free-living, although some major parasitic lineages are of great public health importance. Despite their diversity in nature, few species parasitize humans with any regularity. Two major clades of human parasites are Dorylaimia and Chromadoria, which vary greatly

in their biology, route of transmission, and morphology. Most nematodes have six developmental stages: egg, four larval stages (L1–L4), and adult (L5); some larviparous species do not lay eggs. For most parasitic nematodes of humans, the infectious stage is the L3 larva, this fact is often called the ‘Rule of the Infective Third State’. Notable exceptions are members of the Trichinellida, for which the infectious stage is the L1 larva, and *Eustrongylides* for which the infectious stage is believed to be the precocious L4 (Anderson 2000). Most intestinal nematodes infect the human host after ingestion of fully embryonated eggs in food, water, or fomites contaminated with feces, or after ingestion of an infected intermediate or paratenic host. L3 larvae of some species, such as *Strongyloides* and the hookworms, can penetrate intact human skin during walking on contaminated soil. Filarial nematodes (Spirurida: Filarioidea) are vectored by sanguivorous arthropods (Ash and Orihel 2007; Adl and Mathison 2019; Carroll et al. 2019).

The arthropods are the largest group of Metazoa, but relatively few species are adapted to parasitism on humans. Two main clades are the Chelicerata and Mandibulata, which can be broadly separated by the structure of their mouthparts. The Chelicerata includes the mites and ticks. The Mandibulata includes Pancrustacea, which contains the clades Crustacea (crustaceans) and Hexapoda (insects). The only parasitic crustaceans of humans are zoonotic pentastomids (tongueworms). The pancrustacean clade Hexapoda includes parasitic insects such as lice, fleas, bed bugs, triatomine (kissing) bugs, and myiasis-causing flies. Arthropods are characterized by a jointed exoskeleton containing chitin. Development is variable and can be gradual (hemimetabolous) or complete (holometabolous). Holometabolous insects have morphologically and biologically markedly different life cycle stages including larvae, pupae, and adults (Mathison and Pritt 2014; Mathison and Pritt 2015; Adl and Mathison 2019).

Here we provide an updated checklist of the parasitic protozoans, helminths, annelids, and arthropods of humans, nearly 20 years since the last such endeavor (Ashford and Crewe 2003), while providing an updated higher taxonomy based on advances in eukaryotic taxonomy (Adl et al. 2019). Taxonomy and systematics are largely ignored in the education of medical and public health professionals. There are frequent updates to taxonomic changes in Medical Parasitology (Simner 2017; Adl and Mathison 2019; Mathison and Pritt 2019; Mathison and Pritt 2020; Mathison et al. 2021), but taxonomy remains a challenging discipline in the medical and public health realms. We also provide annotations on the geographic distribution, host range, route of infection, and anatomic site of infection for human eukaryotic parasites.

Materials and methods

The bulk of this manuscript is an annotated checklist of protozoan, helminthic, annelid, and arthropod parasites reported from the human host, based on extensive literature searches through April 2021. A systematic literature search of the PubMed database (U.S. National Library of Medicine National Institutes of Health; <https://www.ncbi.nlm.nih.gov/pubmed>), Google Scholar (<https://scholar.google.com/>) and

Google (<https://www.google.com>) was also performed using the keyword search phrases using various taxa in combination with human infection and case reports. We aimed to include all protozoan and helminthic parasites known to be reported from humans. The leeches and arthropods, however, provided a special challenge, as an exhaustive list of every sanguivorous species that may feed on a human host would be large and beyond the scope of the audience. For the arthropods, the focus is on ectoparasites that spend prolonged time on the human host (lice, sarcoptid and demodecid mites, hard ticks, myiasis-causing flies, *Tunga* fleas) and visceral parasites (pentastomids). For leeches and arthropods that are short-term blood feeders or biters, such as non-*Tunga* fleas, soft ticks, zoonotic biting mites, bed bugs, and triatomine bugs, the focus is on those species that are commonly associated with humans and/or vector medically-important pathogens. Sanguivorous flies (mosquitoes, black flies, deer flies, etc.) are not included, nor are those flies and other insects (e.g., cockroaches) that may passively transmit pathogens. Arthropods that are considered medically important due to envenomation, stings, or urticarial and allergic reactions are also not included.

Suprageneric taxa are listed in a hierarchal system lacking formal designations, following Adl et al. (Adl et al. 2019). Each suprageneric taxon is indicated by a series of icons (●) rather than using traditional Linnaean categories such as order, family, etc. Each taxon has one more icon than the preceding taxon in which it belongs. The number of clades presented varies based on the diversity within a given clade in relation to human parasites.

For organisms traditionally classified as protozoans (Amoebozoa, Excavates, and SAR), clades above the traditional 'family' level follow Adl and colleagues (Adl et al. 2019). The family-group taxa (typically ending in -idae) are presented based on the current classifications of the given organisms. For the helminths and arthropods, the organisms considered 'true' animals (Opisthokonta), the classification follows Adl and colleagues (Adl et al. 2019) to the level Holozoa, after which clades follow the current classifications for the given organisms. Genera within a family-group taxon, and species within each genus, are presented alphabetically.

Species, and genera for which isolates from human cases have not been characterized at the species level, have at least four bullet-points of information: 1) Geographic distribution, 2) Natural hosts, 3) Route of infection, and 4) Site in/on human host. In addition, many of the arthropods and leeches also have an additional bullet-point, 5) Vectored pathogens.

Geographic distribution. This is the known geographic distribution of a given species or genus. For widespread organisms, the distribution is presented generally (e.g., Worldwide, North America, Circumtropical). For organisms that are more geographically restricted, the information is presented at the country level.

Natural hosts. These are the hosts that are part of the parasite's natural life cycle. For multi-host parasites, the hosts are presented in developmental order, starting with the first intermediate host and finishing with the definitive host. Hosts for species or genera with broad host ranges are presented generally (e.g., mammals, carnivores,

birds), while hosts for parasites that are more host specific are presented at a more granular level (e.g., mosquitoes in the genus *Anopheles*).

Route of infection. This is route through which the human host becomes infected or colonized.

Site in/on human host. This is where the parasite resides in the human host, and includes both the typical site as well as common sites of ectopic infection.

Vectored pathogens. This bullet-point is for all the arthropods, except for the pentastomids and most of the myiasis-causing fly larvae, and leeches. It provides information on infectious agents (bacteria, viruses, parasites) that are pathogenic for the human host that are transmitted by the given arthropod. It includes agents that are believed to be vectored by a given arthropod in a natural setting and does not include agents that have simply been detected in an arthropod using molecular surveillance or in experimental models.

For all taxonomic levels, select synonyms are presented, especially for names that show up frequently in the literature regarding human cases.

Lastly, a list of excluded species is presented. These are species that have been reported as human parasites but are either not believed to be true parasites in the human host, represent data based on misidentifications, or are invalid names. The excluded species are presented alphabetically.

Checklist of eukaryotic human parasites

Amoebozoa Lühe, 1913, amend Cavalier-Smith, 1998

- *Tubulinea* Smirnov et al., 2005
- *Echinamoebida* Cavalier-Smith in Cavalier-Smith et al. 2004

Genus *Vermamoeba* Smirnov et al., 2011

Vermamoeba vermiformis (Page, 1967)

Geographic distribution. Worldwide (Scheid 2019).

Natural host. None; occurs in natural freshwater environments, surface water, soil, and biofilms. Humans are incidental hosts (Scheid 2019; Scheid et al. 2019; Siddiqui et al. 2021).

Route of infection. Presumably environmental exposure into eyes, mucus membranes, and wounds (Abedkhozasteh et al. 2013; Scheid 2019; Scheid et al. 2019).

Site in human host. Skin and soft tissues, eye (Scheid et al. 2019).

Notes. Previous records of *Harmannella* from human clinical specimens probably refer to this species.

- Elardia Kang et al., 2017
- Euamoebida Lepši, 1960 sensu Smirnov et al., 2011
- Hartmannellidae Volkonsky, 1931

Genus *Hartmannella* Alexeiff, 1912

Geographic distribution. Worldwide.

Natural host. None (environmental); humans are incidental hosts (Bradbury 2014).

Route of infection. Unknown; presumed environmental exposure and contamination (Bradbury 2014).

Site in human host. Intestinal tract (Bradbury 2014).

Notes. Most extraintestinal reports of human infection with *Hartmannella* apply to *H. vermiformis* Page, which is now in the genus *Vermamoeba*.

- Evosea Kang et al., 2017
- Archamoeba Cavalier-Smith, 1993 sensu Cavalier-Smith et al., 2004
- Entamoebida Cavalier-Smith, 1993
- Entamoebidae Chatton, 1925

Genus *Endolimax* Kuenen & Swellengrebel, 1913

Endolimax nana (Wenyon et O'Connor, 1917)

Geographic distribution. Worldwide (Poulsen and Stensvold 2016).

Natural host. Humans (Poulsen and Stensvold 2016).

Route of infection. Ingestion of mature cysts in fecally contaminated water, food, fomites (Poulsen and Stensvold 2016).

Site in human host. Large intestine, cecum, colon (Poulsen and Stensvold 2016).

Genus *Entamoeba* Casagrandi & Barbagallo, 1895

Entamoeba bangladeshi Royer et al., 2012

Geographic distribution. Southeast Asia, sub-Saharan Africa (Hooshyar et al. 2015; Ngobeni et al. 2017).

Natural host. Humans (Hooshyar et al. 2015).

Route of infection. Ingestion of mature cysts in fecally contaminated water, food, fomites (Ali 2015).

Site in human host. Large intestine, cecum, colon (Ali 2015).

***Entamoeba coli* (Grassi, 1879)**

Entamoeba hominis Casagrandi & Barbagallo, 1897

Entamoeba loeschi Lesage, 1908

Councilmania lafleuri Kofoid & Swezy, 1921

Geographic distribution. Worldwide (Hooshyar et al. 2015).

Natural hosts. Humans, monkeys (Hooshyar et al. 2015).

Route of infection. Ingestion of mature cysts in fecally contaminated water, food, fomites (Ali 2015).

Site in human host. Large intestine, cecum, colon (Ali 2015).

***Entamoeba chattoni* Swellengrebel, 1914**

Geographic distribution. Africa (Verweij et al. 2001; Hooshyar et al. 2015).

Natural hosts. Non-human primates. Zoonotic in humans (Sargeaunt et al. 1992; Hooshyar et al. 2015).

Route of infection. Ingestion of mature cysts in fecally contaminated water, food, fomites (Sargeaunt et al. 1992).

Site in human host. Large intestine, cecum, colon (Sargeaunt et al. 1992).

***Entamoeba dispar* Brumpt, 1925**

Geographic distribution. Worldwide (Ali 2015; Hooshyar et al. 2015).

Natural hosts. Humans, chimpanzee, baboons, macaques (Ali 2015; Hooshyar et al. 2015).

Route of infection. Ingestion of mature cysts in fecally contaminated water, food, fomites (Ali 2015).

Site in human host. Large intestine, cecum, colon (Ali 2015).

***Entamoeba gingivalis* (Gros, 1849)**

Amoeba buccalis Steinberg, 1862

Amoeba dentalis Grassi, 1879

Amoeba kartulis Doflein, 1901

Entamoeba maxillaris Kartulis, 1906

Entamoeba canibuccalis Smith, 1938

Endamoeba confusa Craig, 1916

Entamoeba equibuccalis Simitich, 1938

Entamoeba suigingivalis Tumka, 1959

Geographic distribution. Worldwide (Hooshyar et al. 2015).

Natural hosts. Humans, horses, pigs, cats, monkeys (Hooshyar et al. 2015).

Route of infection. Person-to-person contact (Bonner et al. 2018; Bradbury et al. 2019b).

Site in human host. Oral cavity; ectopic colonization of urogenital tract (Cembranelli et al. 2013; Bonner et al. 2018; Bradbury et al. 2019b; Dubar et al. 2020).

***Entamoeba hartmanni* Von Prowazek, 1912**

Entamoeba histolytica, small race

Entamoeba minuta Woodcock & Penfold, 1916

Entamoeba tenuis Kuenen & Swellengrebel, 1917

Entamoeba minutissima Brug, 1917

Geographic distribution. Worldwide (Hooshyar et al. 2015).

Natural host. Humans (Hooshyar et al. 2015).

Route of infection. Ingestion of mature cysts in fecally contaminated water, food, fomites (Ali 2015).

Site in human host. Large intestine, cecum, colon (Ali 2015).

***Entamoeba histolytica* Schaudinn, 1903**

Amoeba coli Lösch, 1875

Amoeba dysenteriae Councilman & Laffleur, 1891

Entamoeba africana Hartmann & Von Prowazek, 1907

Entamoeba schaudinii Lesage, 1908

Entamoeba minuta Elmassian, 1909

Entamoeba nipponica Koidzumi, 1909

Entamoeba brasiliensis Aragao, 1912

Entamoeba venaticum Darling, 1915

Entamoeba caudata Carini & Reichenow, 1949

Geographic distribution. Worldwide; endemic areas of clinical amebiasis include Central and South America, Africa, Southeast Asia (Ali 2015; Shirley et al. 2018).

Natural host. Humans (Hooshyar et al. 2015).

Route of infection. Ingestion of mature cysts in fecally contaminated water, food, fomites; oral-anal sex (Shirley et al. 2018).

Site in human host. Large intestine, cecum, colon; ectopic colonization of liver, skin, lungs, brain (Ali 2015; Mathison and Pritt 2018; Shirley et al. 2018).

***Entamoeba moshkovskii* (Tshalaia, 1941)**

Entamoeba histolytica, Laredo strain

Entamoeba histolytica, Huff strain

Geographic distribution. Presumed worldwide; high areas of endemicity include Australia, India, Bangladesh, Tanzania, Pakistan, Iran (Ali 2015; Kyany'a et al. 2019).

Natural host. Humans (Hooshyar et al. 2015; Lopez et al. 2015).

Route of infection. Ingestion of mature cysts in fecally contaminated water, food, fomites (Ali 2015).

Site in human host. Large intestine, cecum, colon (Ali 2015).

***Entamoeba nuttalli* (Castellani, 1908)**

Loeschia duboscqi Mathis, 1913

Amoeba ateles Eichhorn & Gallagher, 1916

Loeschia cynomolgi Brug, 1923

Geographic distribution. Native to Southeast Asia, probably worldwide in zoos and from the animal trade (Tanaka et al. 2019).

Natural hosts. Non-human primates. Zoonotic in humans (Hooshyar et al. 2015; Levecke et al. 2015).

Route of infection. Ingestion of mature cysts in fecally contaminated water, food, fomites.

Site in human host. Large intestine, cecum, colon (Levecke et al. 2015)

***Entamoeba polecki* (Von Prowazek, 1912)**

Entamoeba deblickei Nieschulz, 1923

Geographic distribution. Worldwide; spots of high endemicity include Venezuela, Iran, Southeast Asia, Papua New Guinea (Verweij et al. 2001; Stensvold et al. 2018).

Natural hosts. Pigs, monkeys. Zoonotic in humans (Hooshyar et al. 2015).

Route of infection. Ingestion of mature cysts in fecally contaminated water, food, fomites (Ali 2015).

Site in human host. Large intestine, cecum, colon (Ali 2015).

Genus *Iodamoeba* Dobell, 1919***Iodamoeba buetschlii* (Von Prowazek, 1912)**

Geographic distribution. Worldwide (Ash and Orihel 2007).

Natural host. Human (Ash and Orihel 2007).

Route of infection. Ingestion of mature cysts in fecally contaminated water, food, fomites (Ash and Orihel 2007).

Site in human host. Large intestine, cecum, colon (Ash and Orihel 2007).

- **Discosea Cavalier-Smith et al., 2004**
- **Flabellinia Smirnov et al., 2005**
- **Thecamoebida Schaeffer, 1926, emend. Smirnov et al., 2011**
- **Thecamoebidae Schaeffer, 1926, emend. Smirnov et al., 2011**

Genus *Sappinia* Dangeard, 1896

***Sappinia pedata* Dangeard, 1896**

Geographic distribution. Presumed worldwide (Visvesvara 2013).

Natural host. None (environmental); humans are incidental hosts

Route of infection. Presumably by the entry of cysts or trophozoites through mucous membranes (e.g., nasal passages) or broken skin (Qvarnstrom et al. 2009; Visvesvara 2013).

Site in human host. Central nervous system (CNS) (Qvarnstrom et al. 2009; Visvesvara 2013).

- **Centramoebia Cavalier-Smith et al., 2016**
- **Acanthopodida Page, 1976**
- **Acanthamoebidae Sawyer & Griffin, 1975**

Genus *Acanthamoeba* Volonsky, 1931

Notes. Clinically, *Acanthamoeba* species are usually characterized by their genotypes rather than traditional Linnaean binominal nomenclature (Booton et al. 2005).

***Acanthamoeba* Genotype T1**

Geographic distribution. Presumed worldwide (Booton et al. 2005).

Natural host. None (environmental); humans are incidental hosts.

Route of infection. Entry of environmental trophozoites or cysts through mucous membranes (e.g., eye, nasal passages) or broken skin (Booton et al. 2005; Khan 2006).

Site in human host. CNS (Booton et al. 2005; Khan 2006).

***Acanthamoeba* Genotype T2**

Geographic distribution. Presumed worldwide (Booton et al. 2005).

Natural host. None (environmental); humans are incidental hosts.

Route of infection. Entry of environmental trophozoites or cysts through mucous membranes (e.g., eye, nasal passages) (Booton et al. 2005; Khan 2006).

Site in human host. Eye (Khan 2006).

Acanthamoeba Genotype T3

Geographic distribution. Presumed worldwide (Booton et al. 2005).

Natural host. None (environmental); humans are incidental hosts.

Route of infection. Entry of environmental trophozoites or cysts through mucous membranes (e.g., eye, nasal passages) (Booton et al. 2005; Khan 2006).

Site in human host. Eye (Booton et al. 2005; Khan 2006).

Acanthamoeba Genotype T4

Geographic distribution. Worldwide (Booton et al. 2005).

Natural host. None (environmental); humans are incidental hosts.

Route of infection. Entry of environmental trophozoites or cysts through mucous membranes (e.g., eye, nasal passages) or broken skin (Booton et al. 2005; Khan 2006).

Site in human host. Eye, skin, lung, brain, sinus cavity, disseminated infection (Booton et al. 2005; Khan 2006).

Acanthamoeba Genotype T5

Geographic distribution. Presumed worldwide (Booton et al. 2005).

Natural host. None (environmental); humans are incidental hosts.

Route of infection. Entry of environmental trophozoites or cysts through mucous membranes (e.g., eye, nasal passages) (Ledee et al. 2009).

Site in human host. Eye (Ledee et al. 2009).

Acanthamoeba Genotype T6

Geographic distribution. Presumed worldwide (Booton et al. 2005).

Natural host. None (environmental); humans are incidental hosts.

Route of infection. Entry of environmental trophozoites or cysts through mucous membranes (e.g., eye, nasal passages) (Booton et al. 2005; Khan 2006).

Site in human host. Eye (Booton et al. 2005; Khan 2006).

Acanthamoeba Genotype T8

Geographic distribution. Presumed worldwide (Booton et al. 2005).

Natural host. None (environmental); humans are incidental hosts.

Route of infection. Entry of environmental trophozoites or cysts through mucous membranes (e.g., eye, nasal passages) (Orosz et al. 2018).

Site in human host. Eye (Orosz et al. 2018).

Acanthamoeba Genotype T10

Geographic distribution. Presumed worldwide (Booton et al. 2005).

Natural host. None (environmental); humans are incidental hosts

Route of infection. Entry of environmental trophozoites or cysts through mucous membranes (e.g., eye, nasal passages) or broken skin (Booton et al. 2005; Khan 2006; Nuprasert et al. 2010).

Site in human host. CNS, eye (Booton et al. 2005; Khan 2006; Nuprasert et al. 2010).

Acanthamoeba Genotype T11

Geographic distribution. Presumed worldwide (Booton et al. 2005).

Natural host. None (environmental); humans are incidental hosts.

Route of infection. Entry of environmental trophozoites or cysts through mucous membranes (e.g., eye, nasal passages) (Booton et al. 2005; Khan 2006).

Site in human host. Eye (Booton et al. 2005; Khan 2006).

Acanthamoeba Genotype T12

Geographic distribution. Presumed worldwide (Booton et al. 2005).

Natural host. None (environmental); humans are incidental hosts.

Route of infection. Entry of environmental trophozoites or cysts through mucous membranes (e.g., eye, nasal passages) or broken skin (Booton et al. 2005; Khan 2006).

Site in human host. CNS (Booton et al. 2005; Khan 2006).

Acanthamoeba Genotype T13

Geographic distribution. Presumed worldwide (Booton et al. 2005).

Natural host. None (environmental); humans are incidental hosts.

Route of infection. Entry of environmental trophozoites or cysts through mucous membranes (e.g., eye, nasal passages) (Grün et al. 2014)

Site in human host. Eye (Grün et al. 2014).

Acanthamoeba Genotype T15

Geographic distribution. Presumed worldwide (Booton et al. 2005).

Natural host. None (environmental); humans are incidental hosts.

Route of infection. Entry of environmental trophozoites or cysts through mucous membranes (e.g., eye, nasal passages) (Di Cave et al. 2009).

Site in human host. Eye (Di Cave et al. 2009).

●●●● **Balamuthiidae Cavalier-Smith et al., 2004**

Genus *Balamuthia* Visvesvara et al., 1993

***Balamuthia mandrillaris* Visvesvara et al., 1993**

Geographic distribution. Worldwide (Visvesvara 2013; Cope et al. 2019).

Natural host. None (environmental); humans are incidental hosts.

Route of infection. Entry of environmental trophozoites or cysts through mucous membranes (e.g., eye, nasal passages) or broken skin (Visvesvara 2013; Cope et al. 2019).

Site in human host. Disseminated infection with predilection for CNS (Visvesvara 2013; Cope et al. 2019).

SAR (Stramenopiles-Alveolata-Rhizaria) Burki et al., 2008, emend. Adl et al., 2012

● **Stramenopiles Patterson 1989, emend. Adl et al., 2005**

●● **Bigyra Cavalier-Smith, 1998, emend. Cavalier-Smith et al., 2006**

●●● **Opalozoa Cavalier-Smith, 1991, emend. Cavalier-Smith et al., 2006**

●●●● **Opalinata Wenyon, 1926, emend. Cavalier-Smith, 1997**

Genus *Blastocystis* Alexeieff, 1911

Notes. As the molecular epidemiology is becoming better understood, it is becoming less common to recognize *Blastocystis* by traditional Linnaean binominal nomenclature but rather by their genetic subtypes (Stensvold et al. 2007; Tan 2008).

***Blastocystis* subtype 1**

Geographic distribution. Worldwide (Stensvold et al. 2007; Tan 2008).

Natural hosts. Various mammals. Zoonotic in humans (Stensvold et al. 2007; Tan 2008).

Route of infection. Ingestion of cyst-forms and/or vacuolar-forms in contaminated food, water, fomites (Tan 2008).

Site in human host. Large intestine (Tan 2008).

***Blastocystis* subtype 2**

Geographic distribution. Worldwide (Stensvold et al. 2007; Tan 2008).

Natural hosts. Non-human primates, pigs. Zoonotic in humans (Stensvold et al. 2007; Tan 2008).

Route of infection. Ingestion of cyst-forms and/or vacuolar-forms in contaminated food, water, fomites (Tan 2008).

Site in human host. Large intestine (Tan 2008).

Blastocystis subtype 3

Geographic distribution. Worldwide (Stensvold et al. 2007; Tan 2008).

Natural hosts. Humans (Stensvold et al. 2007; Tan 2008).

Route of infection. Ingestion of cyst-forms and/or vacuolar-forms in contaminated food, water, fomites (Tan 2008).

Site in human host. Large intestine (Tan 2008).

Blastocystis subtype 4

Geographic distribution. Worldwide (Stensvold et al. 2007; Tan 2008).

Natural hosts. Rodents. Zoonotic in humans (Stensvold et al. 2007; Tan 2008).

Route of infection. Ingestion of cyst-forms and/or vacuolar-forms in contaminated food, water, fomites (Tan 2008).

Site in human host. Large intestine (Tan 2008).

Blastocystis subtype 5

Geographic distribution. Worldwide (Stensvold et al. 2007; Tan 2008).

Natural hosts. Pigs, cattle. Zoonotic in humans (Stensvold et al. 2007; Tan 2008).

Route of infection. Ingestion of cyst-forms and/or vacuolar-forms in contaminated food, water, fomites (Tan 2008).

Site in human host. Large intestine (Tan 2008).

Blastocystis subtype 6

Geographic distribution. Worldwide (Stensvold et al. 2007; Tan 2008).

Natural hosts. Birds. Zoonotic in humans (Stensvold et al. 2007; Tan 2008).

Route of infection. Ingestion of cyst-forms and/or vacuolar-forms in contaminated food, water, fomites (Tan 2008).

Site in human host. Large intestine (Tan 2008).

Blastocystis subtype 7

Geographic distribution. Worldwide (Stensvold et al. 2007; Tan 2008).

Natural hosts. Birds. Zoonotic in humans (Stensvold et al. 2007; Tan 2008).

Route of infection. Ingestion of cyst-forms and/or vacuolar-forms in contaminated food, water, fomites (Tan 2008).

Site in human host. Large intestine (Tan 2008).

Blastocystis subtype 8

Geographic distribution. Worldwide (Stensvold et al. 2007; Tan 2008).

Natural hosts. Monkeys, birds. Zoonotic in humans (Stensvold et al. 2007; Tan 2008).

Route of infection. Ingestion of cyst-forms and/or vacuolar-forms in contaminated food, water, fomites (Tan 2008).

Site in human host. Large intestine (Tan 2008).

Blastocystis subtype 9

Geographic distribution. Worldwide (Stensvold et al. 2007; Tan 2008).

Natural hosts. Unknown; presumed zoonotic in humans (Stensvold et al. 2007; Tan 2008).

Route of infection. Ingestion of cyst-forms and/or vacuolar-forms in contaminated food, water, fomites (Tan 2008).

Site in human host. Large intestine (Tan 2008).

Blastocystis subtype 12

Geographic distribution. Worldwide (Stensvold et al. 2007; Tan 2008).

Natural hosts. Goats, cattle. Zoonotic in humans (Stensvold et al. 2007; Tan 2008; Udonsom et al. 2018; Jiménez et al. 2019).

Route of infection. Ingestion of cyst-forms and/or vacuolar-forms in contaminated food, water, fomites (Tan 2008).

Site in human host. Large intestine (Tan 2008).

- Alveolata Cavalier-Smith, 1991
- Colpodellida Cavalier-Smith, 1993, amend. Adel et al. 2005, 2019
- Colpodellidae Simpson & Patterson, 1996

Genus *Colpodella* Cienkowski, 1865

Geographic distribution. Worldwide (Getty et al. 2021).

Natural hosts. None; free-living predators of other protozoans. Humans are incidental hosts (Getty et al. 2021).

Route of infection. Unknown, proposed transmission via the bite of ticks in the genus *Ixodes* (Yuan et al. 2012; Jiang et al. 2018).

Site in human host. Blood (Yuan et al. 2012; Jiang et al. 2018).

Notes. In nature, *Colpodella* species are free-living predators of other protozoans. *Copodella*-like organisms have been diagnosed twice in patients from China, one from blood (Yuan et al. 2012) and once from CSF (Jiang et al. 2018); only the former also demonstrated morphologic evidence of a potential parasite. In both cases, the identifications were made by sequencing analyses. The species-level identification was not made in either case, and the organisms in at least one of the cases (Yuan et al. 2012) might represent an as-of-yet undescribed genus. Further work is

needed to confirm the ability of *Colpodella* and related organisms to cause parasitic infections in humans.

●● **Apicomplexa** Levine, 1970, emend. Adl et al. 2005

●●● **Aconoidasida** Mehlhorn et al., 1980

Hematazoa Vivier, 1982

●●●● **Haemospororida** Danilewsky, 1885

●●●●● **Plasmodiidae** Mesnil, 1903

Genus *Plasmodium* Marchiafava & Celli, 1885

Plasmodium brasilianum Gonder et von Berenberg-Gossler, 1908

Geographic distribution. Brazil, Venezuela, Colombia, Panama, Peru (Coatney et al. 1971).

Natural hosts. Intermediate hosts are New World monkeys. Arthropod vector and definitive hosts are mosquitoes in the genus *Anopheles*. Zoonotic in humans as an intermediate host (Coatney et al. 1971; Lalremruata et al. 2015).

Route of infection. Introduction of sporozoites via the bite of infected *Anopheles* mosquito (Coatney et al. 1971).

Site in human host. Liver (initial infection), blood (Coatney et al. 1971).

Notes. It has been suggested that *P. brasilianum* is conspecific with *P. malariae* and that *P. malariae* adapted to non-human primates in Latin America after being introduced from Africa (Guimaraes et al. 2012).

Plasmodium coatneyi Eyles et al., 1962

Geographic distribution. Peninsular Malaysia, Philippines (Coatney et al. 1971).

Natural hosts. Intermediate host is the crab-eating macaque (*Macaca fascicularis*). Arthropod vector and definitive hosts are mosquitoes in the genus *Anopheles*. Zoonotic in humans as an intermediate host (Coatney et al. 1971; Yap et al. 2021).

Route of infection. Introduction of sporozoites via the bite of infected *Anopheles* mosquito (Coatney et al. 1971).

Site in human host. Liver (initial infection), blood (Coatney et al. 1971).

Plasmodium cynomolgi Mayer, 1907

Geographic distribution. Southeast Asia (Coatney et al. 1971).

Natural hosts. Intermediate hosts are macaques. Arthropod vector and definitive hosts are mosquitoes in the genus *Anopheles*. Zoonotic in humans as an intermediate host (Coatney et al. 1971; Hartmeyer et al. 2019; Yap et al. 2021).

Route of infection. Introduction of sporozoites via the bite of infected *Anopheles* mosquito (Coatney et al. 1971).

Site in human host. Liver (initial infection, chronic sequestration), blood (Coatney et al. 1971).

Plasmodium falciparum (Welch, 1897)

Oscillaria malariae Laveran, 1881

Haemamoeba praecox Feletti & Grassi, 1890

Haemamoeba immaculata Grassi, 1891

Haemamoeba laverani Labbe, 1894

Haemosporidium sedecimanae Lewkowitz, 1897

Haemosporidium vigesimotertiana Lewkowitz, 1897

Geographic distribution. Circumtropical; sub-Saharan Africa, Asia, Latin America, Caribbean (Snow et al. 2005; Weiss et al. 2019).

Natural hosts. Intermediate hosts are humans. Arthropod vector and definitive hosts are mosquitoes in the genus *Anopheles* (Coatney et al. 1971).

Route of infection. Introduction of sporozoites via the bite of infected *Anopheles* mosquito (Coatney et al. 1971).

Site in human host. Liver (initial infection), blood, CNS (cerebral malaria) (Coatney et al. 1971; Idro et al. 2010).

Plasmodium inui Helberstaedter & Von Prowazek, 1907

Geographic distribution. Southeast Asia (Coatney et al. 1971).

Natural hosts. Intermediate hosts are Old World monkeys. Arthropod vector and definitive hosts are mosquitoes in the genus *Anopheles*. Zoonotic in humans as an intermediate host (Coatney et al. 1971; Yap et al. 2021).

Route of infection. Introduction of sporozoites via the bite of infected *Anopheles* mosquito (Coatney et al. 1971).

Site in human host. Liver (initial infection), blood (Coatney et al. 1971; Liew et al. 2021).

Plasmodium knowlesi Sinton & Mulligan, 1932

Geographic distribution. Southeast Asia (Coatney et al. 1971).

Natural hosts. Intermediate hosts are macaques. Arthropod vector and definitive hosts are mosquitoes in the genus *Anopheles*. Zoonotic in humans as an intermediate host (Coatney et al. 1971; Singh and Daneshvar 2013; Yap et al. 2021).

Route of infection. Introduction of sporozoites via the bite of infected *Anopheles* mosquito (Coatney et al. 1971).

Site in human host. Liver (initial infection), blood (Coatney et al. 1971).

***Plasmodium malariae* (Grassi & Feletti, 1890)**

Plasmodium quartanae Celli & Sanfelice, 1891

Haematosporidium tertiana Labbe, 1894

Geographic distribution. Sub-Saharan Africa, Southeast Asia, Indonesia, Pacific Islands, Amazonian South America (Coatney et al. 1971; Collins and Jeffery 2007).

Natural hosts. Intermediate hosts are humans; non-human primates in Latin America are reservoir hosts. Arthropod vector and definitive hosts are mosquitoes in the genus *Anopheles* (Coatney et al. 1971; Collins and Jeffery 2007).

Route of infection. Introduction of sporozoites via the bite of infected *Anopheles* mosquito (Coatney et al. 1971; Collins and Jeffery 2007).

Site in human host. Liver (initial infection), blood (Coatney et al. 1971; Collins and Jeffery 2007).

***Plasmodium ovale curtisi* Sutherland et al., 2010**

Geographic distribution. Western sub-Saharan Africa, Southeast Asia (Coatney et al. 1971).

Natural hosts. Intermediate hosts are humans. Arthropod vector and definitive hosts are mosquitoes in the genus *Anopheles* (Coatney et al. 1971; Collins and Jeffery 2005; Oguike et al. 2011).

Route of infection. Introduction of sporozoites via the bite of infected *Anopheles* mosquito (Coatney et al. 1971; Collins and Jeffery 2005).

Site in human host. Liver (initial infection, chronic sequestration), blood (Coatney et al. 1971; Collins and Jeffery 2005).

Notes. With the separation of *P. ovale* into the subspecies *P. ovale curtisi* and *P. o. wallikeri*, the epidemiology of the two species is not well understood. Both subspecies are sympatric at the country level in several African countries, but their individual distributions in Southeast Asia are not well defined (Oguike et al. 2011).

***Plasmodium ovale wallikeri* Sutherland et al., 2010**

Geographic distribution. Western sub-Saharan Africa, Southeast Asia (Coatney et al. 1971).

Natural hosts. Intermediate hosts are humans. Arthropod vector and definitive hosts are mosquitoes in the genus *Anopheles* (Coatney et al. 1971; Collins and Jeffery 2005; Oguike et al. 2011).

Route of infection. Introduction of sporozoites via the bite of infected *Anopheles* mosquito (Coatney et al. 1971; Collins and Jeffery 2005).

Site in human host. Liver (initial infection, chronic sequestration), blood (Coatney et al. 1971; Collins and Jeffery 2005).

Notes. With the separation of *P. ovale* into the subspecies *P. ovale curtisi* and *P. o. wallikeri*, the epidemiology of the two species is not well understood. Both subspecies are sympatric at the country level in several African countries, but their individual distributions in Southeast Asia are not well defined (Oguike et al. 2011).

***Plasmodium schwetzi* Brumpt, 1938**

Geographic distribution. Tropical Africa (Coatney et al. 1971).

Natural hosts. Intermediate hosts are gorillas and chimpanzees. Arthropod vector and definitive hosts are mosquitoes in the genus *Anopheles*. Zoonotic in humans as intermediate hosts (Contacos et al. 1970; Coatney et al. 1971).

Route of infection. Introduction of sporozoites via the bite of infected *Anopheles* mosquito (Coatney et al. 1971).

Site in human host. Liver (initial infection, presumed chronic sequestration), blood (Coatney et al. 1971).

***Plasmodium simiovale* Dissanaïke, Nelson & Garnham, 1965**

Geographic distribution. Sri Lanka, Malaysia (Coatney et al. 1971).

Natural hosts. Intermediate hosts is the toque macaque (*Macaca sinica*). Arthropod vector and definitive hosts are mosquitoes in the genus *Anopheles*. Zoonotic in humans as intermediate hosts (Coatney et al. 1971; Yap et al. 2021).

Route of infection. Introduction of sporozoites via the bite of infected *Anopheles* mosquito (Coatney et al. 1971).

Site in human host. Liver (initial infection, chronic sequestration), blood (Coatney et al. 1971; Yap et al. 2021).

***Plasmodium simium* Fonseca, 1951**

Geographic distribution. Brazil (Coatney et al. 1971).

Natural hosts. Intermediate hosts are howler monkeys and capuchins. Arthropod vector and definitive hosts are mosquitoes in the genus *Anopheles*. Zoonotic in humans as intermediate hosts (Coatney et al. 1971; Brasil et al. 2017).

Route of infection. Introduction of sporozoites via the bite of infected *Anopheles* mosquito (Coatney et al. 1971).

Site in human host. Liver (initial infection, chronic sequestration), blood (Coatney et al. 1971; Brasil et al. 2017).

Notes. It has been suggested *P. simium* is conspecific with *P. vivax*, and that *P. vivax* adapted to non-human primates in Latin America after being introduced by humans (Daron et al. 2020).

***Plasmodium vivax* (Grassi & Feletti, 1890)**

Haemamoeba malariae Feletti & Grassi, 1890, in part

Haemosporidium tertianae Lewkowitz, 1897

Plasmodium camarensis Ziemann, 1915

Geographic distribution. Africa (east, Horn of Africa, and Madagascar), Central and South America, Central Asia, Indian Subcontinent, Southeast Asia, Korean Peninsula (Coatney et al. 1971; Howes et al. 2016).

Natural hosts. Intermediate hosts are humans; non-human primates can serve as reservoir hosts. Arthropod vector and definitive hosts are mosquitoes in the genus *Anopheles* (Coatney et al. 1971).

Route of infection. Introduction of sporozoites via the bite of infected *Anopheles* mosquito (Coatney et al. 1971).

Site in human host. Liver (initial infection, chronic sequestration), blood, CNS (Coatney et al. 1971; Sarkar and Bhattacharya 2008; Mukhtar et al. 2019).

●●●● **Piroplasmorida** Wenyon, 1926

●●●●● **Babesiidae** Poche, 1913

Genus *Babesia* Starcovivi, 1893

Babesia divergens M'Fadyean & Stockman, 1911

Geographic distribution. Europe (Ord and Lobo 2015)

Natural hosts. Intermediate hosts are primarily cattle. Arthropod vector and definitive hosts are ticks in the genus *Ixodes*, primarily *I. ricinus*. Zoonotic in humans as intermediate hosts (Ord and Lobo 2015; Kukina et al. 2018, 2019).

Route of infection. Introduction of sporozoites via the bite of the *Ixodes* tick vector; person-to-person transmission via blood and solid organ transplants (Ord and Lobo 2015; Kukina et al. 2018, 2019).

Site in human host. Blood (Ord and Lobo 2015; Kukina et al. 2018, 2019).

Babesia duncani Conrad et al., 2006

Babesia strain WA1

Babesia strain WA2

Babesia strain CA5

Babesia strain CA6

Geographic distribution. Western and northern North America (Conrad et al. 2006; Ord and Lobo 2015).

Natural hosts. Intermediate hosts are suspected as being mule deer (*Odocoileus hemionus*). Arthropod vector and definitive host is believed to be the tick *Dermacentor albipictus*. Zoonotic in humans (Swei et al. 2019).

Route of infection. Introduction of sporozoites via the bite of the presumptive tick vector; possibly also via blood transfusion and solid organ transplants (Conrad et al. 2006).

Site in human host. Blood.

***Babesia microti* (França, 1910)**

Geographic distribution. Northeastern North America, Australia (Senanayake et al. 2012; Ord and Lobo 2015; Westblade et al. 2017).

Natural hosts. Intermediate hosts are rodents, primarily white-footed mice (*Peromyscus leucopus*), and deer, primarily white-tailed deer (*Odocoileus virginianus*). Arthropod vector and definitive hosts are ticks in the genus *Ixodes*, primarily *I. scapularis*. Zoonotic in humans as intermediate hosts (Westblade et al. 2017).

Route of infection. Introduction of sporozoites via the bite of infected tick, blood transfusion, and solid organ transplants (Ord and Lobo 2015; Westblade et al. 2017).

Site in human host. Blood (Ord and Lobo 2015; Westblade et al. 2017).

***Babesia motasi* Wenyou, 1926**

Babesia sp. Strain KO-1

Babesia ovis-like

Geographic distribution. Eurasia (Kim et al. 2007; Hong et al. 2019).

Natural hosts. Intermediate hosts are sheep. Definitive hosts are ticks in the genus *Haemaphysalis*. Zoonotic in humans as intermediate hosts (Kim et al. 2007; Hong et al. 2019; Wang et al. 2020).

Route of infection. Introduction of sporozoites via the bite of infected tick (Kim et al. 2007; Hong et al. 2019; Wang et al. 2020).

Site in human host. Blood (Hong et al. 2019).

***Babesia odocoilei* Emerson & Wright, 1970**

Geographic distribution. Eastern North America (Pattullo et al. 2013).

Natural hosts. Intermediate hosts are cervids, primarily white-tailed deer (*Odocoileus virginianus*). Arthropod vector and definitive host is the tick *Ixodes scapularis*; birds may serve as reservoir hosts for larvae or nymphs. Zoonotic in humans as intermediate hosts (Pattullo et al. 2013; Scott et al. 2021).

Route of infection. Introduction of sporozoites via the bite of infected tick (Scott et al. 2021).

Site in human host. Blood (Scott et al. 2021).

***Babesia venatorum* Herwaldt et al., 2003**

Babesia strain EU1

Geographic distribution. Europe (Herwaldt et al. 2003).

Natural hosts. Intermediate hosts are red deer (*Cervus elaphus*); roe deer, sheep, dogs, cats, and other mammals may serve as reservoir hosts. Arthropod vectors and definitive hosts are ticks in the genus *Ixodes*, primarily *I. ricinus*. Zoonotic in humans as intermediate hosts (Herwaldt et al. 2003; Gray et al. 2019).

Route of infection. Introduction of sporozoites via the bite of infected *Ixodes* tick; possibly also via blood transfusion and solid organ transplants (Herwaldt et al. 2003).

Site in human host. Blood (Herwaldt et al. 2003).

***Babesia* sp. Strain MO-1**

Babesia divergens-like

Geographic distribution. Central and Pacific Northwest USA (Burgess et al. 2017).

Natural hosts. Unknown; presumed intermediate hosts are unknown and presumed arthropod vector and definitive hosts are ticks in the genus *Ixodes*. Zoonotic in humans as intermediate hosts (Burgess et al. 2017; Herc et al. 2018).

Route of infection. Introduction of sporozoites via the bite of infected tick; possibly also via blood transfusion and solid organ transplantation (Burgess et al. 2017; Herc et al. 2018).

Site in human host. Blood (Burgess et al. 2017; Herc et al. 2018).

Notes. This strain was originally described from Missouri, USA. It has since been described from the states of Washington, Kentucky, and Arkansas (Burgess et al. 2017). Genetically it is very similar to the Palearctic *B. divergens* and is often referred to as *Babesia divergens*-like (Herc et al. 2018).

***Babesia* sp. crassa-like**

Geographic distribution. Undefined; human cases known from China and Slovenia (Jia et al. 2018; Strasek-Smrđel et al. 2020).

Natural hosts. Unknown; presumed intermediate hosts are sheep, goats, presumed arthropod vector and definitive hosts are ticks in the genera *Haemaphysalis* and/or *Ixodes*. Zoonotic in humans as intermediate hosts (Jia et al. 2018; Strasek-Smrđel et al. 2020).

Route of infection. Introduction of sporozoites via the bite of infected tick; possibly also via blood transfusion and solid organ transplantation (Jia et al. 2018; Strasek-Smrđel et al. 2020).

Site in human host. Blood (Jia et al. 2018; Strasek-Smrđel et al. 2020).

●●●●● Anthemosomatidae Levine, 1981**Genus *Anthemosoma* Landau et al., 1969*****Anthemosoma garnhami* Landau et al., 1969**

Geographic distribution. Sub-Saharan Africa (Chavatte et al. 2018).

Natural hosts. Intermediate hosts are rodents, including spiny mice (*Acomys*). Arthropod vector and definitive hosts are unknown but presumed to be ixodid ticks. Zoonotic in humans (Chavatte et al. 2018; Stead et al. 2021).

Route of infection. Presumably by introduction of sporozoites via the bite of an infected tick (Stead et al. 2021).

Site in human host. Blood (Stead et al. 2021).

●●● Conoidasida Levine, 1988**●●●● Coccidia Leuckart, 1879****●●●●● Eimeriorina Léger, 1911****●●●●●● Eimeriidae Minchin, 1903****Genus *Cyclospora* Schneider, 1881*****Cyclospora cayetanensis* Ortega et al., 1994**

Geographic distribution. Nearly worldwide, hot spots of endemicity include Latin America, the Caribbean, Middle East, and Southeast Asia (Ortega and Sanchez 2010; Almeria et al. 2019).

Natural host. Human (Ortega and Sanchez 2010; Almeria et al. 2019).

Route of infection. Ingestion of sporulated oocysts in fecally contaminated produce and water (Ortega and Sanchez 2010; Cama and Mathison 2015; Almeria et al. 2019).

Site in human host. Small intestine (Ortega and Sanchez 2010; Almeria et al. 2019).

●●●●●● Sarcocystidae Poche, 1913**Genus *Cystoisospora* Frenkel, 1977*****Cystoisospora belli* (Wenyon, 1923)**

Geographic distribution. Nearly worldwide, more prevalent in tropics and subtropics (Cama and Mathison 2015; Dubey and Almeria 2019).

Natural host. Human (Cama and Mathison 2015; Dubey and Almeria 2019).

Route of infection. Ingestion of sporulated oocysts in fecally contaminated water, food, fomites (Cama and Mathison 2015; Dubey and Almeria 2019)

Site in human host. Small intestine (Cama and Mathison 2015; Dubey and Almeria 2019; Rowan et al. 2020)

Genus *Sarcocystis* Lankester, 1882

***Sarcocystis heydorni* Dubey et al., 2015**

Geographic distribution. Europe, China (Dubey et al. 2015).

Natural hosts. Cattle are intermediate hosts. Humans are the definitive hosts (Dubey 2015; Dubey et al. 2015).

Route of infection. Ingestion of cysts (bradyzoites) in undercooked beef (Dubey 2015; Dubey et al. 2015).

Site in human host. Small intestine (Dubey 2015; Dubey et al. 2015)

***Sarcocystis hominis* (Raillet & Lucet, 1891)**

Sarcocystis bovi-hominis Heydorn et al., 1975

Geographic distribution. Nearly worldwide, more prevalent in tropics and subtropics (Dubey 2015).

Natural hosts. Intermediate hosts are cattle. Definitive hosts are humans and non-human primates (Fayer 2004; Dubey 2015; Fayer et al. 2015).

Route of infection. Ingestion of cysts (bradyzoites) in undercooked beef (Fayer 2004; Dubey 2015; Fayer et al. 2015).

Site in human host. Small intestine (Fayer 2004; Dubey 2015; Fayer et al. 2015).

***Sarcocystis nesbitti* Mandour, 1969**

Geographic distribution. Southeast Asia (Abubakar et al. 2013; Fayer et al. 2015).

Natural hosts. Unknown. Natural intermediate host is unknown, non-human primates can serve as intermediate or reservoir hosts. Definitive hosts are presumed to be snakes or other reptiles. Zoonotic in humans as incidental intermediate hosts (Abubakar et al. 2013).

Route of infection. Ingestion of fully-sporulated oocysts in contaminated food, water, fomites (Abubakar et al. 2013; Fayer et al. 2015).

Site in human host. Skeletal muscle (Abubakar et al. 2013; Fayer et al. 2015).

***Sarcocystis suihominis* Heydorn, 1977**

Geographic distribution. Nearly worldwide, more prevalent in tropics and subtropics (Dubey 2015).

Natural hosts. Intermediate hosts are pigs. Definitive hosts are humans and non-human primates (Fayer 2004; Dubey 2015; Fayer et al. 2015).

Route of infection. Ingestion of cysts (bradyzoites) in undercooked pork (Fayer 2004; Dubey 2015; Fayer et al. 2015).

Site in human host. Small intestine (Fayer 2004; Dubey 2015; Fayer et al. 2015).

Genus *Toxoplasma* Nicolle & Manceaux, 1909

Toxoplasma gondii (Nicolle & Manceaux, 1908)

Geographic distribution. Worldwide (Robert-Gangneux and Darde 2012).

Natural hosts. Wild and domestic felids. Many mammals and birds can serve as reservoir hosts; zoonotic in humans as dead-end hosts (Attias et al. 2020).

Route of infection. Ingestion of fully sporulated oocysts in food, water, fomites contaminated with cat feces; ingestion of cysts (bradyzoites) in infected paratenic hosts; blood transfusion or solid organ transplantation; transplacentally from mother to fetus (Montoya and Remington 2008; Robert-Gangneux and Darde 2012; Attias et al. 2020).

Site in human host. Disseminated infection, common sites are skeletal muscle, myocardium, brain, and eyes; can be transmitted transplacentally from mother to fetus (Montoya and Remington 2008; Robert-Gangneux and Darde 2012).

●●●● Gregarinasina Dufour, 1828

●●●●● Cryptogregarinorida Cavalier-Smith, 2014, emend. Adl et al., 2019

●●●●●● Cryptosporidiidae Leger, 1911

Genus *Cryptosporidium* Tyzzer, 1910

Cryptosporidium andersoni Lindsay et al., 2000

Geographic distribution. Worldwide (Feng et al. 2011).

Natural hosts. Cattle, camels, sheep, goats, horses. Zoonotic in humans (Feng et al. 2011; Liu et al. 2015).

Route of infection. Ingestion of sporulated oocysts in water, food, fomites contaminated with animal feces (Jiang et al. 2014).

Site in human host. Unknown, presumed stomach.

Cryptosporidium bovis Barker & Carbonell, 1974

Geographic distribution. Worldwide (Feng et al. 2007).

Natural hosts. Cattle. Zoonotic in humans (Ng et al. 2012; Das et al. 2019).

Route of infection. Ingestion of sporulated oocysts in water, food, fomites contaminated with cattle feces (Ng et al. 2012; Das et al. 2019).

Site in human host. Duodenum and small intestine.

***Cryptosporidium canis* Fayer et al., 2001**

Geographic distribution. Worldwide (Fayer et al. 2001).

Natural hosts. Dogs. Zoonotic in humans (Fayer et al. 2001; Ryan et al. 2016).

Route of infection. Ingestion of sporulated oocysts in water, food, fomites contaminated with dog feces (Fayer et al. 2001; Ryan et al. 2016).

Site in human host. Duodenum and small intestine

***Cryptosporidium cuniculus* Inman & Takeuchi, 1979**

Geographic distribution. Undefined, most human cases from Europe and Australia (Koehler et al. 2014; Puleston et al. 2014).

Natural hosts. Rabbits, kangaroos. Zoonotic in humans (Koehler et al. 2014; Puleston et al. 2014).

Route of infection. Ingestion of sporulated oocysts water, food, fomites contaminated with rabbit/animal feces (Koehler et al. 2014; Puleston et al. 2014; Ryan et al. 2016).

Site in human host. Duodenum and small intestine.

***Cryptosporidium ditrichi* Čondlová et al., 2018**

Geographic distribution. Europe (Condlova et al. 2018).

Natural hosts. Field mice (*Apodemus* spp.). Zoonotic in humans (Condlova et al. 2018; Beser et al. 2020).

Route of infection. Ingestion of sporulated oocysts in water, food, fomites contaminated with rodent feces (Beser et al. 2020; Condlova et al. 2018).

Site in human host. Duodenum and small intestine

***Cryptosporidium erinacei* Kváč et al., 2014**

Cryptosporidium hedgehog genotype

Geographic distribution. Europe (Kváč et al. 2014).

Natural hosts. Hedgehogs, horses. Zoonotic in humans (Kváč et al. 2014; Zahedi et al. 2016).

Route of infection. Ingestion of sporulated oocysts in water, food, fomites contaminated with animal feces (Kváč et al. 2014; Zahedi et al. 2016).

Site in human host. Duodenum and small intestine

***Cryptosporidium fayeri* Ryan et al., 2008**

Geographic distribution. Australia (Ryan et al. 2008; Waldron et al. 2010).

Natural hosts. Marsupials. Zoonotic in humans (Waldron et al. 2010).

Route of infection. Ingestion of sporulated oocysts in water, food, fomites contaminated with marsupial feces (Waldron et al. 2010).

Site in human host. Duodenum and small intestine

***Cryptosporidium felis* Iseki, 1979**

Geographic distribution. Worldwide (Jiang et al. 2020).

Natural hosts. Cats, foxes, horses, cattle. Zoonotic in humans (Caccio et al. 2002; Jiang et al. 2020).

Route of infection. Ingestion of sporulated oocysts in fecally contaminated water, food, fomites contaminated with cat/animal feces (Caccio et al. 2002; Jiang et al. 2020).

Site in human host. Duodenum and small intestine

***Cryptosporidium hominis* Morgan-Ryan et al., 2002**

Geographic distribution. Worldwide (Cama and Mathison 2015).

Natural hosts. Humans, non-human primates (Cama and Mathison 2015; Ryan et al. 2016; Chen et al. 2019).

Route of infection. Ingestion of sporulated oocysts in water, food, fomites contaminated with human feces (Cama and Mathison 2015).

Site in human host. Duodenum and small intestine, disseminated infections in immunocompromised patients (Bouzid et al. 2013; Cama and Mathison 2015).

***Cryptosporidium meleagridis* Slavin, 1955**

Geographic distribution. Worldwide

Natural hosts. Birds. Zoonotic in humans (Akiyoshi et al. 2003).

Route of infection. Ingestion of sporulated oocysts in fecally contaminated water, food, fomites contaminated with bird feces (Akiyoshi et al. 2003).

Site in human host. Duodenum and small intestine, disseminated infections in immunocompromised patients (Akiyoshi et al. 2003; Kopacz et al. 2019).

***Cryptosporidium muris* Tyzzer, 1910**

Geographic distribution. Worldwide (Chappell et al. 2015).

Natural hosts. Rodents, ruminants, horses, non-human primates, carnivores. Zoonotic in humans (Chappell et al. 2015).

Route of infection. Ingestion of sporulated oocysts in water, food, fomites contaminated with animal feces (Chappell et al. 2015).

Site in human host. Unknown, presumed stomach (Chappell et al. 2015).

***Cryptosporidium occultus* Kváč et al., 2018**

Cryptosporidium parvum VF383

Geographic distribution. Worldwide (Kvac et al. 2018).

Natural hosts. Rodents. Zoonotic in humans (Kvac et al. 2018; Xu et al. 2020).

Route of infection. Ingestion of sporulated oocysts in water, food, fomites contaminated with rodent feces (Kvac et al. 2018; Xu et al. 2020).

Site in human host. Duodenum and small intestine

Cryptosporidium parvum Tyzzer, 1912

Geographic distribution. Worldwide (Cama and Mathison 2015; Ryan et al. 2016).

Natural hosts. Cattle, sheep, goats, and other ruminants, rodents. Zoonotic in humans (Cama and Mathison 2015; Ryan et al. 2016).

Route of infection. Ingestion of sporulated oocysts in water, food, fomites contaminated with human and animal feces (Cama and Mathison 2015; Ryan et al. 2016).

Site in human host. Duodenum and small intestine, disseminated infections in immunocompromised patients (Cama and Mathison 2015; Ryan et al. 2016).

Cryptosporidium ryanae Fayer et al., 2008

Cryptosporidium deer-like genotype

Geographic distribution. XX (Fayer et al. 2008).

Natural hosts. Cattle. Zoonotic in humans (Chako et al. 2010; Das et al. 2019).

Route of infection. Ingestion of sporulated oocysts in water, food, fomites contaminated with cattle feces (Das et al. 2019).

Site in human host. Duodenum and small intestine.

Cryptosporidium scrofarum Kváč et al., 2013

Cryptosporidium pig genotype II

Geographic distribution. Worldwide (Kváč et al. 2013).

Natural hosts. Pigs. Zoonotic in humans (Ryan et al. 2016).

Route of infection. Ingestion of sporulated oocysts in water, food, fomites contaminated with pig feces (Ryan et al. 2016).

Site in human host. Duodenum and small intestine.

Cryptosporidium suis Ryan et al., 2004

Geographic distribution. Worldwide (Ryan et al. 2004).

Natural hosts. Pigs. Zoonotic in humans (Xiao et al. 2002).

Route of infection. Ingestion of sporulated oocysts in water, food, fomites contaminated with pig feces (Xiao et al. 2002; Nemejc et al. 2013)

Site in human host. Duodenum and small intestine

Cryptosporidium tyzzeri* Ren et al., 2012Cryptosporidium* mouse genotype I**Geographic distribution.** Presumed worldwide (Raskova et al. 2013).**Natural hosts.** Rodents, horses may serve as reservoir hosts. Zoonotic in humans (Raskova et al. 2013; Wagnerová et al. 2015).**Route of infection.** Ingestion of sporulated oocysts in water, food, fomites contaminated with rodent feces (Raskova et al. 2013).**Site in human host.** Duodenum and small intestine***Cryptosporidium ubiquitum* Fayer et al., 2010***Cryptosporidium* cervine genotype**Geographic distribution.** Worldwide (Li et al. 2014).**Natural hosts.** Many mammal species, including ruminants, rodents, and carnivores; presumed zoonotic in humans (Li et al. 2014).**Route of infection.** Ingestion of sporulated oocysts in water, food, fomites contaminated with animal feces (Li et al. 2014).**Site in human host.** Duodenum and small intestine***Cryptosporidium viatorum* Elwin et al., 2012****Geographic distribution.** Worldwide (Koehler et al. 2018).**Natural hosts.** Rodents. Zoonotic in humans (Koehler et al. 2018).**Route of infection.** Ingestion of sporulated oocysts in water, food, fomites contaminated with rodent and human feces (Koehler et al. 2018).**Site in human host.** Duodenum and small intestine.

●● Ciliophora Doflein, 1901

●●● Postodesmatophora Gerassimova & Seravin, 1976

●●●● SAL (Spirotrichea-Lamellicorticata-Armophorea) Gentekaki et al., 2014

●●●●● Litostomatea Small & Lynn, 1981

●●●●●● Trichostomatia Bütschli, 1889

●●●●●●● Balantidiidae Reichenow in Doflein & Reichenow, 1929

Genus *Balantioides* Alexeieff, 1931*Neobalantidium* Pomajbíková et al., 2013

***Balantioides coli* (Malmsten, 1857)**

Geographic distribution. Worldwide; high areas of endemicity include the Altiplano region of Bolivia, Philippines, New Guinea, Middle East (Schuster and Ramirez-Avila 2008).

Natural hosts. Pigs, rodents. Zoonotic in humans (Schuster and Ramirez-Avila 2008).

Route of infection. Ingestion of mature cysts in fecally contaminated water, food, fomites (Schuster and Ramirez-Avila 2008).

Site in human host. Large intestine (Schuster and Ramirez-Avila 2008).

Excavates Cavalier-Smith, 2002, emend. Simpson, 2003.

- **Metamonada** Grassé, 1952
- **Fornicata** Simpson, 2003
- **Diplomonadida** Wenyon, 1926
- **Hexamitidae** Kent, 1880

Genus *Enteromonas* da Fonseca, 1915***Enteromonas hominis* da Fonseca, 1915**

Tricercomonas intestinalis Wenyon et O'Connor, 1917

Geographic distribution. Worldwide (Ash and Orihel 2007).

Natural host. Humans (Ash and Orihel 2007).

Route of infection. Ingestion of mature cysts in fecally contaminated water, food, fomites (Spriegel et al. 1989; Ash and Orihel 2007).

Site in human host. Large intestine (Ash and Orihel 2007).

Genus *Giardia* Künstler, 1882***Giardia duodenalis* Stiles, 1902**

Cercomonas intestinalis Lambl, 1859

Giardia lamblia Kofoid & Christiansen, 1915

Giardia enterica Kofoid & Christiansen, 1920

Geographic distribution. Worldwide (Adam 2001; Cama and Mathison 2015).

Natural hosts. Two genetic assemblages implicated in human disease: A and B. Both have been found in humans and numerous non-human mammals and birds (Ryan and Caccio 2013; Heyworth 2016).

Route of infection. Ingestion of mature cysts in fecally contaminated water, food, fomites; sexual transmission via oral-anal route (Adam 2001; Escobedo et al. 2014; Cama and Mathison 2015).

Site in human host. Duodenum, small intestine (Adam 2001; Cama and Mathison 2015).

●●● Retortamonadida Grassé, 1952

●●●● Retortamonadidae Wenrich, 1932

Genus *Chilomastix* Alexeieff, 1912

Chilomastix mesnili (Wenyon, 1910)

Geographic distribution. Worldwide (Ash and Orihel 2007).

Natural hosts. Humans, non-human primates (Ash and Orihel 2007).

Route of infection. Ingestion of mature cysts in fecally contaminated water, food, fomites (Morimoto et al. 1996; Ash and Orihel 2007).

Site in human host. Large intestine (Ash and Orihel 2007).

Genus *Retortamonas* Grassi, 1879

Retortamonas intestinalis (Wenyon et O'Connoer, 1917)

Geographic distribution. Worldwide (Ash and Orihel 2007).

Natural host. Humans (Ash and Orihel 2007).

Route of infection. Ingestion of mature cysts in fecally contaminated water, food, fomites (Mendez et al. 2002; Ash and Orihel 2007).

Site in human host. Large intestine (Ash and Orihel 2007).

●● Parabasalia Honigberg, 1973

●●● Trichonomadida Kirby, 1947

●●●● Monocercomonadidae Kirby, 1944

Genus *Dientamoeba* Jepps & Dobell, 1918

Dientamoeba fragilis Jepps & Dobell, 1918

Geographic distribution. Worldwide (Stark et al. 2016).

Natural hosts. Humans, non-human primates, pigs (Stark et al. 2016).

Route of infection. Ingestion of trophozoites in fecally contaminated water, food, fomites. A putative cyst stage has been described, but it is not yet widely accepted among protozoologists (Munasinghe et al. 2013; Stark et al. 2014, 2016).

Site in human host. Large intestine (Stark et al. 2016).

●●●● **Trichomonadidae Chalmers & Pekkola, 1918, emend. Hampl et al., 2004**

Genus *Pentatrichomonas* Mensil, 1914

***Pentatrichomonas hominis* (Davaine, 1860)**

Geographic distribution. Worldwide (Ash and Orihel 2007).

Natural hosts. Humans, non-human primates, dogs, cats, foxes, rabbits, cattle, sheep, goats, deer (Li et al. 2016, 2017, 2018).

Route of infection. Ingestion of trophozoites in fecally contaminated fomites (Ash and Orihel 2007).

Site in human host. Large intestine (Ash and Orihel 2007).

Genus *Trichomonas* Donné, 1837

***Trichomonas tenax* (Müller, 1773)**

Geographic distribution. Worldwide (Dubar et al. 2020).

Natural host. Humans (Dubar et al. 2020).

Route of infection. Direct person-to-person contact (Dubar et al. 2020).

Site in human host. Oral cavity; ectopic colonization of pleural cavity (Bellanger et al. 2008; Marty et al. 2017; Dubar et al. 2020).

***Trichomonas vaginalis* (Donné, 1836)**

Geographic distribution. Worldwide (Schwebke and Burgess 2004; Kissinger 2015).

Natural host. Humans (Schwebke and Burgess 2004; Kissinger 2015).

Route of infection. Sexual contact (Schwebke and Burgess 2004; Kissinger 2015).

Site in human host. Urogenital tract; rare ectopic colonization of CSF (Schwebke and Burgess 2004; Hamilton et al. 2018).

Genus *Tritrichomonas* Kofoid, 1920

***Tritrichomonas foetus* Reidmüller, 1928**

***Trichomonas suis* Gruby & Delafond, 1843**

Geographic distribution. Worldwide (Suzuki et al. 2016).

Natural host. Cattle, pigs, cats. Zoonotic and opportunistic in immunocompromised humans (Maritz et al. 2014; Yao and Köster 2015; Suzuki et al. 2016).

Route of infection. Unknown.

Site in human host. Respiratory tract, CNS, urogenital tract (Duboucher et al. 2006; Zalonis et al. 2011; Suzuki et al. 2016).

Genus *Tetratrichomonas* Parisi, 1910

Tetratrichomonas sp. cf. *gallinarum*

Geographic distribution. Unknown, presumed worldwide

Natural host. Unknown, presumed birds. Zoonotic in humans (Mantini et al. 2009; Maritz et al. 2014).

Route of infection. Unknown

Site in human host. Respiratory tract, possibly associated with bacterial infection (Mantini et al. 2009; Maritz et al. 2014).

Notes. This species has yet to be formally described. It was first isolated from pleural fluid in a patient with empyema. Bacterial cultures of the pleural fluid grew two *Streptococcus* species and *Prevotella* (Mantini et al. 2009). The protozoan may have been associated with bacteria and not an agent of human disease.

Tetratrichomonas sp. (undescribed species)

Tetratrichomonas empyemagena Lopez-Escamilla et al., 2013

Geographic distribution. Unknown, presumed worldwide

Natural hosts. Unknown

Route of infection. Unknown

Site in human host. Respiratory tract (Lopez-Escamilla et al. 2013)

Notes. This species was reported from two patients from Mexico (Lopez-Escamilla et al. 2013). The species was originally named *T. empyemagena*, but because it was not described under the criteria of the ICZN, its name should be considered *nomen nudum*. This isolate might be conspecific with that described by Mantini et al. (2009).

●●●● Simplicimonadidae Čepička et al., 2010

Genus *Simplicimonas* Čepička et al., 2010

Simplicimonas similis Čepička et al., 2010

Geographic distribution. Undefined, possibly worldwide (Dimasuay et al. 2013).

Natural hosts. Several vertebrates; previously recorded from lizards, cattle, and chickens (Dimasuay et al. 2013).

Route of infection. Unknown.

Site in human host. Intestinal tract (Greigert et al. 2018).

Notes. *Simplicimonas similis* was detected by molecular methods in four human patients in Madagascar (Greigert et al. 2018). The authors acknowledged that the parasite was not detected by microscopy and the finding may be based on spurious passage after consuming insects or another host, and further studies are required to determine if humans can serve as a reliable host (Greigert et al. 2018).

- *Discoba* Simpson in Hampl et al., 2009
- *Heterolobosea* Page & Blanton, 1985
- *Tetramitia* Cavalier-Smith, 1993
- *Vahlkampfiidae* Jollos, 1917

Genus *Naegleria* Alexeieff, 1912

***Naegleria fowleri* Carter, 1970**

Geographic distribution. Worldwide (Gharpure et al. 2020).

Natural host. None (environmental); humans are incidental hosts (Visvesvara 2013).

Route of infection. Exposure to trophozoites and cysts in contaminated fresh-water via the nasal cavity; improper use of nasal irrigation devices (Yoder et al. 2012; Visvesvara 2013).

Site in human host. CNS (Visvesvara 2013; Yoder et al. 2012; Mathison and Pritt 2018).

Genus *Paravahlkampfia* Brown et de Jonckheere, 1999

***Paravahlkampfia francinae* Visvesvara et al., 2009**

Geographic distribution. Unknown (single case from Ohio, USA) (Visvesvara et al. 2009).

Natural host. None (environmental); humans are incidental hosts (Visvesvara et al. 2009).

Route of infection. Unknown; presumed environmental exposure.

Site in human host. CNS (Visvesvara et al. 2009).

Genus *Tetramitus* Perty, 1852

***Tetramitus entericus* (Page, 1974)**

Geographic distribution. Worldwide (Brown and De Jonckheere 1999).

Natural host. None (environmental); humans are incidental hosts (Walochnik et al. 2000).

Route of infection. Presumed environmental exposure.

Site in human host. Eye (Walochnik et al. 2000).

***Tetramitus jugosus* (Page, 1967)**

Geographic distribution. Worldwide (Enzien et al. 1989).

Natural host. None (environmental); humans are incidental hosts (Dua et al. 1998).

Route of infection. Presumed environmental exposure (Dua et al. 1998).

Site in human host. Eye (Dua et al. 1998).

Genus *Vahlkampfia* Chatton et Lalung-Bonnaire, 1912

Geographic distribution. Worldwide (Walochnik et al. 2000).

Natural host. None (environmental); humans are incidental hosts (Walochnik et al. 2000).

Route of infection. Unknown; presumed environmental exposure.

Site in human host. Eye (Walochnik et al. 2000).

Notes. Most reports of *Vahlkampfia* from human clinical specimens are not characterized at the species level and may represent species currently assigned to other genera such as *Tetramitus*, *Allovahlkampfia*, and *Paravahlkampfia*.

●●●● Acrasidae Poche, 1913

Genus *Allovahlkampfia* Walochnik & Mulec, 2009

Allovahlkampfia spelaea Walochnik & Mulec, 2009

Geographic distribution. Not fully known; described from caves in Slovenia with single human case from Egypt (Walochnik 2009; Tolba et al. 2016).

Natural host. None (environmental); humans are incidental hosts (Tolba et al. 2016).

Route of infection. Unknown; presumed environmental exposure.

Site in human host. Eye (Tolba et al. 2016).

●● Euglenozoa Cavalier-Smith, 1981, emend. Simpson, 1997

●●● Kinetoplastea Honigberg, 1963

●●●● Metakinetoplastina Vickerman in Moreira et al., 2004

●●●●● Trypanosomatida Kent 1880, emend. Vickerman in Moreira et al. 2004

●●●●●● Trypanosomatidae Doflein, 1901

Genus *Crithidia* Léger, 1902

Geographic distribution. Worldwide (McGhee and Cosgrove 1980).

Natural hosts. Arthropods, mainly insects. Zoonotic in humans (McGhee and Cosgrove 1980).

Route of infection. Introduction of promastigotes via the bite of infected arthropod vector (Ghobakhloo et al. 2019; Maruyama et al. 2019).

Site in human host. Skin, bone marrow (Ghobakhloo et al. 2019; Maruyama et al. 2019).

Notes. *Crithidia* isolates from human clinical specimens have yet to be characterized at the species level. To date, cases are known from Brazil (Maruyama et al. 2019) and Iran (Ghobakhloo et al. 2019). The vectors in the human cases are also unknown.

Genus *Endotrypanum* Mesnil & Brimont, 1908***Endotrypanum colombiensis* (Kreuter et al., 1991)**

Geographic distribution. Colombia (Kreutzer et al. 1991).

Natural hosts. Mammalian hosts are sloths; zoonotic in humans. Arthropod vectors are sand flies in the genus *Lutzomyia* (Kreutzer et al. 1991).

Route of infection. Introduction of promastigotes via the bite of infected *Lutzomyia* sand fly (Kreutzer et al. 1991).

Site in human host. Skin (Kreutzer et al. 1991)

***Endotrypanum equatoriensis* (Grimaldi-Junior et al., 1992)**

Geographic distribution. Colombia, Ecuador (Grimaldi Junior et al. 1992).

Natural hosts. Mammalian hosts are sloths and squirrels; zoonotic in humans. Arthropod vectors are sand flies in the genus *Lutzomyia* (Grimaldi Junior et al. 1992).

Route of infection. Introduction of promastigotes via the bite of infected *Lutzomyia* sand fly (Grimaldi Junior et al. 1992).

Site in human host. Skin (Grimaldi Junior et al. 1992).

Genus *Leishmania* Ross, 1903***Euleishmania* Cupolillo et al., 2000****Subgenus *Leishmania* Ross, 1903*****Leishmaniai* (*L.*) *aethiopica* Bray et al., 1973**

Geographic distribution. Africa (Ethiopia, Kenya, Uganda), Yemen (van Henten et al. 2018).

Natural hosts. Mammalian hosts include humans, rock hyraxes, and other wild mammals. Arthropod vectors are sand flies in the genus *Phlebotomus*, mainly *P. longipes* and *P. pedifer* (Bray et al. 1973).

Route of infection. Introduction of promastigotes via the bite of infected *Phlebotomus* sand fly (van Henten et al. 2018).

Site in human host. Skin (van Henten et al. 2018).

Leishmania* (*L.*) *amazonensis* Lainson & Shaw, 1972**Leishmania garnhami* Scorza et al., 1978**

Geographic distribution. Amazonian South America (Brazil, Venezuela, Bolivia) (Torres-Guerrero et al. 2017).

Natural hosts. Mammals and birds, including humans, rodents, dogs, chickens. Arthropod vectors are sand flies in the genus *Lutzomyia* (Torres-Guerrero et al. 2017; Valdivia et al. 2017).

Route of infection. Introduction of promastigotes via the bite of infected *Lutzomyia* sand fly (Torres-Guerrero et al. 2017; Valdivia et al. 2017).

Site in human host. Skin, mucosal membranes of nose, throat, and mouth (Torres-Guerrero et al. 2017).

***Leishmania (L.) donovani* (Laveran & Mesnil, 1903)**

Leishmania archibaldi Castellani & Chalmers, 1919

Geographic distribution. Indian subcontinent, China, Central Africa; hot spots of endemicity include India, Bangladesh, Nepal, Sudan (Lukes et al. 2007; Ready 2014).

Natural hosts. Mammalian hosts include humans, dogs, foxes, marsupials, and rodents. Arthropod vectors are sand flies in the genera *Phlebotomus* (Old World) and *Lutzomyia* (New World) (Lukes et al. 2007; Ready 2014).

Route of infection. Introduction of promastigotes via the bite of infected sand fly (Torres-Guerrero et al. 2017).

Site in human host. Skin, spleen, liver, small intestine, lymph nodes (Torres-Guerrero et al. 2017).

***Leishmania (L.) infantum* Nicolle, 1908**

Leishmania chagasi da Cunha & Chagas, 1937

Geographic distribution. Mediterranean Europe and Africa, southeastern Europe, Middle East, Central Asia, Central and South America (Mexico, Venezuela, Brazil, Bolivia) (Pratlong et al. 2013; Steverding 2017).

Natural hosts. Mammalian hosts include dogs, cats, foxes, lagomorphs, rodents, marsupials, and others; zoonotic in humans. Arthropod vectors are sand flies in the genera *Phlebotomus* (Old World) and *Lutzomyia* (New World) (Millan et al. 2014).

Route of infection. Introduction of promastigotes via the bite of infected sand fly (Torres-Guerrero et al. 2017).

Site in human host. Skin, spleen, liver, small intestine, lymph nodes (Torres-Guerrero et al. 2017).

***Leishmania (L.) major* Yakimoff & Schokhor, 1914**

Geographic distribution. Northern and Central Africa, Middle East, India, China (Guerrant et al. 2006; Steverding 2017).

Natural hosts. Humans; gerbils and birds may serve as reservoir hosts; dogs can become infected but are not adequate reservoir hosts for human disease. Arthropod vectors are sand flies in the genus *Phlebotomus* (Al-Bajalan et al. 2018).

Route of infection. Introduction of promastigotes via the bite of infected *Phlebotomus* sand fly (Torres-Guerrero et al. 2017).

Site in human host. Skin (Torres-Guerrero et al. 2017).

***Leishmania (L.) mexicana* Biagi, 1953, emend. Garnham, 1962**

Leishmania pifanoi Medina & Romero, 1962

Geographic distribution. Texas, Mexico, Belize, Guatemala, Brazil, Ecuador, Peru, Venezuela (Steverding 2017).

Natural hosts. Mammalian hosts include humans, dogs, rodents, and bats. Arthropod vectors are sand flies in the genus *Lutzomyia* (Berzunza-Cruz et al. 2015).

Route of infection. Introduction of promastigotes via the bite of infected sand fly (Steverding 2017; Torres-Guerrero et al. 2017).

Site in human host. Skin (Steverding 2017; Torres-Guerrero et al. 2017).

***Leishmania (L.) tropica* Wright, 1903**

Geographic distribution. North and Central Africa, Middle East, Central Asia, India (Steverding 2017).

Natural hosts. Mammalian hosts include humans, rock hyraxes, gundi, and dogs (although dogs are not an adequate reservoir host for human disease). Arthropod vectors are sand flies in the genus *Phlebotomus* (Ajaoud et al. 2013).

Route of infection. Introduction of promastigotes via the bite of infected sand fly (Steverding 2017; Torres-Guerrero et al. 2017).

Site in human host. Skin (Steverding 2017; Torres-Guerrero et al. 2017).

***Leishmania (L.) venezuelensis* (Bonfante-Garrido, 1980)**

Geographic distribution. Venezuela, northern South America (Steverding 2017).

Natural hosts. Mammalian hosts include humans, cats. Arthropod vectors are sand flies in the genus *Lutzomyia* (Rivas et al. 2018).

Route of infection. Introduction of promastigotes via the bite of infected *Lutzomyia* sand fly (Steverding 2017; Torres-Guerrero et al. 2017).

Site in human host. Skin (Steverding 2017; Torres-Guerrero et al. 2017).

***Leishmania (L.) waltoni* Shaw et al., 2015**

Geographic distribution. Dominican Republic (Shaw et al. 2015).

Natural hosts. Mammalian hosts unknown, rats may serve as reservoir hosts; presumed zoonotic in humans. Arthropod vectors are sand flies in the genus *Lutzomyia* (Shaw et al. 2015).

Route of infection. Introduction of promastigotes via the bite of infected *Lutzomyia* sand fly (Shaw et al. 2015).

Site in human host. Skin (Shaw et al. 2015).

Subgenus *Mundinia* Shaw et al. in Espinosa et al., 2016

Leishmania (M.) martiniquensis Desbois et al., 2014

Geographic distribution. Martinique, Thailand (Desbois et al. 2014; Pothirat et al. 2014).

Natural hosts. Mammalian hosts are humans. Rats may serve as reservoir hosts. Arthropod vectors are sand flies, possibly in the genera *Sergentomyia* (Old World) and *Lutzomyia* (New World) (Leelayoova et al. 2017).

Route of infection. Introduction of promastigotes via the bite of infected sand fly

Site in human host. Skin, mucosal membranes of nose, throat, and mouth (Desbois et al. 2014; Liautaud et al. 2015).

Leishmania (M.) orientalis Jariyapan et al., 2018

Leishmania siamensis Sukmee et al., 2008

Geographic distribution. Thailand (Sukmee et al. 2008; Jariyapan et al. 2018).

Natural hosts. Mammalian hosts are humans. Arthropod vectors are sand flies. (Sukmee et al. 2008; Jariyapan et al. 2018).

Route of infection. Introduction of promastigotes via the bite of infected sand fly (genus unknown).

Site in human host. Skin (Sukmee et al. 2008; Jariyapan et al. 2018).

Subgenus *Sauroleishmania* Ranque, 1973

Leishmania (S.) tarentolae Weynon, 1921

Geographic distribution. North Africa, southern Europe, Middle East (Klatt et al. 2019).

Natural hosts. Definitive hosts are lizards. Arthropod vectors are sand flies in the genus *Sergentomyia*. Zoonotic in humans (Klatt et al. 2019).

Route of infection. Introduction of promastigotes in bite of infected *Sergentomyia* sand fly (Klatt et al. 2019; Pombi et al. 2020).

Site in human host. Blood (Pombi et al. 2020).

Notes. The single human case was diagnosed by molecular detection (only) in blood (Pombi et al. 2020).

Subgenus *Viannia* Lainson & Shaw, 1987

Leishmania (V.) braziliensis Vianna, 1911

Geographic distribution. Amazonian South America (Brazil, Bolivia, Venezuela, Peru), Guatemala (Castro et al. 2007; Steverding 2017).

Natural hosts. Mammalian hosts include humans, dogs, rodents, and marsupials. Arthropod vectors are sand flies in the genus *Lutzomyia* (Castro et al. 2007).

Route of infection. Introduction of promastigotes via the bite of infected *Lutzomyia* sand fly (Steverding 2017; Torres-Guerrero et al. 2017).

Site in human host. Skin, mucosal membranes of nose, throat, and mouth (Steverding 2017; Torres-Guerrero et al. 2017).

***Leishmania (V.) guyanensis* Floch, 1954**

Geographic distribution. Northern South America (French Guiana, Suriname, Brazil, Bolivia) (Guerra et al. 2011; Steverding 2017).

Natural hosts. Mammalian hosts include sloths, anteaters, rodents, marsupials, and humans. Arthropod vectors are sand flies in the genus *Lutzomyia* (Guerra et al. 2011).

Route of infection. Introduction of promastigotes via the bite of infected *Lutzomyia* sand fly (Steverding 2017; Torres-Guerrero et al. 2017).

Site in human host. Skin, mucosal membranes of nose, throat, and mouth (Steverding 2017; Torres-Guerrero et al. 2017).

***Leishmania (V.) lainsoni* Silveira et al., 1987**

Geographic distribution. Brazil, Bolivia, Ecuador, Peru, Suriname, French Guiana (Kato et al. 2016; Mans et al. 2017; Steverding 2017).

Natural hosts. Mammalian host is lowland paca; zoonotic in humans. Arthropod vectors are sand flies in the genus *Lutzomyia* (Mans et al. 2017).

Route of infection. Introduction of promastigotes via the bite of infected *Lutzomyia* sand fly (Steverding 2017; Torres-Guerrero et al. 2017).

Site in human host. Skin (Steverding 2017; Torres-Guerrero et al. 2017).

***Leishmania (V.) lindenbergi* Silveira et al., 2002**

Geographic distribution. Brazil (Cantanhêde et al. 2019).

Natural hosts. Mammalian hosts unknown; presumed zoonotic in humans. Arthropod vectors are sand flies in the genus *Lutzomyia* (Cantanhêde et al. 2019).

Route of infection. Introduction of promastigotes via the bite of infected *Lutzomyia* sand fly (Steverding 2017; Torres-Guerrero et al. 2017).

Site in human host. Skin (Steverding 2017; Torres-Guerrero et al. 2017).

***Leishmania (V.) naiffi* Lainson & Shaw, 1989**

Geographic distribution. Brazil, Bolivia, Peru (Lainson and Shaw 1989; Fagundes-Silva et al. 2015).

Natural hosts. Mammalian hosts are armadillos; zoonotic in humans. Arthropod vectors are sand flies in the genus *Lutzomyia* (Lainson and Shaw 1989; Fagundes-Silva et al. 2015).

Route of infection. Introduction of promastigotes via the bite of infected *Lutzomyia* sand fly (Steverding 2017; Torres-Guerrero et al. 2017).

Site in human host. Skin (Steverding 2017; Torres-Guerrero et al. 2017).

Leishmania (V.) panamensis Lainson & Shaw, 1972

Geographic distribution. Central and South America (Steverding 2017; Torres-Guerrero et al. 2017).

Natural hosts. Mammalian hosts include sloths, procyonids, rodents, humans, and non-human primates. Arthropod vectors are sand flies in the genus *Lutzomyia* (Steverding 2017; Torres-Guerrero et al. 2017).

Route of infection. Introduction of promastigotes via the bite of infected *Lutzomyia* sand fly (Steverding 2017; Torres-Guerrero et al. 2017).

Site in human host. Skin, mucosal membranes of nose, throat, and mouth (Steverding 2017; Torres-Guerrero et al. 2017).

Leishmania (V.) peruviana Velez, 1913

Geographic distribution. Peru, Bolivia (Steverding 2017; Torres-Guerrero et al. 2017).

Natural hosts. Mammalian hosts include humans, rodents, marsupials, and domestic dogs. Arthropod vectors are sand flies in the genera *Lutzomyia* and possibly *Pintomyia* (Kato et al. 2021).

Route of infection. Introduction of promastigotes via the bite of infected sand fly (Steverding 2017; Torres-Guerrero et al. 2017).

Site in human host. Skin, mucosal membranes of nose, throat, and mouth (Steverding 2017; Torres-Guerrero et al. 2017).

Leishmania (V.) shawi Lainson et al., 1989

Geographic distribution. Brazil (Lainson et al. 1989)

Natural hosts. Mammalian hosts include non-human primates, sloths, coati; zoonotic in humans. Arthropod vectors are sand flies in the genus *Lutzomyia* (Lainson et al. 1989).

Route of infection. Introduction of promastigotes via the bite of infected *Lutzomyia* sand fly (Steverding 2017; Torres-Guerrero et al. 2017).

Site in human host. Skin (Steverding 2017; Torres-Guerrero et al. 2017).

Genus *Leptomonas* Kent, 1880

Leptomonas seymouri (Wallace, 1877)

Geographic distribution. Worldwide (Kraeva et al. 2015).

Natural hosts. Insects. Zoonotic in humans (Ghosh et al. 2012; Kraeva et al. 2015; Thakur et al. 2020).

Route of infection. Presumed introduction of promastigotes via the bite of infected *Phlebotomus* sand flies (Ghosh et al. 2012; Kraeva et al. 2015; Thakur et al. 2020).

Site in human host. Skin, spleen, bone marrow (Ghosh et al. 2012; Kraeva et al. 2015; Thakur et al. 2020)

Notes. Human cases of *L. seymouri* to date have been found in Southeast Asia associated with co-infection with *Leishmania donovani*, suggesting incidental human infection after the bite of an infected sand fly vector.

Genus *Trypanosoma* Gruby, 1843

Subgenus *Herpetosoma* Doflein, 1901

Trypanosoma (H.) lewisi Laveran & Mesnil, 1901

Geographic distribution. Cosmopolitan (Desquesnes et al. 2002).

Natural hosts. Mammalian hosts are rats (*Rattus* spp.); zoonotic in humans. Arthropod vectors are rat fleas *Nosophyllus fasciatus* and *Xenopsylla cheopis* (Desquesnes et al. 2002).

Route of infection. Ingestion of infected fleas or their feces (Desquesnes et al. 2002).

Site in human host. Blood (Verma et al. 2011; de Sousa 2014).

Trypanosoma (H.) rangeli Tejera, 1920

Trypanosoma saimirii Rodhain, 1941

Geographic distribution. South America (Guhl and Vallejo 2003).

Natural hosts. Mammalian hosts include coati, sloths, tamandua, and opossums; zoonotic in humans. Arthropod vectors are triatomine bugs, most-notably members of the genus *Rhodnius* (Miles et al. 1983; Guhl and Vallejo 2003; Ferreira L de et al. 2015).

Route of infection. Introduction of trypomastigotes via the bite of infected triatomine bug (Guhl and Vallejo 2003; Ferreira L de et al. 2015).

Site in human host. Blood (Guhl and Vallejo 2003; Ferreira L de et al. 2015).

Subgenus *Schizotrypanum* Chagas, 1909

Trypanosoma (S.) cruzi Chagas, 1909

Geographic distribution. Southern United States, Central and South America (Bern et al. 2011; Kirchhoff 2011; Connors et al. 2016).

Natural hosts. Mammalian hosts include humans and many domestic and wild mammals. Arthropod vectors are triatomine bugs, most-notably members of the genera *Triatoma*, *Panstrongylus*, and *Rhodnius* (Kirchhoff 2011).

Route of infection. Introduction of metacyclic trypomastigotes in the feces of infected triatomine bug when rubbed into wounds or mucus membranes; ingestion of food contaminated with bugs or their feces; transplacentally from mother to fetus; via blood transfusion and organ transplantation (Coura 2015; Conners et al. 2016; Lidani et al. 2019).

Site in human host. Blood (acute infection, reactivation); heart muscle, esophagus, large intestine, peripheral nervous system, CNS, other organs (chronic infection) (Coura 2015; Conners et al. 2016; Lidani et al. 2019).

Subgenus *Trypanozoon* Lühe, 1906

***Trypanosoma (T.) brucei gambiense* (Dutton, 1902)**

Geographic distribution. West-central Africa; hot spots of endemicity include Angola, Democratic Republic of Congo, South Sudan, Central African Republic, Uganda (Brun et al. 2010; Franco et al. 2014).

Natural hosts. Mammalian hosts include humans, non-human primates, and ungulates. Arthropod vectors are tsetse flies in the genus *Glossina* (Brun et al. 2010; Franco et al. 2014).

Route of infection. Introduction of trypomastigotes via the bite of infected *Glossina* tsetse fly (Brun et al. 2010).

Site in human host. Blood, lymphatics, CNS (Brun et al. 2010).

***Trypanosoma (T.) brucei rhodesiense* (Stephens & Fantham, 1910)**

Geographic distribution. East and southeast Africa; hot spots of endemicity include Malawi, Zambia, Tanzania, Botswana, Kenya (Brun et al. 2010; Franco et al. 2014).

Natural hosts. Mammalian hosts include livestock and wild ungulates; zoonotic in humans. Arthropod vectors are tsetse flies in the genus *Glossina* (Brun et al. 2010; Franco et al. 2014).

Route of infection. Introduction of trypomastigotes via the bite of infected *Glossina* tsetse fly (Brun et al. 2010).

Site in human host. Blood, lymphatics, CNS (Brun et al. 2010).

***Trypanosoma (T.) evansi* Balbiani, 1888**

Geographic distribution. Equatorial regions of Africa, Asia, Latin America (Van Vinh Chau et al. 2016).

Natural hosts. Mammalian hosts include horses, camels, bovines; zoonotic in humans. Arthropod vectors are biting flies in the genera *Stomoxys* and *Tabanus* (Van Vinh Chau et al. 2016).

Route of infection. Introduction of trypomastigotes via the bite of infected flies; possibly direct contact with blood/wounds of infected animals (Joshi et al. 2005; Powar et al. 2006; Van Vinh Chau et al. 2016).

Site in human host. Blood (Joshi et al. 2005; Powar et al. 2006; Van Vinh Chau et al. 2016).

Opisthokonta Cavalier-Smith, 1987, emend. Adl et al., 2005

- Holozoa Lang et al., 2002
- Metazoa Haeckel, 1874, emend. Adl et al. 2005
- Protostomia Grobden, 1908
- Spiralia Edgecombe et al., 2011
- Acanthocephala Koelreuther, 1771
- Oligacanthorhynchida Petrotschenko, 1956
- Oligacanthorhynchidae Southwell & Macfie, 1925

Genus *Macracanthorhynchus* Travassos, 1917

Macracanthorhynchus hirudinaceus (Pallas, 1781)

Taenia haeruca Pallas, 1776

Echinorhynchus gigas Bloch, 1782

Geographic distribution. Worldwide (Mathison et al. 2016).

Natural hosts. Intermediate hosts are insects, primarily scarabaeoid beetles. Definitive hosts are pigs; dogs and other mammals can serve as reservoir hosts. Zoonotic in humans (Mathison et al. 2016).

Route of infection. Ingestion of cystacanths in infected insect intermediate host (Mathison et al. 2016).

Site in human host. Small intestine (Mathison et al. 2016).

Macracanthorhynchus ingens (von Linstow, 1879)

Geographic distribution. Eastern North America (Elkins and Nickol 1983; Richardson et al. 2017).

Natural hosts. Intermediate hosts are millipedes. Definitive hosts are raccoons (*Procyon lotor*) and American black bear (*Ursus americanus*); many other mammals, especially carnivores, can serve as definitive hosts. Snakes and other animals may serve as paratenic hosts. Zoonotic in humans (Elkins and Nickol 1983; Richardson et al. 2017).

Route of infection. Ingestion of cystacanths in infected arthropod intermediate host or paratenic hosts (Mathison et al. 2016; Chancey et al. 2021).

Site in human host. Small intestine (Mathison et al. 2016; Chancey et al. 2021).

●●●●●● **Moniliformida** Schmidt, 1972

●●●●●● **Moniliformidae** Van Cleave, 1924

Genus *Moniliformis* Travassos, 1915

Moniliformis moniliformis (Bremser, 1811)

Echinorhynchus grassi Railliet, 1893

Echinorhynchus canis Porta, 1914

Echinorhynchus belgicus Railliet, 1918

Moniliformis moniliformis siciliensis Meyer, 1932

Moniliformis moniliformis aegypticus Meyer, 1932

Moniliformis dubius Meyer, 1932

Geographic distribution. Worldwide (Mathison et al. 2016).

Natural hosts. Intermediate hosts are insects, primarily beetles and cockroaches. Definitive hosts are primarily rodents and carnivores. Zoonotic in humans (Mathison et al. 2016).

Route of infection. Ingestion of cystacanths in infected insect intermediate host (Mathison et al. 2016).

Site in human host. Small intestine (Mathison et al. 2016).

●●●●●● **Echinorhynchida** Petrotschenko, 1956

●●●●●● **Echinorhynchidae** Cobbold, 1876

Genus *Acanthocephalus* Koelreuther, 1771

Acanthocephalus rauschi (Schmidt, 1969)

Geographic distribution. Alaska (Schmidt 1969).

Natural hosts. Alaskan grayling (*Thymallus arcticus*). Otters can serve as paratenic hosts. Zoonotic in humans (Schmidt 1969).

Route of infection. Unknown; presumed ingestion of cystacanths in infected intermediate host.

Site in human host. Peritoneum (Schmidt 1969, 1971).

Genus *Pseudoacanthocephalus* Petrotschenko, 1958

Pseudoacanthocephalus bufonis (Shiple, 1903)

Acanthocephalus sinensis Van Cleave, 1937

Geographic distribution. Southeast Asia, Hawaii (Barton and Pichelin 1999; Bush et al. 2009).

Natural hosts. Toads. Zoonotic in humans (Bush et al. 2009).

Route of infection. Unknown; presumed ingestion of cystacanths in infected intermediate host.

Site in human host. Small intestine (Schmidt 1971).

●●●●●● **Polymorphida Petrotschenko, 1956**

●●●●●● **Polymorphidae Meyer, 1931**

Genus *Bolbosoma* Porta, 1908

***Bolbosoma nipponicum* Yamaguti, 1939**

Geographic distribution. North Atlantic (Yamaguti 1963).

Natural hosts. Marine crustaceans serve as intermediate hosts. Fish and cephalopods may serve as paratenic hosts. Whales and seals serve as definitive hosts. Zoonotic in humans (Yamaguti 1963).

Route of infection. Unknown; presumed ingestion of cystacanths in infected fish paratenic hosts.

Site in human host. Small intestine (Yamamoto et al. 2018).

***Bolbosoma* sp. cf. *capitatum* (von Linstow, 1880)**

Geographic distribution. Worldwide (Amin and Margolis 1998).

Natural hosts. Marine crustaceans serve as intermediate hosts. Whales serve as definitive hosts. Zoonotic in humans (Amin and Margolis 1998).

Route of infection. Unknown; presumed ingestion of cystacanths in infected fish paratenic hosts.

Site in human host. Small intestine (Arizono et al. 2012).

Notes. The single case was reported from Japan as *Bolbosoma* cf. *capitatum* based on molecular and histological examination (Arizono et al. 2012).

Genus *Corynosoma* Lühe, 1904

***Corynosoma strumosum* (Rudolphi, 1802)**

Geographic distribution. Worldwide (Aznar et al. 2006; Leidenberger et al. 2020).

Natural hosts. Intermediate hosts are marine amphipods. Several groups of fish may serve as paratenic hosts. Definitive hosts are seals, walrus, beluga whale. Zoonotic in humans (Aznar et al. 2006; Leidenberger et al. 2020; Nickol et al. 2002).

Route of infection. Ingestion of cystacanths in infected fish paratenic host (Schmidt 1971).

Site in human host. Small intestine (Schmidt 1971).

***Corynosoma villosum* Van Cleave, 1953**

Geographic distribution. North Pacific (Moles 1982).

Natural hosts. Intermediate hosts are marine amphipods. Several groups of fish may serve as paratenic hosts. Definitive hosts are pinnipeds and whales. Zoonotic in humans (Moles 1982).

Route of infection. Ingestion of cystacanths in infected fish paratenic host (Fujita et al. 2016).

Site in human host. Small intestine (Fujita et al. 2016).

***Corynosoma* sp. cf. *validum* Van Cleave, 1953**

Geographic distribution. North Pacific (Ionita et al. 2008).

Natural hosts. Intermediate hosts are marine amphipods. Several groups of fish may serve as paratenic hosts. Definitive host is the northern fur seal (*Callorhinus ursinus*), and other pinnipeds. Zoonotic in humans (Ionita et al. 2008).

Route of infection. Ingestion of cystacanths in infected fish paratenic host (Takahashi et al. 2016).

Site in human host. Small intestine (Takahashi et al. 2016).

Notes. The single case was reported from Japan as *Corynosoma* cf. *validum* based on DNA sequencing (Takahashi et al. 2016).

●●●●● Annelida Lamarck, 1802

●●●●●● Hirudinea Lamarck, 1818

●●●●●●● Rhynchobdellida Blanchard, 1894

●●●●●●●● Glossiphoniidae Vaillant, 1890

Genus *Haementeria* de Filippi, 1849***Haementeria acuecuyetzin* Ocegüera-Figueroa, 2008**

Geographic distribution. Mexico (Ocegüera-Figueroa 2008).

Natural hosts. Various vertebrates. Zoonotic on humans (Ocegüera-Figueroa 2008; Charruau et al. 2020).

Route of infection. Exposure to fresh water (Ocegüera-Figueroa 2008; Charruau et al. 2020).

Site in human host. Skin (Ocegüera-Figueroa 2008).

Vectored pathogens. None.

***Haementeria ghilianii* de Filippi, 1849**

Geographic distribution. Amazon South America (Sawyer et al. 1981).

Natural hosts. Various mammals. Zoonotic on humans (Sawyer et al. 1981; Farrar 2014).

Route of infection. Exposure to fresh water (Farrar 2014).

Site in human host. Skin (Farrar 2014).

Vectored pathogens. None.

Genus *Parabdella* Autrum, 1936

***Parabdella quadrioculata* (Moore, 1930)**

Geographic distribution. East Asia (Moore 1930).

Natural hosts. Freshwater turtles and frogs. Zoonotic on humans (Moore 1930).

Route of infection. Exposure to fresh water (Yamauchi et al. 2013).

Site in human host. Skin (Yamauchi et al. 2013).

Vectored pathogens. None.

Genus *Placobdella* Blanchard, 1893

***Placobdella costata* (Müller, 1846)**

Placobdella catenigera Moquin-Tandon, 1846

Geographic distribution. Mediterranean Region, Central Europe (Wilkialis 1973).

Natural hosts. Freshwater turtles. Zoonotic on humans (Wilkialis 1973).

Route of infection. Exposure to fresh water (Aloto 2018).

Site in human host. Skin (Aloto 2018).

Vectored pathogens. None.

●●●●●●● **Hirudiniformes Caballero, 1952**

●●●●●●● **Xerobdellidae Moore, 1946**

Genus *Diestecostoma* Vaillant, 1890

Heterobdella Baird, 1869

Hygrobdella Caballero, 1940

***Diestecostoma mexicana* (Baird, 1869)**

Geographic distribution. Central America (Oceguera-Figueroa and León-Règagnon 2014).

Natural hosts. Various mammals. Zoonotic on humans (Farrar 2014).

Route of infection. Environmental exposure (Farrar 2014).

Site in human host. Skin (Farrar 2014).

Vectored pathogens. None.

●●●●●●●● **Haemadipsidae Blanchard, 1896**

Genus *Haemadipsa* Tennent, 1859

***Haemadipsa hainana* Song, 1977**

Geographic distribution. Southeast Asia (Tessler et al. 2018).

Natural hosts. Various mammals and birds. Zoonotic on humans (Tessler et al. 2018).

Route of infection. Environmental exposure (Tessler et al. 2018).

Site in human host. Skin (Tessler et al. 2018).

Vectored pathogens. None.

***Haemadipsa interrupta* Moore, 1935**

Geographic distribution. Southeast Asia (Tessler et al. 2018).

Natural hosts. Various mammals and birds. Zoonotic on humans (Tessler et al. 2018).

Route of infection. Environmental exposure (Tessler et al. 2018).

Site in human host. Skin (Tessler et al. 2018).

Vectored pathogens. None.

***Haemadipsa japonica* Whitman, 1886**

Geographic distribution. East Asia (Aizawa and Morishima 2018).

Natural hosts. Various mammals, birds, amphibians. Zoonotic on humans (Hanya et al. 2019).

Route of infection. Environmental exposure (Ji-Tuan 1997).

Site in human host. Skin, mucous membranes or the rectum and genital tract (Ji-Tuan 1997).

Vectored pathogens. None.

***Haemadipsa picta* Moore, 1929**

Geographic distribution. Southeast Asia (Lai et al. 2011).

Natural hosts. Various large mammals, including humans (Lai et al. 2011).

Route of infection. Environmental exposure (Fogden and Proctor 1985; Lai et al. 2011).

Site in human host. Skin (Fogden and Proctor 1985; Lai et al. 2011).

Vectored pathogens. None.

***Haemadipsa rjukjuana* Oka, 1910**

Geographic distribution. East Asia (Lai et al. 2011; Won et al. 2014).

Natural hosts. Various mammals, including humans, and birds (Lai et al. 2011).

Route of infection. Environmental exposure (Lai et al. 2011; Won et al. 2014).

Site in human host. Skin (Lai et al. 2011; Won et al. 2014).

Vectored pathogens. None

***Haemadipsa sylvestris* Blanchard, 1894**

Geographic distribution. Southeast Asia (Farrar 2014).

Natural hosts. Various mammals. Zoonotic on humans (Farrar 2014).

Route of infection. Environmental exposure (Farrar 2014).

Site in human host. Skin (Farrar 2014).

Vectored pathogens. None.

***Haemadipsa trimaculosa* Ngamprasertwong et al., 2007**

Geographic distribution. Southeast Asia (Ngamprasertwong et al. 2007; Tessler et al. 2018).

Natural hosts. Various mammals and birds. Zoonotic on humans (Ngamprasertwong et al. 2007).

Route of infection. Environmental exposure (Tessler et al. 2018).

Site in human host. Skin (Tessler et al. 2018).

Vectored pathogens. None

***Haemadipsa zeylanica* (Moquin-Tandon, 1826)**

Geographic distribution. Japan, Southeast Asia (Fogden and Proctor 1985).

Natural hosts. Various amphibians and mammals. Zoonotic on humans (Fogden and Proctor 1985).

Route of infection. Environmental exposure (Fogden and Proctor 1985).

Site in human host. Skin (Fogden and Proctor 1985).

Vectored pathogens. None.

Genus *Phytobdella* Blanchard, 1894

***Phytobdella catenifera* Moore, 1942**

Geographic distribution. Malaysia (Farrar 2014).

Natural hosts. Reptiles. Zoonotic on humans (Farrar 2014).

Route of infection. Environmental exposure (Farrar 2014).

Site in human host. Skin (Farrar 2014).

Vectored pathogens. None.

●●●●●●●● **Hirudinidae Whitman, 1886**

Genus *Hirudo* Linnaeus, 1758

***Hirudo medicinalis* Linnaeus, 1758**

Geographic distribution. Europe, Asia (Kutschera and Elliott 2014).

Natural hosts. Immature stages usually on amphibians. Adults on mammals. Zoonotic on humans (Kutschera and Elliott 2014).

Route of infection. Exposure to fresh water (Kutschera and Elliott 2014).

Site in human host. Skin (Kutschera and Elliott 2014).

Vectored pathogens. *Aeromonas* spp. (Graf 1999).

***Hirudo nipponia* Whitman, 1886**

Geographic distribution. Russia, Japan, Southeast Asia (Yang 1996).

Natural hosts. Immature stages usually on amphibians. Adults on mammals. Zoonotic on humans (Yang 1996).

Route of infection. Exposure to fresh water (Oda et al. 1984).

Site in human host. Skin, Eye (Oda et al. 1984).

Vectored pathogens. None.

***Hirudo orientalis* Utevsky & Trontelj, 2005**

Geographic distribution. Central Asia, Eastern Europe (Utevsky and Trontelj 2005).

Natural hosts. Immature stages usually on amphibians. Adults on mammals. Zoonotic on humans (Utevsky and Trontelj 2005).

Route of infection. Exposure to fresh water (Utevsky and Trontelj 2005).

Site in human host. Skin, Eye (Utevsky and Trontelj 2005).

Vectored pathogens. *Aeromonas* spp. (Laufer et al. 2008).

***Hirudo troctina* Johnson, 1816**

Geographic distribution. North Africa, Iberian Peninsula (Marrone and Canale 2019).

Natural hosts. Immature stages usually on amphibians. Adults on mammals. Zoonotic on humans (Trontelj and Utevsky 2005).

Route of infection. Exposure to fresh water (Trontelj and Utevsky 2005).

Site in human host. Skin, Eye (Trontelj and Utevsky 2005).

Vectored pathogens. None.

***Hirudo verbena* Carena, 1820**

Geographic distribution. Mediterranean Europe (Marrone and Canale 2019).

Natural hosts. Immature stages usually on amphibians. Adults on mammals. Zoonotic on humans (Kutschera and Elliott 2014).

Route of infection. Exposure to fresh water (Kutschera and Elliott 2014).

Site in human host. Skin (Kutschera and Elliott 2014).

Vectored pathogens. *Aeromonas* spp. (Laufer et al. 2008).

Genus *Hirudinaria* Whitman, 1886

Hirudinaria manillensis (Lesson, 1842)

Hirudo multistriata Schmarda, 1861

Hirudo luzoniae Kinberg, 1866

Hirudo maculosa Grube, 1868

Hirudo maculata Baird, 1869

Limnatis granulosa Blanchard, 1893

Hirudo boyntoni Wharton, 1913

Geographic distribution. Southeast Asia (Jeratthitikul et al. 2020).

Natural hosts. Mammals, including buffalo, cattle. Zoonotic in humans (Jeratthitikul et al. 2020).

Route of infection. Exposure to fresh water (Hii et al. 1978).

Site in human host. Upper respiratory tract (Hii et al. 1978).

Vectored pathogens. None

Genus *Limnatis* Moquin-Tandon, 1827

Limnatis nilotica (Savigny, 1822)

Geographic distribution. Southern Europe, North Africa, Middle East (Arfuso et al. 2019).

Natural hosts. Mammals, including sheep, cattle, dogs, and donkeys. Zoonotic on humans (Arfuso et al. 2019).

Route of infection. Exposure to fresh water

Site in human host. Upper respiratory tract (Almallah 1968; Ağin et al. 2008).

Vectored pathogens. None.

Genus *Poecilobdella* Blanchard, 1893

Poecilobdella granulosa (Savigny, 1822)

Geographic distribution. India, Sri Lanka, Southeast Asia (Nesemann and Sharma 2001).

Natural hosts. Amphibians, Mammals. Zoonotic in humans (Nesemann and Sharma 2001).

Route of infection. Exposure to fresh water (Zainuddin et al. 2016).

Site in human host. Skin, upper respiratory tract (Zainuddin et al. 2016).

Vectored pathogens. None.

Poecilobdella viridis Moore in Harding & Moore, 1927

Geographic distribution. India, Sri Lanka (Kulkarni et al. 1977).

Natural hosts. Amphibians, Mammals. Zoonotic in humans (Kulkarni et al. 1977).

Route of infection. Exposure to fresh water (Zainuddin et al. 2016).

Site in human host. Upper respiratory tract (Zainuddin et al. 2016).

Vectored pathogens. None.

●●●●●●●● Macrobdellidae Richardson, 1969

Genus *Macrobdella* Verrill, 1872

Macrobdella decora (Say, 1824)

Geographic distribution. Eastern North America, Mexico (Phillips et al. 2016, 2019).

Natural hosts. Various vertebrate animals. Zoonotic on humans (Phillips et al. 2016, 2019).

Route of infection. Exposure to fresh water (Phillips et al. 2016, 2019).

Site in human host. Skin (Phillips et al. 2016, 2019).

Vectored pathogens. None.

Macrobdella diploertia Meyer, 1975

Geographic distribution. East-central United States (Phillips et al. 2016, 2019).

Natural hosts. Various vertebrate animals. Zoonotic on humans (Phillips et al. 2016, 2019).

Route of infection. Exposure to fresh water (Phillips et al. 2016, 2019).

Site in human host. Skin (Phillips et al. 2016, 2019).

Vectored pathogens. None.

Macrobdella ditetra Moore, 1953

Geographic distribution. Southeastern United States (Phillips et al. 2016, 2019).

Natural hosts. Various vertebrate animals. Zoonotic on humans (Phillips et al. 2016, 2019).

Route of infection. Exposure to fresh water (Phillips et al. 2016, 2019).

Site in human host. Skin (Phillips et al. 2016, 2019).

Vectored pathogens. None.

***Macrobodella mimicus* Phillips, 2019**

Geographic distribution. Northeastern United States (Phillips et al. 2019).

Natural hosts. Various vertebrate animals. Zoonotic on humans (Phillips et al. 2019).

Route of infection. Exposure to fresh water (Phillips et al. 2019).

Site in human host. Skin (Phillips et al. 2019).

Vectored pathogens. None.

***Macrobodella sestertia* Whitman, 1886**

Geographic distribution. Northeastern United States (Phillips et al. 2016, 2019).

Natural hosts. Various vertebrate animals. Zoonotic on humans (Phillips et al. 2016, 2019).

Route of infection. Exposure to fresh water (Phillips et al. 2016, 2019).

Site in human host. Skin (Phillips et al. 2016, 2019).

Vectored pathogens. None.

●●●●●●●● Praeobdellidae Sawyer, 1986**Genus *Dinobdella* Moore in Harding & Moore, 1927*****Dinobdella ferox* (Blanchard, 1896)**

Geographic distribution. India, Southeast Asia (Lai 2019).

Natural hosts. Various mammals. Zoonotic on humans (Lai 2019).

Route of infection. Exposure to fresh water (Makiya et al. 1988).

Site in human host. Nasal mucosa (Makiya et al. 1988).

Vectored pathogens. None.

Genus *Myxobdella* Oka, 1917***Myxobdella africana* Moore, 1939**

Geographic distribution. Sub-Saharan Africa (Cundall et al. 1986; Estambale et al. 1992).

Natural hosts. Mammals, including dogs. Zoonotic in humans (Cundall et al. 1986; Estambale et al. 1992).

Route of infection. Exposure to fresh water (Cundall et al. 1986; Estambale et al. 1992).

Site in human host. Upper respiratory tract (Cundall et al. 1986; Estambale et al. 1992).

Vectored pathogens. None.

Genus *Tyrannobdella* Philipps et al., 2010

Tyrannobdella rex Phillips et al., 2010

Geographic distribution. Amazon South America (Phillips et al. 2010).

Natural hosts. Unknown; presumed zoonotic on humans (Phillips et al. 2010).

Route of infection. Exposure to fresh water (Phillips et al. 2010).

Site in human host. Nasopharynx (Phillips et al. 2010).

Vectored pathogens. None.

●●●●● Cestoda Rudolphi, 1809

●●●●●● Eucestoda Southwell, 1930

●●●●●●● Diphyllobothriidea Kuchta et al., 2008

●●●●●●●● Diphyllobothriidae Lühe, 1910

Genus *Adenocephalus* Nybelin, 1931

Adenocephalus pacificus Nybelin, 1931

Adenocephalus septentrionalis Nybelin, 1931

Diphyllobothrium arctocephali Drummond, 1937

Diphyllobothrium krotovi Delyamure, 1955

Dibothriocephalus atlanticum Delyamure & Parukhin, 1968

Geographic distribution. Coastal Pacific waters and southern Africa (Hernandez-Orts et al. 2015; Kuchta et al. 2015).

Natural hosts. The first arthropod intermediate host is unknown, but presumed to be marine copepods. The second intermediate hosts are various marine fish. Definitive hosts are eared seals and sea lions; dogs and jackals can serve as reservoir hosts. Zoonotic in humans (Hernandez-Orts et al. 2015; Kuchta et al. 2015).

Route of infection. Ingestion of plerocercoids in infected fish (Kuchta et al. 2015).

Site in human host. Small intestine (Kuchta et al. 2015).

Genus *Dibothriocephalus* Lühe, 1899

Dibothriocephalus dalliae (Rausch, 1956)

Geographic distribution. Northern Pacific (Alaska and northeastern Siberia) (Scholz and Kuchta 2016).

Natural hosts. First intermediate hosts are freshwater copepods. Second intermediate hosts are freshwater fish, primarily Alaska blackfish (*Dallia pectoralis*). Definitive hosts are Arctic fox, with domestic dogs as reservoir hosts. Zoonotic in humans (Scholz and Kuchta 2016).

Route of infection. Ingestion of plerocercoids in infected fish (Scholz and Kuchta 2016).

Site in human host. Small intestine (Scholz and Kuchta 2016).

***Dibothriocephalus dendriticus* (Nitzsch, 1824)**

Dibothriocephalus fissiceps Creplin, 1829

Dibothriocephalus cordiceps Leidy, 1872

Dibothriocephalus exile Linton, 1892

Sparganum sebago Ward, 1910

Dibothriocephalus minor Cholodkovsky, 1916

Diphyllobothrium canadense Cooper, 1921

Diphyllobothrium strictum Talysin, 1932

Diphyllobothrium obdoriense Piotnikoff, 1933

Diphyllobothrium laruei Vergeer, 1934

Diphyllobothrium nenzi Petrov, 1938

Diphyllobothrium oblongatum Thomas, 1946

Diphyllobothrium medium Fahmy, 1954

Diphyllobothrium microcordiceps Szidat & Soria, 1957

Diphyllobothrium norvegicum Vik, 1957

Geographic distribution. Holarctic (Scholz and Kuchta 2016).

Natural hosts. The first intermediate hosts are freshwater copepods. The second intermediate and paratenic hosts are many different freshwater fish. Definitive hosts are fish-eating birds, primarily gulls, and mammals, including wild and domestic canids, bears, and otters. Zoonotic in humans (Scholz and Kuchta 2016).

Route of infection. Ingestion of plerocercoids in infected fish (Kuchta et al. 2013).

Site in human host. Small intestine (Kuchta et al. 2013).

***Dibothriocephalus hians* Diesing, 1850**

Geographic distribution. Mediterranean and Circumboreal (Scholz and Kuchta 2016).

Natural hosts. The first intermediate hosts are presumed to be marine copepods. The second intermediate hosts are unknown, but presumed to be marine fish. Definitive hosts are seals. Zoonotic in humans (Scholz and Kuchta 2016).

Route of infection. Ingestion of plerocercoids in infected fish (Scholz and Kuchta 2016).

Site in human host. Small intestine (Scholz and Kuchta 2016).

***Dibothriocephalus latus* (Linnaeus, 1758)**

Diphyllobothrium americanum Hall & Wigdor, 1918

Diphyllobothrium tungussicum Podyapolskaya & Gnedina, 1932

Diphyllobothrium skrjabini Plotnikoff, 1933

Geographic distribution. Holarctic, South America (Scholz and Kuchta 2016).

Natural hosts. The first intermediate hosts are freshwater copepods. The second intermediate and paratenic hosts are freshwater fish, including perch, pike, burbot, walleye, and pikeperch. Definitive hosts are fish-eating carnivores, including wild and domestic dogs and cats, bears, and mustelids. Zoonotic in humans (Scholz and Kuchta 2016).

Route of infection. Ingestion of plerocercoids in infected fish (Scholz et al. 2009).

Site in human host. Small intestine (Scholz et al. 2009).

Dibothriocephalus nihonkaiensis (Yamane et al., 1886)*Diphyllobothrium giljadicum* Rutkevich, 1937*Diphyllobothrium klebanovskii* Muratov & Posokov, 1988

Geographic distribution. Northern Pacific (Scholz and Kuchta 2016; Ikuno et al. 2018).

Natural hosts. First intermediate hosts are marine copepods. Second intermediate hosts are salmonid fish. Definitive hosts are bears, wild canids, and mink. Zoonotic in humans.

Route of infection. Ingestion of plerocercoids in infected fish (Scholz and Kuchta 2016).

Site in human host. Small intestine (Scholz and Kuchta 2016).

Dibothriocephalus ursi (Rausch, 1954)*Diphyllobothrium gonodo* Yamaguti, 1942

Geographic distribution. Northern North America (Scholz and Kuchta 2016).

Natural hosts. First intermediate hosts are marine copepods. Second intermediate hosts are salmonid fish, primarily sockeye salmon (*Oncorhynchus nerka*). Definitive hosts are bears. Zoonotic in humans (Scholz and Kuchta 2016).

Route of infection. Ingestion of plerocercoids in infected fish (Scholz and Kuchta 2016).

Site in human host. Small intestine (Scholz and Kuchta 2016).

Genus *Diphyllobothrium* Cobbold, 1858*Diphyllobothrium balaenopterae* (Lönnerberg, 1892)*Krabbea grandis* Blanchard, 1894*Diplogonoporus fukuokaensis* Kamo & Miyazaki, 1970

Geographic distribution. Worldwide (Scholz and Kuchta 2016).

Natural hosts. First intermediate hosts are marine copepods. Second intermediate hosts are marine fish, including Japanese anchovy, Japanese sardine, and skipjack tuna. Definitive hosts are baleen whales. Zoonotic in humans (Scholz and Kuchta 2016).

Route of infection. Ingestion of plerocercoids in infected fish (Scholz and Kuchta 2016).

Site in human host. Small intestine (Scholz and Kuchta 2016).

***Diphyllobothrium stemmacephalum* Cobbold, 1858**

Diphyllobothrium ponticum Delyamure, 1971

Diphyllobothrium yonagoense Yamane et al., 1981

Geographic distribution. Northern Hemisphere (Scholz and Kuchta 2016).

Natural hosts. First intermediate hosts are presumed to be marine copepods. Second intermediate hosts are unknown, but presumed to be marine fish. Definitive hosts are dolphins and porpoises. Zoonotic in humans (Scholz and Kuchta 2016).

Route of infection. Ingestion of plerocercoids in infected fish (Scholz and Kuchta 2016).

Site in human host. Small intestine (Scholz and Kuchta 2016).

Diphyllobothrium, incertae sedis:

***Diphyllobothrium cordatum* (Leuckart, 1863)**

Geographic distribution. Circumpolar (Scholz and Kuchta 2016).

Natural hosts. First intermediate hosts are presumed to be marine copepods. Second intermediate hosts are unknown, but presumed to be marine fish. Definitive hosts are seals and walrus. Zoonotic in humans (Scholz and Kuchta 2016).

Route of infection. Ingestion of plerocercoids in infected fish (Scholz and Kuchta 2016).

Site in human host. Small intestine (Scholz and Kuchta 2016).

***Diphyllobothrium lanceolatum* (Krabbe, 1865)**

Geographic distribution. Circumpolar (Scholz and Kuchta 2016).

Natural hosts. First intermediate hosts are marine copepods. Second intermediate hosts are marine fish, including sardine cisco (*Coregonus sardinella*). Definitive hosts are seals. Zoonotic in humans (Scholz and Kuchta 2016).

Route of infection. Ingestion of plerocercoids in infected fish (Scholz and Kuchta 2016).

Site in human host. Small intestine (Scholz and Kuchta 2016).

Genus *Spirometra* Faust et al., 1929

Spirometra decepiens (Diesing, 1850)

Geographic distribution. North and South America, Southeast Asia (Jeon et al. 2015; Jeon et al. 2018b).

Natural hosts. First intermediate hosts are believed to be freshwater copepods. Second intermediate hosts are believed to be freshwater fish and/or amphibians. Definitive hosts are wild and domestic cats; dogs may serve as reservoir hosts. Zoonotic in humans as dead-end hosts that harbor plerocercoid larvae (Jeon et al. 2015; Jeon et al. 2018b).

Route of infection. Ingestion of plerocercoids in infected paratenic or intermediate host; ingestion of water containing copepods infected with proceroids (Jeon et al. 2015; Jeon et al. 2018b).

Site in human host. Skin and soft tissues, pleural and peritoneal cavities, abdominal viscera, eye, CNS (Jeon et al. 2015; Jeon et al. 2018b).

Spirometra erinaceieuropaei (Rudolphi, 1819)

Geographic distribution. Europe, Asia, and Australia (Tang et al. 2017; Kuchta et al. 2021).

Natural hosts. First intermediate hosts are freshwater copepods. Second intermediate hosts are amphibians, reptiles and mammals. Definitive hosts are believed to be wild and domestic canids and felids. Zoonotic in humans as dead-end hosts that harbor plerocercoid larvae (Tang et al. 2017; Kuchta et al. 2021).

Route of infection. Ingestion of plerocercoids in infected paratenic or intermediate host; ingestion of water containing copepods infected with proceroids (Lee et al. 1984; Tang et al. 2017).

Site in human host. Skin and soft tissues, pleural and peritoneal cavities, abdominal viscera, eye, CNS as plerocercoid larvae (spargana); rarely as adults in small intestine (Lee et al. 1984; Tang et al. 2017).

Spirometra mansoni (Joyeux & Houdemer, 1928)

Geographic distribution. Southeast Asia (Hong et al. 2016).

Natural hosts. First intermediate hosts are freshwater copepods. Second intermediate hosts are amphibians, while reptiles, birds, and pigs may serve as paratenic hosts. Definitive hosts are felids and canids. Zoonotic in humans as dead-end hosts that harbor plerocercoid larvae (Hong et al. 2016).

Route of infection. Ingestion of plerocercoids in infected paratenic or intermediate host; ingestion of water containing copepods infected with proceroids (Galan-Puchades 2019).

Site in human host. Skin and soft tissues, pleural and peritoneal cavities, abdominal viscera, eye, CNS (Galan-Puchades 2019).

***Spirometra mansonoides* (Mueller, 1935)**

Geographic distribution. North America (McHale et al. 2020).

Natural hosts. First intermediate hosts are freshwater copepods. Second intermediate hosts are amphibians, reptiles, birds, small mammals. Definitive hosts are felids and canids. Zoonotic in humans as dead-end hosts that harbor plerocercoid larvae (McHale et al. 2020).

Route of infection. Ingestion of plerocercoids in infected paratenic or intermediate host; ingestion of water containing copepods infected with proceroids (Landero et al. 1991).

Site in human host. Skin and soft tissues, pleural and peritoneal cavities, abdominal viscera, eye, CNS (Landero et al. 1991).

***Spirometra ranarum* (Gastaldi, 1854)**

Geographic distribution. Africa, Southeast Asia (Jeon et al. 2018a; Saksirisampant et al. 2020).

Natural hosts. First intermediate hosts are freshwater copepods. Second intermediate hosts are amphibians and reptiles. Definitive hosts are wild and domestic felids and canids. Zoonotic in humans as dead-end hosts that harbor plerocercoid larvae (Jeon et al. 2018a; Saksirisampant et al. 2020).

Route of infection. Ingestion of infected intermediate or paratenic host (Saksirisampant et al. 2020).

Site in human host. Skin and soft tissues, pleural and peritoneal cavities, abdominal viscera, eye, CNS (Saksirisampant et al. 2020).

***Spirometra* spp.**

Geographic distribution. East Africa (Eberhard et al. 2015).

Natural hosts. Unknown (Eberhard et al. 2015).

Route of infection. Either ingestion of plerocercoids in undercooked intermediate or paratenic hosts, or water containing copepods infected with proceroids (Eberhard et al. 2015).

Site in human host. Skin and soft tissues (Eberhard et al. 2015).

Notes. Several infections with *Spirometra* sp. have been confirmed in South Sudan and sporadically in other East African countries. Molecular data have confirmed distinctiveness from other known zoonotic *Spirometra* spp., but no specific identification has been determined (Eberhard et al. 2015).

Diphyllobothriidae, incertae sedis:***Sparganum proliferum* (Ilima, 1905)**

Geographic distribution. Undefined (Mueller and Strano 1974).

Natural hosts. Unknown, currently known only from humans in which it is believed to be zoonotic (Mueller and Strano 1974).

Route of infection. Unknown, but presumed to be ingestion of infected intermediate or paratenic host (Mueller and Strano 1974).

Site in human host. Skin and soft tissues, pleural and peritoneal cavities, abdominal viscera, eye, CNS (Mueller and Strano 1974).

Notes. The name *Sparganum proliferum* is the name given to an enigmatic parasite of unknown affinities. To date, it is only known from its larval form from humans.

●●●●●●●● **Cyclophyllidea van Beneden in Braun, 1990**

●●●●●●●● **Dipylidiidae Stiles, 1896**

Genus *Dipylidium* Leuckart, 1863

Microtaenia Sedgwick in Claus & Sedgwick, 1884

***Dipylidium caninum* (Linnaeus, 1758)**

Geographic distribution. Worldwide (Sapp and Bradbury 2020).

Natural hosts. Intermediate hosts are insects, primarily fleas (*Ctenocephalides, Pulex*) and lice (*Trichodectes*). Definitive hosts are wild and domestic felids and canids. Zoonotic in humans (Sapp and Bradbury 2020).

Route of infection. Ingestion of cysticercoids in infected arthropod intermediate host (Cabello et al. 2011; Sapp and Bradbury 2020).

Site in human host. Small intestine (Cabello et al. 2011; Sapp and Bradbury 2020).

●●●●●●●● **Hymenolepididae Ariola, 1899**

Genus *Hymenolepis* Weinland, 1858

***Hymenolepis diminuta* (Rudolphi, 1819)**

Taenia flavopunctata Weinland, 1858

Geographic distribution. Worldwide (Arai 1980).

Natural hosts. Intermediate hosts are insects, commonly granary beetles. Definitive hosts are rodents (primarily rats). Zoonotic in humans (Arai 1980).

Route of infection. Ingestion of cysticercoids infected insect intermediate host (Tiwari et al. 2014).

Site in human host. Small intestine (Tiwari et al. 2014).

***Hymenolepis hibernia* Montgomery et al., 1987**

Geographic distribution. Europe, Asia (Sargison et al. 2018).

Natural hosts. Intermediate hosts are beetles. Definitive hosts are mice in the genus *Apodemus*. Zoonotic in humans (Sargison et al. 2018).

Route of infection. Presumed ingestion of cysticercoids in infected insect intermediate host (Nkouawa et al. 2016).

Site in human host. Small intestine (Nkouawa et al. 2016).

Genus *Rodentolepis* Spasskii, 1954***Rodentolepis microstoma* (Dujardin, 1845)**

Geographic distribution. Worldwide (Cunningham and Olson 2010).

Natural hosts. Intermediate hosts are insects, commonly granary beetles and fleas. Definitive hosts are fleas. Zoonotic in humans (Andreassen et al. 2004; Cunningham and Olson 2010).

Route of infection. Presumably, ingestion of cysticercoids in infected insect intermediate host (Macnish et al. 2003).

Site in human host. Small intestine (Macnish et al. 2003).

Notes. The few cases of *H. microstoma* in humans have been diagnosed by molecular detection in stool (Macnish et al. 2003).

Rodentolepis nana* (Bilharz, 1851)**Hymenolepis fraterna* Stiles, 1906**

Geographic distribution. Worldwide (Sataeva et al. 2018).

Natural hosts. Intermediate hosts are insects, commonly granary beetles and fleas. Definitive hosts are rodents. Zoonotic in humans, although human-to-human transmission can occur (Sataeva et al. 2018).

Route of infection. Ingestion of cysticercoids in infected insect intermediate host; ingestion of eggs on fecally contaminated fomites (Thompson 2015; Sataeva et al. 2018).

Site in human host. Small intestine (Thompson 2015; Sataeva et al. 2018).

●●●●●●●● Taeniidae Ludwig, 1886**Genus *Echinococcus* Rudolphi, 1801*****Echinococcus canadensis* Webster & Cameron, 1961*****Echinococcus granulosus borealis* Williams & Sweatman, 1863**

Echinococcus G6, camel-pig strain

Echinococcus G7, camel-pig strain

Echinococcus G8, American cervid strain

Echinococcus G10, Fennoscandian strain

Geographic distribution. North America, Europe, Middle East, China, east Africa (Romig et al. 2015; Hua et al. 2019).

Natural hosts. Intermediate hosts are camels, pigs, goats, and deer. Definitive hosts are wild and domestic canids. Zoonotic in humans as dead-end hosts harboring larval cysts (Romig et al. 2015; Hua et al. 2019).

Route of infection. Ingestion of eggs in food, water, fomites contaminated with dog feces (Romig et al. 2015).

Site in human host. Disseminated infection, common sites of colonization are liver, lung, brain, bone (Romig et al. 2015).

***Echinococcus equinus* Willams & Sweatman, 1963**

Echinococcus G4, horse strain

Geographic distribution. Europe, North Africa, Middle East (Romig et al. 2015).

Natural hosts. Intermediate hosts are horses. Definitive hosts are wild and domestic canids. Zoonotic in humans as dead-end hosts harboring larval cysts (Romig et al. 2015).

Route of infection. Ingestion of eggs in food, water, fomites contaminated with canid feces (Romig et al. 2015; Kim et al. 2020).

Site in human host. Disseminated infection, common sites of colonization are liver, lung, brain, bone (Kim et al. 2020).

***Echinococcus granulosus* (Batsch, 1796)**

Echinococcus minimus Cameron, 1926

Echinococcus longimanubrius Cameron, 1926

Echinococcus cameroni Ortlepp, 1934

Echinococcus lycaontis Ortlepp, 1934

Echinococcus intermedius Lopez-Neyra & Soler Planas, 1943

Echinococcus patagonicus Szidat, 1960

Echinococcus cepananoi Szidat, 1971

Echinococcus G1, sheep-buffalo strain

Echinococcus G2, sheep-buffalo strain

Echinococcus G3, sheep-buffalo strain

Geographic distribution. Worldwide (Eckert and Deplazes 2004; Romig et al. 2015).

Natural hosts. Intermediate hosts are primarily sheep, goats, cattle, and buffalo. Definitive hosts are wild and domestic canids. Zoonotic in humans as dead-end hosts harboring larval cysts (Eckert and Deplazes 2004; Romig et al. 2015).

Route of infection. Ingestion of eggs in food, water, fomites contaminated with canid feces (Eckert and Deplazes 2004; Romig et al. 2015).

Site in human host. Disseminated infection, common sites of colonization are liver, lung, brain, bone (Eckert and Deplazes 2004; Romig et al. 2015).

***Echinococcus multilocularis* (Leuckart, 1863)**

Echinococcus sibiricensis Rausch & Schiller, 1964

Echinococcus russicensis Tang et al., 2007

Geographic distribution. Europe, Asia, North America (Eckert and Deplazes 2004; Conraths and Deplazes 2015).

Natural hosts. Intermediate hosts are rodents, primarily voles. Definitive hosts are wild canids, primarily red fox (*Vulpes vulpes*); raccoon dogs, domestic dogs, and domestic cats may serve as reservoir hosts. Zoonotic in humans as dead-end hosts harboring larval cysts (Conraths and Deplazes 2015; Eckert and Deplazes 2004).

Route of infection. Ingestion of eggs in food, water, fomites contaminated with canid feces (Eckert and Deplazes 2004; Conraths and Deplazes 2015).

Site in human host. Liver, with dissemination to other organs, such as lung, heart, brain, bone (Eckert and Deplazes 2004; Conraths and Deplazes 2015).

***Echinococcus oligarthra* Diesing, 1863**

Echinococcus cruzi Brumpt & Joyeux, 1924

Echinococcus pampeanus Szidat, 1967

Geographic distribution. South America (D'Alessandro and Rausch 2008).

Natural hosts. Intermediate hosts are rodents and marsupials. Definitive hosts are wild felids. Zoonotic in humans as dead-end hosts harboring larval cysts (D'Alessandro and Rausch 2008).

Route of infection. Ingestion of eggs in food, water, fomites contaminated with felid feces (D'Alessandro and Rausch 2008).

Site in human host. Liver, peritoneal and pleural cavities, with dissemination to other organs (D'Alessandro and Rausch 2008).

***Echinococcus ortleppi* Lopez-Neyra & Soler Planas, 1943**

Echinococcus G5, cattle strain

Geographic distribution. Africa, Europe, India, South America (Romig et al. 2015).

Natural hosts. Intermediate hosts are primarily bovids. Definitive hosts are wild and domestic canids. Zoonotic in humans as dead-end hosts harboring larval cysts (Romig et al. 2015).

Route of infection. Ingestion of eggs in food, water, fomites contaminated with canid feces (Romig et al. 2015).

Site in human host. Disseminated infection, common sites of colonization are liver, lung, brain, bone (Romig et al. 2015).

Echinococcus vogeli Rausch & Berstein, 1972

Geographic distribution. South America (D’Alessandro and Rausch 2008).

Natural hosts. Intermediate hosts are rodents, primarily paca (*Cuniculus paca*). Definitive hosts are bush dogs (*Speothos venaticus*). Zoonotic in humans as dead-end hosts harboring larval cysts (D’Alessandro and Rausch 2008).

Route of infection. Ingestion of eggs in food, water, fomites contaminated with bush dog feces (D’Alessandro and Rausch 2008).

Site in human host. Liver, peritoneal and pleural cavities, with dissemination to other organs (D’Alessandro and Rausch 2008).

Genus *Hydatigera* Lamarck, 1816

Hydatigera taeniaeformis (Batsch, 1786)

Cysticercus fasciolaris Rudolphi, 1808

Geographic distribution. Worldwide (Nakao et al. 2013; Guo 2020).

Natural hosts. Intermediate hosts are primarily rodents and lagomorphs. Definitive hosts are felids. Zoonotic in humans (Nakao et al. 2013; Guo 2020).

Route of infection. Unknown

Site in human host. Small intestine, liver (Stěrba and Barus 1976; Guo 2020).

Genus *Taenia* Linnaeus, 1758

Taenia brauni Setti, 1897

Geographic distribution. Africa (Deplazes et al. 2019).

Natural hosts. Intermediate hosts are rodents and non-human primates. Definitive hosts are canids and genets. Zoonotic in humans as dead-end hosts harboring larval cysts (Deplazes et al. 2019).

Route of infection. Ingestion of eggs in food, water, fomites contaminated with canid feces (Lescano and Zunt 2013).

Site in human host. Skin and soft tissues, eyes (Vanderick et al. 1964; Lescano and Zunt 2013; Deplazes et al. 2019).

Notes. *Taenia brauni* is often considered a subspecies or synonym of *T. serialis*.

***Taenia crassiceps* (Zeder, 1800)**

Taenia hyperborea von Linstow, 1905

Geographic distribution. Northern Hemisphere (Ntoukas et al. 2013).

Natural hosts. Intermediate hosts are rodents. Definitive hosts are canids, primarily foxes. Zoonotic in humans as dead-end hosts harboring larval cysts (Freeman 2011; Ntoukas et al. 2013).

Route of infection. Ingestion of eggs in food, water, fomites contaminated with canid feces (Ntoukas et al. 2013; Tappe et al. 2016a).

Site in human host. Skin and soft tissues, eyes, CNS (Ntoukas et al. 2013; Tappe et al. 2016a).

***Taenia glomeratus* (Railliet & Henry, 1915)**

Geographic distribution. Africa (Deplazes et al. 2019).

Natural hosts. Intermediate hosts are rodents. Definitive hosts are canids and genets. Zoonotic in humans as dead-end hosts harboring larval cysts (Lescano and Zunt 2013; Deplazes et al. 2019).

Route of infection. Ingestion of eggs in food, water, fomites contaminated with canid feces (Lescano and Zunt 2013; Deplazes et al. 2019).

Site in human host. Eyes (Lescano and Zunt 2013; Deplazes et al. 2019).

Notes. *Taenia glomeratus* is often considered a subspecies or synonym of *T. serialis*.

***Taenia martis* (Zeder, 1803)**

Geographic distribution. North America, Europe (Brunet et al. 2015).

Natural hosts. Intermediate hosts include rodents. Definitive hosts are mustelids and foxes. Zoonotic in humans has a dead-end host harboring larval cysts (Eberwein et al. 2013; Brunet et al. 2015; Deplazes et al. 2019).

Route of infection. Ingestion of eggs in food, water, fomites contaminated with feces of definitive host (Brunet et al. 2015; Deplazes et al. 2019).

Site in human host. Skin and soft tissues, CNS, eyes (Eberwein et al. 2013; Brunet et al. 2015; Deplazes et al. 2019).

***Taenia multiceps* Leske, 1790**

Geographic distribution. Worldwide (Varcasia et al. 2015).

Natural hosts. Intermediate hosts are ungulates, primarily sheep. Definitive hosts are canids, primarily foxes. Zoonotic in humans as dead-end hosts harboring larval cysts (Varcasia et al. 2015; Deplazes et al. 2019).

Route of infection. Ingestion of eggs in food, water, fomites contaminated with canid feces (Deplazes et al. 2019).

Site in human host. Skin and soft tissues, CNS (Deplazes et al. 2019).

Taenia saginata Goeze, 1782

Taenia confusa Ward, 1896

Taenia africana von Linstow, 1900

Taenia hominis von Linstow, 1904

Geographic distribution. Worldwide (Hoberg 2002; Braae et al. 2018).

Natural hosts. Intermediate hosts are cattle. Definitive hosts are humans (Braae et al. 2018; Hoberg 2002).

Route of infection. Ingestion of cysticercoids in infected cattle (Braae et al. 2018).

Site in human host. Small intestine (Braae et al. 2018).

Taenia serialis Gervais, 1847

Multiceps radians Joyeux et al., 1922

Geographic distribution. Worldwide; highest prevalence in Africa (Schneider-Crease et al. 2017).

Natural hosts. Intermediate hosts include rabbits, rodents, cattle, sheep, goats. Definitive hosts are wild and domestic canids. Zoonotic in humans as dead-end hosts harboring larval cysts (Schneider-Crease et al. 2017).

Route of infection. Ingestion of eggs in food, water, fomites contaminated with canid feces (Tappe et al. 2016a).

Site in human host. CNS (Tappe et al. 2016a).

Taenia solium Linnaeus, 1758

Cysticercus cellulosae Gmelin, 1790

Geographic distribution. Worldwide (Garcia et al. 2003; Nakao et al. 2002; Coral-Almeida et al. 2015).

Natural hosts. Intermediate hosts are pigs. Definitive hosts are humans, although in cases of cysticercosis humans function as dead-end intermediate hosts (Garcia et al. 2003; Coral-Almeida et al. 2015).

Route of infection. Ingestion of cysticercoids in infected pigs (taeniasis); ingestion of eggs in food, water, fomites contaminated with human feces (cysticercosis) (Garcia et al. 2003).

Site in human host. Small intestine (taeniasis); disseminated infection with predilection for CNS (cysticercosis) (Garcia et al. 2003).

***Taenia sui hominis* Mathison et al., 2021**

Taenia asiatica Eom & Rim, 1993

Taenia saginata asiatica Eom & Rim, 1993

Taenia asiaticus Eom et al., 2020

Geographic distribution. East Asia (Ale et al. 2014).

Natural hosts. Intermediate hosts are pigs. Definitive hosts are humans (Eom and Rim 1993; Ale et al. 2014).

Route of infection. Ingestion of cysticerci in infected pigs (Ale et al. 2014; Eom and Rim 1993).

Site in human host. Small intestine (Eom and Rim 1993; Ale et al. 2014).

Genus *Versteria* Nakao et al., 2013

Geographic distribution. Europe, Central Asia, North America (Nakao et al. 2013; Deplazes et al. 2019).

Natural hosts. Intermediate hosts are rodents; non-human primates can be incidental hosts. Definitive hosts are mustelids. Zoonotic in humans (Nakao et al. 2013; Deplazes et al. 2019).

Route of infection. Presumably by the ingestion of eggs in food, water, fomites contaminated with feces of definitive host (Deplazes et al. 2019).

Site in human host. Various organs, disseminated infection (Barkati et al. 2018; Lehman et al. 2019).

Notes. Human and primate *Versteria* infections have not been identified to species level but bear close genetic resemblance to isolates from North American mink and are somewhat similar to isolates of *V. mustelae* (Gmelin, 1790) from European mink (Lee et al. 2016; Lehman et al. 2019).

●●●●●●● Anoplocephalidae Cholodkovsky, 1902**Genus *Bertiella* Stiles & Hassall, 1902**

Bertia Blanchard, 1891

***Bertiella mucronotata* (Meyner, 1895)**

Geographic distribution. South America, Caribbean (Sapp and Bradbury 2020).

Natural hosts. First intermediate hosts are oribatid mites. Definitive hosts are New World monkeys. Zoonotic in humans (Sapp and Bradbury 2020).

Route of infection. Ingestion of cysticercoids in infected mite intermediate host (Sapp and Bradbury 2020).

Site in human host. Small intestine (Sapp and Bradbury 2020).

***Bertiella studeri* (Blanchard, 1891)**

Geographic distribution. Africa, Asia (Sapp and Bradbury 2020).

Natural hosts. First intermediate hosts are oribatid mites. Definitive hosts are Old World monkeys and non-human apes. Zoonotic in humans (Sapp and Bradbury 2020).

Route of infection. Ingestion of cysticercoids in infected mite intermediate host (Sapp and Bradbury 2020).

Site in human host. Small intestine (Sapp and Bradbury 2020).

Genus *Inermicapsifer* Janicki, 1910***Inermicapsifer madagascariensis* (Davaine, 1870)**

Acanthocephala arvicanthidis Kofend, 1917

Raillietina cubensis Kouri, 1938

Geographic distribution. Sub-Saharan Africa, South America, West Indies, Southeast Asia (Goldsmid and Muir 1972; Sapp and Bradbury 2020).

Natural hosts. Intermediate hosts are not known, but presumed to be terrestrial arthropods. Definitive hosts are rodents and hyraxes. Zoonotic in humans (Goldsmid and Muir 1972; Sapp and Bradbury 2020).

Route of infection. Ingestion of cysticercoids in infected arthropod intermediate host (Sapp and Bradbury 2020).

Site in human host. Small intestine (Sapp and Bradbury 2020).

Genus *Mathevotaenia* Akhumyan, 1946***Mathevotaenia symmetrica* (Baylis, 1927)**

Geographic distribution. Europe, Asia (Lamon and Greer 1986).

Natural hosts. Intermediate hosts are suspected as being arthropods. Definitive hosts are rodents. Zoonotic in humans (Lamon and Greer 1986).

Route of infection. Presumably, ingestion of cysticercoids in infected arthropod intermediate host (Lamon and Greer 1986).

Site in human host. Small intestine (Lamon and Greer 1986).

Genus *Moniezia* Blanchard, 1891***Moniezia expansa* (Rudolphi, 1810)**

Geographic distribution. Worldwide (single human case from Egypt) (Elliott 1986; el-Shazly et al. 2004).

Natural hosts. Intermediate hosts are mites. Definitive hosts are sheep, goats, cattle. Zoonotic in humans (Elliott 1986; el-Shazly et al. 2004).

Route of infection. Presumably, ingestion of cysticercoids in infected arthropod intermediate host (el-Shazly et al. 2004).

Site in human host. Small intestine (el-Shazly et al. 2004).

●●●●●●●● **Davaineidae Braun, 1900**

Genus *Raillietina* Fuhrman, 1920

***Raillietina celebensis* (Janicki, 1902)**

Taenia asiatica von Linstow, 1891

Taenia formosana Akashi, 1916

Raillietina funerebris Meggitt & Subramanian, 1927

Raillietina garrisoni Tubanguai, 1931

Raillietina sinensis Hsu, 1935

Raillietina murium Joyeux & Baer, 1938

Geographic distribution. Asia, Australia (Baer and Sandars 1956; Sapp and Bradbury 2020).

Natural hosts. Intermediate hosts are invertebrates, including insects (primarily ants and beetles) and terrestrial mollusks. Definitive hosts are rodents and shrews. Zoonotic in humans (Baer and Sandars 1956; Sapp and Bradbury 2020).

Route of infection. Ingestion of cysticercoids in infected arthropod intermediate host (Baer and Sandars 1956; Sapp and Bradbury 2020).

Site in human host. Small intestine (Baer and Sandars 1956; Sapp and Bradbury 2020).

***Raillietina demerariensis* (Daniels, 1895)**

Raillietina quitensis Leon, 1935

Raillietina brumpti Doilfus, 1939

Raillietina equatoriensis Dollfus, 1939

Raillietina leoni Dollfus, 1939

Raillietina luisaleoni Dollfus, 1939

Raillietina halli Perez-Vigueras, 1934

Geographic distribution. South America, Caribbean (Baer and Sandars 1956; Sapp and Bradbury 2020).

Natural hosts. Intermediate hosts are invertebrates, including insects (primarily ants and beetles) and terrestrial mollusks. Definitive hosts are rodents; non-human primates may serve as reservoir hosts. Zoonotic in humans (Baer and Sandars 1956; Sapp and Bradbury 2020).

Route of infection. Ingestion of cysticercooids in infected arthropod intermediate host (Baer and Sandars 1956; Sapp and Bradbury 2020).

Site in human host. Small intestine (Baer and Sandars 1956; Sapp and Bradbury 2020).

Raillietina siriraji Chandler & Pradatsundarasar, 1957

Geographic distribution. Thailand (Sapp and Bradbury 2020).

Natural hosts. Intermediate hosts are invertebrates, including insects (primarily ants and beetles). Definitive hosts are rodents. Zoonotic in humans (Sapp and Bradbury 2020).

Route of infection. Ingestion of cysticercooids in infected arthropod intermediate host (Sapp and Bradbury 2020).

Site in human host. Small intestine (Sapp and Bradbury 2020).

●●●●●●● Mesocestoididae Fuhrmann, 1907

Genus *Mesocestoides* Valliant, 1863

Mesocestoides lineatus (Goeze, 1782)

Geographic distribution. Europe and Asia (Sapp and Bradbury 2020).

Natural hosts. The complete life cycle and presumed number of hosts is not completely understood. The presumed first intermediate host is believed to be an arthropod. The second intermediate hosts are a variety of vertebrates, including small mammals, birds, reptiles, and amphibians. Definitive hosts are carnivores, including foxes, wolves, raccoon dogs, and European badgers. Zoonotic in humans (Padgett and Boyce 2004; Sapp and Bradbury 2020).

Route of infection. Unknown; presumed to be ingestion of tetrathyridia in meat and viscera of infected second intermediate hosts (Sapp and Bradbury 2020).

Site in human host. Small intestine (Sapp and Bradbury 2020).

Mesocestoides variabilis Mueller, 1928

Geographic distribution. North America (Sapp and Bradbury 2020).

Natural hosts. The complete life cycle and presumed number of hosts is not completely understood. The presumed first intermediate host is believed to be an arthropod. The second intermediate hosts are a variety of vertebrates, including small mammals, birds, reptiles, and amphibians. Definitive hosts include wild and domestic canids and felids, mustelids, and opossums. Zoonotic in humans (Padgett and Boyce 2004; Sapp and Bradbury 2020).

Route of infection. Unknown; presumed to be ingestion of tetrathyridia in meat and viscera of infected second intermediate hosts (Sapp and Bradbury 2020).

Site in human host. Small intestine (Sapp and Bradbury 2020).

- Trematoda Rudolphi, 1808
- Digenea Carus, 1863
- Diplostomida Olson et al., 2003
- Diplostomata Olson et al., 2003
- Brachylaeimidae Joyeux & Foley, 1930

Genus *Brachylaeima* Dujardin, 1843

Brachylaeima cribbi Butcher & Grove, 2001

Geographic distribution. Australia (Chai and Jung 2020).

Natural hosts. First intermediate hosts are terrestrial snails in the genus *Theba*. Second intermediate hosts are land snails in the genus *Cerनुualla*. Definitive hosts include a variety of birds, mammals, and reptiles. Zoonotic in humans (Chai and Jung 2020).

Route of infection. Ingestion of metacercariae in infected snail intermediate hosts (Chai and Jung 2020).

Site in human host. Small intestine (Chai and Jung 2020).

- Cyathocotylidae Mühling, 1898

Genus *Prohemistomum* Odhner, 1913

Prohemistomum vivax (Sonsino, 1892)

Geographic distribution. Europe, North Africa, Middle East (Nasr 1941; Chai and Jung 2020).

Natural hosts. First intermediate hosts are freshwater snails in the genus *Cleopatra*. Second intermediate hosts are various brackish water and freshwater fish. Definitive hosts are felids, canids, and birds. Zoonotic in humans (Nasr 1941; Chai and Jung 2020).

Route of infection. Ingestion of metacercariae in infected fish intermediate hosts (Nasr 1941; Chai and Jung 2020)

Site in human host. Small intestine (Nasr 1941; Chai and Jung 2020).

- Diplostomidae Poirier, 1886

Genus *Alaria* Schrank, 1788

Alaria americana Hall & Wigdor, 1918

Geographic distribution. North America (Möhl et al. 2009).

Natural hosts. First intermediate hosts are snails in the genera *Planorbis*, *Heliosoma*, *Lymnea*, and *Anisus*. Second intermediate hosts are amphibians; many vertebrate

animals can serve as paratenic hosts. Definitive hosts are carnivores. Zoonotic in humans (Möhl et al. 2009).

Route of infection. Ingestion of metacercariae in undercooked game meat (Möhl et al. 2009).

Site in human host. Subcutaneous tissues, eyes, lungs, disseminated infections (Freeman et al. 1976; Möhl et al. 2009).

Genus *Fibricola* Dubois, 1932

Fibricola cratera (Barker & Noll, 1915)

Geographic distribution. North America (Hoffman 1955; Chai and Jung 2020).

Natural hosts. First intermediate hosts are freshwater snails in the genus *Physa*. Second intermediate hosts are amphibians. Definitive hosts are rodents and shrews. Zoonotic in humans (Hoffman 1955; Chai and Jung 2020).

Route of infection. Ingestion of metacercariae in infected amphibian intermediate hosts (Chai and Jung 2020).

Site in human host. Small intestine (Chai and Jung 2020).

Genus *Neodiplostomum* Railliet, 1919

Neodiplostomum seoulense Seo et al., 1964

Geographic distribution. China, South Korea (Chai and Jung 2020).

Natural hosts. First intermediate hosts are freshwater snails in the genera *Hippeutis* and *Segmentina*. Second intermediate hosts are amphibians; snakes can serve as paratenic hosts. Definitive hosts are rodents and humans (Chai and Jung 2020).

Route of infection. Ingestion of metacercariae in infected reptile and amphibian intermediate and paratenic hosts (Chai and Jung 2020).

Site in human host. Small intestine (Chai and Jung 2020).

●●●●●●●● Strigeidae Railliet, 1919

Genus *Cotylurus* Szidat, 1928

Cotylurus japonicus Ishii, 1932

Geographic distribution. Central and Southeast Asia, Russia, Japan (Chai and Jung 2020).

Natural hosts. First intermediate hosts are unknown but hypothesized to be freshwater snails. Second intermediate hosts are unknown but presumed to be freshwater snails. Definitive hosts are ducks. Zoonotic in humans (Chai and Jung 2020).

Route of infection. Ingestion of metacercariae in infected insect intermediate hosts (Chai and Jung 2020).

Site in human host. Small intestine (Chai and Jung 2020).

●●●●●●●● **Schistosomatidae Stiles & Hassall, 1898**

Genus *Schistosoma* Weinland, 1858

***Schistosoma bovis* (Bilharz, 1852)**

Geographic distribution. Southern Europe, Middle East, South Asia, Africa (Christensen et al. 1983).

Natural hosts. First intermediate hosts are snails in the genus *Bulinus*. Definitive hosts are ruminants, primarily cattle. Zoonotic in humans (Christensen et al. 1983).

Route of infection. Penetration of skin by free-swimming cercariae

Site in human host. Mesenteric veins of the large intestine (Chunge et al. 1986; Mouchet et al. 1988)

Notes. Human infections with hybrids of *S. bovis* and *S. haematobium* have been confirmed using molecular methods (Oleaga et al. 2019).

***Schistosoma guineensis* Pagès et al., 2003**

Schistosoma intercalatum, Guinea strain

Geographic distribution. West Africa (Webster et al. 2006).

Natural hosts. First intermediate hosts are snails in the genus *Bulinus*. Definitive hosts are humans, rodents, ruminants (Webster et al. 2006).

Route of infection. Penetration of skin by free-swimming cercariae (Webster et al. 2006).

Site in human host. Mesenteric veins of the large intestine (Webster et al. 2006).

***Schistosoma haematobium* (Bilharz, 1852)**

Geographic distribution. Africa, isolated regions of the Middle East (Colley et al. 2014; Gautret et al. 2015).

Natural hosts. First intermediate hosts are snails in the genus *Bulinus*. Definitive hosts are humans (Colley et al. 2014).

Route of infection. Penetration of skin by free-swimming cercariae (Colley et al. 2014).

Site in human host. Venous plexus of the urinary bladder (Colley et al. 2014).

Notes. Human infection of hybrids of *S. haematobium* and *S. bovis* have been documented in Corsica, France (Oleaga et al. 2019).

***Schistosoma intercalatum* Fisher, 1934**

Geographic distribution. Democratic Republic of the Congo (Webster et al. 2006; Colley et al. 2014).

Natural hosts. First intermediate hosts are snails in the genus *Bulinus*. Definitive hosts are humans, rodents, ruminants (Colley et al. 2014).

Route of infection. Penetration of skin by free-swimming cercariae (Colley et al. 2014).

Site in human host. Mesenteric veins of the large intestine (Colley et al. 2014).

***Schistosoma japonicum* (Katsurada, 1904)**

Geographic distribution. China, Philippines, Indonesia (Sulawesi) (Colley et al. 2014).

Natural hosts. First intermediate hosts are snails in the genus *Oncomelania*. Definitive hosts include a variety of mammals, including canids, felids, pigs, ruminants, rodents, and humans (Colley et al. 2014).

Route of infection. Penetration of skin by free-swimming cercariae (Colley et al. 2014).

Site in human host. Mesenteric veins of the small and large intestine (Colley et al. 2014).

***Schistosoma mansoni* Sambon, 1907**

Geographic distribution. Sub-Saharan Africa, South America, Caribbean, parts of the Arabian Peninsula (Colley et al. 2014).

Natural hosts. First intermediate hosts are snails in the genus *Biomphalaria*. Definitive hosts are humans, occasionally non-human primates (Colley et al. 2014).

Route of infection. Penetration of skin by free-swimming cercariae (Colley et al. 2014).

Site in human host. Mesenteric veins of the large intestine (Colley et al. 2014).

***Schistosoma mattheei* Veglia & LeRoux, 1929**

Geographic distribution. Africa, Middle East (Christensen et al. 1983).

Natural hosts. First intermediate hosts are snails in the genus *Bulinus*. Definitive hosts are wild and domestic ruminants. Zoonotic in humans (Christensen et al. 1983).

Route of infection. Penetration of skin by free-swimming cercariae (Christensen et al. 1983).

Site in human host. Presumed to be venous plexus of the urinary bladder (Wolmarans et al. 1990; Cnops et al. 2020).

Notes. Reports of patent, egg-producing human infections with *S. mattheei* are most likely a result of hybridization of this species with *S. haematobium*. Dead-end acute infections with *S. mattheei* × *S. haematobium* have been confirmed using molecular methods (Kruger and Evans 2009; Cnops et al. 2020).

***Schistosoma mekongi* Voge et al., 1978**

Geographic distribution. Mekong River Valley of Cambodia and Laos (Colley et al. 2014).

Natural hosts. First intermediate host is the snail *Neotricula aperta*. Definitive hosts include a variety of mammals, including canids, felids, pigs, ruminants, rodents, and humans (Colley et al. 2014).

Route of infection. Penetration of skin by free-swimming cercariae (Colley et al. 2014).

Site in human host. Mesenteric veins of the large intestine (Colley et al. 2014).

Cercarial dermatitis

Many species of avian, and some mammalian, schistosomes cause a limited cutaneous reaction in humans referred to as cercarial dermatitis (or “swimmer’s itch”). The identity of causative agents is typically inferred via environmental investigations following outbreaks and may or may not be identified to species level; only the major genera are listed here:

Genus *Anserobilharzia* Brandt et al., 2013

Geographic distribution. Northern Hemisphere (Brant et al. 2013).

Natural hosts. Intermediate hosts are freshwater snails in the family Planorbidae. Definitive hosts are charadriiform birds. Zoonotic in humans (Brant et al. 2013; Horák et al. 2015).

Route of infection. Penetration of skin by free-swimming cercariae (Horák et al. 2015).

Site in human host. Skin (Horák et al. 2015).

Genus *Austroilharzia* Johnston, 1917

Geographic distribution. Northern Hemisphere, Australia (Farley 1971).

Natural hosts. Intermediate hosts are freshwater snails in the family Planorbidae. Definitive hosts are charadriiform birds. Zoonotic in humans (Farley 1971).

Route of infection. Penetration of skin by free-swimming cercariae (Horák et al. 2015).

Site in human host. Skin (Horák et al. 2015).

Genus *Bilharziella* Looss, 1899

Geographic distribution. Europe (Prüter et al. 2017).

Natural hosts. Intermediate hosts are freshwater snails in the family Planorbidae. Definitive hosts are waterfowl and other wading birds. Zoonotic in humans (Prüter et al. 2017).

Route of infection. Penetration of skin by free-swimming cercariae (Horák et al. 2015).

Site in human host. Skin (Horák et al. 2015).

Genus *Bivitellobilharzia* Vogel & Minning, 1940

Geographic distribution. Africa, Asia (Devkota et al. 2014).

Natural hosts. Snail intermediate hosts are unknown. Definitive hosts are elephants and rhinoceroses. Zoonotic in humans (Devkota et al. 2014).

Route of infection. Penetration of skin by free-swimming cercariae (Horák et al. 2015).

Site in human host. Skin (Horák et al. 2015).

Genus *Gigantobilharzia* Odhner, 1910

Geographic distribution. North America (McDonald 1981).

Natural hosts. Intermediate hosts are freshwater snails in the family Physidae. Definitive hosts are passerine birds. Zoonotic in humans (McDonald 1981).

Route of infection. Penetration of skin by free-swimming cercariae (Horák et al. 2015).

Site in human host. Skin (Horák et al. 2015).

Genus *Heterobilharzia* Price, 1929

Geographic distribution. North America (Graham et al. 2021).

Natural hosts. Intermediate hosts are freshwater snails in the family Lymnaeidae. Definitive hosts are various mammals, including carnivores and marsupials. Zoonotic in humans (Graham et al. 2021).

Route of infection. Penetration of skin by free-swimming cercariae (Horák et al. 2015).

Site in human host. Skin (Horák et al. 2015).

Genus *Ornithobilharzia* Odhner, 1912

Geographic distribution. Worldwide (Farley 1971)

Natural hosts. Intermediate hosts are marine snails in the family Batillariidae. Definitive hosts are various birds and mammals. Zoonotic in humans (Farley 1971).

Route of infection. Penetration of skin by free-swimming cercariae (Horák et al. 2015).

Site in human host. Skin (Horák et al. 2015).

Genus *Trichobilharzia* Skrjabin & Zakharow, 1920

Geographic distribution. Worldwide (Horák et al. 2002).

Natural hosts. Intermediate hosts are freshwater snails in the families Lymnaeidae and Physidae. Definitive hosts are waterfowl. Zoonotic in humans (Horák et al. 2002).

Route of infection. Penetration of skin by free-swimming cercariae (Horák et al. 2015).

Site in human host. Skin (Horák et al. 2015).

●●●●●●●● Plagiorchiida La Rue, 1957

●●●●●●●● Bucephalata La Rue, 1926

●●●●●●●● Gymnophallidae Odhner, 1905

Genus *Gymnophalloides* Fujita, 1925

Gymnophalloides seoi Chai et al., 2003

Geographic distribution. South Korea (Lee and Chai 2001; Chai and Jung 2020).

Natural hosts. First intermediate hosts are unknown. Second intermediate hosts are oysters in the genus *Crassostrea*. Definitive hosts are oystercatchers (*Haematopus*). Zoonotic in humans (Lee and Chai 2001; Chai and Jung 2020).

Route of infection. Ingestion of metacercariae in infected oyster intermediate hosts (Lee and Chai 2001; Chai and Jung 2020),

Site in human host. Small intestine (Lee and Chai 2001; Chai and Jung 2020).

●●●●●●●● Echinostomata La Rue, 1926

●●●●●●●● Himasthlidae Odhner, 1910

Genus *Acanthoparyphium* Dietz, 1909

Acanthoparyphium tyosenense Yamaguti, 1939

Geographic distribution. Korea, Japan (Toledo and Esteban 2016; Chai and Jung 2020).

Natural hosts. First intermediate hosts are marine gastropods. The second intermediate hosts are brackish water bivalves. Definitive hosts are ducks. Zoonotic in humans (Toledo and Esteban 2016; Chai and Jung 2020).

Route of infection. Ingestion of metacercariae in infected bivalves (Toledo and Esteban 2016; Chai and Jung 2020).

Site in human host. Small intestine (Toledo and Esteban 2016; Chai and Jung 2020).

Genus *Himasthla* Dietz, 1909

Himasthla muehlensi Vogel, 1933

Geographic distribution. North America, Colombia (Toledo and Esteban 2016; Chai and Jung 2020).

Natural hosts. First intermediate hosts are unknown, presumed to be marine gastropods. Second intermediate hosts are presumed to be marine clams and mussels. Definitive hosts are birds. Zoonotic in humans (Toledo and Esteban 2016; Chai and Jung 2020).

Route of infection. Ingestion of metacercariae in infected mollusk intermediate hosts (Toledo and Esteban 2016; Chai and Jung 2020).

Site in human host. Small intestine (Toledo and Esteban 2016; Chai and Jung 2020).

●●●●●●●● Echinochasmidae Odhner, 1910

Genus *Echinochasmus* Dietz, 1909

Echinochasmus caninus (Verma, 1935)

Geographic distribution. Southeast Asia, India (Toledo and Esteban 2016; Chai and Jung 2020).

Natural hosts. First intermediate hosts are unknown. Second intermediate hosts are freshwater fish in the genus *Macropodus*. Definitive hosts are wild and domestic canids. Zoonotic in humans (Toledo and Esteban 2016; Chai and Jung 2020).

Route of infection. Ingestion of metacercariae in infected fish intermediate host (Toledo and Esteban 2016; Chai and Jung 2020).

Site in human host. Small intestine (Toledo and Esteban 2016; Chai and Jung 2020).

Echinochasmus fujianensis Chen, in Chen et al., 1992

Geographic distribution. China (Toledo and Esteban 2016; Chai and Jung 2020).

Natural hosts. First intermediate hosts are freshwater snails in the genus *Bellamya*. Second intermediate hosts are freshwater fish in the genera *Pseudorasbora* and *Cyprinus*. Definitive hosts are wild and domestic canids and felids, pigs, and rodents. Zoonotic in humans (Toledo and Esteban 2016; Chai and Jung 2020).

Route of infection. Ingestion of metacercariae in infected fish intermediate host (Toledo and Esteban 2016; Chai and Jung 2020).

Site in human host. Small intestine (Toledo and Esteban 2016; Chai and Jung 2020).

Echinochasmus japonicus Tanabe, 1926

Geographic distribution. Southeast Asia, Japan (Toledo and Esteban 2016; Chai and Jung 2020).

Natural hosts. First intermediate hosts are freshwater snails, including the genus *Parafossarulis*. Second intermediate hosts are various freshwater fish. Definitive hosts are felids, insectivores, and birds. Zoonotic in humans (Toledo and Esteban 2016; Chai and Jung 2020).

Route of infection. Ingestion of metacercariae in infected fish intermediate host (Toledo and Esteban 2016; Chai and Jung 2020).

Site in human host. Small intestine (Toledo and Esteban 2016; Chai and Jung 2020).

***Echinochasmus jiufoensis* Liang & Ke, 1988**

Geographic distribution. China (Toledo and Esteban 2016; Chai and Jung 2020).

Natural hosts. First intermediate hosts are unknown. Second intermediate hosts are presumed to be freshwater fish or mollusks. Definitive hosts are felids, canids, and pigs. Zoonotic in humans (Toledo and Esteban 2016; Chai and Jung 2020).

Route of infection. Ingestion of metacercariae in infected intermediate host (Toledo and Esteban 2016; Chai and Jung 2020).

Site in human host. Small intestine (Toledo and Esteban 2016; Chai and Jung 2020).

***Echinochasmus liliputanus* (Looss, 1896)**

Geographic distribution. China, Middle East, North Africa (Toledo and Esteban 2016; Chai and Jung 2020).

Natural hosts. First intermediate hosts are freshwater snails in the genus *Parafossarulus*. Second intermediate hosts are freshwater fish in the genera *Pseudorasbora* and *Carassius*. Definitive hosts are felids, canids, raccoons, and badgers. Zoonotic in humans (Toledo and Esteban 2016; Chai and Jung 2020).

Route of infection. Ingestion of metacercariae in infected intermediate fish host or water contaminated with metacercariae (Toledo and Esteban 2016; Chai and Jung 2020).

Site in human host. Small intestine (Toledo and Esteban 2016; Chai and Jung 2020).

***Echinochasmus perfoliatus* (Ratz, 1908)**

Geographic distribution. Southeast Asia, Russia, Europe, North Africa (Toledo and Esteban 2016; Chai and Jung 2020).

Natural hosts. First intermediate hosts are various freshwater snails (*Parafossarulus*, *Bithynia*, *Lymnaea*). Second intermediate hosts are various freshwater fish (*Zacco*, *Carassius*, *Pseudorasbora*). Definitive hosts are felids, canids, pigs, and birds. Zoonotic in humans (Toledo and Esteban 2016; Chai and Jung 2020).

Route of infection. Ingestion of metacercariae in infected fish intermediate host (Toledo and Esteban 2016; Chai and Jung 2020).

Site in human host. Small intestine (Toledo and Esteban 2016; Chai and Jung 2020).

●●●●●●●● **Echinostomatidae Looss, 1899**

Genus *Artyfechinostomum* Lane, 1915

***Artyfechinostomum malayanum* (Leiper, 1911)**

Geographic distribution. Southeast Asia (Toledo and Esteban 2016; Chai and Jung 2020).

Natural hosts. First intermediate hosts are freshwater snails in the genera *Indoplanorbis* and *Gyraulus*. Second intermediate hosts are several freshwater snails. Definitive hosts include pigs, rodents, shrews, felids, canids. Zoonotic in humans (Toledo and Esteban 2016; Chai and Jung 2020).

Route of infection. Ingestion of metacercariae in infected snail intermediate hosts (Toledo and Esteban 2016; Chai and Jung 2020).

Site in human host. Small intestine (Toledo and Esteban 2016; Chai and Jung 2020).

Artyfechinostomum oraoni Bandyopadhyay et al., 1989

Geographic distribution. India (Toledo and Esteban 2016; Chai and Jung 2020).

Natural hosts. First intermediate hosts are snails in the genus *Lymnaea*. Second intermediate hosts and definitive hosts are unknown; presumed zoonotic in humans (Toledo and Esteban 2016; Chai and Jung 2020).

Route of infection. Ingestion of metacercariae in infected intermediate hosts (Toledo and Esteban 2016; Chai and Jung 2020).

Site in human host. Small intestine (Toledo and Esteban 2016; Chai and Jung 2020).

Artyfechinostomum sufratyfex Lane, 1915

Cercaria mehrai Faruqui, 1930

Geographic distribution. India, Vietnam (Toledo and Esteban 2016; Chai and Jung 2020).

Natural hosts. First intermediate hosts are snails in the genera *Indoplanorbis* and *Lymnaea*. Second intermediate hosts include freshwater snails, fish, and amphibians. Definitive hosts are pigs, canids, and rodents. Zoonotic in humans (Toledo and Esteban 2016; Chai and Jung 2020).

Route of infection. Ingestion of metacercariae in infected intermediate hosts (Toledo and Esteban 2016; Chai and Jung 2020).

Site in human host. Small intestine (Toledo and Esteban 2016; Chai and Jung 2020).

Genus *Echinoparyphium* Dietz, 1909

Echinoparyphium recurvatum (von Linstow, 1873)

Geographic distribution. North America, Europe, North Africa, Middle East, Central and Southeast Asia (Toledo and Esteban 2016; Chai and Jung 2020).

Natural hosts. First intermediate hosts are snails in the genera *Physa* and *Lymnaea*. Second intermediate hosts include freshwater snails and amphibians. Definitive hosts are canids and rodents. Zoonotic in humans (Toledo and Esteban 2016; Chai and Jung 2020).

Route of infection. Ingestion of metacercariae in infected intermediate hosts (Toledo and Esteban 2016; Chai and Jung 2020).

Site in human host. Small intestine (Toledo and Esteban 2016; Chai and Jung 2020).

Genus *Echinostoma* Rudolphi, 1809

***Echinostoma aegyptica* Khalil & Abaza, 1924**

Geographic distribution. North Africa, Middle East, Southeast Asia, Japan (Toledo and Esteban 2016; Chai and Jung 2020).

Natural hosts. First and second intermediate hosts are unknown. Definitive hosts are rats. Zoonotic in humans (Toledo and Esteban 2016; Chai and Jung 2020).

Route of infection. Ingestion of metacercariae in infected intermediate hosts (Toledo and Esteban 2016; Chai and Jung 2020).

Site in human host. Small intestine (Toledo and Esteban 2016; Chai and Jung 2020).

***Echinostoma angustitestis* Wang, 1977**

Geographic distribution. China (Toledo and Esteban 2016; Chai and Jung 2020).

Natural hosts. First intermediate hosts are presumed to be freshwater snails. Second intermediate hosts are freshwater fish. Definitive hosts are dogs and livestock. Zoonotic in humans (Toledo and Esteban 2016; Chai and Jung 2020).

Route of infection. Ingestion of metacercariae in infected fish intermediate hosts (Toledo and Esteban 2016; Chai and Jung 2020).

Site in human host. Small intestine (Toledo and Esteban 2016; Chai and Jung 2020).

***Echinostoma cinetorchis* Ando & Ozaki, 1923**

Geographic distribution. Southeast Asia, Japan (Toledo and Esteban 2016; Chai and Jung 2020).

Natural hosts. First intermediate hosts are snails in the genera *Hippeutis* and *Segmentina*. Second intermediate hosts include freshwater snails, fish (*Misgurnus*), and amphibians. Definitive hosts are rats. Zoonotic in humans (Toledo and Esteban 2016; Chai and Jung 2020).

Route of infection. Ingestion of metacercariae in infected intermediate hosts (Toledo and Esteban 2016; Chai and Jung 2020).

Site in human host. Small intestine (Toledo and Esteban 2016; Chai and Jung 2020).

***Echinostoma ilocanum* (Garrison, 1908)**

Geographic distribution. Central and Southeast Asia (Toledo and Esteban 2016; Chai and Jung 2020).

Natural hosts. First intermediate hosts are snails in the genera *Gyraulus* and *Hippeutis*. Second intermediate hosts include freshwater snails in the genera *Pila* and *Vivaparus*. Definitive hosts are rats and canids. Zoonotic in humans (Toledo and Esteban 2016; Chai and Jung 2020).

Route of infection. Ingestion of metacercariae in infected snail intermediate hosts (Toledo and Esteban 2016; Chai and Jung 2020).

Site in human host. Small intestine (Toledo and Esteban 2016; Chai and Jung 2020).

***Echinostoma lindoense* Sandground & Bonne, 1940**

Geographic distribution. Europe, Southeast Asia, South America (Toledo and Esteban 2016; Chai and Jung 2020).

Natural hosts. First intermediate hosts are several genera of freshwater snails. Second intermediate hosts include freshwater snails and mussels. Definitive hosts include a variety of mammals and birds. Zoonotic in humans (Toledo and Esteban 2016; Chai and Jung 2020).

Route of infection. Ingestion of metacercariae in infected mollusk intermediate hosts (Toledo and Esteban 2016; Chai and Jung 2020).

Site in human host. Small intestine (Toledo and Esteban 2016; Chai and Jung 2020).

***Echinostoma macrorchis* Ando & Ozaki, 1923**

Geographic distribution. Southeast Asia, Japan (Toledo and Esteban 2016; Chai and Jung 2020).

Natural hosts. First intermediate hosts are snails in the genera *Segmentina* and *Gyraulus*. Second intermediate hosts include freshwater snails (*Segmentina*) and amphibians. Definitive hosts are rodents and wading birds. Zoonotic in humans (Toledo and Esteban 2016; Chai and Jung 2020).

Route of infection. Ingestion of metacercariae in infected intermediate hosts (Toledo and Esteban 2016; Chai and Jung 2020).

Site in human host. Small intestine (Toledo and Esteban 2016; Chai and Jung 2020).

***Echinostoma paraensei* Lie & Basch, 1967**

Geographic distribution. Australia, Brazil (Toledo and Esteban 2016; Chai and Jung 2020).

Natural hosts. First and second intermediate hosts are snails in the genus *Biomphalaria*. Definitive hosts are rats. Zoonotic in humans (Toledo and Esteban 2016; Chai and Jung 2020).

Route of infection. Ingestion of metacercariae in infected snail intermediate hosts (Toledo and Esteban 2016; Chai and Jung 2020).

Site in human host. Small intestine (Toledo and Esteban 2016; Chai and Jung 2020).

***Echinostoma revolutum* (Froelich, 1802)**

Geographic distribution. Europe, Russia, Middle East, Central and Southeast Asia, North America (Toledo and Esteban 2016; Chai and Jung 2020).

Natural hosts. First intermediate hosts include several genera of freshwater snails. Second intermediate hosts include freshwater snails and clams, and amphibians. Definitive hosts are felids, canids, rodents, and birds. Zoonotic in humans (Toledo and Esteban 2016; Chai and Jung 2020).

Route of infection. Ingestion of metacercariae in infected intermediate hosts (Toledo and Esteban 2016; Chai and Jung 2020).

Site in human host. Small intestine (Toledo and Esteban 2016; Chai and Jung 2020).

Notes. Numerous nominal species are considered part of the “*E. revolutum* complex” or the “37-collar-spined *Echinostoma* complex”; taxonomic resolution of species within this group is an area of ongoing investigation.

Genus *Hypoderaeum* Dietz, 1909***Hypoderaeum conoideum* (Bloch, 1782)**

Geographic distribution. Southeast Asia, Japan, Europe, North and Central America (Toledo and Esteban 2016; Chai and Jung 2020).

Natural hosts. First intermediate hosts include several genera of freshwater snails. Second intermediate hosts include freshwater snails and amphibians. Definitive hosts are birds. Zoonotic in humans (Toledo and Esteban 2016; Chai and Jung 2020).

Route of infection. Ingestion of metacercariae infected intermediate hosts (Toledo and Esteban 2016; Chai and Jung 2020).

Site in human host. Small intestine (Toledo and Esteban 2016; Chai and Jung 2020).

Genus *Isthmiophora* Lühe, 1909***Isthmiophora hortensis* (Asada, 1926)**

Geographic distribution. China, Japan, Korea (Toledo and Esteban 2016; Chai and Jung 2020).

Natural hosts. First intermediate hosts include freshwater snails in the genera *Lymnaea* and *Radix*. Second intermediate hosts include several genera of freshwater fish. Definitive hosts are felids, canids, and rodents. Zoonotic in humans (Toledo and Esteban 2016; Chai and Jung 2020).

Route of infection. Ingestion of metacercariae in infected fish intermediate hosts (Toledo and Esteban 2016; Chai and Jung 2020).

Site in human host. Small intestine (Toledo and Esteban 2016; Chai and Jung 2020).

***Isthmiophora melis* (Schrank, 1788)**

Geographic distribution. Europe, USA, Taiwan (Toledo and Esteban 2016; Chai and Jung 2020).

Natural hosts. First intermediate hosts freshwater snails in the genus *Lymnaea*. Second intermediate hosts are freshwater fish and amphibians. Definitive hosts are canids, mustelids, hedgehogs, and rodents. Zoonotic in humans (Toledo and Esteban 2016; Chai and Jung 2020).

Route of infection. Ingestion of metacercariae in infected intermediate hosts (Toledo and Esteban 2016; Chai and Jung 2020).

Site in human host. Small intestine (Toledo and Esteban 2016; Chai and Jung 2020).

●●●●●●●● Fasciolidae Railliet, 1895**Genus *Fasciola* Linnaeus, 1758*****Fasciola gigantica* Cobbold, 1855**

Geographic distribution. Africa, Middle East, Southeast Asia, Japan (Tolan 2011; Phalee et al. 2015).

Natural hosts. First intermediate hosts are freshwater snails in the genus *Lymnaea*. Second intermediate hosts are aquatic plants. Definitive hosts are primarily sheep, also cattle, buffalo, goats, pigs. Zoonotic in humans (olan 2011; Phalee et al. 2015).

Route of infection. Ingestion of metacercariae on contaminated aquatic vegetation (olan 2011; Phalee et al. 2015).

Site in human host. Biliary ducts; ectopic migration to skin, brain (olan 2011; Phalee et al. 2015).

***Fasciola hepatica* Linnaeus, 1758**

Geographic distribution. Worldwide; hot spots of endemicity for human infection include the Bolivian Altiplano, Ecuador, Peru, Cuba, Portugal, Spain, Turkey, North Africa (Nile Delta), Iran, Vietnam (Tolan 2011).

Natural hosts. First intermediate hosts are freshwater snails, particularly members of the genera *Galba*, *Fossaria*, and *Pseudosuccinea*. Second intermediate hosts are aquatic plants. Definitive hosts include primarily cattle and buffalo, also sheep, goats, and deer. Zoonotic in humans (Tolan 2011).

Route of infection. Ingestion of metacercariae on contaminated aquatic vegetation (Tolan 2011).

Site in human host. Biliary ducts; ectopic migration to skin, brain (Tolan 2011).

Genus *Fasciolopsis* Looss, 1899***Fasciolopsis buski* (Lankester in Küchenmeister, 1857)**

Distoma rathouisi Poirier, 1887

Fasciolopsis fuelleborni Rodenwadt, 1909

Fasciolopsis goddardi Ward, 1910

Geographic distribution. Central and Southeast Asia (Sah et al. 2019; Chai and Jung 2020).

Natural hosts. First intermediate hosts are several genera of freshwater snails, especially the genera *Segmentina*, *Hippeutis*, and *Gyraulus*. Second intermediate hosts are freshwater plants. Definitive hosts are pigs, rabbits, and canids. Zoonotic in humans (Ma et al. 2017; Sah et al. 2019; Chai and Jung 2020).

Route of infection. Ingestion of metacercariae on contaminated freshwater vegetation (Sah et al. 2019; Chai and Jung 2020).

Site in human host. Small intestine (Sah et al. 2019; Chai and Jung 2020).

●●●●●●●●●● *Philophthalmidae* Looss, 1899**Genus *Philophthalmus* Looss, 1899*****Philophthalmus gralli* Matthis & Leger, 1910**

Philophthalmus anatinus Sugimoto, 1928

Philophthalmus nyrocae Yamaguti, 1934

Geographic distribution. East and Southeast Asia, North America, South America (Nollen and Kanev 1995).

Natural hosts. Intermediate hosts are snails in the genus *Melanooides*. Definitive hosts are birds, primarily water birds and fowl. Zoonotic in humans (Nollen and Kanev 1995).

Route of infection. Ingestion of metacercariae in water or on submerged substrates (e.g., mollusk shells, plants); also direct exposure of eyes to free-swimming cercariae in water (Nollen and Kanev 1995).

Site in human host. Eyes (Nollen and Kanev 1995).

***Philophthalmus palpebrarum* Looss, 1899**

Geographic distribution. Middle East, North Africa (Nollen and Kanev 1995).

Natural hosts. Intermediate hosts are snails in the genus *Melanooides*. Definitive hosts are birds, primarily water birds and fowl. Zoonotic in humans (Nollen and Kanev 1995).

Route of infection. Ingestion of metacercariae in water or on submerged substrates (e.g., mollusk shells, plants); also direct exposure of eyes to free-swimming cercariae in water (Nollen and Kanev 1995).

Site in human host. Eyes (Nollen and Kanev 1995).

***Philophthalmus lacrymosus* Braun, 1902**

Geographic distribution. Central and South America (Nollen and Kanev 1995).

Natural hosts. Intermediate hosts are snails in the genus *Melanoides*. Definitive hosts are birds, primarily water birds and fowl. Zoonotic in humans (Nollen and Kanev 1995).

Route of infection. Ingestion of metacercariae in water or on submerged substrates (e.g., mollusk shells, plants); also direct exposure of eyes to free-swimming cercariae in water (Nollen and Kanev 1995).

Site in human host. Eyes (Nollen and Kanev 1995).

●●●●●●●● **Hemiurata Skrjabin & Guschanskaja, 1954**

●●●●●●●● **Isoparorchiiidae Travassos, 1922**

Genus *Isoparorchis* Southwell, 1913

Leptolecithum Kobayashi, 1915

***Isoparorchis hypselobagri* (Billet, 1898)**

Geographic distribution. Central and Southeast Asia, Russia, Australia (Chai and Jung 2020).

Natural hosts. First intermediate hosts are freshwater snails in the genus *Melanoides*. Second intermediate hosts are catfish. Definitive hosts are predator fish. Zoonotic in humans (Chai and Jung 2020).

Route of infection. Ingestion of metacercariae in infected fish intermediate hosts (Chai and Jung 2020).

Site in human host. Small intestine (Chai and Jung 2020).

●●●●●●●● **Opisthorchiata La Rue, 1957**

●●●●●●●● **Heterophyidae Leiper, 1909**

Genus *Acanthotrema* Travassos, 1928

Acanthotrema felis Sohn et al., 2003

Geographic distribution. Korea (Chai and Jung 2020).

Natural hosts. First intermediate hosts are unknown. Second intermediate hosts are brackish water fish in the genus *Acanthogobius*. Definitive hosts are felids. Zoonotic in humans (Chai and Jung 2020).

Route of infection. Ingestion of metacercariae in infected fish intermediate host (Chai and Jung 2020).

Site in human host. Small intestine (Chai and Jung 2020).

Genus *Apophallus* Lühe, 1909

***Apophallus donicus* (Skrjabin & Lindtrop, 1919)**

Geographic distribution. North America (Chai and Jung 2020).

Natural hosts. First intermediate hosts are freshwater snails in the genus *Flumenicola*. Second intermediate hosts are various freshwater fish. Definitive hosts are carnivores and lagomorphs. Zoonotic in humans (Chai and Jung 2020).

Route of infection. Ingestion of metacercariae in infected fish intermediate host (Chai and Jung 2020).

Site in human host. Small intestine (Chai and Jung 2020).

Genus *Ascocotyle* Looss, 1899

***Ascocotyle longa* Ransom, 1920**

Geographic distribution. North America, Europe (Chai and Jung 2020).

Natural hosts. First intermediate hosts are brackish water snails in the genus *Heleobia*. Second intermediate hosts are brackish water fish in the genus *Mugil*. Definitive hosts are canids. Zoonotic in humans (Chai and Jung 2020).

Route of infection. Ingestion of metacercariae in infected fish (Chai and Jung 2020).

Site in human host. Small intestine (Chai and Jung 2020).

Genus *Centrocestus* Looss, 1899

***Centrocestus armatus* (Tanabe, 1922)**

Geographic distribution. Korea, Japan (Chai and Jung 2020).

Natural hosts. First intermediate hosts are freshwater snails in the genus *Semisulcospira*. Second intermediate hosts are freshwater fish in the genus *Zacco*. Definitive hosts are wild and domestic canids and felids, rabbits, rats, and birds. Zoonotic in humans (Chai and Jung 2020).

Route of infection. Ingestion of metacercariae in infected fish (Chai and Jung 2020).

Site in human host. Small intestine (Chai and Jung 2020).

***Centrocestus cuspidatus* (Looss, 1896)**

Geographic distribution. Southeast Asia, North Africa, Middle East (Chai and Jung 2020).

Natural hosts. First intermediate hosts are freshwater snails in the genus *Oncomelania*. Second intermediate hosts are freshwater fish in the genus *Gambusia*. Definitive hosts are carnivores, rodents, and birds. Zoonotic in humans (Chai and Jung 2020).

Route of infection. Ingestion of metacercariae in infected fish (Chai and Jung 2020).

Site in human host. Small intestine (Chai and Jung 2020).

***Centrocestus formosanus* Nishigori, 1924**

Centrocestus caninus Leiper, 1913

Geographic distribution. Southeast Asia, Central and South America, North Africa, Middle East (Chai and Jung 2020).

Natural hosts. First intermediate hosts are freshwater snails in the genus *Stenomelania*. Second intermediate hosts are freshwater fish, including *Cyclocheilichthys* and *Puntius*. Definitive hosts are wild and domestic canids, chickens, and ducks. Zoonotic in humans (Chai and Jung 2020).

Route of infection. Ingestion of metacercariae in infected fish (Chai and Jung 2020).

Site in human host. Small intestine (Chai and Jung 2020).

***Centrocestus kurokawai* (Kurowawa, 1935)**

Geographic distribution. Japan (Chai and Jung 2020).

Natural hosts. First intermediate hosts are unknown. Second intermediate hosts are unknown but presumed to be freshwater fish. Definitive hosts unknown; presumed zoonotic in humans (Chai and Jung 2020).

Route of infection. Ingestion of metacercariae in infected intermediate host (Chai and Jung 2020).

Site in human host. Small intestine (Chai and Jung 2020).

Genus *Cryptocotyle* Lühe, 1899***Cryptocotyle lingua* (Creplin, 1825)**

Geographic distribution. North America, Europe, Russia, Japan (Chai and Jung 2020).

Natural hosts. First intermediate hosts are brackish water snails in the genus *Littorina*. Second intermediate hosts are brackish and freshwater fish in the genus *Gobius*. Definitive hosts are carnivores, rodents, and birds. Zoonotic in humans (Chai and Jung 2020).

Route of infection. Ingestion of metacercariae in infected fish (Chai and Jung 2020).

Site in human host. Small intestine (Chai and Jung 2020).

Genus *Haplorchis* Looss, 1899***Haplorchis pumilio* (Looss, 1896)**

Geographic distribution. Central and Southeast Asia, Australia, Middle East, North Africa, Central and South America (Chai and Jung 2020).

Natural hosts. First intermediate hosts are snails in the genus *Melania*. Second intermediate hosts are various freshwater fish. Definitive hosts are carnivores. Zoonotic in humans (Chai and Jung 2020).

Route of infection. Ingestion of metacercariae in infected freshwater fish (Chai and Jung 2020).

Site in human host. Small intestine (Chai and Jung 2020).

***Haplorchis taichui* (Nishigori, 1924)**

Monorchotrema microrchia Katsuta, 1932

Geographic distribution. Central and Southeast Asia, Hawaii, Middle East (Chai and Jung 2020).

Natural hosts. First intermediate hosts are freshwater snails in the genera *Melania* and *Melanooides*. Second intermediate hosts are various freshwater fish. Definitive hosts are carnivores and birds. Zoonotic in humans (Chai and Jung 2020).

Route of infection. Ingestion of metacercariae in infected freshwater fish (Chai and Jung 2020).

Site in human host. Small intestine (Chai and Jung 2020).

***Haplorchis vanissimus* Africa, 1938**

Geographic distribution. Australia, Philippines (Chai and Jung 2020).

Natural hosts. First intermediate hosts are unknown. Second intermediate hosts are unknown but presumed to be freshwater fish. Definitive hosts are wild and domestic canids and felids and birds. Zoonotic in humans (Chai and Jung 2020).

Route of infection. Ingestion of metacercariae in infected fish (Chai and Jung 2020).

Site in human host. Small intestine (Chai and Jung 2020).

***Haplorchis yokogawai* (Katsuta, 1932)**

Geographic distribution. Central and Southeast Asia, Australia, Middle East, North Africa (Chai and Jung 2020).

Natural hosts. First intermediate hosts include several genera of freshwater snails. Second intermediate hosts are various freshwater fish. Definitive hosts are mammals, including cattle and carnivores. Zoonotic in humans (Chai and Jung 2020).

Route of infection. Ingestion of metacercariae in infected freshwater fish (Chai and Jung 2020).

Site in human host. Small intestine (Chai and Jung 2020).

Genus *Heterophyes* Cobbold, 1866

***Heterophyes heterophyes* (Siebold, 1853)**

Heterophyes aegyptiaca Cobbold, 1866

Geographic distribution. Central and Southeast Asia, Japan, Middle East, North Africa (Chai and Jung 2020).

Natural hosts. First intermediate hosts are various freshwater snails. Second intermediate hosts are various freshwater and brackish water fish. Definitive hosts are carnivores. Zoonotic in humans (Chai and Jung 2020).

Route of infection. Ingestion of metacercariae in infected fish (Chai and Jung 2020).

Site in human host. Small intestine (Chai and Jung 2020).

***Heterophyes nocens* Onji & Nishio, 1916**

Heterophyes katuradai Ozaki & Asada, 1926

Geographic distribution. Korea, Japan China (Chai and Jung 2020).

Natural hosts. First intermediate hosts are freshwater snails in the genus *Cerithidea*. Second intermediate hosts are freshwater fish in the genus *Acanthogobius*. Definitive hosts are wild and domestic and felids. Zoonotic in humans (Chai and Jung 2020).

Route of infection. Ingestion of metacercariae in infected fish (Chai and Jung 2020).

Site in human host. Small intestine (Chai and Jung 2020).

Genus *Heterophyopsis* Tubangui & Africa, 1938

***Heterophyopsis continua* (Onki & Nishio, 1916)**

Geographic distribution. Korea, Japan (Chai and Jung 2020).

Natural hosts. First intermediate hosts are unknown. Second intermediate hosts are various freshwater fish. Definitive hosts are carnivores and birds. Zoonotic in humans (Chai and Jung 2020).

Route of infection. Ingestion of metacercariae in infected fish (Chai and Jung 2020).

Site in human host. Small intestine (Chai and Jung 2020).

Genus *Metagonimus* Katsurada, 1912***Metagonimus katsuradai* Izumi, 1935**

Geographic distribution. Japan, Russia (Chai and Jung 2020).

Natural hosts. First intermediate hosts are freshwater snails in the genus *Semisulcospira*. Second intermediate hosts are freshwater fish in the genus *Tanakia*. Definitive hosts are carnivores. Zoonotic in humans (Chai and Jung 2020).

Route of infection. Ingestion of metacercariae in infected fish (Chai and Jung 2020).

Site in human host. Small intestine (Chai and Jung 2020).

***Metagonimus minutus* Katsuta, 1932**

Geographic distribution. Taiwan (Chai and Jung 2020).

Natural hosts. First intermediate hosts are unknown. Second intermediate hosts are brackish water fish in the genus *Mugil*. Definitive hosts are unknown; presumed zoonotic in humans (Chai and Jung 2020).

Route of infection. Ingestion of metacercariae in infected fish (Chai and Jung 2020).

Site in human host. Small intestine (Chai and Jung 2020).

***Metagonimus miyatai* Saito et al., 1997**

Geographic distribution. Korea, Japan (Chai and Jung 2020).

Natural hosts. First intermediate hosts are freshwater snails in the genus *Semisulcospira*. Second intermediate hosts are various freshwater fish. Definitive hosts are carnivores, rodents, and birds. Zoonotic in humans (Chai and Jung 2020).

Route of infection. Ingestion of metacercariae in infected fish (Chai and Jung 2020).

Site in human host. Small intestine (Chai and Jung 2020).

***Metagonimus takahashii* Takahashi, 1929**

Geographic distribution. Korea, Japan (Chai and Jung 2020).

Natural hosts. First intermediate hosts are freshwater snails in the genera *Semisulcospira* and *Koreanomelania*. Second intermediate hosts are various freshwater fish. Definitive hosts are carnivores and birds. Zoonotic in humans (Chai and Jung 2020).

Route of infection. Ingestion of metacercariae in infected fish (Chai and Jung 2020).

Site in human host. Small intestine (Chai and Jung 2020).

***Metagonimus yokogawai* (Katsurada, 1912)**

Geographic distribution. Central and Southeast Asia, Russia, Japan, Europe (Chai and Jung 2020).

Natural hosts. First intermediate hosts are freshwater snails in the genus *Semio-sulcospira*. Second intermediate hosts are various freshwater fish. Definitive hosts are carnivores, rodents, and birds. Zoonotic in humans (Chai and Jung 2020).

Route of infection. Ingestion of metacercariae in infected fish (Chai and Jung 2020).

Site in human host. Small intestine (Chai and Jung 2020).

Genus *Procerovum* Onji & Nishio, 1916

***Procerovum calderoni* (Africa & Garcia, 1935)**

Geographic distribution. Egypt, Southeast Asia (Chai and Jung 2020).

Natural hosts. First intermediate hosts are brackish water snails in the genus *Sermyla*. Second intermediate hosts are freshwater fish in the genus *Ophiocephalis*. Definitive hosts are wild and domestic canids and felids. Zoonotic in humans (Chai and Jung 2020).

Route of infection. Ingestion of metacercariae in infected fish (Chai and Jung 2020).

Site in human host. Small intestine (Chai and Jung 2020).

***Procerovum varium* Onji & Nishio, 1916**

Geographic distribution. Korea, Japan (Chai and Jung 2020).

Natural hosts. First intermediate hosts are brackish water snails in the genus *Melanoides*. Second intermediate hosts are brackish water fish in the genus *Mugil*. Definitive hosts are wild and domestic canids and felids. Zoonotic in humans (Chai and Jung 2020).

Route of infection. Ingestion of metacercariae in infected fish (Chai and Jung 2020).

Site in human host. Small intestine (Chai and Jung 2020).

Genus *Pygidiopsis* Looss, 1907

***Pygidiopsis genata* Looss, 1907**

Geographic distribution. North Africa, Middle East, Eastern Europe, Philippines (Chai and Jung 2020).

Natural hosts. First intermediate hosts are freshwater snails in the genera *Melanoides* and *Melanopsis*. Second intermediate hosts are various freshwater fish. Definitive hosts are carnivores, rodents, shrews, and birds. Zoonotic in humans (Chai and Jung 2020).

Route of infection. Ingestion of metacercariae in infected fish (Chai and Jung 2020).

Site in human host. Small intestine (Chai and Jung 2020).

***Pygidiopsis summa* Onji & Nishio, 1916**

Geographic distribution. Korea, Japan, Vietnam (Chai and Jung 2020).

Natural hosts. First intermediate hosts are unknown. Second intermediate hosts are brackish water fish in the genera *Mugil* and *Acnathogonius*. Definitive hosts are wild and domestic felids. Zoonotic in humans (Chai and Jung 2020).

Route of infection. Ingestion of metacercariae in infected fish (Chai and Jung 2020).

Site in human host. Small intestine (Chai and Jung 2020).

Genus *Stellantchasmus* Onji & Nishio, 1916

Stellantchasmus falcatus Onji & Nishio, 1916

Geographic distribution. Central and Southeast Asia, Japan, Middle East, Australia, Hawaii (Chai and Jung 2020).

Natural hosts. First intermediate hosts are freshwater snails in the genera *Stenomelania* and *Thiara*. Second intermediate hosts are mullets. Definitive hosts are wild and domestic canids and felids, rabbits, rats, and birds. Zoonotic in humans (Chai and Jung 2020).

Route of infection. Ingestion of metacercariae in infected fish (Chai and Jung 2020).

Site in human host. Small intestine (Chai and Jung 2020).

Genus *Stictodora* Looss, 1899

Stictodora fuscata (Onji & Nishio, 1916)

Geographic distribution. Korea, Japan, Kuwait (Chai and Jung 2020).

Natural hosts. First intermediate hosts are unknown. Second intermediate hosts are brackish water fish in the genera *Acanthogobius* and *Pseudorasbora*. Definitive hosts are wild and domestic canids and felids. Zoonotic in humans (Chai and Jung 2020).

Route of infection. Ingestion of metacercariae in infected fish (Chai and Jung 2020).

Site in human host. Small intestine (Chai and Jung 2020).

Stictodora lari Yamaguti, 1939

Geographic distribution. Southeast Asia, Japan, Russia, Australia (Chai and Jung 2020).

Natural hosts. First intermediate hosts are brackish water snails, including the genus *Velacumantus*. Second intermediate hosts are various brackish water fish. Definitive hosts are wild and domestic canids and felids. Zoonotic in humans (Chai and Jung 2020).

Route of infection. Ingestion of metacercariae in infected fish (Chai and Jung 2020).

Site in human host. Small intestine (Chai and Jung 2020).

●●●●●●●●●● **Opisthorchiidae Looss, 1899**

Genus *Amphimerus* Barker, 1911

Geographic distribution. North, Central, and South America (Calvopiña et al. 2011, 2015).

Natural hosts. Intermediate hosts are various freshwater snails. Second intermediate hosts are freshwater fish. Natural hosts are a variety of mammals, birds, and reptiles, including domestic dogs and cats. Zoonotic in humans (Calvopiña et al. 2011, 2015).

Route of infection. Ingestion of metacercariae in infected fish (Calvopiña et al. 2011).

Site in human host. Liver (Calvopiña et al. 2011).

Notes. The natural hosts and species identification have not been confirmed from human isolates, most of which have been documented in Ecuador.

Genus *Clonorchis* Looss, 1907

***Clonorchis sinensis* (Cobbold, 1875)**

Geographic distribution. East Asia; hot spots of endemicity for human disease include Korea, China, Taiwan, and Vietnam (Qian et al. 2012).

Natural hosts. First intermediate hosts are various genera of freshwater snails. Second intermediate host are a variety of freshwater fish, primarily cyprinids. Definitive hosts are fish-eating mammals, including wild and domestic canids and felids, mustelids, and pigs. Zoonotic in humans (Qian et al. 2012).

Route of infection. Ingestion of metacercariae in infected fish (Qian et al. 2012).

Site in human host. Biliary ducts (Qian et al. 2012).

Genus *Metorchis* Looss, 1899

***Metorchis bilis* (Braun, 1790)**

Distoma albidus Braun, 1893

Distoma crassiusculus Rudolphi, 1809

Geographic distribution. Central and Eastern Europe, Russia (Mordvinov et al. 2012).

Natural hosts. First intermediate hosts include snails of the genus *Bithynia*. Second intermediate hosts are typically cyprinid fishes (e.g., carps, minnows). Natural definitive hosts are various fish-eating mammals and birds. Zoonotic in humans (Mordvinov et al. 2012).

Route of infection. Ingestion of metacercariae in infected fish (Mordvinov et al. 2012).

Site in human host. Biliary ducts (Mordvinov et al. 2012).

***Metorchis conjunctus* (Cobbold, 1860)**

Geographic distribution. Northern North America (Behr et al. 1998).

Natural hosts. First intermediate hosts are freshwater snails in the genus *Amnicola*. Second intermediate hosts are freshwater fish, primarily members of the genus *Catostomus*. Definitive hosts are carnivores. Zoonotic in humans (Behr et al. 1998).

Route of infection. Ingestion of metacercariae in infected fish (MacLean et al. 1996; Behr et al. 1998).

Site in human host. Biliary ducts (MacLean et al. 1996; Behr et al. 1998).

***Metorchis orientalis* Tanabe, 1921**

Geographic distribution. East Asia (Na et al. 2016).

Natural hosts. First intermediate hosts include snails of the genera *Bithynia* and *Parafossarulus*. Second intermediate hosts are typically cyprinid fishes (e.g., carps, minnows). Natural definitive hosts are various fish-eating mammals and birds. Zoonotic in humans (Na et al. 2016).

Route of infection. Ingestion of metacercariae in infected fish (Na et al. 2016).

Site in human host. Biliary ducts (Na et al. 2016).

***Metorchis taiwanensis* Morishita & Tsuchimochi, 1952**

Geographic distribution. East Asia (Zhan et al. 2017).

Natural hosts. First intermediate hosts include snails of the genus *Bithynia*. Second intermediate hosts are typically cyprinid fishes (e.g., carps, minnows). Natural definitive hosts are various fish-eating mammals and birds. Zoonotic in humans (Zhan et al. 2017).

Route of infection. Ingestion of metacercariae in infected fish (Zhan et al. 2017).

Site in human host. Biliary ducts (Zhan et al. 2017).

Genus *Opisthorchis* Blanchard, 1895***Opisthorchis felineus* (Rivolta, 1884)**

Geographic distribution. Eastern Europe, Russia, Central Asia (Pakharukova and Mordvinov 2016).

Natural hosts. First intermediate hosts are freshwater snails in the genus *Bithynia*. Second intermediate hosts are various freshwater fish. Definitive hosts are mammals, primarily carnivores. Zoonotic in humans (Pakharukova and Mordvinov 2016).

Route of infection. Ingestion of metacercariae in infected fish (Pakharukova and Mordvinov 2016).

Site in human host. Biliary ducts (Pakharukova and Mordvinov 2016).

***Opisthorchis viverrini* (Poirier, 1886)**

Geographic distribution. Southeast Asia, with high prevalence of human infection in Thailand, Laos, Vietnam, and Cambodia (Kaewpitoon et al. 2008; Suwannatrai et al. 2018).

Natural hosts. First intermediate hosts are freshwater snails in the genus *Bithynia*. Second intermediate hosts are various freshwater fish. Definitive hosts are wild and domestic canids and felids. Zoonotic in humans (Kaewpitoon et al. 2008; Suwannatrai et al. 2018).

Route of infection. Ingestion of metacercariae in infected fish (Kaewpitoon et al. 2008).

Site in human host. Biliary ducts (Kaewpitoon et al. 2008).

Genus *Pseudamphistomum* Lühe, 1908***Pseudamphistomum truncatum* (Rudolphi, 1819)**

Geographic distribution. Europe, Asia (Neimanis et al. 2016).

Natural hosts. First intermediate hosts are freshwater and brackish snails. Second intermediate host are freshwater, brackish, and marine fish, primarily cyprinids. Definitive hosts are carnivores, including cats, pinnipeds, and mustelids. Zoonotic in humans (Neimanis et al. 2016).

Route of infection. Ingestion of metacercariae in infected fish (Neimanis et al. 2016).

Site in human host. Biliary ducts (Neimanis et al. 2016).

●●●●●●●● Pronocephalata Olson et al., 2003**●●●●●●●● Paramphistomidae Fiscoeder, 1901****Genus *Fiscoederius* Stiles & Goldberger, 1910*****Fiscoederius elongatus* (Poirier, 1883)**

Geographic distribution. Europe, Russia, Central and Southeast Asia, Japan (Chai and Jung 2020).

Natural hosts. First intermediate hosts are freshwater snails in the genus *Lymnaea*. Second intermediate hosts are freshwater plants. Definitive hosts are cattle, deer, goats, and sheep. Zoonotic in humans (Chai and Jung 2020).

Route of infection. Ingestion of metacercariae on contaminated aquatic vegetation (Chai and Jung 2020)

Site in human host. Small intestine (Chai and Jung 2020).

Genus *Gastrodiscoides* Leiper, 1913***Gastrodiscoides hominis* (Lewis & McConnell, 1876)**

Geographic distribution. Europe, Central and Southeast Asia (Mas-Coma 2006; Chai and Jung 2020).

Natural hosts. First intermediate hosts are freshwater snails in the genus *Helicorbis*. Second intermediate hosts are freshwater plants, amphibians, and crayfish. Definitive hosts are pigs and humans (Mas-Coma 2006; Chai and Jung 2020).

Route of infection. Ingestion of metacercariae in infected intermediate hosts (Mas-Coma 2006; Chai and Jung 2020).

Site in human host. Small intestine (Mas-Coma 2006; Chai and Jung 2020).

Genus *Watsonius* Stiles & Goldberger, 1910***Watsonius watsoni* (Conyngham, 1904)**

Geographic distribution. Sub-Saharan Africa, Vietnam (Chai and Jung 2020).

Natural hosts. First intermediate hosts are freshwater snails in the genus *Physa*. Second intermediate hosts are presumed to be freshwater vegetation. Definitive hosts are non-human primates. Zoonotic in humans (Chai and Jung 2020).

Route of infection. Ingestion of metacercariae on contaminated aquatic vegetation (Chai and Jung 2020).

Site in human host. Small intestine (Chai and Jung 2020).

●●●●●●●● **Xiphidiata Olson et al., 2003**●●●●●●●● **Dicrocoeliidae Looss, 1899****Genus *Dicrocoelium* Dujardin, 1845*****Dicrocoelium dendriticum* (Rudolphi, 1819)**

Dicrocoelium lanceolatum Stiles & Hassall, 1898

Geographic distribution. North America, Africa, Asia, Europe, Middle East (Cengiz et al. 2010).

Natural hosts. First intermediate hosts are various genera of terrestrial snails. Second intermediate hosts are ants (primarily genus *Formica*). Definitive hosts are ungulates, including cattle, sheep, and goats. Zoonotic in humans (Cengiz et al. 2010).

Route of infection. Ingestion of metacercariae in infected ants (Cengiz et al. 2010).

Site in human host. Bile ducts, gall bladder (Cengiz et al. 2010).

***Dicrocoelium hospes* Looss, 1907**

Geographic distribution. West Africa (Odei 1966).

Natural hosts. First intermediate hosts are snails in the genera *Achatina* and *Limicolaria*. Second intermediate hosts are ants in the genera *Dorylus* and *Crematogaster*. Definitive hosts are ungulates, including cattle and sheep. Zoonotic in humans (Odei 1966).

Route of infection. Ingestion of infected intermediate hosts (Odei 1966).

Site in human host. Bile ducts (Odei 1966)

Notes. It is uncertain whether *D. hospes* causes true infection in humans. Most reported cases of the finding of eggs in stool are believed to be spurious after the consumption of infected beef liver (Wolfe 2007). The two cases from Ghana believed to be true infections are in children who denied eating beef liver (Odei 1966). True infection should be confirmed by the examination of follow-up stool specimens (Wolfe 2007).

●●●●●●●● **Lecithodendriidae Lühe, 1901**

Genus *Caprimolgorchis* Jha, 1943***Caprimolgorchis molenkampii* Lie Kian Joe, 1961**

Geographic distribution. Southeast Asia (Chai and Jung 2020).

Natural hosts. First intermediate hosts are unknown, but hypothesized to be freshwater snails in the genera *Bithynia* or *Zebrina*. Second intermediate hosts are dragonflies and damselflies. Definitive hosts are rodents and bats. Zoonotic in humans (Chai and Jung 2020).

Route of infection. Ingestion of infected insects (Chai and Jung 2020).

Site in human host. Small intestine (Chai and Jung 2020).

●●●●●●●● **Microphalidae Ward, 1901**

Genus *Gynaecotyla* Yamaguti, 1939***Gynaecotyla squatarolae* (Yamaguti, 1934)**

Geographic distribution. Japan, South Korea, Taiwan (Chai and Jung 2020).

Natural hosts. First intermediate hosts are brackish water snails in the genus *Batillaria*. Second intermediate hosts are brackish water crabs in the genus *Macrophthalmus*. Definitive hosts are shorebirds. Zoonotic in humans (Chai and Jung 2020).

Route of infection. Ingestion of metacercariae in infected crabs (Chai and Jung 2020).

Site in human host. Small intestine (Chai and Jung 2020).

Genus *Microphallus* Ward, 1901***Microphallus brevicaeca* (Africa & Garcia, 1935)**

Geographic distribution. Papua New Guinea, Philippines (Chai and Jung 2020).

Natural hosts. First intermediate hosts are unknown. Second intermediate hosts are brackish water crustaceans in the genera *Carcinus* and *Macrobrachium*. Definitive hosts are birds and mammals, especially non-human primates. Zoonotic in humans (Chai and Jung 2020).

Route of infection. Ingestion of metacercariae in infected crustaceans (Chai and Jung 2020).

Site in human host. Small intestine (Chai and Jung 2020).

●●●●●●●● **Phaenerosolidae Mehra, 1935****Genus *Phaneropsolus* Looss, 1899*****Phaneropsolus bonnei* Lie Kian Joe, 1951**

Geographic distribution. Central and Southeast Asia (Chai and Jung 2020).

Natural hosts. First intermediate hosts are unknown, but hypothesized to be freshwater snails in the genus *Bithynia*. Second intermediate hosts are dragonflies and damselflies. Definitive hosts are non-human primates. Zoonotic in humans (Chai and Jung 2020).

Route of infection. Ingestion of infected insects (Chai and Jung 2020).

Site in human host. Small intestine (Chai and Jung 2020).

***Phaneropsolus spinicirrus* Kaewkes et al., 1991**

Geographic distribution. Thailand (Chai and Jung 2020).

Natural hosts. First intermediate hosts are unknown. Second intermediate hosts are unknown, but presumed to be dragonflies and damselflies. Definitive hosts are unknown; presumed zoonotic in humans (Chai and Jung 2020).

Route of infection. Ingestion of infected intermediate hosts (Chai and Jung 2020).

Site in human host. Small intestine (Chai and Jung 2020).

●●●●●●●● **Plagiorchiidae Lühe, 1901****Genus *Plagiorchis* Lühe, 1899*****Plagiorchis javensis* Sandground, 1940**

Geographic distribution. Indonesia (Chai and Jung 2020).

Natural hosts. First intermediate hosts are unknown. Second intermediate hosts include snails, insects, and possibly fish. Definitive hosts are birds and bats. Zoonotic in humans (Chai and Jung 2020).

Route of infection. Ingestion of metacercariae in infected intermediate hosts (Chai and Jung 2020).

Site in human host. Small intestine (Chai and Jung 2020).

***Plagiorchis muris* (Tanabe, 1922)**

Geographic distribution. Europe, Central and Southeast Asia, Japan, North and Central America (Chai and Jung 2020).

Natural hosts. First intermediate hosts are freshwater snails in the genus *Lymnaea*. Second intermediate hosts are freshwater insects, crustaceans, and fish. Definitive hosts are canids, felids, raccoons, rodents, bats, and birds. Zoonotic in humans (Chai and Jung 2020).

Route of infection. Ingestion of metacercariae in infected intermediate hosts (Chai and Jung 2020).

Site in human host. Small intestine (Chai and Jung 2020).

***Plagiorchis philippinensis* Africa & Garcia, 1937**

Geographic distribution. Philippines (Chai and Jung 2020).

Natural hosts. First intermediate hosts are unknown. Second intermediate hosts are presumed to be aquatic insects. Definitive hosts rodents and possibly birds. Zoonotic in humans (Chai and Jung 2020).

Route of infection. Ingestion of metacercariae in infected intermediate hosts (Chai and Jung 2020).

Site in human host. Small intestine (Chai and Jung 2020).

***Plagiorchis vespertilionis* (Müller, 1784)**

Geographic distribution. Europe, North Africa, Central and Southeast Asia, Japan, Madagascar, North America (Chai and Jung 2020).

Natural hosts. First intermediate hosts are freshwater snails in the genus *Lymnaea*. Second intermediate hosts are freshwater insects. Definitive hosts are rodents and bats. Zoonotic in humans (Chai and Jung 2020).

Route of infection. Ingestion of metacercariae in infected insect intermediate hosts (Chai and Jung 2020).

Site in human host. Small intestine (Chai and Jung 2020).

●●●●●●●● Troglotremata Schell, 1890

●●●●●●●● Nanophyetidae Dollfus, 1939

Genus *Nanophyetus* Chapin, 1928***Nanophyetus salmincola* (Chapin, 1926)**

Geographic distribution. Northern North America (Chai and Jung 2020).

Natural hosts. First intermediate hosts are snails in the genus *Oxytrema*. Second intermediate hosts are several general of freshwater fish, especially salmonids. Definitive hosts canids, felids, raccoons, and birds. Zoonotic in humans (Chai and Jung 2020).

Route of infection. Ingestion of metacercariae in infected intermediate hosts (Chai and Jung 2020).

Site in human host. Small intestine (Chai and Jung 2020).

***Nanophyetus schikhobalowi* Skrjabin & Podiapolskaia, 1931**

Geographic distribution. Northern Eurasia (Voronova et al. 2017; Voronova and Chelomina 2018).

Natural hosts. First intermediate hosts are snails in the genus *Oxytrema*. Second intermediate hosts are several general of freshwater fish, especially salmonids. Definitive hosts canids, felids, raccoons, and birds. Zoonotic in humans (Voronova et al. 2017; Voronova and Chelomina 2018).

Route of infection. Ingestion of metacercariae in infected intermediate hosts (Voronova et al. 2017; Voronova and Chelomina 2018).

Site in human host. Small intestine (Voronova et al. 2017; Voronova and Chelomina 2018).

●●●●●●●● Paragonimidae Dollfus, 1939**Genus *Paragonimus* Braun, 1899*****Paragonimus africanus* Volker & Vogel, 1965**

Geographic distribution. West Africa (Blair et al. 1999; Cumberlidge et al. 2018).

Natural hosts. First intermediate hosts are presumed to be freshwater or brackish snails, but the specific species have yet to be identified. Second intermediate hosts are freshwater crabs in the family Potamidae. Definitive hosts are carnivores and non-human primates. Zoonotic in humans (Blair et al. 1999; Cumberlidge et al. 2018).

Route of infection. Ingestion of metacercariae in infected intermediate or paratenic hosts (Blair et al. 1999; Cumberlidge et al. 2018).

Site in human host. Lungs, possibly ectopic infection at other sites (Blair et al. 1999; Cumberlidge et al. 2018).

***Paragonimus heterotremus* Chen & Hsia, 1964**

Paragonimus tuanshansis Chung et al., 1964

Paragonimus pseudoheterotremus Waikagul et al., 2007

Geographic distribution. Southeast Asia (Yoshida et al. 2019).

Natural hosts. First intermediate hosts are freshwater snails in the superfamily Rissooidea. Second intermediate hosts are freshwater crabs in the families Potamidae and Parathelphusidae. Definitive hosts are carnivores and non-human primates. Wild boar, rodents, and deer can serve as paratenic hosts. Zoonotic in humans (Yoshida et al. 2019).

Route of infection. Ingestion of metacercariae in infected intermediate or paratenic hosts (Yoshida et al. 2019).

Site in human host. Lungs, possibly ectopic infection at other sites (Yoshida et al. 2019).

Notes. *Paragonimus heterotremus* is often referred to as a species complex; the status of many members as distinct and valid species is an area of ongoing investigation.

***Paragonimus kellicotti* Ward, 1908**

Geographic distribution. Eastern and Midwestern North America (Procop 2009).

Natural hosts. First intermediate hosts are freshwater snails in the family Pomatiopsidae. Second intermediate hosts are freshwater crayfish, primarily of the family Astacidae. Definitive hosts are carnivores and marsupials. Zoonotic in humans (Procop 2009).

Route of infection. Ingestion of metacercariae in infected intermediate and paratenic hosts (Procop 2009).

Site in human host. Lungs, possibly ectopic infection at other sites (Procop 2009).

***Paragonimus mexicanus* Miyazaki & Ishii, 1968**

Paragonimus peruvianus Miyazaki et al., 1969

Paragonimus ecuadorensis Voekler & Arzube, 1979

Geographic distribution. Central and South America (Blair et al. 1999).

Natural hosts. First intermediate hosts are freshwater and brackish snails in the families Hydrobiidae and Pomatiopsidae. Second intermediate hosts are freshwater crabs in the families Pseudothelphusidae and Trichodactylidae. Definitive hosts are carnivores and marsupials. Zoonotic in humans (Blair et al. 1999).

Route of infection. Ingestion of metacercariae in infected crustaceans (Blair et al. 1999).

Site in human host. Lungs, possibly ectopic infection at other sites (Blair et al. 1999).

Paragonimus ohirai* Miyazaki, 1939Paragonimus iloktsuensis* Chen, 1940*Paragonimus sadoensis* Miyazaki et al., 1968**Geographic distribution.** East Asia (Blair et al. 1999; Yoshida et al. 2019).**Natural hosts.** First intermediate hosts are freshwater and brackish snails in the families Assimineidae and Pomatiopsidae. Second intermediate hosts are freshwater crabs in the family Potamidae and brackish crabs in the family Grapsidae. Definitive hosts are carnivores. Zoonotic in humans (Blair et al. 1999; Yoshida et al. 2019).**Route of infection.** Ingestion of metacercariae in infected intermediate or paratenic hosts (Blair et al. 1999; Yoshida et al. 2019).**Site in human host.** Lungs, possibly ectopic infection at other sites (Blair et al. 1999; Yoshida et al. 2019).**Notes.** *Paragonimus ohirai* is often referred to as a species complex; the status of many members as distinct and valid species is an area of ongoing investigation.***Paragonimus skrjabini* Chen, 1959***Paragonimus miyazakii* Kamo et al., 1961*Paragonimus szechuanensis* Chung & Tsao, 1962*Paragonimus hueitungensis* Chung et al., 1975*Paragonimus veocularis* Chen & Li, 1979**Geographic distribution.** East and Southeast Asia, Indian subcontinent (Blair et al. 1999; Yoshida et al. 2019).**Natural hosts.** First intermediate hosts are freshwater and brackish snails in the superfamily Rissooidea. Second intermediate hosts are freshwater crabs in the families Potamidae and Parathelphusidae. Definitive hosts are carnivores. Wild boar, rodents, and deer may serve as paratenic hosts. Zoonotic in humans (Blair et al. 1999; Yoshida et al. 2019).**Route of infection.** Ingestion of metacercariae in infected intermediate and paratenic hosts (Blair et al. 1999; Yoshida et al. 2019).**Site in human host.** Lungs, possibly ectopic infection at other sites (Blair et al. 1999; Yoshida et al. 2019).**Notes.** *Paragonimus skrjabini* is often referred to as a species complex; the status of many members as distinct and valid species is an area of ongoing investigation.***Paragonimus uterobilateralis* Volker & Vogel, 1965****Geographic distribution.** West Africa (Blair et al. 1999; Cumberlidge et al. 2018).**Natural hosts.** First intermediate hosts are presumed to be freshwater or brackish snails, but the specific species have yet to be identified. Second intermediate hosts are

freshwater crabs in the family Potamidae. Definitive hosts are carnivores. Zoonotic in humans (Blair et al. 1999; Cumberlidge et al. 2018).

Route of infection. Ingestion of metacercariae in infected intermediate or paratenic hosts (Blair et al. 1999; Cumberlidge et al. 2018).

Site in human host. Lungs, possibly ectopic infection at other sites (Blair et al. 1999; Cumberlidge et al. 2018).

***Paragonimus westermani* (Kerbert, 1878)**

Distoma pulmonalis Baelz, 1880

Distoma ringeri Cobbold, 1880

Paragonimus edwardsi Gulati, 1926

Paragonimus macacae Sandosham, 1953

Paragonimus asymmetricus Chen, 1977

Paragonimus filipinus Miyazaki, 1978

Paragonimus philippinensis Ito et al., 1978

Geographic distribution. East and Southeast Asia, Indian subcontinent, Siberia (Blair et al. 1999; YBlair 2019; oshida et al. 2019).

Natural hosts. First intermediate hosts are freshwater and brackish snails in the superfamily Cerithioidea. Second intermediate hosts are freshwater crabs in the families Potamidae and Parathelphusidae. Definitive hosts are carnivores and non-human primates. Wild boar, rodents, and deer may serve as paratenic hosts. Zoonotic in humans (Blair et al. 1999; Blair 2019; Yoshida et al. 2019).

Route of infection. Ingestion of metacercariae in infected intermediate or paratenic hosts (Blair et al. 1999; Blair 2019; Yoshida et al. 2019).

Site in human host. Lungs, possibly ectopic infection at other sites (Blair et al. 1999; Blair 2019; Yoshida et al. 2019).

Notes. *Paragonimus westermani* is often referred to as a species complex; the status of many members as distinct and valid species is an area of ongoing investigation.

●●●●●●●● **Orchipedidae Skrjabin, 1913**

Achillurbainiidae Dollfus, 1939

Genus *Achillurbainia* Dollfus, 1939

Poikilorchis Fain & Vandepitte, 1957

***Achillurbainia congolensis* (Fain & Vandepitte, 1957)**

Geographic distribution. Sub-Saharan Africa (Fain and Vandepitte 1957; Nieuwenhuyse and Gatti 1968).

Natural hosts. Unknown; presumed zoonotic in humans (Fain and Vandepitte 1957; Nieuwenhuysen and Gatti 1968).

Route of infection. Unknown.

Site in human host. Subcutaneous cysts (usually near the ear), mastoid, middle ear (Fain and Vandepitte 1957; Nieuwenhuysen and Gatti 1968).

***Achillurbainia nouveli* Dolffus, 1939**

Geographic distribution. Southeast Asia (human case from China) (Kannangara 1971).

Natural hosts. First intermediate hosts are unknown. Second intermediate hosts are freshwater crabs (*Paratelphusa*). Definitive hosts are leopards. Zoonotic in humans (Kannangara 1971; Kwo and Lim 1968).

Route of infection. Unknown, presumed ingestion of metacercariae in under-cooked crabs (Kannangara 1971).

Site in human host. Subcutaneous cysts (near the ear) (Kannangara 1971).

***Achillurbainia recondita* Travassos, 1942**

Geographic distribution. Central and South America (Beaver et al. 1977).

Natural hosts. Intermediate hosts unknown. Definitive hosts are opossums (*Didelphis marsupialis*). Zoonotic in humans (Beaver et al. 1977).

Route of infection. Unknown.

Site in human host. Peritoneal cavity (Beaver et al. 1977).

- Ecdysozoa Anguinaldo et al., 1997
- Panarthropoda Nielsen, 1995
- Arthropoda von Siebold, 1848
- Chelicerata Heymons, 1901
- Euchelicerata Weygoldt & Paulus, 1979
- Arachnida Lamarck, 1801
- Acari Leach, 1817
- Acariformes Zakhvatkin, 1952
- Sarcoptiformes Reuter, 1909
- Astigmatina Canestrini, 1891
- Acaridia Latreille, 1802
- Histiotomatidae Berlese, 1897

Histiotomatidae, incertae sedis

Geographic distribution. Unknown (single human case from Saudi Arabia) (Al-Arfaj et al. 2007).

Natural hosts. Unknown

Natural hosts. Rodents and insectivores. Zoonotic on humans as incidental hosts (Diaz 2009; Elliott et al. 2019).

Route of infection. Exposure to larvae in the environment (Diaz 2009; Elliott et al. 2019).

Site in human host. Skin.

Vectored pathogens. *Orientia tsutsugamushi* (Oriental scrub typhus) (Diaz 2009; Elliott et al. 2019).

Leptotrombidium arenicola (Traub, 1960)

Geographic distribution. Malaysia (Diaz 2009; Elliott et al. 2019).

Natural hosts. Rodents and insectivores. Zoonotic on humans as incidental hosts (Diaz 2009; Elliott et al. 2019).

Route of infection. Exposure to larvae in the environment (Diaz 2009; Elliott et al. 2019).

Site in human host. Skin.

Vectored pathogens. *Orientia tsutsugamushi* (Oriental scrub typhus) (Diaz 2009; Elliott et al. 2019).

Leptotrombidium deliense (Walch, 1922)

Geographic distribution. Southeast Asia, Japan, Philippines, Australia, Pacific Islands (Lv et al. 2018).

Natural hosts. Rodents and insectivores. Zoonotic on humans as incidental hosts (Lv et al. 2018).

Route of infection. Exposure to larvae in the environment (Lv et al. 2018).

Site in human host. Skin.

Vectored pathogens. *Orientia tsutsugamushi* (Oriental scrub typhus) (Lv et al. 2018).

Leptotrombidium fletcheri (Womersley & Heaslip, 1943)

Geographic distribution. Malaysia (Diaz 2009; Elliott et al. 2019).

Natural hosts. Rodents and insectivores. Zoonotic on humans as incidental hosts (Diaz 2009; Elliott et al. 2019).

Route of infection. Exposure to larvae in the environment (Diaz 2009; Elliott et al. 2019).

Site in human host. Skin.

Vectored pathogens. *Orientia tsutsugamushi* (Oriental scrub typhus) (Diaz 2009; Elliott et al. 2019).

Leptotrombidium pallidum (Nagayo et al., 1919)

Geographic distribution. Japan (Diaz 2009; Elliott et al. 2019).

Natural hosts. Rodents and insectivores. Zoonotic on humans as incidental hosts (Diaz 2009; Elliott et al. 2019).

Route of infection. Exposure to larvae in the environment (Diaz 2009; Elliott et al. 2019).

Site in human host. Skin

Vectored pathogens. *Orientia tsutsugamushi* (Oriental scrub typhus) (Diaz 2009; Elliott et al. 2019).

Leptotrombidium pavlovskyi (Schluger, 1948)

Geographic distribution. Eastern Russia (Diaz 2009; Elliott et al. 2019).

Natural hosts. Rodents and insectivores. Zoonotic on humans as incidental hosts (Diaz 2009; Elliott et al. 2019).

Route of infection. Exposure to larvae in the environment (Diaz 2009; Elliott et al. 2019).

Site in human host. Skin

Vectored pathogens. *Orientia tsutsugamushi* (Oriental scrub typhus) (Diaz 2009; Elliott et al. 2019).

Leptotrombidium scutellaris (Nagayo et al., 1921)

Geographic distribution. Japan (Diaz 2009; Elliott et al. 2019).

Natural hosts. Rodents and insectivores. Zoonotic on humans as incidental hosts (Diaz 2009; Elliott et al. 2019).

Route of infection. Exposure to larvae in the environment (Diaz 2009; Elliott et al. 2019).

Site in human host. Skin

Vectored pathogens. *Orientia tsutsugamushi* (Oriental scrub typhus) (Diaz 2009; Elliott et al. 2019).

Genus *Neotrombicula* Hirst, 1925

Neotrombicula autumnalis (Shaw, 1790)

Geographic distribution. Europe (Schöler et al. 2006).

Natural hosts. Mammals and birds. Zoonotic on humans as incidental hosts (Schöler et al. 2006).

Route of infection. Exposure to larvae in the environment (Diaz 2009).

Site in human host. Skin.

Vectored pathogens. None.

Site in human host. Skin (Rosen et al. 2002).

Vectored pathogens. None.

Pyemotes ventricosus (Newport, 1850)

Geographic distribution. Southern Europe (Moser 1975).

Natural hosts. Insects. Zoonotic on humans as incidental hosts (Moser 1975).

Route of infection. Environmental exposure (Diaz 2009).

Site in human host. Skin (Diaz 2009).

Vectored pathogens. None.

●●●●●●●●●● Parazitiformes Leach, 1815

●●●●●●●●●● Ixodoidea Leach, 1815

●●●●●●●●●● Ixodidae Koch, 1844

Genus *Amblyomma* Koch, 1844

Amblyomma americanum (Linnaeus, 1758)

Ixodes unipunctata Packard, 1869

Geographic distribution. Eastern North America (Cooley 1944; Guglielmo and Robbins 2018).

Natural hosts. Many birds and mammals, including humans (Cooley 1944; Guglielmo and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. *Ehrlichia chaffeensis* (human monocytic ehrlichiosis), *E. ewingii* (human granulocytic ehrlichiosis), *Francisella tularensis* (tularemia), Heartland virus, Bourbon virus; also implicated in alpha-gal syndrome and Southern Tick-Associated Rash Illness (STARI) (Madison-Antenucci et al. 2020).

Amblyomma aureolatum (Pallas, 1772)

Amblyomma striatum Koch, 1844

Geographic distribution. South America (Guglielmo and Robbins 2018).

Natural hosts. Mammals, including carnivores and rodents, and birds. Zoonotic on humans (Guglielmo and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. *Rickettsia rickettsii* (Rocky Mountain Spotted Fever) (Szabó et al. 2013).

Amblyomma babirussae Schülze, 1933

Geographic distribution. Indonesia (Guglielmone and Robbins 2018).

Natural hosts. Mammals, including pigs, bovids, deer, carnivores, and rodents. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Amblyomma brasiliense Aragão, 1908

Geographic distribution. South America (Guglielmone and Robbins 2018).

Natural hosts. Several mammals and birds. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Amblyomma breviscutatum Neumann, 1899

Amblyomma cyprium Koch & Neumann, 1899

Amblyomma cyprium aeratipes Schülze, 1932

Geographic distribution. Southeast Asia, Australia, Pacific Islands (Guglielmone and Robbins 2018).

Natural hosts. Immature stages primarily on rodents and small birds. Adults are primarily on mammals, including bovids, deer, pigs, horses. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Amblyomma cajennense (Fabricius, 1787)

Geographic distribution. Southern United States, Central and South America, Caribbean (Cooley 1944; Martins et al. 2016; Guglielmone and Robbins 2018).

Natural hosts. Many mammals, including humans, and birds (Cooley 1944; Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Amblyomma calcartum* Neumann, 1899**

Geographic distribution. Central and South America, Caribbean (Guglielmone and Robbins 2018).

Natural hosts. Mammals, primarily anteaters; also birds. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Amblyomma clypeolatum* Neumann, 1899**

Geographic distribution. India, Sri Lanka, Myanmar (Liyanaarachchi et al. 2015).

Natural hosts. Reptiles. Zoonotic on humans (Liyanaarachchi et al. 2015; Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Amblyomma coelebs* Neumann, 1899**

Geographic distribution. Central and South America (Guglielmone and Robbins 2018).

Natural hosts. Several groups of mammals and birds. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Amblyomma cobraerens* Dönitz, 1909**

Geographic distribution. Sub-Saharan Africa (Guglielmone and Robbins 2018).

Natural hosts. Primarily bovids, but also other mammals and birds. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Amblyomma cordiferum* Neumann, 1899**

Geographic distribution. Southeast Asia (Audy 1960).

Natural hosts. Immature stages are primarily on rodents. Adults are on reptiles and ungulates. Zoonotic on humans (Audy 1960).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Notes. The single record of *A. cordiferum* from a human in Malaysia (Audy 1960) is considered a tentative identification (Guglielmono and Robbins 2018).

***Amblyomma dissimile* Koch, 1844**

Geographic distribution. North, Central, and South America and Caribbean (Cooley 1944; Guglielmono and Robbins 2018).

Natural hosts. Primarily reptiles and amphibians, but also a variety of mammals and birds. Zoonotic on humans (Cooley 1944; Guglielmono and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Amblyomma dubitatum* Neumann, 1899**

Amblyomma cooperi Nuttall & Warburton, 1908

Geographic distribution. South America (Guglielmono and Robbins 2018).

Natural hosts. Primarily rodents, but also other mammals and birds. Zoonotic on humans (Guglielmono and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Amblyomma falsomarmoreum* Tonelli-Rondelli, 1935**

Geographic distribution. Sub-Saharan Africa (Guglielmono and Robbins 2018).

Natural hosts. Reptiles, occasionally other mammals. Zoonotic on humans (Guglielmono and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Amblyomma fuscum* Neumann, 1907**

Geographic distribution. Brazil (Guglielmono and Robbins 2018).

Natural hosts. Immature stages on mammals, including rodents, carnivores, and marsupials. Adults usually on reptiles and amphibians. Zoonotic on humans (Guglielmono and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Amblyomma gemma* Dönitz, 1909**

Geographic distribution. Sub-Saharan Africa (Guglielmone and Robbins 2018).

Natural hosts. Immature stages on small mammals and birds. Adults primarily on bovids, pigs, camels, and horses, but also other mammals, large birds, and reptiles. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Amblyomma geomydae* (Cantor, 1847)**

Amblyomma malayanum Neumann, 1906

Geographic distribution. Southeast Asia (Guglielmone and Robbins 2018).

Natural hosts. Immature stages on a variety of small mammals, birds, and reptiles. Adults primarily on tortoises, but also several mammals, including pigs, bovids, deer. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Amblyomma hadanii* Nava et al., 2014**

Geographic distribution. Argentina (Guglielmone and Robbins 2018).

Natural hosts. Primarily bovids, tapirs, horses, also carnivores. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Amblyomma hebraeum* Koch, 1844**

Geographic distribution. Sub-Saharan Africa (Guglielmone and Robbins 2018).

Natural hosts. Mammals, including bovids, goats, sheep, other ruminants, and humans; occasionally birds and reptiles (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. *Rickettsia africae* (African tick bite fever) (Jongejan et al. 2020).

***Amblyomma incisum* Neumann, 1906**

Geographic distribution. South America (Guglielmone and Robbins 2018).

Natural hosts. Mammals, primarily tapirs, but also rodents, deer, and carnivores. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Amblyomma inoratum (Banks, 1909)

Geographic distribution. North and Central America (Guglielmone and Robbins 2018).

Natural hosts. Several groups of mammals and birds. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Amblyomma integrum Karsch, 1879

Geographic distribution. India, Sri Lanka (Guglielmone and Robbins 2018).

Natural hosts. Primarily bovids, but also other mammals and birds. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin, ears (Dilrukshi et al. 2004).

Vectored pathogens. None.

Amblyomma javanense (Supino, 1897)

Aponomma sublaeve Neumann, 1899

Geographic distribution. Central, Southeastern Asia (Guglielmone and Robbins 2018).

Natural hosts. Mammals, primarily pangolins, but also rodents, and carnivores, and reptiles. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Amblyomma latepunctatum Tonelli-Rondelli, 1939

Geographic distribution. South America (Guglielmone and Robbins 2018).

Natural hosts. Mammals, primarily tapirs and peccaries, also marsupials and rodents. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Amblyomma latum* Koch, 1844**

Geographic distribution. Sub-Saharan Africa (Guglielmone and Robbins 2018).

Natural hosts. Primarily reptiles, also amphibians, rodents, and shrews. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Amblyomma lepidum* Dönitz, 1909**

Geographic distribution. Sub-Saharan Africa, Middle East (Guglielmone and Robbins 2018).

Natural hosts. Primarily bovids, but also other mammals and birds. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Amblyomma limbatum* Neumann, 1899**

Geographic distribution. Australia (Guglielmone and Robbins 2018).

Natural hosts. Reptiles. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Amblyomma loculosum* Neumann, 1907**

Geographic distribution. Sub-Saharan Africa, Australia, several islands in the Indian and South Pacific Oceans (Guglielmone and Robbins 2018).

Natural hosts. Primarily birds and reptiles, occasionally bovids. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Amblyomma longirostre* (Koch, 1844)**

Geographic distribution. Central and South America (Guglielmone and Robbins 2018).

Natural hosts. Primarily birds (immature stages) and New World porcupines (adults), but also a variety of mammals and birds. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.
Site in human host. Skin.
Vectored pathogens. None.

***Amblyomma maculatum* Koch, 1844**

Geographic distribution. Southern USA, Central and South America (Cooley 1944; Guglielmone and Robbins 2018; Lado et al. 2018).

Natural hosts. Many reptiles, birds, and mammals, including humans (Cooley 1944; Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. *Rickettsia parkeri* (tidewater spotted fever) (Sumner et al. 2007; Lee et al. 2019).

***Amblyomma marmoreum* Koch, 1844**

Geographic distribution. Sub-Saharan Africa (Guglielmone and Robbins 2018).

Natural hosts. Immature stages primarily on birds and mammals. Adults on tortoises. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Amblyomma mixtum* Koch, 1844**

Geographic distribution. Southern USA, Central and South America (Cooley 1944; Guglielmone and Robbins 2018).

Natural hosts. Variety of mammals, including humans, also birds, amphibians, reptiles (Cooley 1944; Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. *Rickettsia rickettsii* (Rocky Mountain spotted fever) (Bermudez and Troyo 2018).

***Amblyomma moreliae* (Koch, 1867)**

Geographic distribution. Australia (Roberts 1970).

Natural hosts. Reptiles. Zoonotic on humans (Roberts 1970).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Amblyomma naponense* (Packard, 1869)**

Amblyomma mantiquirens Aragão, 1908

Geographic distribution. Central and South America (Guglielmone and Robbins 2018).

Natural hosts. Primarily mammals, including peccaries, carnivores, anteaters, marsupials, and rodents, and birds. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Amblyomma neumanni* Ribaga, 1902**

Amblyomma furcula Dönitz, 1909

Geographic distribution. Argentina, Colombia (Guglielmone and Robbins 2018).

Natural hosts. Mammals, including bovids, deer, carnivores, pigs, peccaries, and humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Amblyomma nuttalli* Dönitz, 1909**

Geographic distribution. Sub-Saharan Africa (Guglielmone and Robbins 2018).

Natural hosts. Immature stages on a variety of mammals, birds, and reptiles. Adults primarily on tortoises. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Amblyomma oblongoguttatum* Koch, 1844**

Geographic distribution. Central and South America (Guglielmone and Robbins 2018).

Natural hosts. A variety of mammals, including deer, rodents, carnivores, and marsupials, and birds. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Amblyomma ovale* Koch, 1844**

Amblyomma fozsum Neumann, 1899

Geographic distribution. Southern USA, Central and South America (Cooley 1944; Guglielmone and Robbins 2018).

Natural hosts. Mammals, primarily carnivores, rodents, tapirs, and humans. Occasionally marsupials, birds (Cooley 1944; Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Amblyomma pacae Aragão, 1911

Geographic distribution. Central and South America (Guglielmone and Robbins 2018).

Natural hosts. Primarily mammals, including rodents and marsupials, but also carnivores and birds. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Amblyomma parkeri Fonseca & Aragão, 1951

Geographic distribution. Brazil (Guglielmone and Robbins 2018).

Natural hosts. Primarily rodents, also carnivores, marsupials, monkeys, and birds. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Amblyomma parvum Aragão, 1908

Geographic distribution. Central and South America (Guglielmone and Robbins 2018).

Natural hosts. Wide variety of mammals, including humans, and birds (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Amblyomma paulopunctatum Neumann, 1899

Geographic distribution. Sub-Saharan Africa (Guglielmone and Robbins 2018).

Natural hosts. Mammals, primarily pigs, but also hippopotamuses, carnivores, and rodents, and birds. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Amblyomma pecarium* Dunn, 1933**

Geographic distribution. Central and South America (Guglielmone and Robbins 2018).

Natural hosts. Peccaries, also deer. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Amblyomma personatum* Neumann, 1901**

Geographic distribution. Sub-Saharan Africa (Guglielmone and Robbins 2018).

Natural hosts. Rhinoceroses, bovids. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Amblyomma postoculatum* Neumann, 1899**

Geographic distribution. Australia (Roberts 1970).

Natural hosts. Marsupials. Zoonotic on humans (Roberts 1970).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Amblyomma pseudoconcolor* Aragão, 1908**

Geographic distribution. South America (Guglielmone and Robbins 2018).

Natural hosts. Primarily armadillos, also marsupials and birds. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Amblyomma pseudoparvum* Guglielmone et al., 1990**

Geographic distribution. Argentina (Guglielmone and Robbins 2018).

Natural hosts. Caviomorph rodents. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Amblyomma romitii* Tonelli-Rondelli, 1939**

Amblyomma tasquei Floch & Adonnenc, 1940

Geographic distribution. South America (Guglielmone and Robbins 2018).

Natural hosts. Rodents, marsupials. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Amblyomma rotundum* Koch, 1844**

Geographic distribution. Southern USA, Central and South America, Pacific Islands (Cooley 1944; Guglielmone and Robbins 2018).

Natural hosts. Reptiles and amphibians, occasionally birds. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Amblyomma sabanerae* Stoll, 1890**

Geographic distribution. Central and South America (Guglielmone and Robbins 2018).

Natural hosts. Primarily reptiles, including tortoises, also amphibians, rodents, and marsupials. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Amblyomma sculpturatum* Neumann, 1906**

Geographic distribution. Central and South America (Guglielmone and Robbins 2018).

Natural hosts. Mammals, including tapirs, pigs, anteaters, peccaries, carnivores, and rodents. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Amblyomma sculptum* (Berlese, 1888)**

Geographic distribution. South America (Martins et al. 2016; Guglielmone and Robbins 2018).

Natural hosts. Several mammals, including capybaras and humans, and birds (Osava et al. 2016; Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment

Site in human host. Skin.

Vectored pathogens. *Rickettsia rickettsii* (Rocky Mountain Spotted Fever) (Ramírez-Hernández et al. 2020).

Amblyomma sparsum Neumann, 1899

Geographic distribution. Sub-Saharan Africa (Guglielmone and Robbins 2018).

Natural hosts. Several groups of mammals, birds, and reptiles. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Amblyomma tapirellum Dunn, 1933

Geographic distribution. Central and South America (Guglielmone and Robbins 2018).

Natural hosts. Tapirs, peccaries, marsupials, rodents. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Amblyomma tenellum Koch, 1844

Amblyomma imitator Kohls, 1958

Geographic distribution. Southern USA, Central America (Guglielmone and Robbins 2018).

Natural hosts. Mammals, including bovids, peccaries, horses, marsupials, carnivores, and rodents, and birds. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Amblyomma testudinarum Koch, 1844

Geographic distribution. Central and Southeast Asia, Japan (Guglielmone and Robbins 2018).

Natural hosts. Several groups of mammals, including humans, birds, and reptiles (Guglielmone and Robbins 2018),

Route of infection. Exposure to larval, nymph, or adult ticks in the environment

Site in human host. Skin, ear (Nakao et al. 2017).

Vectored pathogens. Dabie bandavirus (severe fever with thrombocytopenia syndrome Virus, SFTSV) (Sato et al. 2021).

Amblyomma tholloni Neumann, 1899

Geographic distribution. Sub-Saharan Africa (Guglielmone and Robbins 2018).

Natural hosts. Several groups of mammals, primarily elephants, and birds and reptiles. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Amblyomma tigrinum Koch, 1844

Geographic distribution. South America (Guglielmone and Robbins 2018).

Natural hosts. Primarily carnivores, also rodents and ungulates. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Amblyomma tonelliae Nava et al., 2014

Geographic distribution. South America (Guglielmone and Robbins 2018).

Natural hosts. Mammals, primarily bovids and equids. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. *Rickettsia rickettsii* (Rocky Mountain spotted fever) (Tarragona et al. 2016).

Amblyomma triguttatum Koch, 1844

Geographic distribution. Australia (Guglielmone and Robbins 2018).

Natural hosts. Primarily marsupials, also horses, bovids, carnivores, lagomorphs, rodents, humans, and reptiles (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Amblyomma triste* Koch, 1844**

Geographic distribution. Southern USA, Central and South America (Cooley 1944; Guglielmo and Robbins 2018).

Natural hosts. Several groups of mammals, primarily bovids, deer, and carnivores, and birds. Zoonotic on humans (Cooley 1944; Guglielmo and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. *Rickettsia parkeri* (Tidewater spotted fever) (Silveira et al. 2007).

***Amblyomma tuberculatum* Marx in Hubbard, 1894**

Geographic distribution. Southeastern USA (Cooley 1944; Guglielmo and Robbins 2018).

Natural hosts. Tortoises. Zoonotic on humans (Cooley 1944; Guglielmo and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Amblyomma variegatum* (Fabricius, 1794)**

Geographic distribution. Africa, Middle East, Caribbean (Guglielmo and Robbins 2018).

Natural hosts. Several groups of mammals, primarily bovids, also humans, birds, and reptiles (Guglielmo and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. *Rickettsia africae* (African tick bite fever) (Dantas-Torres et al. 2012).

***Amblyomma varium* Koch, 1844**

Geographic distribution. Central and South America (Guglielmo and Robbins 2018).

Natural hosts. Mammals, primarily sloths, also marsupials, tapirs, carnivores, and rodents, also reptiles and birds. Zoonotic on humans (Guglielmo and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Genus *Bothriocroton* Keirans et al., 1994***Bothriocroton auruginans* (Schülze, 1938)**

Geographic distribution. Australia (Guglielmone and Robbins 2018).

Natural hosts. Marsupials, occasionally carnivores. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Bothriocroton hydrosauri* (Denny, 1843)**

Geographic distribution. Australia (Guglielmone and Robbins 2018).

Natural hosts. Reptiles, occasionally bovids. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Genus *Dermacentor* Koch, 1844***Dermacentor albipictus* (Packard, 1869)**

Ixodes nigrolineatus Packard, 1869

Geographic distribution. North and Central America (Guglielmone and Robbins 2018).

Natural hosts. Mammals, primarily cattle, deer, and horses. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. *Babesia duncani* (babesiosis) (Swei et al. 2019).

***Dermacentor andersoni* Stiles, 1908**

Dermacentor venustus Marx in Neumann 1897

Geographic distribution. Rocky Mountain region of North America (Guglielmone and Robbins 2018).

Natural hosts. Mammals, including humans (Eisen 2007; Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment

Site in human host. Skin

Vectored pathogens. *Rickettsia rickettsii* (Rocky Mountain spotted fever), *Francisella tularensis* (tularemia), Colorado tick fever virus; also implicated in tick paralysis (Eisen 2007).

***Dermacentor atrosignatus* Neumann, 1906**

Geographic distribution. Southeast Asia (Guglielmone and Robbins 2018).

Natural hosts. Pigs. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin, ear (Mariana et al. 2008).

Vectored pathogens. None.

***Dermacentor auratus* Supino, 1897**

Geographic distribution. Southeast Asia (Guglielmone and Robbins 2018).

Natural hosts. Primarily pigs, but also a variety of other mammals, birds, and reptiles. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin, ear (Ariyaratne et al. 2011).

Vectored pathogens. *Rickettsia sibirica* (Siberian tick typhus) (Dantas-Torres et al. 2012).

***Dermacentor bellulus* (Schülze, 1933)**

Geographic distribution. Southeast Asia (Guglielmone and Robbins 2018).

Natural hosts. Immature stages primarily on rodents, also tree shrews, carnivores, lagomorphs, and birds. Adults are primarily on pigs, also carnivores. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Dermacentor circumguttatus* Neumann, 1897**

Geographic distribution. Sub-Saharan Africa (Guglielmone and Robbins 2018).

Natural hosts. Primarily elephants, also duikers, pigs. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Dermacentor compactus* Neumann, 1901**

Geographic distribution. Southeast Asia (Guglielmone and Robbins 2018).

Natural hosts. Primarily pigs, but also a variety of other mammals and reptiles. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin, ear (Mariana et al. 2008).

Vectored pathogens. None.

***Dermacentor hunteri* Bishopp, 1912**

Geographic distribution. Southeast Asia (Guglielmone and Robbins 2018).

Natural hosts. Primarily bovids, also deer, lagomorphs, and rodents. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Dermacentor imitans* Warburton, 1933**

Geographic distribution. Central and South America (Guglielmone and Robbins 2018).

Natural hosts. Peccaries, deer. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Dermacentor latus* Cooley, 1937**

Geographic distribution. Central America (Guglielmone and Robbins 2018).

Natural hosts. Primarily tapirs, also carnivores. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Dermacentor limbooliati* Apanaskevich & Apanaskevich, 2015**

Geographic distribution. Malaysia, Vietnam (Apanaskevich and Apanaskevich 2015).

Natural hosts. Pigs. Zoonotic on humans (Apanaskevich and Apanaskevich 2015).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Dermacentor marginatus* (Sulzer, 1776)**

Geographic distribution. Europe, western and central Asia, northern Africa (Rubel et al. 2016; Guglielmone and Robbins 2018).

Natural hosts. Many mammals, including humans, and birds (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. *Rickettsia raoultii*, Omsk hemorrhagic fever virus (Dantas-Torres et al. 2012).

Dermacentor nitens Neumann, 1897

Geographic distribution. Southern USA, Central and South America, Caribbean (Guglielmone and Robbins 2018).

Natural hosts. Horses, occasionally other mammals, reptiles, and amphibians. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Dermacentor niveus Neumann, 1897

Geographic distribution. Central Asia, China, Russia (Guglielmone and Robbins 2018).

Natural hosts. Mammals, including ungulates, rodents, lagomorphs, hedgehogs, and carnivores. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Dermacentor nuttalli Olenov, 1929

Geographic distribution. Eastern Russia, Mongolia, China (Guglielmone and Robbins 2018).

Natural hosts. Immature stages on rodents, lagomorphs, hedgehogs. Adults primarily on ungulates and horses. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. *Rickettsia sibirica* (Siberian tick typhus), *Francisella tularensis* (tularemia), Crimean-Congo hemorrhagic fever virus (Gui et al. 2021; Kulakova et al. 2014).

Dermacentor occidentalis Marx in Curtice, 1892

Geographic distribution. Pacific coastal areas of Mexico and USA (Guglielmone and Robbins 2018).

Natural hosts. Several groups of mammals. Immature stages primarily on rodents and lagomorphs. Adults on cattle, deer, and humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. *Rickettsia phlipii* (Pacific Coast tick fever) (Padgett et al. 2016).

***Dermacentor parumapertus* Neumann, 1901**

Geographic distribution. North America (Guglielmone and Robbins 2018).

Natural hosts. Lagomorphs, rodents, bovids, deer, carnivores, and birds. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Dermacentor raskemensis* Pomerantsev, 1946**

Geographic distribution. Central and Southeast Asia (Guglielmone and Robbins 2018).

Natural hosts. Bovids, carnivores, rodents, lagomorphs. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Dermacentor reticulatus* (Fabricius, 1794)**

Dermacentor pictus Hermann, 1804

Geographic distribution. Europe and western Asia (Rubel et al. 2016; Guglielmone and Robbins 2018).

Natural hosts. Immature stages primarily on rodents, lagomorphs, hedgehogs, rarely birds and reptiles, and amphibians. Adults primarily on canids, also ungulates, and humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. *Rickettsia slovaca* (tick-borne lymphadenopathy, TIBOLA), *Rickettsia raoultii*, Omsk hemorrhagic fever virus (Földvári et al. 2016).

***Dermacentor rhinocerinus* (Denny, 1843)**

Geographic distribution. Sub-Saharan Africa (Guglielmone and Robbins 2018).

Natural hosts. Primarily rhinoceroses, occasionally other mammals and reptiles. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Dermacentor silvarum* Olenev, 1931**

Dermacentor asiaticus Emel'yanova & Kozlovskaya, 1967

Geographic distribution. Eastern Russia, China, Mongolia (Guglielmone and Robbins 2018).

Natural hosts. Mammals, including bovids, deer, sheep, dogs, and humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. *Rickettsia sibirica* (Siberian tick typhus), *Rickettsia raoultii* (Cao et al. 2008).

***Dermacentor similis* Lado, Glon, et Klompen, 2021**

Geographic distribution. North America, west of the Rocky Mountains (Lado et al. 2021).

Natural hosts. Several groups of mammals, including humans, and birds (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. *Rickettsia rickettsii* (Rocky Mountain spotted fever), *Francisella tularensis* (tularemia); also implicated in tick paralysis (Wikswow et al. 2014; Moraru 2019; Mullen and Durden 2019).

***Dermacentor steini* (Schülze, 1933)**

Geographic distribution. Southeast Asia (Guglielmone and Robbins 2018).

Natural hosts. Primarily pigs, but also a variety of other mammals and reptiles. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin, ear (Mariana et al. 2008).

Vectored pathogens. None.

***Dermacentor tamokensis* Apanaskevich & Apanaskevich, 2016**

Geographic distribution. Southeast Asia (Apanaskevich and Apanaskevich 2016; Guglielmone and Robbins 2018).

Natural hosts. Pigs. Zoonotic on humans (Guglielmone and Robbins 2018)

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Dermacentor variabilis* (Say, 1821)**

Dermacentor electus Koch, 1844

Geographic distribution. Eastern North America (Guglielmone and Robbins 2018; Lado et al. 2021).

Natural hosts. Several groups of mammals, including humans, and birds. (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin, ear (Grady et al. 2011).

Vectored pathogens. *Rickettsia rickettsii* (Rocky Mountain spotted fever), *Francisella tularensis* (tularemia); also implicated in tick paralysis (Ammerman et al. 2004; Dergousoff and Chilton 2012; Moraru 2019; Mullen and Durden 2019).

Genus *Haemaphysalis* Koch, 1844***Haemaphysalis aculeata* Lavarra, 1904**

Geographic distribution. India, Sri Lanka (Guglielmone and Robbins 2018).

Natural hosts. Chevrotains, bovids, deer. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. Kyasanur Forest Disease virus (Sharma et al. 2019).

***Haemaphysalis anomala* Warburton, 1913**

Geographic distribution. Southeast Asia (Guglielmone and Robbins 2018),

Natural hosts. Immature stages on rodents and small birds. Adults on bovids, deer, carnivores. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Haemaphysalis aponommoides* Warburton, 1913**

Geographic distribution. China, India, Nepal (Guglielmone and Robbins 2018).

Natural hosts. Immature stages on rodents, shrews, birds. Adults on mammals, primarily bovids. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Haemaphysalis bancrofti* Nuttall & Warburton, 1915**

Geographic distribution. Australia, Indonesia, Papua New Guinea (Guglielmone and Robbins 2018).

Natural hosts. Primarily marsupials, but also a variety of other mammals and birds. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Haemaphysalis birmaniae* Supino, 1897**

Geographic distribution. Southeast Asia (Guglielmone and Robbins 2018).

Natural hosts. Bovids, deer, pigs, carnivores, and rodents. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Haemaphysalis bispinosa* Neumann, 1897**

Geographic distribution. Central and Southeast Asia (Guglielmone and Robbins 2018).

Natural hosts. Mammals, primarily bovids and carnivores, and birds. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. Kyasanur Forest disease virus (Sharma et al. 2019).

***Haemaphysalis campanulata* Warburton, 1908**

Geographic distribution. Central and Southeast Asia, Japan (Guglielmone and Robbins 2018).

Natural hosts. Immature stages primarily on rodents. Adults on mammals, primarily carnivores. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Haemaphysalis caucasica* Olenov, 1928**

Geographic distribution. Eastern Europe, Central Asia (Guglielmone and Robbins 2018).

Natural hosts. Mammals, including lagomorphs, carnivores, rodents, and bovids, and birds. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Haemaphysalis celebensis Hoogstral et al., 1965

Geographic distribution. Indonesia (Guglielmone and Robbins 2018).

Natural hosts. Primarily pigs, bovids, deer, and horses. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Haemaphysalis chordeilis (Packard, 1869)

Geographic distribution. North America (Guglielmone and Robbins 2018).

Natural hosts. Primarily birds, also mammals including bovids, horses, rodents. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Haemaphysalis colasbelcouri (Santos Dias, 1958)

Geographic distribution. China, Vietnam, Laos (Guglielmone and Robbins 2018).

Natural hosts. Primarily bovids and deer. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Haemaphysalis concinna Koch, 1844

Geographic distribution. Palearctic (Guglielmone and Robbins 2018; Kiewra et al. 2019).

Natural hosts. Small mammals, birds, and reptiles. Zoonotic on humans (Guglielmone and Robbins 2018; Kiewra et al. 2019).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. *Francisella tularensis* (tularemia), tick-borne encephalitis viruses (TBE) (Kiewra et al. 2019).

***Haemaphysalis cornigera* Neumann, 1897**

Geographic distribution. Southeast Asia, Japan (Guglielmone and Robbins 2018).

Natural hosts. Mammals, including bovids, deer, carnivores, shrews, and tree shrews. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. *Rickettsia japonica* (Japanese spotted fever) (Li et al. 2019).

***Haemaphysalis cuspidata* Warburton, 1910**

Geographic distribution. India, Sri Lanka (Guglielmone and Robbins 2018).

Natural hosts. Many groups of mammals, including ungulates, carnivores, rodents, lagomorphs, shrews, and non-human primates, and birds. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. Kyasanur Forest disease virus (Sharma et al. 2019).

***Haemaphysalis darjeeling* Hoogstraal & Dhanda, 1970**

Geographic distribution. Southeast Asia (Guglielmone and Robbins 2018).

Natural hosts. Mammals, including bovids, deer, pigs, and carnivores. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Haemaphysalis doenitzi* Warburton & Nuttall, 1909**

Geographic distribution. Southeast Asia, Japan, Russia, Australia (Guglielmone and Robbins 2018).

Natural hosts. Primarily birds, occasionally lagomorphs and reptiles. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Haemaphysalis elliptica* (Koch, 1844)**

Geographic distribution. Sub-Saharan Africa (Guglielmone and Robbins 2018).

Natural hosts. Primarily carnivores, also rodents, elephant shrews, and birds. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.
Site in human host. Skin.
Vectored pathogens. None.

Haemaphysalis elongata Neumann, 1897

Geographic distribution. Madagascar (Guglielmone and Robbins 2018).
Natural hosts. Primarily tenrecs, also carnivores and hedgehogs. Zoonotic on humans (Guglielmone and Robbins 2018).
Route of infection. Exposure to larval, nymph, or adult ticks in the environment.
Site in human host. Skin.
Vectored pathogens. None.

Haemaphysalis erinacei Pavesi, 1884

Haemaphysalis numidiana Neumann, 1905
Haemaphysalis erinacei taurica Pospelova-Shtrom, 1939

Geographic distribution. Europe, northern Africa, Russia (Guglielmone and Robbins 2018).
Natural hosts. Mammals, primarily hedgehogs and carnivores, also lagomorphs and bovids, birds, and reptiles. Zoonotic on humans (Guglielmone and Robbins 2018).
Route of infection. Exposure to larval, nymph, or adult ticks in the environment.
Site in human host. Skin.
Vectored pathogens. None.

Haemaphysalis flava Neumann, 1897

Geographic distribution. East Asia (Guglielmone and Robbins 2018).
Natural hosts. Several groups of mammals and birds. Zoonotic on humans (Guglielmone and Robbins 2018).
Route of infection. Exposure to larval, nymph, or adult ticks in the environment.
Site in human host. Skin.
Vectored pathogens. *Rickettsia japonica* (Japanese spotted fever) (Li et al. 2019).

Haemaphysalis heinrichi Schülze, 1939

Geographic distribution. Southeast Asia (Guglielmone and Robbins 2018).
Natural hosts. Mammals, including bovids, carnivores, rodents, and shrews. Zoonotic on humans (Guglielmone and Robbins 2018).
Route of infection. Exposure to larval, nymph, or adult ticks in the environment.
Site in human host. Skin.
Vectored pathogens. None.

***Haemaphysalis hirsuta* Hoogstraal et al., 1966**

Geographic distribution. Indonesia (Guglielmone and Robbins 2018).

Natural hosts. Mammals, including bovids, pigs, carnivores, and rodents. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Haemaphysalis hoodi* Warburton & Nuttall, 1909**

Geographic distribution. Sub-Saharan Africa (Guglielmone and Robbins 2018).

Natural hosts. Primarily birds, also mammals including bovids, non-human primates, rodents, carnivores, and lagomorphs. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Haemaphysalis humerosa* Warburton & Nuttall, 1909**

Geographic distribution. Indonesia, Australia (Guglielmone and Robbins 2018).

Natural hosts. Primarily marsupials, also monotremes, rodents, and birds. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Haemaphysalis hylobatis* Schülze, 1933**

Geographic distribution. Southeast Asia, Japan (Guglielmone and Robbins 2018).

Natural hosts. Several groups of mammals and birds. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Haemaphysalis hystricis* Supino, 1897**

Haemaphysalis nishiyamai Sugimoto, 1935

Haemaphysalis trispinosa Tounamoff, 1941

Geographic distribution. Southeast Asia, Japan (Guglielmone and Robbins 2018).

Natural hosts. Several groups of mammals and birds. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. *Rickettsia japonica* (Japanese spotted fever) (Li et al. 2019).

Haemaphysalis indoflava Dhanda & Bhat, 1968

Geographic distribution. Southeast Asia, Japan (Guglielmone and Robbins 2018).

Natural hosts. Bovids, pigs, carnivores. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Haemaphysalis inermis Birula, 1895

Geographic distribution. Southern Europe, Middle East (Guglielmone and Robbins 2018).

Natural hosts. Primarily birds, also rodents, shrews, and reptiles. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Haemaphysalis intermedia Warburton & Nuttall, 1909

Geographic distribution. Central Asia (Guglielmone and Robbins 2018).

Natural hosts. Several groups of mammals and birds. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Haemaphysalis japonica Warburton, 1908

Haemaphysalis japonica douglasi Nuttall & Warburton, 1915

Geographic distribution. Southeast Asia, Japan, Russia (Guglielmone and Robbins 2018).

Natural hosts. Mammals, including deer, carnivores, lagomorphs, pigs, horses, bovines, and hedgehogs, and birds. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Haemaphysalis juxtakochi* Cooley, 1946**

Haemaphysalis kohlsi Aragão & Fonseca, 1951

Geographic distribution. USA, Central and South America (Guglielmone and Robbins 2018).

Natural hosts. Mammals, including deer, peccaries, bovids, horses, carnivores, marsupials, rodents, and lagomorphs, and birds. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Haemaphysalis kitaokai* Hoogstraal, 1969**

Geographic distribution. Southeast Asia, Japan (Guglielmone and Robbins 2018).

Natural hosts. Mammals, including bovids, deer, horses, and rodents, and birds. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Haemaphysalis koningsbergeri* Warburton & Nuttall, 1909**

Geographic distribution. Southeast Asia (Guglielmone and Robbins 2018).

Natural hosts. Mammals, primarily carnivores and rodents. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Haemaphysalis lagrangei* Larrousse, 1925**

Haemaphysalis hystricis indochinensis Phan Trong, 1977

Geographic distribution. Southeast Asia (Guglielmone and Robbins 2018).

Natural hosts. Several groups of mammals and birds. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Haemaphysalis leachi* (Audouin, 1826)**

Geographic distribution. Africa (Apanaskevich et al. 2007; Guglielmono and Robbins 2018).

Natural hosts. Mammals, primarily carnivores, bovids, pigs, non-human primates, hedgehogs, and rodents, and birds. Zoonotic on humans (Apanaskevich et al. 2007; Guglielmono and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. *Rickettsia conorii* (boutonneuse fever) (Apanaskevich et al. 2007).

***Haemaphysalis leporispalustris* Packard, 1869**

Geographic distribution. North, Central, and South America (Guglielmono and Robbins 2018; Sánchez-Montes et al. 2020).

Natural hosts. Primarily rabbits, also other mammals and birds. Zoonotic on humans (Guglielmono and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Haemaphysalis longicornis* Neumann, 1901**

Geographic distribution. East Asia, Australia, New Zealand, Pacific Islands, eastern North America (Beard et al. 2018; Guglielmono and Robbins 2018; Wormser et al. 2020).

Natural hosts. Mammals, including sheep, deer, and horses, and birds. Zoonotic on humans (Guglielmono and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin, ear (Choi et al. 2018).

Vectored pathogens. Dabie bandavirus (severe fever with thrombocytopenia syndrome virus, SFTSV), *Rickettsia japonica* (Japanese spotted fever) (Li et al. 2019; Sato et al. 2021).

***Haemaphysalis mageshimaensis* Saito & Hoogstraal, 1973**

Geographic distribution. Southeast Asia, Japan (Guglielmono and Robbins 2018).

Natural hosts. Mammals, including bovids, deer, carnivores, pigs, rodents, and birds. Zoonotic on humans (Guglielmono and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Haemaphysalis megaspinosa Saito, 1969

Geographic distribution. Southeast Asia, Japan (Guglielmone and Robbins 2018).

Natural hosts. Deer, carnivores, horses. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Haemaphysalis mjoebergi Warburton, 1926

Geographic distribution. Indonesia (Guglielmone and Robbins 2018).

Natural hosts. Deer, bovids. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Haemaphysalis montgomeryi Nuttall, 1912

Geographic distribution. Central and Southeast Asia, Japan (Guglielmone and Robbins 2018).

Natural hosts. Mammals, including bovids, rodents, camels, deer, horses, and hedgehogs, and birds. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Haemaphysalis nadchatrami Hoogstraal et al., 1965

Geographic distribution. Southeast Asia (Guglielmone and Robbins 2018).

Natural hosts. Rodents, bovids, deer, pigs, carnivores, horses, tapirs. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Haemaphysalis nepalensis Hoogstraal, 1962

Geographic distribution. Central and Southeast Asia (Guglielmone and Robbins 2018).

Natural hosts. Bovids, carnivores, rodents. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Haemaphysalis novaeguineae* Hirst, 1914**

Geographic distribution. Australia, Papua New Guinea (Guglielmone and Robbins 2018).

Natural hosts. Mammals, including bovids, pigs, marsupials, carnivores, horses, and rodents, and birds. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Haemaphysalis obesa* Larrousse, 1925**

Geographic distribution. Southeast Asia (Guglielmone and Robbins 2018).

Natural hosts. Several groups of mammals. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Haemaphysalis papuana* Thorell, 1883**

Geographic distribution. Southeast Asia (Guglielmone and Robbins 2018).

Natural hosts. Mammals, including pigs, deer, musk deer, carnivores, and rodents, and birds. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Haemaphysalis paraleachi* Camicas et al. 1983**

Geographic distribution. Sub-Saharan Africa (Guglielmone and Robbins 2018).

Natural hosts. Carnivores, rodents, bovids, non-human primates. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Haemaphysalis parmata* Neumann, 1905**

Geographic distribution. Sub-Saharan Africa (Guglielmone and Robbins 2018).

Natural hosts. Mammals, primarily bovids, rarely reptiles and birds. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Haemaphysalis parva* (Neumann, 1897)**

Haemaphysalis otophila Schülze, 1919

Geographic distribution. Europe, northern Africa, western Asia (Guglielmone and Robbins 2018).

Natural hosts. Mammals, including hedgehogs, carnivores, lagomorphs, rodents, and humans, birds, and reptiles (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Haemaphysalis punctata* Canestrini & Fanzago, 1878**

Haemaphysalis punctata autumnalis Schülze, 1919

Geographic distribution. Europe, northern Africa, western and central Asia (Guglielmone and Robbins 2018).

Natural hosts. Several groups of mammals, including humans, birds, and reptiles (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Haemaphysalis qinghaiensis* Teng, 1980**

Geographic distribution. China (Guglielmone and Robbins 2018).

Natural hosts. Bovids, horses, lagomorphs. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Haemaphysalis ramachandrai* Dhanda et al., 1970**

Geographic distribution. India, Nepal (Guglielmone and Robbins 2018).

Natural hosts. Deer, bovids, carnivores. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Haemaphysalis roubaudi* Toumanoff, 1940**

Geographic distribution. Vietnam (Guglielmone and Robbins 2018).

Natural hosts. Deer. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Haemaphysalis semermis* Neumann, 1901**

Geographic distribution. Southeast Asia (Guglielmone and Robbins 2018).

Natural hosts. Chevrotains, deer, pigs, carnivores, tapirs, rodents, tree shrews. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Haemaphysalis shimoga* Trapido & Hoogstraal, 1964**

Haemaphysalis cornigera vietnama Phan Trong, 1977

Geographic distribution. Southeast Asia (Guglielmone and Robbins 2018).

Natural hosts. Mammals, including bovids, pigs, deer, and rodents. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Haemaphysalis silacea* Robinson, 1912**

Geographic distribution. Sub-Saharan Africa (Guglielmone and Robbins 2018)

Natural hosts. Mammals, including bovids, elephant shrews, rhinoceroses, carnivores, lagomorphs, and rodents, and birds. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Haemaphysalis spinigera* Neumann, 1897**

Geographic distribution. Southeast Asia (Guglielmone and Robbins 2018).

Natural hosts. Several groups of mammals, including humans, and birds (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. Kyasanur Forest Disease virus (Varma et al. 1960; Sharma et al. 2019).

Haemaphysalis sulcata Canestrini & Fanzago, 1878

Haemaphysalis cholodkovskiyi Olenov. 1928

Geographic distribution. Europe, North Africa, Asia (Guglielmone and Robbins 2018).

Natural hosts. Mammals, primarily bovids, but also carnivores, rodents, lagomorphs, bats, hedgehogs, and humans, and reptiles (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Haemaphysalis turturis Nuttall & Warburton, 1915

Geographic distribution. India, Sri Lanka (Guglielmone and Robbins 2018).

Natural hosts. Several groups of mammals, birds, and reptiles. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. Kyasanur Forest disease virus (Sharma et al. 2019).

Haemaphysalis wellingtoni Nuttall & Warburton, 1908

Geographic distribution. Southeast Asia (Guglielmone and Robbins 2018).

Natural hosts. Primarily birds, occasionally mammals. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. Kyasanur Forest disease virus (Sharma et al. 2019).

Genus *Hyalomma* Koch, 1844

Hyalomma aegyptium (Linnaeus, 1758)

Geographic distribution. Europe, North Africa, Middle East, Central Asia (Guglielmone and Robbins 2018).

Natural hosts. Primarily tortoises, but also other reptiles and mammals. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. *Borrelia turcica* (tick-borne relapsing fever) (Güner et al. 2004).

***Hyalomma albiparmatum* Schülze, 1919**

Geographic distribution. East Africa (Apanaskevich and Horak 2008; Guglielmone and Robbins 2018).

Natural hosts. Many mammals, primarily wild ungulates, also lagomorphs and birds. Zoonotic on humans (Apanaskevich and Horak 2008; Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. *Rickettsia conorii* (boutonneuse fever) (Apanaskevich and Horak 2008).

***Hyalomma anatolicum* Koch, 1844**

Geographic distribution. North Africa, Middle East, Central Asia (Vatansever 2017a; Guglielmone and Robbins 2018).

Natural hosts. Mammals, including cattle, horses, camels, sheep, goats, and humans, and birds (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. Crimean-Congo hemorrhagic fever virus (Kassari et al. 2020).

***Hyalomma asiaticum* Schülze & Schlottke, 1930**

Hyalomma asiaticum kozlovi Olenov, 1931

Geographic distribution. Central Asia, Middle East (Guglielmone and Robbins 2018; Vatansever 2017b).

Natural hosts. Mammals, including bovids, camels, rodents, lagomorphs, and hedgehogs, birds, and reptiles. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. *Rickettsia sibirica mongolitimonae*, *Coxiella burnetti* (Q fever), and possibly Crimean-Congo Hemorrhagic Fever virus (Ramos et al. 2013; Batu et al. 2020; Kassari et al. 2020).

***Hyalomma brevipunctatum* Sharif, 1928**

Geographic distribution. Southeast and Central Asia (Guglielmone and Robbins 2018).

Natural hosts. Mammals, including bovids, deer, camels, horses, rodents, carnivores, shrews, and birds. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Hyalomma dromedarii Koch, 1844

Ixodes camelinus Fischer von Waldheim, 1823

Ixodes arenicola Eichwald, 1830

Ixodes trilineatus Lucas, 1844

Ixodes cinctus Lucas, 1844

Hyalomma yakimovi Olenov, 1931

Hyalomma yakimovi persiacum Olenov, 1931

Hyalomma delphy Schülze and Gossel in Schülze, 1936

Geographic distribution. Africa, Middle East, Central and Southeast Asia (Guglielmone and Robbins 2018).

Natural hosts. Primarily camels, but also other mammals, birds, and reptiles. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. Crimean-Congo hemorrhagic fever virus (Champour et al. 2016).

Hyalomma excavatum Koch, 1844

Geographic distribution. North Africa, Mediterranean Europe, Middle East, Central Asia (Guglielmone and Robbins 2018).

Natural hosts. Mammals, including bovids, camels, lagomorphs, rodents, hedgehogs, and humans, birds, and reptiles (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Hyalomma glabrum Delpy, 1949

Geographic distribution. South Africa (Guglielmone and Robbins 2018).

Natural hosts. Immature stages on lagomorphs, carnivores, hyrax, rodents, and birds. Adults on bovids, occasionally carnivores. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Hyalomma hussaini* Sharif, 1928**

Geographic distribution. India, Pakistan, Myanmar (Guglielmone and Robbins 2018).

Natural hosts. Immature stages on rodents and shrews. Adults on bovids, camels, pigs, carnivores, and horses. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Hyalomma impeltatum* Schülze & Schlottke, 1930**

Geographic distribution. Africa, Middle East, Central Asia (Guglielmone and Robbins 2018).

Natural hosts. Mammals, including bovids, camels, horses, rhinoceroses, carnivores, and rodents, birds, and reptiles. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Hyalomma isaaci* Sharif, 1928**

Geographic distribution. Central and Southeast Asia (Guglielmone and Robbins 2018).

Natural hosts. Immature stages on lagomorphs, deer, carnivores, and birds. Adults primarily on bovids. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Hyalomma lustanicum* Koch, 1844**

Geographic distribution. Mediterranean Europe and Africa (Guglielmone and Robbins 2018).

Natural hosts. Mammals, including bovids and lagomorphs, and birds. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Hyalomma marginatum* Koch, 1844**

Geographic distribution. Europe, North Africa, Middle East, Central Asia (Estrada-Peña et al. 2012; Guglielmone and Robbins 2018).

Natural hosts. Several groups of mammals, including humans, and birds. Immature stages are primarily on small mammals and birds. Adults primarily on ungulates (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host: Skin.

Vectored pathogens. Crimean-Congo hemorrhagic fever virus (Gargili et al. 2013; Kassari et al. 2020).

Hyalomma rufipes Koch, 1844

Geographic distribution. Africa, Middle East (Guglielmone and Robbins 2018).

Natural hosts. Several groups of mammals and birds. Immature stages are primarily on birds and lagomorphs. Adults are primarily on bovids. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. Crimean-Congo hemorrhagic fever virus (Hoogstraal 1979).

Hyalomma schulzei Olenev, 1931

Geographic distribution. Middle East, Central Asia (Guglielmone and Robbins 2018).

Natural hosts. Immature stages primarily on rodents, hedgehogs, and lagomorphs. Adults primarily on camels, occasionally bovids. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Hyalomma scupense Schülze, 1919

Hyalomma detritum Schülze, 1919

Hyalomma detritum dardanicum Schülze & Schlottke, 1930

Hyalomma uralense Schülze & Schlottke, 1930

Hyalomma volgense Schülze & Schlottke, 1930

Geographic distribution. Europe, North Africa, Middle East, Central and East Asia (Guglielmone and Robbins 2018).

Natural hosts. Bovids, camels, horses, pigs, deer, and humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Hyalomma truncatum* Koch, 1844Hyalomma transiens* Schülze, 1919

Geographic distribution. Africa, Saudi Arabia, Yemen (Apanaskevich and Horak 2008; Guglielmone and Robbins 2018).

Natural hosts. Many groups of mammals, birds, and reptiles. Immature stages primarily on rodents and lagomorphs. Adults primarily on domestic and wild ungulates. Zoonotic on humans (Apanaskevich and Horak 2008; Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment

Site in human host. Skin

Vectored pathogens. Crimean-Congo hemorrhagic fever virus, *Rickettsia conorii* (boutonneuse fever), *Coxiella burnetii* (Q fever) (Apanaskevich and Horak 2008).

***Hyalomma turanicum* Pomerantzev, 1946**

Geographic distribution. North Africa, Middle East, Central and East Asia (Guglielmone and Robbins 2018).

Natural hosts. Mammals, including bovids, lagomorphs, camels, pigs, horses, humans, and birds (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Genus *Ixodes* Latreille, 1795***Ixodes acuminatus* Neumann, 1901**

Geographic distribution. Europe, Middle East (Guglielmone and Robbins 2018).

Natural hosts. Mammals, primarily rodents, but also other mammals and birds. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Ixodes acutitarsus* (Karsch, 1880)Ixodes gigas* Warburton, 1910

Geographic distribution. Southeast Asia, Japan, India (Guglielmone and Robbins 2018).

Natural hosts. Immature stages are typically on rodents; adults on ungulates. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Ixodes angustus Neumann, 1899

Geographic distribution. North America, Russia, Japan (Spencer 1963; Guglielmone and Robbins 2018).

Natural hosts. Mammals, primarily rodents, but also other mammals and birds. Zoonotic on humans (Spencer 1963; Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Ixodes apronophorus Schülze, 1924

Geographic distribution. Europe, Asia (Guglielmone and Robbins 2018).

Natural hosts. Mammals, primarily rodents, but also other mammals, reptiles, and birds. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Ixodes asanumai Kitaoka, 1973

Geographic distribution. Japan (Okono 2010).

Natural hosts. Reptiles, rarely mammals and birds. Zoonotic on humans (Okono 2010).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Ixodes australiensis Neumann, 1904

Geographic distribution. Australia (Kwak 2018).

Natural hosts. Marsupials, dogs, bovids. Zoonotic on humans (Kwak 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Ixodes baergi Cooley & Kohls, 1942

Geographic distribution. North America (Walker et al. 1998).

Natural hosts. Passerine birds. Zoonotic on humans (Walker et al. 1998).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Ixodes banksi Bishopp, 1911

Geographic distribution. North America (Guglielmone and Robbins 2018).

Natural hosts. Rodents, carnivores. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Ixodes boliviensis Neumann, 1904

Ixodes bicornis Neumann, 1906

Geographic distribution. Central and South America (Guglielmone and Robbins 2018).

Natural hosts. Several groups of mammals and birds. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Ixodes brunneus Koch, 1844

Ixodes ricinus californicus Banks, 1908

Geographic distribution. Western Hemisphere (Guglielmone and Robbins 2018).

Natural hosts. Birds. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Ixodes canisuga Johnston, 1849

Geographic distribution. Europe (Guglielmone and Robbins 2018).

Natural hosts. Carnivores; immature stages occasionally on birds. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Ixodes cavipalpus* Nuttall & Warburton, 1908**

Ixodes rubicundus limbatus Neumann, 1908

Geographic distribution. Sub-Saharan Africa (Guglielmone and Robbins 2018).

Natural hosts. Ungulates, primarily cattle. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Ixodes columnae* Takada & Fujita, 1992**

Geographic distribution. Japan (Takada and Fujita 1992).

Natural hosts. Rodents, birds. Zoonotic on humans (Takada and Fujita 1992).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Ixodes confusus* Roberts, 1960**

Geographic distribution. Australia, Papua New Guinea (Roberts 1970).

Natural hosts. Ungulates, equids, marsupials. Zoonotic on humans (Roberts 1970).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Ixodes cookei* Packard, 1869**

Ixodes cruciarius Fitch, 1872

Geographic distribution. North America (Guglielmone and Robbins 2018).

Natural hosts. Several groups of mammals, including humans, and birds (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. Powassan virus lineage I (Khan et al. 2019).

***Ixodes cornuatus* Roberts, 1960**

Geographic distribution. Australia (Guglielmone and Robbins 2018).

Natural hosts. Mammals, primarily carnivores and rodents, and birds. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Ixodes crenulatus Koch, 1844

Geographic distribution. Europe, Asia (Guglielmone and Robbins 2018).

Natural hosts. Mammals, primarily rodents, carnivores, lagomorphs. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Ixodes cumulatimpunctatus Schülze, 1943

Ixodes pseudorasus Arthur & Burrow, 1957

Geographic distribution. Sub-Saharan Africa (Guglielmone and Robbins 2018).

Natural hosts. Several groups of mammals and birds. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Ixodes dentatus Marx in Neumann, 1899

Geographic distribution. North America (Guglielmone and Robbins 2018).

Natural hosts. Immature stages on a variety of mammals and birds; adults on mammals, primarily lagomorphs and carnivores. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Ixodes eichhorni Nuttall, 1916

Geographic distribution. Southeast Asia, Australia, Pacific Islands (Guglielmone and Robbins 2018).

Natural hosts. Birds. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Ixodes feicalis Warburton & Nuttall, 1909

Geographic distribution. Australia, Papua New Guinea (Domrow and Derrick 1964).

Natural hosts. Marsupials. Zoonotic on humans (Domrow and Derrick 1964).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment

Site in human host. Skin

Vectored pathogens. None

Ixodes festai Rondelli, 1926

Geographic distribution. Southern Europe, northern Africa (Guglielmone and Robbins 2018).

Natural hosts. Birds. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Ixodes frontalis (Panzer, 1798)

Geographic distribution. Europe, Middle East (Guglielmone and Robbins 2018).

Natural hosts. Birds. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Ixodes gibbosus Nuttall, 1916

Geographic distribution. Southern, eastern Europe, Middle East (Guglielmone and Robbins 2018).

Natural hosts. Ungulates, primarily bovids, rarely birds. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Ixodes granulatus Supino, 1897

Geographic distribution. Southeast Asia (Guglielmone and Robbins 2018).

Natural hosts. Mammals, birds, reptiles. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Ixodes hexagonus* Leach, 1815**

Geographic distribution. Europe (Guglielmone and Robbins 2018).

Natural hosts. Mammals, primarily hedgehogs, also birds. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Ixodes holocyclus* Neumann, 1899**

Geographic distribution. East coastal Australia (Guglielmone and Robbins 2018).

Natural hosts. Marsupials, primarily bandicoots, also livestock, cats, dogs, and humans, and birds (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. *Rickettsia australis* (Queensland tick typhus); also implicated in tick paralysis (Hall-Mendelin et al. 2011; Stewart et al. 2017).

***Ixodes kaschmiricus* Pomerantsev, 1948**

Geographic distribution. Central and East Asia (Guglielmone and Robbins 2018).

Natural hosts. Mammals, primarily bovids, carnivores, rodents. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Ixodes kazakstani* Olenov & Sorokoumov, 1934**

Geographic distribution. Central Asia (Guglielmone and Robbins 2018).

Natural hosts. Several mammals and birds. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Ixodes kingi* Bishopp, 1911**

Geographic distribution. North America (Guglielmone and Robbins 2018).

Natural hosts. Mammals, including carnivores, rodents, lagomorphs. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Ixodes kohlsi Arthur, 1955

Geographic distribution. Australia (Roberts 1970).

Natural hosts. Birds. Zoonotic on humans (Roberts 1970).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Ixodes laguri Olenov, 1929

Geographic distribution. Central and Eastern Europe, Asia (Guglielmone and Robbins 2018).

Natural hosts. Mammals, primarily rodents, lagomorphs, shrews, hedgehogs, carnivores. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Ixodes marxi Banks, 1908

Geographic distribution. North America (Guglielmone and Robbins 2018).

Natural hosts. Mammals, primarily rodents; also birds. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. Powassan virus lineage I (Khan et al. 2019).

Ixodes monospinosus Saito, 1968

Geographic distribution. Japan (Guglielmone and Robbins 2018).

Natural hosts. Mammals, including bovids, deer, rodents, carnivores, shrews. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Ixodes muniensis Arthur & Burrow, 1957

Geographic distribution. Sub-Saharan Africa (Guglielmone and Robbins 2018).

Natural hosts. Mammals, including bovids, pigs, giraffes, hyrax, carnivores, rodents. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.
Vectored pathogens. None.

Ixodes muris Bishopp & Smith, 1937

Geographic distribution. North America (Guglielmone and Robbins 2018).

Natural hosts. Mammals, primarily rodents. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Ixodes myrmecobii Roberts, 1962

Geographic distribution. Australia (Guglielmone and Robbins 2018).

Natural hosts. Marsupials, rarely other mammals, birds. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Ixodes nipponensis Kitaoka & Saito, 1967

Geographic distribution. Southeast Asia, Japan, Russia (Guglielmone and Robbins 2018).

Natural hosts. Mammals, primarily carnivores, bovids, deer, lagomorphs. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Ixodes ovatus Neumann, 1899

Ixodes japonensis Neumann, 1904

Ixodes frequens Ogura & Takada, 1927

Ixodes carinatus Kishida, 1930

Ixodes lindbergi Santos Dias, 1959

Geographic distribution. Central, Eastern, Southeastern Asia (Guglielmone and Robbins 2018).

Natural hosts. Mammals, including rodents, lagomorphs, carnivores, shrews, humans, and birds (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin, ear (Iwasaki et al. 2007).

Vectored pathogens. *Rickettsia japonica* (Japanese spotted fever) (Li et al. 2019).

***Ixodes pacificus* Cooley & Kohls, 1943**

Ixodes californicus Banks, 1904

Geographic distribution. Western North America (Padgett and Lane 2001; Guglielmone and Robbins 2018).

Natural hosts. Small mammals, birds, and reptiles. Zoonotic on humans (Padgett and Lane 2001; Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. *Borrelia burgdorferi* (Lyme disease), *B. miyamotoi* (tick-borne relapsing fever), *Anaplasma phagocytophilum* (human granulocytic anaplasmosis) (Padgett and Lane 2001).

***Ixodes pararicinus* Keirans & Clifford in Keirans et al. 1985**

Geographic distribution. Argentina, Colombia, Peru (Saracho-Bottero et al. 2018).

Natural hosts. Immature stages are primarily on small mammals and birds. Adults are on ungulates, including bovids, peccaries, and deer. Zoonotic on humans (Saracho-Bottero et al. 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Ixodes pavlovskyi* Pomerantzev, 1946**

Geographic distribution. Central Asia, Russia, Japan (Guglielmone and Robbins 2018).

Natural hosts. Birds and mammals, including rodents, lagomorphs, carnivores, hedgehogs, shrews, and humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Ixodes persulcatus* (Schülze, 1930)**

Geographic distribution. Europe, Northern Asia (Guglielmone and Robbins 2018).

Natural hosts. Mammals, including dogs, deer, rodents, lagomorphs, and humans, and ground-nesting birds (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin, ear (Kogoashiwa et al. 2009),

Vectored pathogens. *Borrelia miyamotoi* (tick-borne relapsing fever), *B. garinii*, tick-borne encephalitis viruses (TBE) (Jaenson et al. 2016).

***Ixodes petauristae* Warburton, 1933**

Geographic distribution. India, Sri Lanka (Guglielmone and Robbins 2018).

Natural hosts. Mammals, primarily rodents. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Ixodes pilosus* Koch, 1844**

Geographic distribution. Southern Africa (Guglielmone and Robbins 2018).

Natural hosts. Mammals, primarily ungulates. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Ixodes rageaui* Arthur, 1958**

Geographic distribution. Sub-Saharan Africa (Guglielmone and Robbins 2018).

Natural hosts. Non-human primates. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Ixodes rarus* Neumann, 1899**

Geographic distribution. Sub-Saharan Africa (Guglielmone and Robbins 2018).

Natural hosts. Mammals, including ungulates, carnivores, rodents, and hyrax, and birds, Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Ixodes redikorzevi* Olenev, 1927**

Ixodes theodori Warburton, 1927

Geographic distribution. Europe, North Africa, Middle East, Asia (Guglielmone and Robbins 2018).

Natural hosts. Mammals, including rodents, hedgehogs, lagomorphs, and carnivores, birds, and reptiles. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Ixodes ricinus (Linnaeus, 1758)

Acarus reduvius Linnaeus, 1758

Geographic distribution. Europe, North Africa, Middle East, northern Asia (Guglielmone and Robbins 2018).

Natural hosts. Many mammals, including humans, and birds. Immature stages are typically on small mammals, reptiles, and birds. Adults are on ungulates and dogs (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. *Borrelia burgdorferi* sensu lato (Lyme disease), *B. miyamotoi* (tick-borne relapsing fever), *Coxiella burnetii* (Q fever), *Babesia divergens* (babesiosis), tick-borne encephalitis viruses (TBE), louping ill virus (Davidson et al. 1991; Gern 2005; Rizzoli et al. 2014).

Ixodes rubicundus Neumann, 1904

Geographic distribution. South Africa (Guglielmone and Robbins 2018).

Natural hosts. Mammals, including ungulates, rodents, lagomorphs, and carnivores. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Ixodes rugosus Bishopp, 1911

Geographic distribution. North America (Guglielmone and Robbins 2018).

Natural hosts. Mammals, primarily carnivores, but also opossums and rodents. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Ixodes scapularis Say, 1821

Ixodes ozarkus Cooley, 1944

Ixodes dammini Spielman et al., 1979

Geographic distribution. Eastern and Midwestern North America (Guglielmone and Robbins 2018).

Natural hosts. Immature stages are primarily on rodents and other small mammals, reptiles, small birds. Adults are on ungulates (primarily deer) and dogs. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. *Borrelia burgdorferi* sensu lato and *B. mayonii* (Lyme disease), *B. miyamotoi* (tick-borne relapsing fever), *Babesia microti* (babesiosis), *Anaplasma phagocytophilum* (human granulocytic anaplasmosis), deer tick virus (Powassan virus lineage II) (Nelder et al. 2016; Westblade et al. 2017; Khan et al. 2019; Wolf et al. 2020).

***Ixodes schillingsi* Neumann, 1901**

Geographic distribution. Sub-Saharan Africa (Guglielmone and Robbins 2018).

Natural hosts. Non-human primates. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Ixodes sculptus* Neumann, 1904**

Geographic distribution. Central Asia, Russia, Japan (Guglielmone and Robbins 2018).

Natural hosts. Primarily rodents and lagomorphs, rarely ungulates and birds. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Ixodes sinensis* Teng, 1977**

Geographic distribution. China (Guglielmone and Robbins 2018).

Natural hosts. Mammals, including ungulates, rodents, and lagomorphs, and birds. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Ixodes soricis* Gregson, 1942**

Geographic distribution. North America (Spencer 1963).

Natural hosts. Shrews and rodents. Zoonotic on humans (Spencer 1963).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Ixodes spinicoxalis Neumann, 1899

Geographic distribution. Southeast Asia (Guglielmone and Robbins 2018).

Natural hosts. Mammals, tree shrews, rodents, and carnivores, and birds. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Ixodes spinipalpis Hawden et Nuttall in Nuttall, 1916

Geographic distribution. North America (Guglielmone and Robbins 2018).

Natural hosts. Mammals, primarily rodents, but also lagomorphs and carnivores, and birds. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Ixodes tancitarius Cooley & Kohls, 1942

Geographic distribution. Mexico (Guglielmone and Robbins 2018).

Natural hosts. Rodents. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Notes. The single human record from Mexico is considered tentative (Guglielmone and Robbins 2018).

Ixodes tanuki Saito, 1964

Geographic distribution. Southeast Asia, Japan (Guglielmone and Robbins 2018).

Natural hosts. Mammals, including carnivores, rodents, and ungulates. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Ixodes tasmani Neumann, 1899

Geographic distribution. Southeast Asia, Japan (Guglielmone and Robbins 2018).

Natural hosts. Mammals, including marsupials, carnivores, and rodents. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Ixodes texanus Banks, 1909

Geographic distribution. North America (Guglielmone and Robbins 2018).

Natural hosts. Mammals, primarily carnivores. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Ixodes trainguliceps Birula, 1895

Geographic distribution. Europe, northwestern Asia (Guglielmone and Robbins 2018).

Natural hosts. Mammals, primarily shrews, but also rodents and carnivores; birds, lizards. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Ixodes turdus Nakatsudi, 1942

Geographic distribution. Southeast Asia, Japan (Guglielmone and Robbins 2018).

Natural hosts. Birds, rarely rodents. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Ixodes uriae White, 1852

Hyalomma puta Pickard-Cambridge, 1876

Geographic distribution. North America, Europe, Russia, southern Africa, Australia (Guglielmone and Robbins 2018).

Natural hosts. Predominately birds, but also mammals, including carnivores, rodents, and humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Ixodes vanidicus* Schülze, 1943**

Geographic distribution. Sub-Saharan Africa (Estrada-Pena and Jongejan 1999).

Natural hosts. Mammals, including carnivores and elephant shrews. Zoonotic on humans (Estrada-Pena and Jongejan 1999).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Ixodes ventalloi* Gil Collado, 1936**

Geographic distribution. Europe, northern Africa (Guglielmone and Robbins 2018).

Natural hosts. Mammals, including carnivores, rodents, and lagomorphs; birds, and reptiles. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Ixodes vespertilionis* Koch, 1844**

Geographic distribution. Europe, Africa, Asia (Guglielmone and Robbins 2018).

Natural hosts. Bats. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Ixodes woodi* Bishopp, 1911**

Geographic distribution. North America (Guglielmone and Robbins 2018).

Natural hosts. Mammals, primarily rodents, also carnivores. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Genus *Nosomma* Schülze, 1919***Nosomma monstrosum* (Nuttall & Warburton, 1908)**

Geographic distribution. Central and Southeast Asia (Guglielmone and Robbins 2018).

Natural hosts. Immature stage primarily on rodents and shrews. Adults primarily on bovids, deer, pigs, horses, and carnivores. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Genus *Rhipicephalus* Koch, 1844

***Rhipicephalus annulatus* (Say, 1821)**

Rhipicephalus calcaratus Birula, 1895

Geographic distribution. North America, Africa, Europe, Middle East, Central Asia (Guglielmone and Robbins 2018).

Natural hosts. Primarily cattle, but also other mammals, birds, and reptiles. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Rhipicephalus appendiculatus* Neumann, 1901**

Geographic distribution. Sub-Saharan Africa (Perry et al. 1990).

Natural hosts. Mammals, including cattle, buffalo, antelope, warthogs, equids, lagomorphs, and dogs. Zoonotic on humans (Perry et al. 1990).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Rhipicephalus armatus* Pocock, 1900**

Geographic distribution. East Africa (Guglielmone and Robbins 2018).

Natural hosts. Carnivores, bovids, and lagomorphs. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Rhipicephalus aurantiacus* Neumann, 1907**

Geographic distribution. Sub-Saharan Africa (Guglielmone and Robbins 2018).

Natural hosts. Bovids, pigs. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Rhipicephalus australis* Fuller, 1899.**

Geographic distribution. Southeast Asia, Australia, Pacific Islands (Guglielmone and Robbins 2018).

Natural hosts. Mammals, primarily bovids. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Rhipicephalus bequaerti* Zumpt, 1950**

Geographic distribution. Sub-Saharan Africa (Guglielmone and Robbins 2018).

Natural hosts. Bovids, pigs. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Rhipicephalus bursa* Canestrini et Fanzago, 1878**

Geographic distribution. Southern Europe, North Africa, Middle East, Central and East Asia (Guglielmone and Robbins 2018).

Natural hosts. Mammals, primarily bovids, and birds. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Rhipicephalus carnivoralis* Walker, 1966**

Geographic distribution. East Africa (Guglielmone and Robbins 2018).

Natural hosts. Primarily felids, also other mammals including bovids and hyrax. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Rhipicephalus complanatus* Neumann, 1911**

Geographic distribution. West Africa (Guglielmone and Robbins 2018).

Natural hosts. Pigs, bovids, carnivores, and rodents. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Rhipicephalus compositus* Neumann, 1897**

Geographic distribution. Sub-Saharan Africa (Guglielmone and Robbins 2018).

Natural hosts. Immature stages primarily on rodents. Adults primarily on bovids, pigs, carnivores, horses, rhinoceroses. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Rhipicephalus decoloratus* Koch, 1844**

Geographic distribution. Sub-Saharan Africa (Guglielmone and Robbins 2018).

Natural hosts. Primarily bovids, also other mammals, birds, and tortoises. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Rhipicephalus distinctus* Bedford, 1932**

Geographic distribution. Sub-Saharan Africa (Guglielmone and Robbins 2018).

Natural hosts. Primarily hyrax, also lagomorphs, carnivores, rodents, and elephant shrews. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Rhipicephalus evertsi* Neumann, 1897**

Geographic distribution. Sub-Saharan Africa (Guglielmone and Robbins 2018).

Natural hosts. Several groups of mammals and birds. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Rhipicephalus follis* Dönitz, 1910**

Geographic distribution. Sub-Saharan Africa (Guglielmone and Robbins 2018).

Natural hosts. Mammals, including bovids, pigs, carnivores, horses, rhinoceroses, hyrax, and rodents. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Rhipicephalus fulvus Neumann, 1913

Geographic distribution. Central and northern Africa (Guglielmone and Robbins 2018).

Natural hosts. Bovids, camels, rodents. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Rhipicephalus gertrudae Feldman-Muhsam, 1960

Geographic distribution. Southern Africa (Guglielmone and Robbins 2018).

Natural hosts. Several groups of mammals, birds, and reptiles. Immature stages primarily on rodents. Adults primarily on bovids. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Rhipicephalus glabroscutatus Du Toit, 1941

Geographic distribution. South Africa (Guglielmone and Robbins 2018).

Natural hosts. Immature stages primarily on carnivores, rodents, birds. Adults primarily on lagomorphs, horses, and hyrax. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Rhipicephalus guilhoni Morel & Vassiliades, 1963

Geographic distribution. Africa (Guglielmone and Robbins 2018).

Natural hosts. Several groups of mammals and birds. Immature stages primarily on rodents, lagomorphs. Adults primarily on bovids. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Rhipicephalus haemaphysaloides* Supino, 1897**

Geographic distribution. Central and Southeast Asia (Guglielmone and Robbins 2018).

Natural hosts. Mammals, including carnivores, rodents, bovids, and shrews; birds. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin, ear (Dilrukshi et al. 2004).

Vectored pathogens. None.

***Rhipicephalus humeralis* Tonelli-Rondelli, 1926**

Geographic distribution. East Africa (Guglielmone and Robbins 2018).

Natural hosts. Various groups of mammals. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Rhipicephalus hurti* Wilson, 1954**

Geographic distribution. East Africa (Guglielmone and Robbins 2018).

Natural hosts. Mammals, including bovids, pigs, carnivores, rhinoceroses, and rodents. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Rhipicephalus jeanneli* Neumann, 1913**

Geographic distribution. East Africa (Guglielmone and Robbins 2018).

Natural hosts. Mammals, including bovids, pigs, horses, rhinoceroses, and carnivores, and birds. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Rhipicephalus kochi* Dönitz, 1905**

Rhipicephalus neavi Warburton, 1912

Geographic distribution. Sub-Saharan Africa (Guglielmone and Robbins 2018).

Natural hosts. Mammals, including bovids, elephant shrews, lagomorphs, and pigs; birds. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Rhipicephalus longus Neumann, 1907

Geographic distribution. Sub-Saharan Africa (Guglielmone and Robbins 2018).

Natural hosts. Mammals, primarily bovids and pigs. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Rhipicephalus lunulatus Neumann, 1907

Geographic distribution. Sub-Saharan Africa (Guglielmone and Robbins 2018).

Natural hosts. Mammals, including bovids, lagomorphs, elephant shrews, and rodents; birds. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Rhipicephalus maculatus Neumann, 1901

Geographic distribution. Southern and East Africa (Guglielmone and Robbins 2018).

Natural hosts. Mammals, including bovids, rhinoceroses, carnivores, horses, and elephants; rarely reptiles. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Rhipicephalus microplus (Canestrini, 1888)

Geographic distribution. Circumtropical (Guglielmone and Robbins 2018).

Natural hosts. Several groups of mammals, especially livestock; birds and reptiles. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Rhipicephalus muehlensi Zumpt, 1943

Geographic distribution. Africa (Guglielmone and Robbins 2018).

Natural hosts. Several groups of mammals. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Rhipicephalus pilans Schülze, 1935

Geographic distribution. East Timor, Indonesia, Philippines (Guglielmone and Robbins 2018).

Natural hosts. Several groups of mammals. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Rhipicephalus planus Neumann, 1907

Rhipicephalus reichenowi Zumpt, 1943

Geographic distribution. Sub-Saharan Africa (Guglielmone and Robbins 2018).

Natural hosts. Several groups of mammals. Immature stages primarily on rodents and lagomorphs. Adults primarily on bovids and pigs. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Rhipicephalus praetextatus Gerstäcker, 1873

Geographic distribution. Africa, Yemen (Guglielmone and Robbins 2018).

Natural hosts. Several groups of mammals and birds. Immature stages primarily on rodents. Adults primarily on bovids. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Rhipicephalus parvus Dönitz, 1910

Geographic distribution. East Africa (Guglielmone and Robbins 2018).

Natural hosts. Several groups of mammals, birds, and reptiles. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Rhipicephalus pulchellus* (Gerstäcker, 1873)**

Geographic distribution. East Africa (Guglielmone and Robbins 2018).

Natural hosts. Several groups of mammals, especially bovids, and birds. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Rhipicephalus pumilio* Schülze, 1935**

Geographic distribution. Central and East Asia (Guglielmone and Robbins 2018).

Natural hosts. Several groups of mammals, including hedgehogs, rodents, and lagomorphs, and birds. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Rhipicephalus pusillus* Gil Collado, 1936**

Geographic distribution. Mediterranean Europe and Africa (Guglielmone and Robbins 2018).

Natural hosts. Several groups of mammals and birds. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Rhipicephalus rossicus* Yakimov et Kohl-Yakimova, 1911**

Geographic distribution. Eastern Europe, Asia (Guglielmone and Robbins 2018).

Natural hosts. Several groups of mammals, birds, and reptiles. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Rhipicephalus sanguineus* (Latreille, 1806)**

Geographic distribution. Worldwide (Guglielmone and Robbins 2018).

Natural hosts. Several groups of mammals, especially dogs. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. *Rickettsia rickettsii* (Rocky Mountain spotted fever), *R. conorii* (boutonnoise fever) (Dantas-Torres 2010; Dantas-Torres et al. 2012).

***Rhipicephalus schulzei* Olenev, 1929**

Geographic distribution. Central and East Asia (Guglielmone and Robbins 2018).

Natural hosts. Mammals, including rodents, lagomorphs, and carnivores; birds. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Rhipicephalus senegalensis* Koch, 1844**

Geographic distribution. Sub-Saharan Africa (Guglielmone and Robbins 2018).

Natural hosts. Several groups of mammals and birds. Immature stages primarily on carnivores, elephant shrews, and rodents. Adults primarily on bovids, especially domestic cattle, and other mammals. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Rhipicephalus simus* Koch, 1844**

Geographic distribution. Southern and East Africa (Guglielmone and Robbins 2018).

Natural hosts. Several groups of mammals. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Rhipicephalus sulcatus* Neumann, 1908**

Geographic distribution. Sub-Saharan Africa (Guglielmone and Robbins 2018).

Natural hosts. Several groups of mammals. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Rhipicephalus supertritus* Neumann, 1907**

Geographic distribution. Sub-Saharan Africa (Guglielmone and Robbins 2018).

Natural hosts. Several groups of mammals, especially bovids. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Rhipicephalus turanicus* Pomerantzev, 1936**

Geographic distribution. Hard to define but generally considered broadly distributed in Africa and Asia (Guglielmone and Robbins 2018).

Natural hosts. Several groups of mammals, including humans, birds, and reptiles (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. *Rickettsia conorii* (boutonneuse fever) (Dantas-Torres et al. 2012).

***Rhipicephalus warburtoni* Walker et Horak in Walker et al., 2000**

Geographic distribution. South Africa (Guglielmone and Robbins 2018).

Natural hosts. Mammals, including bovids, elephant shrews, lagomorphs, carnivores, horses, and rodents; birds. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Rhipicephalus zambeziensis* Walker et al., 1981**

Geographic distribution. Sub-Saharan Africa (Guglielmone and Robbins 2018).

Natural hosts. Many groups of mammals and birds. Immature stages primarily on lagomorphs. Adults primarily on bovids. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Rhipicephalus ziemanni* Neumann, 1904**

Geographic distribution. Sub-Saharan Africa (Guglielmone and Robbins 2018).

Natural hosts. Mammals, primarily carnivores and rodents. Zoonotic on humans (Guglielmone and Robbins 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. *Borrelia hispanica* (tick-borne relapsing fever) (Boinas et al. 2014; Talagrand-Reboul et al. 2018).

Ornithodoros graingeri Heisch & Guggisberg, 1953

Geographic distribution. Kenya (Heisch and Harvey 1953).

Natural hosts. Rodents. Zoonotic on humans as incidental hosts (Heisch and Harvey 1953; Talagrand-Reboul et al. 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. *Borrelia graingeri* (tick-borne relapsing fever) (Talagrand-Reboul et al. 2018).

Ornithodoros hermsi Wheeler, 1935

Geographic distribution. Western North America (Sage et al. 2017).

Natural hosts. Rodents. Zoonotic on humans as incidental hosts (Sage et al. 2017; Talagrand-Reboul et al. 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. *Borrelia hermsii* (tick-borne relapsing fever) (Sage et al. 2017; Talagrand-Reboul et al. 2018).

Ornithodoros maroccanus Velu, 1919

Geographic distribution. Mediterranean Region (Vial 2009).

Natural hosts. Many mammals, including rodents, insectivores, canids, mustelids, bats. Zoonotic on humans as incidental hosts (Vial 2009; Talagrand-Reboul et al. 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. *Borrelia hispanica* (tick-borne relapsing fever) (Talagrand-Reboul et al. 2018).

Ornithodoros moubata Murray, 1877

Geographic distribution. Sub-Saharan Africa (Vial 2009).

Natural hosts. Pigs, poultry. Zoonotic on humans as incidental hosts (Vial 2009; Talagrand-Reboul et al. 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. *Borrelia duttoni* (tick-borne relapsing fever) (Talagrand-Reboul et al. 2018).

***Ornithodoros parkeri* Cooley, 1936**

Geographic distribution. Western North America (Lopez et al. 2016).

Natural hosts. Rodents, horses. Zoonotic on humans as incidental hosts (Lopez et al. 2016; Talagrand-Reboul et al. 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. *Borrelia parkeri* (tick-borne relapsing fever) (Lopez et al. 2016; Talagrand-Reboul et al. 2018).

***Ornithodoros rudis* Karsch, 1880**

Ornithodoros venezuelensis Brumpt, 1921

Geographic distribution. Panama and South America (Faccini-Martínez Á and Botero-García 2016).

Natural hosts. Birds, primarily poultry. Zoonotic on humans as incidental hosts (Faccini-Martínez Á and Botero-García 2016; Talagrand-Reboul et al. 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. *Borrelia venezuelensis* (tick-borne relapsing fever) (Faccini-Martínez Á and Botero-García 2016; Talagrand-Reboul et al. 2018).

***Ornithodoros sonrai* Sautet & Witkowski, 1944**

Geographic distribution. Western and northern Africa (Vial 2009).

Natural hosts. Rodents, insectivores. Zoonotic on humans as incidental hosts (Vial 2009; Talagrand-Reboul et al. 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. *Borrelia crociduræ* (tick-borne relapsing fever) (Talagrand-Reboul et al. 2018).

***Ornithodoros talajae* (Guérin-Méneville, 1849)**

Geographic distribution. Southern United States, Central and South America (Cooley 1945; Faccini-Martínez Á and Botero-García 2016).

Natural hosts. Many mammals. Zoonotic on humans as incidental hosts (Cooley 1945; Talagrand-Reboul et al. 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. *Borrellia mazzottii* (tick-borne relapsing fever) (Elelu 2018; Talagrand-Reboul et al. 2018; Moraru 2019).

Ornithodoros tholozani Laboulbène & Mégnin, 1882

Geographic distribution. Middle East, North Africa, Central Asia, India (Vial 2009).

Natural hosts. Rodents, insectivores. Zoonotic on humans as incidental hosts (Vial 2009; Talagrand-Reboul et al. 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. *Borrelia persica* (tick-borne relapsing fever) (Talagrand-Reboul et al. 2018).

Ornithodoros turicata (Dugès, 1876)

Geographic distribution. Western and Midwestern North America, Mexico (Cooley 1945).

Natural hosts. Rodents, dogs. Zoonotic on humans as incidental hosts (Cooley 1945; Talagrand-Reboul et al. 2018).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. *Borrelia turicatae* (tick-borne relapsing fever) (Talagrand-Reboul et al. 2018).

Ornithodoros verrucosus Olenov et al., 1934

Geographic distribution. Eastern Europe (Vial 2009).

Natural hosts. Rodents. Zoonotic on humans as incidental hosts (Vial 2009).

Route of infection. Exposure to larval, nymph, or adult ticks in the environment.

Site in human host. Skin.

Vectored pathogens. *Borrelia caucasica* (tick-borne relapsing fever) (Talagrand-Reboul et al. 2018).

Genus *Otobius* Banks, 1912

Otobius megnini (Dugès, 1884)

Geographic distribution. Nearly worldwide wherever livestock are raised (Cooley 1945).

Natural hosts. Cattle, sheep, goats, deer, other ruminants, carnivores, rabbits. Zoonotic on humans as incidental hosts (Cooley 1945).

Route of infection. Exposure to larval or nymph ticks in the environment.

- Mandibulata Snodgrass, 1938
- Pancrustacea Zrzavý & Štys, 1997
 - Tetraconata Dohle, 2001
- Crustacea Brünnich, 1772
- Maxillopoda Dahl, 1956
- Pentastomida Diesing, 1836
- Porocephalidae Sambon, 1922

Genus *Armillifer* Sambon, 1922

***Armillifer agkistrodontis* Self & Kuntz, 1966**

Geographic distribution. China (Chen et al. 2010; Christoffersen and Assis 2013; Ioannou and Vamvoukaki 2019).

Natural hosts. Intermediate hosts are rodents. Definitive hosts are snakes. Zoonotic in humans as dead-end hosts (Chen et al. 2010; Christoffersen and Assis 2013; Ioannou and Vamvoukaki 2019).

Route of infection. Ingestion of infected intermediate or reservoir host (Chen et al. 2010; Ioannou and Vamvoukaki 2019).

Site in human host. Disseminated infection, including the mesenteries, peritoneum, lungs (Chen et al. 2010; Ioannou and Vamvoukaki 2019).

***Armillifer armillatus* Wyman, 1845**

Geographic distribution. Central and western Africa (Christoffersen and Assis 2013; Ioannou and Vamvoukaki 2019).

Natural hosts. Intermediate hosts are various mammals. Definitive hosts are pythons. Zoonotic in humans as dead-end hosts (Christoffersen and Assis 2013; Ioannou and Vamvoukaki 2019).

Route of infection. Ingestion of infected intermediate or reservoir host (Ioannou and Vamvoukaki 2019).

Site in human host. Disseminated infection, including the mesenteries, peritoneum, lungs, brain, eye. (Ioannou and Vamvoukaki 2019).

***Armillifer grandis* Hett, 1915**

Geographic distribution. Central and western Africa (Christoffersen and Assis 2013; Ioannou and Vamvoukaki 2019).

Natural hosts. Intermediate hosts are rodents. Definitive hosts are viperid snakes. Zoonotic in humans as dead-end hosts (Christoffersen and Assis 2013; Ioannou and Vamvoukaki 2019).

Route of infection. Ingestion of infected intermediate or reservoir host (Ioannou and Vamvoukaki 2019).

Site in human host. Disseminated infection, including the mesenteries, peritoneum, lungs, brain, and eye (Tappe and Büttner 2009; Ioannou and Vamvoukaki 2019).

Armillifer moniliformis (Diesing, 1835)

Geographic distribution. Southeast Asia (Christoffersen and Assis 2013; Ioannou and Vamvoukaki 2019).

Natural hosts. Intermediate hosts are various mammals. Definitive hosts are snakes. Zoonotic in humans as dead-end hosts (Christoffersen and Assis 2013; Ioannou and Vamvoukaki 2019).

Route of infection. Ingestion of infected intermediate or reservoir host (Ioannou and Vamvoukaki 2019).

Site in human host. Disseminated infection, including the mesenteries, peritoneum, lungs, liver (Latif et al. 2011; Ioannou and Vamvoukaki 2019).

Genus *Porocephalus* Humboldt, 1811

Porocephalus crotali (Humboldt, 1808)

Geographic distribution. New World (Esslinger 1962; Christoffersen and Assis 2013).

Natural hosts. Intermediate hosts are small mammals, including rodents, monkeys. Definitive hosts are snakes, primarily rattlesnakes (*Crotalus*). Zoonotic in humans as dead-end hosts (Esslinger 1962; Christoffersen and Assis 2013).

Route of infection. Ingestion of infected intermediate or reservoir host (Tappe and Büttner 2009).

Site in human host. Disseminated infection, including the mesenteries, peritoneum, lungs (Tappe and Büttner 2009).

Porocephalus subuliferum (Leuckart, 1860)

Geographic distribution. Africa (Sambon 1922; Christoffersen and Assis 2013).

Natural hosts. Intermediate and definitive hosts are snakes. Zoonotic in humans as dead-end hosts (Sambon 1922; Christoffersen and Assis 2013).

Route of infection. Ingestion of infected intermediate host (Sambon 1922).

Site in human host. Viscera (Sambon 1922; Christoffersen and Assis 2013).

Porocephalus taiwana Qui et al., 2005

Geographic distribution. Southeast Asia (Qiu et al. 2005; Yao et al. 2008; Christoffersen and Assis 2013).

Natural hosts. Intermediate hosts are unknown. Definitive hosts are snakes. Zoonotic in humans as dead-end hosts (Christoffersen and Assis 2013).

Route of infection. Ingestion of infected intermediate or reservoir host (Qiu et al. 2005).

Site in human host. Disseminated disease, including greater omentum, small intestine, liver, lungs (Qiu et al. 2005; Yao et al. 2008).

●●●●●●●●●● *Linguatulidae* Haldeman, 1851

Genus *Linguatula* Fröhlich, 1789

Linguatula serrata Fröhlich, 1789

Geographic distribution. Worldwide (Christoffersen and Assis 2013).

Natural hosts. Intermediate hosts are sheep, bovids, rodents. Definitive hosts are felids and canids. Zoonotic in humans as dead-end hosts (Christoffersen and Assis 2013).

Route of infection. Ingestion of infected intermediate or reservoir host (Tappe and Büttner 2009).

Site in human host. Disseminated infection, including the mesenteries, peritoneum, lungs, eyes (Tappe and Büttner 2009).

●●●●●●●●●● *Raillietiellidae* Sambon, 1922

Genus *Raillietiella* Sambon in Vaney & Sambon, 1910

Geographic distribution. Worldwide (Christoffersen and Assis 2013).

Natural hosts. Intermediate hosts are insects. Definitive hosts are reptiles and amphibians. Zoonotic in humans as dead-end hosts (Christoffersen and Assis 2013).

Route of infection. Ingestion of infected intermediate or reservoir host (Tappe et al. 2016b).

Site in human host. Disseminated infection, including the mesenteries, peritoneum, lungs (Tappe et al. 2016b).

Notes. *Raillietiella* isolates from human hosts have not been characterized at the species level. Cases of *R. hemidactyli* Hett, 1934 and *R. gehyrae* Bovien, 1927 from Vietnam were not confirmed morphologically (Tappe et al. 2016b).

●●●●●●●●●● **Sebekidae Sambon, 1922**

Genus *Leiperia* Sambon, 1922

***Leiperia cincinallis* Sambon, in Vaney and Sambon 1910**

Geographic distribution. Africa (Christoffersen and Assis 2013).

Natural hosts. Intermediate hosts are freshwater fish. Definitive hosts are crocodiles and turtles. Zoonotic in humans as dead-end hosts (Christoffersen and Assis 2013).

Route of infection. Presumably ingestion of infected intermediate or paratenic host.

Site in human host. Intestine (?) (Fain 1960, 1961)

Notes. *Leiperia cincinallis* has been reported from the stool of a European woman in Zaire (Fain 1960, 1961), but it is believed this may represent spurious passage after ingesting infected fish (Christoffersen and Assis 2015).

Genus *Sebekia* Sambon, 1922

Geographic distribution. Australia, South America, Southeast Asia, southern Africa (Christoffersen and Assis 2013).

Natural hosts. Intermediate hosts are freshwater fish. Definitive hosts are crocodiles and snakes (Christoffersen and Assis 2013).

Route of infection. Presumably by ingestion of infected intermediate or paratenic host.

Site in human host. Skin (Mairena et al. 1989).

Notes. A single human case of *Sebekia*, manifesting as dermatitis in a woman in Costa Rica (Mairena et al. 1989) was not identified to the species level.

●●●●●●●●●● **Hexapoda Latreille, 1825**

●●●●●●●●●● **Insecta Linnaeus, 1758**

●●●●●●●●●● **Hemiptera Linnaeus, 1758**

●●●●●●●●●● **Reduviidae Latreille, 1807**

Genus *Meccus* Stål, 1859

***Meccus pallidipennis* (Stål, 1872)**

Geographic distribution. Mexico (Franzini-Junior et al. 2018).

Natural hosts. Various mammals. Zoonotic on humans (Franzini-Junior et al. 2018).

Route of infection. Exposure to nymphs and adults in the environment.

Site in human host. Skin.

Vectored pathogens. *Trypanosoma cruzi* (Chagas Disease, American trypanosomiasis) (Franzini-Junior et al. 2018).

Genus *Panstrongylus* Berg, 1879***Panstrongylus geniculatus* (Latreille, 1811)**

Geographic distribution. Central and South America (Melo et al. 2018).

Natural hosts. Armadillos. Zoonotic on humans (Melo et al. 2018).

Route of infection. Exposure to nymphs and adults in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Panstrongylus megistus* (Burmeister, 1835)**

Geographic distribution. Argentina, Bolivia, Brazil, Paraguay, Uruguay (Jurberg and Galvão 2006).

Natural hosts. Various mammals. Zoonotic on humans (Jurberg and Galvão 2006).

Route of infection. Exposure to nymphs and adults in the environment.

Site in human host. Skin.

Vectored pathogens. *Trypanosoma cruzi* (Chagas disease, American trypanosomiasis) (Jurberg and Galvão 2006).

Genus *Paratriatoma* Barber, 1938***Paratriatoma hirsuta* Barber, 1938**

Geographic distribution. Southwestern USA, Mexico (Bern et al. 2011).

Natural hosts. Rodents. Zoonotic on humans (Bern et al. 2011).

Route of infection. Exposure to nymphs and adults in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Genus *Rhodnius* Stål, 1859***Rhodnius prolixus* Stål, 1859**

Geographic distribution. Central and South America (Jurberg and Galvão 2006).

Natural hosts. Various mammals. Zoonotic on humans (Jurberg and Galvão 2006).

Route of infection. Exposure to nymphs and adults in the environment.

Site in human host. Skin.

Vectored pathogens. *Trypanosoma cruzi* (Chagas disease, American trypanosomiasis) (Jurberg and Galvão 2006).

Genus *Triatoma* Laporte de Castelnaud, 1832***Triatoma brasiliensis* Neiva, 1911**

Geographic distribution. Brazil (Jurberg and Galvão 2006).

Natural hosts. Various mammals. Zoonotic on humans (Jurberg and Galvão 2006).

Route of infection. Exposure to nymphs and adults in the environment.

Site in human host. Skin.

Vectored pathogens. *Trypanosoma cruzi* (Chagas disease, American trypanosomiasis) (Jurberg and Galvão 2006).

***Triatoma dimidiata* (Latrielle, 1811)**

Geographic distribution. Central and South America (Jurberg and Galvão 2006).

Natural hosts. Various mammals. Zoonotic on humans (Jurberg and Galvão 2006).

Route of infection. Exposure to nymphs and adults in the environment.

Site in human host. Skin.

Vectored pathogens. *Trypanosoma cruzi* (Chagas disease, American trypanosomiasis) (Jurberg and Galvão 2006).

***Triatoma gerstaeckeri* (Stål, 1859)**

Geographic distribution. Southwestern USA, Mexico (Wozniak et al. 2015).

Natural hosts. Various mammals. Zoonotic on humans (Wozniak et al. 2015).

Route of infection. Exposure to nymphs and adults in the environment.

Site in human host. Skin.

Vectored pathogens. None (*T. gerstaeckeri* has been found to be naturally infected with *Trypanosoma cruzi* but its role as a natural vector for human disease is not fully understood) (Reisenman et al. 2010).

***Triatoma infestans* (Klug in Meyen, 1834)**

Geographic distribution. South America (Jurberg and Galvão 2006).

Natural hosts. Various mammals. Zoonotic on humans (Jurberg and Galvão 2006).

Route of infection. Exposure to nymphs and adults in the environment.

Site in human host. Skin.

Vectored pathogens. *Trypanosoma cruzi* (Chagas disease, American trypanosomiasis) (Jurberg and Galvão 2006).

***Triatoma lectularia* (Stål, 1859)**

Geographic distribution. Southern and southeastern USA, Mexico (Bern et al. 2011).

Natural hosts. Various mammals. Zoonotic on humans (Bern et al. 2011).

Route of infection. Exposure to nymphs and adults in the environment.

Site in human host. Skin.

Vectored pathogens. None (*T. lectularia* has been found to be naturally infected with *Trypanosoma cruzi* but its role as a natural vector for human disease is not fully understood) (Bern et al. 2011).

Triatoma protracta (Uhler, 1894)

Geographic distribution. Southwestern USA, Mexico (Bern et al. 2011).

Natural hosts. Woodrats (*Neotoma*). Zoonotic on humans (Bern et al. 2011).

Route of infection. Exposure to nymphs and adults in the environment.

Site in human host. Skin.

Vectored pathogens. None (*T. protracta* has been found to be naturally infected with *Trypanosoma cruzi* but its role as a natural vector for human disease is not fully understood) (Bern et al. 2011).

Triatoma pseudomaculata Correa & Espínola, 1964

Geographic distribution. Brazil (Ribeiro et al. 2019; Soares et al. 2000).

Natural hosts. Various mammals. Zoonotic on humans (Soares et al. 2000).

Route of infection. Exposure to nymphs and adults in the environment.

Site in human host. Skin.

Vectored pathogens. *Trypanosoma cruzi* (Chagas disease, American trypanosomiasis) (Soares et al. 2000; Ribeiro et al. 2019).

Triatoma recurva (Stål, 1868)

Geographic distribution. Southwestern USA, Mexico (Bern et al. 2011).

Natural hosts. Rodents. Zoonotic on humans (Bern et al. 2011).

Route of infection. Exposure to nymphs and adults in the environment.

Site in human host. Skin.

Vectored pathogens. None (*T. recurva* has been found to be naturally infected with *Trypanosoma cruzi* but its role as a natural vector for human disease is not fully understood) (Bern et al. 2011).

Triatoma rubida (Uhler, 1894)

Geographic distribution. Southwestern USA, Central America (Bern et al. 2011).

Natural hosts. Rodents. Zoonotic on humans (Bern et al. 2011).

Route of infection. Exposure to nymphs and adults in the environment.

Site in human host. Skin.

Vectored pathogens. None (*T. rubida* has been found to be naturally infected with *Trypanosoma cruzi* but its role as a natural vector for human disease is not fully understood) (Reisenman et al. 2010; Bern et al. 2011).

***Triatoma rubrofasciata* (De Geer, 1773)**

Geographic distribution. Eastern North America, Southeast Asia, Pacific Islands (Bern et al. 2011; Huang et al. 2018).

Natural hosts. Rodents. Zoonotic on humans (Bern et al. 2011).

Route of infection. Exposure to nymphs and adults in the environment.

Site in human host. Skin.

Vectored pathogens. None; implicated in anaphylactic shock from bites (Huang et al. 2018).

***Triatoma sanguisuga* (LeConte, 1855)**

Geographic distribution. Eastern United States (Bern et al. 2011).

Natural hosts. Several groups of mammals, birds, reptiles, and amphibians. Zoonotic on humans (Bern et al. 2011).

Route of infection. Exposure to nymphs and adults in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Triatoma sordida* (Stål, 1859)**

Geographic distribution. South America (Crocco and Catalá 1997; Ribeiro et al. 2019).

Natural hosts. Various mammals. Zoonotic on humans (Crocco and Catalá 1997).

Route of infection. Exposure to nymphs and adults in the environment.

Site in human host. Skin

Vectored pathogens. *Trypanosoma cruzi* (Chagas Disease, American trypanosomiasis) (Ribeiro et al. 2019).

●●●●●●●●●● **Cimicidae Latreille, 1802****Genus *Cimex* Linnaeus, 1758*****Cimex adjunctus* Barber, 1939**

Geographic distribution. Eastern North America (Goddard 2012).

Natural hosts. Bats. Zoonotic on humans as incidental hosts (Goddard 2012).

Route of infection. Exposure to nymphs and adults in the environment.

Site in human host. Skin.

Vectored pathogens. None.

***Cimex hemipterus* (Fabricius, 1802)**

Geographic distribution. Circumtropical (Usinger 1966; Mathison and Pritt 2021).

Natural hosts. Humans (Usinger 1966; Mathison and Pritt 2021).

Route of infection. Exposure to nymphs and adults in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Cimex lectularius Linnaeus, 1758

Geographic distribution. Worldwide (Usinger 1966; Mathison and Pritt 2021).

Natural hosts. Humans (Usinger 1966; Mathison and Pritt 2021).

Route of infection. Exposure to nymphs and adults in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Cimex pilosellus (Horváth, 1910)

Geographic distribution. Western North America (Goddard 2012).

Natural hosts. Bats. Zoonotic on humans as incidental hosts (Goddard 2012).

Route of infection. Exposure to nymphs and adults in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Cimex pipistrelli Jenyns, 1839

Geographic distribution. Europe and Central Asia (Goddard 2012).

Natural hosts. Bats. Zoonotic on humans as incidental hosts (Goddard 2012).

Route of infection. Exposure to nymphs and adults in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Genus *Leptocimex* Roubaud, 1913

Leptocimex boueti (Brumpt, 1910)

Geographic distribution. Africa (Miller 2008).

Natural hosts. Bats. Zoonotic on humans as incidental hosts (Miller 2008).

Route of infection. Exposure to nymphs and adults in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Genus *Haemosiphon* Champion, 1900

Haemosiphon inodora (Dugès, 1892)

Geographic distribution. North and Central America (Mullen and Durden 2019).

Natural hosts. Birds, primarily poultry. Zoonotic on humans as incidental hosts (Mullen and Durden 2019).

Route of infection. Exposure to nymphs and adults in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Genus *Hesperocimex* List, 1925

Hesperocimex coloradensis List, 1925

Geographic distribution. North and Central America (Mullen and Durden 2019).

Natural hosts. Birds. Zoonotic on humans as incidental hosts (Mullen and Durden 2019).

Route of infection. Exposure to nymphs and adults in the environment.

Site in human host. Skin.

Vectored pathogens. None.

Genus *Oeciacus* Stål, 1873

Oeciacus vicarius Horváth, 1912

Geographic distribution. North America (Mullen and Durden 2019).

Natural hosts. Birds, primarily chimney swifts (*Chaetura*). Zoonotic on humans as incidental hosts (Mullen and Durden 2019).

Route of infection. Exposure to nymphs and adults in the environment.

Site in human host. Skin.

Vectored pathogens. None.

●●●●●●●●●● Psocodea Hennig, 1966

Anoplura Leach, 1815

Phthiraptera Haeckel, 1896

●●●●●●●●●● Pediculidae Leach, 1817

Genus *Pediculus* Linnaeus, 1758

Pediculus humanus capitis De Geer, 1778

Geographic distribution. Worldwide (Veracx and Raoult 2012; Meister and Ochsendorf 2016).

Natural hosts. Humans (Veracx and Raoult 2012; Bonilla et al. 2013).

Route of infection. Direct person-to-person contact, contaminated fomites (Veracx and Raoult 2012; Bonilla et al. 2013).

Site in human host. Hair shafts, commonly on the scalp (Veracx and Raoult 2012; Bonilla et al. 2013; Meister and Ochsendorf 2016).

Vectored pathogens. None.

***Pediculus humanus humanus* Linnaeus, 1758**

Pediculus corporis De Geer, 1778

Geographic distribution. Worldwide (Veracx and Raoult 2012).

Natural hosts. Humans (Veracx and Raoult 2012; Bonilla et al. 2013).

Route of infection. Direct person-to-person contact, contaminated fomites (Bonilla et al. 2013; Veracx and Raoult 2012).

Site in human host. Skin; *P. h. humanus* primarily lives off the human host and only migrates over to feed (Bonilla et al. 2013).

Vectored pathogens. *Rickettsia prowazekii* (epidemic typhus), *Bartonella quintana* (trench fever), *Borrelia recurrentis* (louse-borne relapsing fever) (Bonilla et al. 2013).

●●●●●●●●●● **Pthiridae Ewing, 1929**

Genus *Pthirus* Leach, 1815

Phthirus, misspelling

***Pthirus pubis* (Linnaeus, 1758)**

Geographic distribution. Worldwide (Anderson and Chaney 2009).

Natural hosts. Humans (Anderson and Chaney 2009).

Route of infection. Direct person-to-person contact (Anderson and Chaney 2009).

Site in human host. Hair shafts, usually coarse hair (pubic hair, eyebrows, eye lashes, facial hair) (Anderson and Chaney 2009).

Vectored pathogens. None.

●●●●●●●●●● **Coleoptera Linnaeus, 1758**

●●●●●●●●●● **Ptinidae Latreille, 1802**

Genus *Stegobium* Motschulsky, 1860

***Stegobium paniceum* (Linnaeus, 1758)**

Geographic distribution. Worldwide (Bousquet 1990).

Natural hosts. None; causes facultative canthariasis (infestation with beetles) in humans (Smadi et al. 2014).

Route of infection. Contamination from the environment (Smadi et al. 2014).

Site in human host. Skin (Smadi et al. 2014).

Vectored pathogens. None.

Notes. This species was reported from the skin of a patient with lupus after treatment for facultative myiasis caused by *Lucilia sericata* (Smadi et al. 2014). This appears to represent facultative canthariasis, with the beetles feeding on dead and/or dried skin.

Genus *Trogoderma* Dejean, 1821

Geographic distribution. Worldwide (Bousquet 1990).

Natural hosts. None; causes facultative canthariasis in humans (Smadi et al. 2014).

Route of infection. Contamination from the environment (Smadi et al. 2014).

Site in human host. Skin (Smadi et al. 2014).

Vectored pathogens. None

Notes. A member of this genus was reported from the skin of a patient with lupus after treatment for facultative myiasis caused by *Lucilia sericata* (Smadi et al. 2014). This appears to represent facultative canthariasis, with the beetles feeding on dead and/or dried skin.

●●●●●●●●●● Tenebrionidae Latreille, 1802

Genus *Tenebrio* Linnaeus, 1758

Tenebrio molitor Linnaeus, 1758

Geographic distribution. Worldwide (Bousquet 1990).

Natural hosts. None; causes facultative canthariasis in humans (Rodriguez-Morales 2018).

Route of infection. Contamination from the environment (Rodriguez-Morales 2018).

Site in human host. Skin ulcer (Rodriguez-Morales 2018).

Vectored pathogens. None.

Notes. *Tenebrio molitor* is a cosmopolitan pest of stored grains and other food-stuffs. A larva of *T. molitor* was discovered in a skin ulcer of an AIDS patient in rural Colombia (Rodriguez-Morales 2018). This case represents facultative canthariasis. Previous reports of *T. molitor* from the human intestinal tract (Palmer 1946) or urinary tract (Aelami et al. 2019) probably represents spurious passage, environmental contamination, or urethral sounding.

●●●●●●●●●● Siphonaptera Latreille, 1824
 ●●●●●●●●●● Heteropsyllidae Baker, 1904

Genus *Tunga* Jarocki, 1838

Tunga penetrans (Linnaeus, 1758)

Geographic distribution. Central and South America, Caribbean, Africa, Madagascar, Central Asia (Feldmeier et al. 2014).

Natural hosts. Many mammals. Zoonotic on humans (Feldmeier et al. 2014).

Route of infection. Contact with contaminated soil (e.g., walking barefoot) (Feldmeier et al. 2014).

Site in human host. Subcutaneous, especially on the bottom of feet and between toes (Feldmeier et al. 2014).

Vectored pathogens. None.

Tunga trimamillata Pampiglione et al., 2002

Geographic distribution. South America (Peru, Bolivia, Brazil) (Feldmeier et al. 2014).

Natural hosts. Mammals, primarily goats, cattle, pigs. Zoonotic on humans (Feldmeier et al. 2014).

Route of infection. Contact with contaminated soil (e.g., walking barefoot) (Feldmeier et al. 2014).

Site in human host. Subcutaneous, especially on the bottom of feet and between toes (Feldmeier et al. 2014).

Vectored pathogens. None.

●●●●●●●●●● Ceratophyllidae Dampf, 1908

Genus *Nodopsyllus* Jordan, 1933

Nodopsyllus fasciatus (Bosc, 1800)

Geographic distribution. Worldwide (Bitam et al. 2010).

Natural hosts. Rodents. Zoonotic on humans (Bitam et al. 2010).

Route of infection. Exposure to adults in the environment.

Site in human host. Skin.

Vectored pathogens. *Hymenolepis nana* (dwarf tapeworm disease), *H. diminuta* (rate tapeworm disease), *Yersinia pestis* (plague) (Bitam et al. 2010).

Genus *Oropsylla* Wagner & Ioff, 1926***Oropsylla montana* (Baker, 1895)**

Geographic distribution. Western North America (Hinnebusch et al. 2017).

Natural hosts. Rodents, primarily prairie dogs and ground squirrels. Zoonotic on humans (Hinnebusch et al. 2017).

Route of infection. Exposure to adults in the environment.

Site in human host. Skin.

Vectored pathogens. *Yersinia pestis* (plague) (Hinnebusch et al. 2017).

●●●●●●●●●● Pulicidae Billberg, 1820**Genus *Ctenocephalides* Stiles & Collins, 1930*****Ctenocephalides canis* (Curtis, 1826)**

Geographic distribution. Worldwide (Mathison and Pritt 2014).

Natural hosts. Wild and domestic canids and felids. Zoonotic on humans (Mathison and Pritt 2014).

Route of infection. Exposure to adults in the environment.

Site in human host. skin.

Vectored pathogens. *Dipylidium caninum* (dog tapeworm disease), *Hymenolepis nana* (dwarf tapeworm disease), *H. diminuta* (rat tapeworm disease) (Mathison and Pritt 2014).

***Ctenocephalides felis* (Bouché, 1835)**

Geographic distribution. Worldwide (Mathison and Pritt 2014).

Natural hosts. Wild and domestic canids and felids. Zoonotic on humans (Mathison and Pritt 2014).

Route of infection. Exposure to adults in the environment.

Site in human host. skin.

Vectored pathogens. *Dipylidium caninum* (dog tapeworm disease), *Hymenolepis nana* (dwarf tapeworm disease), *H. diminuta* (rate tapeworm disease), *Bartonella henselae* (cat-scratch disease), *Rickettsia felis* (feline rickettsiae), *Acanthocheilonema reconditum* (ocular filariasis) (Orihel and Eberhard 1998; Mathison and Pritt 2014).

Genus *Echidnophaga* Olliff, 1886***Echidnophaga gallinacea* (Westwood, 1875)**

Geographic distribution. Worldwide (Carlson and Fox 2009).

Natural hosts. Many birds and mammals. Zoonotic on humans (Carlson and Fox 2009).

Route of infection. Exposure to adults in the environment; exposure to infected birds (Carlson and Fox 2009).

Site in human host. Skin.

Vectored pathogens. None.

Genus *Synopsyllus* Wagner & Roubaud, 1932

***Synopsyllus fonquerniei* Wagner & Roubaud, 1932**

Geographic distribution. Madagascar (Kreppel et al. 2016).

Natural hosts. Rats. Zoonotic on humans (Kreppel et al. 2016).

Route of infection. Exposure to adults in the environment.

Site in human host. Skin.

Vectored pathogens. *Yersinia pestis* (plague) (Kreppel et al. 2016).

Genus *Pulex* Linnaeus, 1758

***Pulex irritans* Linnaeus, 1758**

Geographic distribution. Worldwide (Mathison and Pritt 2014).

Natural hosts. Many mammals, including humans (Mathison and Pritt 2014).

Route of infection. Exposure to adults in the environment.

Site in human host. Skin.

Vectored pathogens. *Dipylidium caninum* (dog tapeworm disease), *Hymenolepis nana* (dwarf tapeworm disease), *H. diminuta* (rate tapeworm disease) (Mathison and Pritt 2014).

Genus *Xenopsylla* Glinkiewicz, 1907

***Xenopsylla brasiliensis* (Baker, 1904)**

Geographic distribution. Africa, South America, India (Miarinjara et al. 2016; Mohammadi et al. 2021).

Natural hosts. Rats. Zoonotic on humans (Miarinjara et al. 2016; Mohammadi et al. 2021).

Route of infection. Exposure to adults in the environment.

Site in human host: Skin.

Vectored pathogens. *Yersinia pestis* (plague) (Miarinjara et al. 2016; Mohammadi et al. 2021).

***Xenopsylla cheopis* (Rothschild, 1903)**

Geographic distribution. Worldwide (Perry and Fetherston 1997).

Natural hosts. Mammals, primarily rodents (rats). Zoonotic on humans (Perry and Fetherston 1997).

Route of infection. Exposure to adults in the environment.

Site in human host. Skin.

Vectored pathogens. *Yersinia pestis* (plague), *Rickettsia typhi* (murine (epidemic) typhus) (Mathison and Pritt 2014; Perry and Fetherston 1997).

●●●●●●●●●● **Diptera Linnaeus, 1758**

●●●●●●●●●● **Phoridae Curtis, 1833**

Genus *Megaselia* Rondani, 1856***Megaselia scalaris* Loew, 1866**

Geographic distribution. Worldwide (Disney 2008).

Natural hosts. None, agent of facultative and incidental myiasis in humans (Disney 2008).

Route of infection. Oviposition on pre-existing wounds (Disney 2008).

Site in human host. Soft tissues, pre-existing wounds (Hira et al. 2004; Disney 2008).

Notes. Although this species is known to cause facultative myiasis in pre-existing wounds of soft tissues and the upper respiratory tract (Hira et al. 2004; Disney 2008), reports from urine (Ghavami and Djalilvand 2015) should be met with caution.

●●●●●●●●●● **Muscidae Latreille, 1802**

Genus *Musca* Linnaeus, 1758***Musca domestica* Linnaeus, 1758**

Musca nebulosa Fabricius, 1794

Musca hottentota Robineau-Desvoidy, 1830

Musca frontalis Macquart, 1843

Musca senegalensis Macquart, 1843

Musca santae-helenae Macquart, 1848

Musca gymnosoma Rondani, 1862

Musca niveisquama Thomson, 1869

Musca curviforceps Saccà & Rivosechi, 1956

Geographic distribution. Worldwide (Francesconi and Lupi 2012).

***Calliphora vicina* Robineau-Desvoidy, 1830**

Geographic distribution. Worldwide (Francesconi and Lupi 2012).

Natural hosts. None; causes facultative myiasis in humans (Francesconi and Lupi 2012).

Route of infection. Oviposition on pre-existing wounds (Francesconi and Lupi 2012).

Site in human host. Skin, soft tissues, pre-existing wounds (Francesconi and Lupi 2012).

Genus *Cochliomyia* Townsend, 1915***Cochliomyia hominivorax* (Coquerel, 1858)**

Geographic distribution. Central and South America, Caribbean (Scott et al. 2017; Parker et al. 2020).

Natural hosts. Most mammals, including humans (Scott et al. 2017; Parker et al. 2020).

Route of infection. Oviposition on pre-existing wounds, ears, nose, oral cavity (Francesconi and Lupi 2012).

Site in human host. Skin, soft tissues, pre-existing wounds, ears, nose, oral cavity (Francesconi and Lupi 2012).

Genus *Chrysomya* Robineau-Desvoidy, 1830***Chrysomya albiceps* (Wiedemann, 1819)**

Geographic distribution. Worldwide (Francesconi and Lupi 2012).

Natural hosts. None; causes facultative myiasis in humans and sheep (Francesconi and Lupi 2012).

Route of infection. Oviposition on pre-existing wounds (Francesconi and Lupi 2012).

Site in human host. Skin, soft tissues, pre-existing wounds (Francesconi and Lupi 2012).

***Chrysomya bezziana* (Villeneuve, 1914)**

Geographic distribution. India, Arabian Peninsula, Indonesia, Philippines, New Guinea (Francesconi and Lupi 2012).

Natural hosts. Sheep. Zoonotic on humans as incidental hosts (Francesconi and Lupi 2012).

Route of infection. Oviposition on pre-existing wounds (Francesconi and Lupi 2012).

Site in human host. Skin, soft tissues, pre-existing wounds (Francesconi and Lupi 2012).

***Chrysomya megacephala* (Fabricius, 1794)**

Geographic distribution. Worldwide (Francesconi and Lupi 2012).

Natural hosts. None; causes facultative myiasis in humans and other mammals (Francesconi and Lupi 2012).

Route of infection. Oviposition on pre-existing wounds (Francesconi and Lupi 2012).

Site in human host. Skin, soft tissues, pre-existing wounds (Francesconi and Lupi 2012).

***Chrysomya rufifacies* (Macquart, 1842)**

Geographic distribution. Worldwide (Francesconi and Lupi 2012).

Natural hosts. None; causes facultative myiasis in humans and other mammals (Francesconi and Lupi 2012).

Route of infection. Oviposition on pre-existing wounds (Francesconi and Lupi 2012).

Site in human host. Skin, soft tissues, pre-existing wounds (Francesconi and Lupi 2012).

Genus *Cordylobia* Gruenberg, 1903***Cordylobia anthropophaga* Blanchard, 1872**

Geographic distribution. East, Central, and West Africa (Francesconi and Lupi 2012).

Natural hosts. Mammals, including rodents, monkeys, ruminants, and humans (Francesconi and Lupi 2012).

Route of infection. Exposure to first-instar larvae on fomites (soil, air-drying laundry) (Francesconi and Lupi 2012).

Site in human host. Skin (Francesconi and Lupi 2012).

***Cordylobia rodhaini* Gedoelst, 1910**

Geographic distribution. Tropical Africa (Pezzi et al. 2015).

Natural hosts. Mammals, primarily rodents. Zoonotic on humans (Pezzi et al. 2015).

Route of infection. Exposure to larvae on fomites (soil, air-drying laundry) (Pezzi et al. 2015).

Site in human host. Skin (Pezzi et al. 2015).

Genus *Lucilia* Robineau-Desvoidy, 1830***Lucilia cuprina* (Wiedemann, 1830)**

Geographic distribution. Nearly worldwide, more common in warm climates (Francesconi and Lupi 2012).

Natural hosts. None, causes facultative myiasis in sheep and humans (Francesconi and Lupi 2012).

Route of infection. Oviposition on pre-existing wounds (Francesconi and Lupi 2012).

Site in human host. Skin, soft tissues, pre-existing wounds (Francesconi and Lupi 2012).

Lucilia sericata (Meigan, 1826)

Geographic distribution. Europe, North, Central, and South America (Francesconi and Lupi 2012).

Natural hosts. None, causes facultative myiasis in sheep and humans (Francesconi and Lupi 2012).

Route of infection. Oviposition on pre-existing wounds (Francesconi and Lupi 2012).

Site in human host. Skin, soft tissues, pre-existing wounds (Francesconi and Lupi 2012).

Vectored pathogens. *Wohlfahrtiimonas chitiniclastica*, *Ignatzschineria indica* (Lysaght et al. 2018).

Genus *Phormia* Robineau-Desvoidy, 1830

Phormia regina (Meigen, 1826)

Geographic distribution. Northern Hemisphere (Francesconi and Lupi 2012).

Natural hosts. None, causes facultative myiasis in humans (Francesconi and Lupi 2012).

Route of infection. Oviposition on pre-existing wounds (Francesconi and Lupi 2012).

Site in human host. Skin, soft tissues, pre-existing wounds (Francesconi and Lupi 2012).

Genus *Protophormia* Townsend, 1908

Protophormia terraenovae Robineau-Desvoidy, 1830

Geographic distribution. Northern Hemisphere (Francesconi and Lupi 2012).

Natural hosts. None, causes facultative myiasis in humans (Francesconi and Lupi 2012).

Route of infection. Oviposition on pre-existing wounds (Francesconi and Lupi 2012).

Site in human host. Skin, soft tissues, pre-existing wounds (Francesconi and Lupi 2012).

***Sarcophaga ruficornis* (Fabricius, 1794)**

Geographic distribution. Worldwide (Suwannayod et al. 2013).

Natural hosts. None, causes facultative myiasis in humans (Suwannayod et al. 2013).

Route of infection. Oviposition on pre-existing wounds (Suwannayod et al. 2013).

Site in human host. Skin and soft tissues (Suwannayod et al. 2013).

***Sarcophaga septentrionalis* (Rohdendorf, 1937)**

Geographic distribution. East Asia (Chigusa et al. 2006).

Natural hosts. None; causes facultative myiasis in humans (Chigusa et al. 2006)

Route of infection. Oviposition on pre-existing wounds (Chigusa et al. 2006).

Site in human host. Oral cavity (Chigusa et al. 2006).

Genus *Wohlfahrtia* Brauer & von Bergenstamm, 1889***Wohlfahrtia magnifica* (Schiner, 1861)**

Geographic distribution. Mediterranean Europe, North Africa, Central and South-east Asia, Russia (Francesconi and Lupi 2012).

Natural hosts. Mammals and birds, including sheep, cattle, poultry, and humans (Francesconi and Lupi 2012).

Route of infection. Larviposition on wounds, tissues (Francesconi and Lupi 2012).

Site in human host. Skin, soft tissues, pre-existing wounds, oral cavity, ears (Francesconi and Lupi 2012).

Vectored pathogens. *Wohlfahrtiimonas chitiniclastica*, *Ignatzschineria indica* (Barker et al. 2014).

***Wohlfahrtia opaca* (Coquillett, 1897)**

Geographic distribution. Western North America (Francesconi and Lupi 2012).

Natural hosts. Wild and domestic canids and felids, rabbits, mustelids. Zoonotic in human as incidental hosts (Francesconi and Lupi 2012).

Route of infection. Larviposition on wounds, tissues (Francesconi and Lupi 2012).

Site in human host. Skin and soft tissues, pre-existing wounds (Francesconi and Lupi 2012).

***Wohlfahrtia vigil* (Walker, 1849)**

Geographic distribution. Eastern North America, southern Europe, Russia, Pakistan (Francesconi and Lupi 2012).

Natural hosts. Wild and domestic canids and felids, rabbits, mustelids. Zoonotic in human as incidental hosts (Francesconi and Lupi 2012).

Route of infection. Larviposition on wounds, tissues (Francesconi and Lupi 2012).

Site in human host. Skin and soft tissues, pre-existing wounds (Francesconi and Lupi 2012).

●●●●●●●●●● **Oestridae Leach, 1815**

Genus *Alouattamyia* Townsend, 1931

Alouattamyia baeri (Shannon & Green, 1926)

Geographic distribution. South America (Francesconi and Lupi 2012).

Natural hosts. Non-human primates. Zoonotic in human as incidental hosts (Francesconi and Lupi 2012).

Route of infection. Exposure to first-instar larvae in the environment (Francesconi and Lupi 2012).

Site in human host. Upper respiratory tract (Francesconi and Lupi 2012).

Genus *Dermatobia* Brauer, 1860

Dermatobia hominis (Linnaeus in Pallas, 1781)

Geographic distribution. Central and South America, Caribbean (Francesconi and Lupi 2012).

Natural hosts. Many mammals, including humans, and birds; vectored by mosquitoes and ticks (Francesconi and Lupi 2012).

Route of infection. Bite of a mosquito or tick, the phoretic host for *D. hominis* eggs (Francesconi and Lupi 2012).

Site in human host. Skin (Francesconi and Lupi 2012).

Genus *Cuterebra* Clark, 1815

Geographic distribution. North and Central America (Francesconi and Lupi 2012).

Natural hosts. Rodents and lagomorphs; cats and other mammals can serve as reservoir hosts. Zoonotic in human as incidental hosts (Francesconi and Lupi 2012).

Route of infection. Exposure to first-instar larvae in contaminated soil, litter, etc.

Site in human host. Skin (second-instar, third-instar larvae), viscera and lungs (third-instar larvae) (Cornet et al. 2003; Delshad et al. 2008; Francesconi and Lupi 2012; Hale et al. 2019).

Notes. Cases of human infestation with *Cuterebra* are usually not identified to the species level.

Genus *Gasterophilus* Leach, 1817

Geographic distribution. Worldwide (Zumpt 1965; Francesconi and Lupi 2012).

Natural hosts. Horses, zebras, elephants, rhinoceroses. Zoonotic in human as incidental hosts (Zumpt 1965; Francesconi and Lupi 2012).

Route of infection. Oviposition on skin and mucus membranes (Francesconi and Lupi 2012).

Site in human host. Skin, eye, oral cavity (Francesconi and Lupi 2012).

Notes. Cases of human infestation with *Gasterophilus* are usually not identified to the species level.

Genus *Hypoderma* Latreille, 1818***Hypoderma bovis* (Linnaeus, 1758)**

Geographic distribution. North America, Europe, Asia, Africa (Zumpt 1965; Francesconi and Lupi 2012).

Natural hosts. Wild and domestic ruminants. Zoonotic in human as incidental hosts (Zumpt 1965; Francesconi and Lupi 2012).

Route of infection. Oviposition on hair or skin, or possibly direct contact with first-instar larvae on infected animals (Francesconi and Lupi 2012).

Site in human host. Skin (Francesconi and Lupi 2012).

***Hypoderma lineatum* (Viller, 1789)**

Geographic distribution. North America, Europe, Asia, Africa (Zumpt 1965; Francesconi and Lupi 2012).

Natural hosts. Wild and domestic ruminants. Zoonotic in human as incidental hosts (Zumpt 1965; Francesconi and Lupi 2012).

Route of infection. Oviposition on hair or skin, or possibly direct contact with first-instar larvae on infected animals (Francesconi and Lupi 2012)

Site in human host. Skin (Francesconi and Lupi 2012).

***Hypoderma tarandi* (Linnaeus, 1758)**

Geographic distribution. Northern North America, Northern Eurasia (Zumpt 1965; Francesconi and Lupi 2012).

Natural hosts. Caribou. Zoonotic in human as incidental hosts (Zumpt 1965; Francesconi and Lupi 2012).

Route of infection. Oviposition on hair, skin, or mucus membranes, or possibly direct contact with first-instar larvae on infected animals (Francesconi and Lupi 2012).

Site in human host. Skin, eyes, oral cavity (Francesconi and Lupi 2012).

Genus *Oestrus* Linnaeus, 1758***Oestrus ovis* Linnaeus, 1758**

Geographic distribution. Worldwide (Francesconi and Lupi 2012).

Natural hosts. Sheep, goats, other ruminants. Zoonotic in human as incidental hosts (Francesconi and Lupi 2012).

Route of infection. Larviposition on eyes and mucus membranes (Francesconi and Lupi 2012).

Site in human host. Eye, throat (Francesconi and Lupi 2012).

●●●●● Nematoda Diesking, 1861

●●●●●● Dorylaimia Inglis, 1983

●●●●●●● Diocetophymatida Baylis & Daubney, 1926

●●●●●●●● Diocetophymidae Railliet, 1915

Genus *Diocetophyme* Collet-Meygret, 1802

Diocetophyma, misspelling

***Diocetophyme renale* (Goeze, 1782)**

Geographic distribution. Worldwide (Angelou et al. 2020)

Natural hosts. First intermediate hosts are annelids. Second intermediate and paratenic hosts are primarily freshwater fish and amphibians. Definitive hosts are carnivores, including mustelids and canids. Zoonotic in humans, harboring L3 larvae or adults (Angelou et al. 2020).

Route of infection. Ingestion of L3 larvae in undercooked paratenic hosts (Orihel and Ash 1995).

Site in human host. Subcutaneous (L3 larvae), kidneys (adults) (Orihel and Ash 1995).

Genus *Eustrongylides* Jägerskiöld, 1909

Geographic distribution. Worldwide (Anderson 2000).

Natural hosts. First intermediate hosts are aquatic annelids. The second intermediate hosts are fish; fish, amphibians, and reptiles can serve as paratenic hosts. Definitive hosts are birds. Zoonotic in humans as dead-end hosts harboring L3 or L4 larvae (Anderson 2000).

Route of Infection. Ingestion of L3 or precocious L4 larvae in infected intermediate or paratenic hosts (Anderson 2000).

Site in human host. Skin, Abdominal and peritoneal cavities, possible ectopic migration to other parts of the body (Orihel and Ash 1995; Eberhard and Ruiz-Tiben 2014).

Notes. There about 20 described species of *Eustrongylides*. Human cases with larval *Eustrongylides* have been reported from North America and East Africa and have not been characterized at the species level.

●●●●●●● **Trichinellida Hall, 1916**

Trichocephalida Spasski, 1954

●●●●●●●● **Anatrichosomatidae Yamaguti, 1961**

Genus *Anatrichosoma* Smith & Chitwood, 1954

Geographic distribution. Worldwide (Eberhard et al. 2010, 2014a).

Natural hosts. The life cycle is not completely understood, and it is not known if there are more than one host. Definitive hosts include non-human primates, tree shrews, rodents, marsupials. Zoonotic in humans as incidental hosts (Eberhard et al. 2010, 2014a).

Route of infection. Unknown, possible ingestion of embryonated eggs or unknown intermediate host (Eberhard et al. 2010, 2014a).

Site in human host. Skin and soft tissues, oral mucosa (Eberhard et al. 2010, 2014a).

Notes. Human cases of anatrichosomiasis have been diagnosed by histopathology. Because *Anatrichosoma* spp. have not been described morphologically by histopathology, it is not currently possible to characterize human isolates at the species level. Given the morphology and epidemiology, cases from the human oral cavity from Mexico and the United States are probably attributable to *A. buccalis* Pence & Little, 1972 (Eberhard et al. 2010, 2014a).

●●●●●●●● **Capillariidae Railliet, 1915**

Genus *Calodium* Dujardin, 1845

***Calodium hepaticum* (Bancroft, 1893)**

Geographic distribution. Worldwide (Li et al. 2010).

Natural hosts. Rodents. Zoonotic in humans as incidental hosts (Li et al. 2010).

Route of infection. Ingestion of embryonated eggs on soil-contaminated fomites (Li et al. 2010).

Site in human host. Liver (Li et al. 2010).

Notes. Eggs of *C. hepaticum* detected in human stool probably represent spurious passage and not true infection.

Genus *Eucoleus* Dujardin, 1845***Eucoleus aerophilus* (Creplin, 1839)**

Geographic distribution. Worldwide (Traversa et al. 2011).

Natural hosts. Paratenic hosts are earthworms. Definitive hosts are carnivores. Zoonotic in humans as incidental hosts (Traversa et al. 2011).

Route of infection. Ingestion of embryonated eggs or paratenic hosts (Traversa et al. 2011).

Site in human host. Lungs (Traversa et al. 2011).

Genus *Paracapillaria* Chitwood et al., 1968***Paracapillaria philippinensis* (Velasquez et al., 1968)**

Geographic distribution. Southeast Asia, Philippines, Egypt, South America (Cross 1992; Attia et al. 2012; Eiras et al. 2018).

Natural hosts. Intermediate hosts are freshwater fish. Definitive hosts are humans, fish-eating birds (Cross 1992).

Route of infection. Ingestion of L1 larvae in infected intermediate host (Cross 1992).

Site in human host. Small intestine (Cross 1992).

●●●●●●●● Trichinellidae Ward, 1907**Genus *Trichinella* Railliet, 1895*****Trichinella britovi* Pozio et al., 1992**

Trichinella strain T3

Geographic distribution. Europe, North America, Asia, West Africa (Gottstein et al. 2009).

Natural hosts. Mammals. Zoonotic in humans (Gottstein et al. 2009).

Route of infection. Ingestion of L1 larvae in infected hosts; common sources of human infection are wild and domestic pigs, horses, foxes, jackals (Gottstein et al. 2009).

Site in human host. Small intestine (adults), skeletal muscle (L1 larvae) (Gottstein et al. 2009).

***Trichinella murelli* Pozio & La Rosa, 2000**

Trichinella strain T5

Geographic distribution. North America (Gottstein et al. 2009; Hall et al. 2012).

Natural hosts. Mammals, primarily carnivores. Zoonotic in humans (Gottstein et al. 2009).

Route of infection. Ingestion of L1 larvae in infected hosts; common sources of human are bears and horses (Gottstein et al. 2009; Hall et al. 2012).

Site in human host. Small intestine (adults), skeletal muscle (L1 larvae) (Gottstein et al. 2009).

Trichinella nativa Britov & Boev, 1972

Trichinella strain T2

Geographic distribution. Circumboreal (Gottstein et al. 2009).

Natural hosts. Mammals, primarily carnivores. Zoonotic in humans (Gottstein et al. 2009).

Route of infection. Ingestion of L1 larvae in infected hosts; common sources of human infection are bears and walrus (Gottstein et al. 2009; Springer et al. 2017).

Site in human host. Small intestine (adults), skeletal muscle (L1 larvae) (Gottstein et al. 2009).

Trichinella nelsoni Britov & Boev, 1972

Trichinella strain T7

Geographic distribution. Europe, North America, Asia, West Africa (Gottstein et al. 2009).

Natural hosts. Mammals. Zoonotic in humans (Gottstein et al. 2009).

Route of infection. Ingestion of L1 larvae in infected hosts; common sources of human infection are wild and domestic pigs, horses, foxes, jackals (Gottstein et al. 2009).

Site in human host. Small intestine (adults), skeletal muscle (L1 larvae) (Gottstein et al. 2009).

Trichinella papuae Pozio et al., 1999

Geographic distribution. Papua New Guinea, Thailand (Gottstein et al. 2009).

Natural hosts. Mammals (primarily pigs), saltwater crocodiles. Zoonotic in humans (Gottstein et al. 2009).

Route of infection. Ingestion of L1 larvae in infected hosts; common sources of human infection are wild and domestic pigs (Gottstein et al. 2009).

Site in human host. Small intestine (adults), skeletal muscle (L1 larvae) (Gottstein et al. 2009).

Trichinella pseudospiralis* Garkavi, 1972Trichinella* strain T4**Geographic distribution.** Worldwide (Gottstein et al. 2009).**Natural hosts.** Mammals and birds. Zoonotic in humans (Gottstein et al. 2009).**Route of infection.** Ingestion of L1 larvae in infected hosts; common sources of human infection are wild and domestic pigs (Gottstein et al. 2009).**Site in human host.** Small intestine (adults), skeletal muscle (L1 larvae) (Gottstein et al. 2009).***Trichinella spiralis* (Owen, 1835)***Trichinella* strain T1**Geographic distribution.** Worldwide (Gottstein et al. 2009).**Natural hosts.** Mammals. Zoonotic in humans (Gottstein et al. 2009).**Route of infection.** Ingestion of L1 larvae in infected hosts; common sources of human infection are wild and domestic pigs (Gottstein et al. 2009).**Site in human host.** Small intestine (adults), skeletal muscle (L1 larvae) (Gottstein et al. 2009).***Trichinella* Strain T6****Geographic distribution.** Northern and montane North America (Gottstein et al. 2009).**Natural hosts.** Mammals, primarily carnivores. Zoonotic in humans (Gottstein et al. 2009).**Route of infection.** Ingestion of L1 larvae in infected hosts; common sources of human infection are bears, wild felids (Gottstein et al. 2009).**Site in human host.** Small intestine (adults), skeletal muscle (L1 larvae) (Gottstein et al. 2009).●●●●●● **Muspiceida** Bain & Chabaud, 1959●●●●●● **Robertdollfusidae** Chabaud & Campana, 1950**Genus *Haycocknema* Spratt et al., 1999*****Haycocknema perplexum* Spratt et al., 1999****Geographic distribution.** Australia (McKelvie et al. 2013; Vos et al. 2016).**Natural hosts.** Unknown, presumed zoonotic in humans (McKelvie et al. 2013; Vos et al. 2016).

Route of infection. Unknown, presumed ingestion of larvae in natural hosts (McKelvie et al. 2013; Vos et al. 2016).

Site in human host. Skeletal muscle (McKelvie et al. 2013; Vos et al. 2016).

●●●●●●●● **Trichuridae Ransom, 1911**

Genus *Trichuris* Roederer, 1761

***Trichuris trichiura* (Linnaeus, 1771)**

Geographic distribution. Worldwide (Betson et al. 2015).

Natural hosts. Humans (Betson et al. 2015).

Route of infection. Ingestion of embryonated eggs in fecally contaminated food, water, fomites (Betson et al. 2015).

Site in human host. Large intestine (Betson et al. 2015).

Notes. The zoonotic potential of *T. trichiura* is not well understood and is unclear whether there are multiple species infecting humans or whether other animals can harbor *T. trichiura* and serve as reservoir hosts for human infection (Betson et al. 2015).

***Trichuris vulpis* Froelich, 1789**

Geographic distribution. Worldwide (Dunn et al. 2002).

Natural hosts. Wild and domestic canids. Zoonotic in humans (Dunn et al. 2002).

Route of infection. Ingestion of embryonated eggs (Dunn et al. 2002).

Site in human host. Large intestine (Dunn et al. 2002).

Notes. Fully-embryonated eggs of *T. vulpis* in human stool specimens may also represent spurious passage and not true infection.

●●●●●● **Chromadoria Inglis, 1983**

●●●●●●● **Spururina Railliet & Henry, 1915**

●●●●●●●● **Ascaridida DeLay & Blaxter, 2002**

●●●●●●●●● **Anisakidae Skrjabin & Karokhin, 1945**

Genus *Anisakis* Dujardin, 1845

Geographic distribution. Worldwide; human infection more common in coastal areas (Audicana and Kennedy 2008; Mathison and da Silva 2018).

Natural hosts. First intermediate hosts are marine microcrustaceans. Paratenic hosts are marine fish and squid. Definitive hosts are fish-eating mammals, primarily cetaceans and pinnipeds. Zoonotic in humans as dead-end hosts harboring L3 larvae (Audicana and Kennedy 2008; Mathison and da Silva 2018).

Route of infection. Ingestion of L3 larvae in infected paratenic hosts (Audicana and Kennedy 2008; Mathison and da Silva 2018).

Site in human host. Stomach, esophagus, oral cavity, large intestine; ectopic colonization of peritoneal cavity, mesenteries, omentum, mesocolic lymph nodes, spleen, pleural cavity, and parametrium (Audicana and Kennedy 2008; Mathison and da Silva 2018).

Notes. While most human cases have been attributed to members of *A. simplex* sensu lato (including *A. simplex* (Rudolphi, 1809), *A. pegreffii* (Campana-Rouget & Biocca, 1955), and *A. berlandi* Mattiucci et al. 2014) or *A. physeteris* (Baylis, 1923), reliable identification human isolates at the species level is difficult and seldom performed due to a lack of species-level morphologic features of the L3 larvae (the stage found in human host) and the lack of a reliable sequencing library for molecular analysis (Umehara et al. 2007).

Genus *Contracaecum* Railliet & Henry, 1912

Geographic distribution. Worldwide; human infection more common on coastal areas (Mathison and da Silva 2018).

Natural hosts. First intermediate hosts are marine microcrustaceans. Paratenic hosts are marine fish and squid. Definitive hosts are fish-eating mammals, primarily cetaceans and pinnipeds, and birds. Zoonotic in humans as dead-end hosts harboring L3 larvae (Mathison and da Silva 2018).

Route of infection. Ingestion of L3 larvae in infected paratenic hosts (Mathison and da Silva 2018).

Site in human host. Stomach, esophagus, oral cavity, large intestine; ectopic colonization of various organs possible (Mathison and da Silva 2018).

Notes. It is not currently possible to reliably identify human isolates of *Contracaecum* at the species level due to a lack of species-level morphologic features of the L3 larvae (the stage found in human host) and the lack of a reliable sequencing library for molecular analysis.

Genus *Pseudoterranova* Mozgovoi, 1951

Phocanema Myers, 1959

Geographic distribution. Worldwide; human infection more common on coastal areas (Mathison and da Silva 2018).

Natural hosts. First intermediate hosts are marine microcrustaceans. Paratenic hosts are marine fish and squid. Definitive hosts are fish-eating mammals, primarily cetaceans and pinnipeds. Zoonotic in humans as dead-end hosts harboring L3 larvae (Mathison and da Silva 2018).

Route of infection. Ingestion of L3 larvae in infected paratenic hosts (Mathison and da Silva 2018).

Site in human host. Stomach, esophagus, oral cavity, large intestine; ectopic colonization of peritoneal cavity, mesenteries, omentum, mesocolic lymph nodes, spleen, pleural cavity, and parametrium (Mathison and da Silva 2018).

Notes. While many human cases have been attributed to the *P. decepiens* (Krabbe, 1878) complex, reliable identification human isolates at the species level is difficult and seldom performed due to a lack of species-level morphologic features of the L3 larvae (the stage found in human host) and the lack of a reliable sequencing library for molecular analysis.

●●●●●●●● Ascarididae Baird, 1853

Genus *Ascaris* Linnaeus, 1758

Ascaris lumbricoides Linnaeus, 1758

Ascaris suum (Goeze, 1782)

Geographic distribution. Worldwide (Betson et al. 2014).

Natural hosts. Pigs. Zoonotic in humans (Betson et al. 2014).

Route of infection. Ingestion of embryonated eggs in fecally contaminated food, water, fomites (Betson et al. 2014).

Site in human host. Lungs (L3 larvae); Small intestine (adults), with ectopic colonization of liver, spleen, gall bladder (Betson et al. 2014).

Notes. The relationship between *A. lumbricoides* and *A. suum* is unresolved. While there are genetic differences among *Ascaris* isolates throughout the world, it is unclear whether such differences represent more than one species with a common ancestor, multiple colonization events between pigs and humans in different geographical areas, or there is one species with geographic variations (Betson et al. 2014). We are maintaining a conservative approach in keeping *A. suum* in synonymy with *A. lumbricoides*, based on ecological, biological, morphological, and molecular data (Leles et al. 2012; Liu et al. 2012).

Genus *Baylisascaris* Sprent, 1968

Baylisascaris procyonis (Stefanski & Zarnowski, 1951)

Geographic distribution. North and Central America, introduced to Europe and Asia (Gavin et al. 2005).

Natural hosts. Raccoons and other procyonids; dogs can serve as reservoir hosts. Zoonotic in humans as dead-end hosts harboring L3 larvae (Gavin et al. 2005).

Route of infection. Ingestion of embryonated eggs in food, water, fomites contaminated with raccoon feces; possibly also ingestion of paratenic hosts (Gavin et al. 2005).

Site in human host. Brain, eye, liver, lungs (Gavin et al. 2005).

Genus *Lagochilascaris* Leiper, 1909***Lagochilascaris minor* Leiper, 1909**

Geographic distribution. Central and South America (Campos et al. 2017).

Natural hosts. Unknown. It has been prosed rodents are intermediate hosts and carnivores are definitive hosts, with humans serving as incidental definitive hosts (Campos et al. 2017).

Route of infection. Unknown, presumed to be ingestion of larvae in infected intermediate or paratenic hosts, possibly ingestion of embryonated eggs (Campos et al. 2017).

Site in human host. Disseminated infection; common sites of infection are subcutaneous, ear, tonsils, nasal sinuses, and mastoid (Campos et al. 2017).

Genus *Toxocara* Stiles, 1905***Toxocara canis* (Werner, 1792)**

Geographic distribution. Worldwide (Despommier 2003).

Natural hosts. Wild and domestic canids. Zoonotic in humans as dead-end hosts harboring L3 larvae (Despommier 2003).

Route of infection. Ingestion of embryonated eggs in food, water, fomites contaminated with canid feces (Despommier 2003).

Site in human host. Disseminated infection; common sites of infection are liver, eye, CNS, and lungs (Despommier 2003).

Toxocara cati* Schrank, 1788**Toxocara mystax* (Zeder, 1800)**

Geographic distribution. Worldwide (Despommier 2003).

Natural hosts. Wild and domestic felids. Zoonotic in humans as dead-end hosts harboring L3 larvae (Despommier 2003).

Route of infection. Ingestion of embryonated eggs in food, water, fomites contaminated with felid feces (Despommier 2003).

Site in human host. Disseminated infection; common sites of infection are liver, eye, CNS, and lungs (Despommier 2003).

●●●●●●●● **Oxyurida Weinland, 1858**

●●●●●●●● **Oxyuridae Cobbold, 1864**

Genus *Enterobius* Baird, 1853

***Enterobius vermicularis* (Linnaeus, 1758)**

Enterobius gregorii Hugot, 1983

Geographic distribution. Worldwide (Wendt et al. 2019).

Natural hosts. Humans (Wendt et al. 2019).

Route of infection. Ingestion of embryonated eggs on contaminated fomites (Wendt et al. 2019).

Site in human host. Cecum, large intestine, appendix; ectopic colonization of female urogenital tract (Wendt et al. 2019).

Genus *Syphacia* Seurat, 1916

***Syphacia oblevata* (Rudolphi, 1802)**

Geographic distribution. Worldwide (Riley 1919).

Natural hosts. Rodents (Riley 1919).

Route of infection. Presumed ingestion of embryonated eggs on contaminated fomites (Riley 1919).

Site in human host. Large intestine (Riley 1919).

●●●●●●●● **Spirudida De Ley & Blaxter, 2002**

●●●●●●●● **Filarioidea Chabaoud & Anderson, 1959**

●●●●●●●● **Onchocercidae Chabaoud & Anderson, 1959**

Genus *Acanthocheilonema* Cobbold, 1870

***Acanthocheilonema reconditum* (Grassi, 1889)**

Geographic distribution. Worldwide (Otranto and Eberhard 2011).

Natural hosts. Vector intermediate hosts are fleas and lice, primarily the cat flea (*Ctenocephalides felis*). Definitive hosts are canids. Zoonotic in humans (Otranto and Eberhard 2011).

Route of infection. Introduction of L3 larvae via the bite of an infected flea (Otranto and Eberhard 2011).

Site in human host. Eyes (Huynh et al. 2001; Otranto and Eberhard 2011).

Notes. Several cases of *Acanthocheilonema*-like nematodes recovered from human eyes have been reported, though not identified further (Otranto and Eberhard 2011).

Genus *Breinlia* Yorke & Maplestone, 1926***Breinlia annulipapillata* (Johnston & Mawson, 1938)**

Geographic distribution. Australia (Koehler et al. 2021).

Natural hosts. Vector intermediate hosts are mosquitoes, although the full range of competent genera is not known. Definitive hosts are kangaroos and wallabies. Zoonotic in humans (Koehler et al. 2021).

Route of infection. Introduction of L3 larvae via the bite of an infected mosquito (Koehler et al. 2021).

Site in human host. Eye (Koehler et al. 2021).

Genus *Brugia* Buckley, 1960***Brugia malayi* (Brug, 1927)**

Geographic distribution. Southeast Asia, including Philippines, Malaysia, Indonesia, South Korea, Vietnam, India (Mak 1987; Dietrich et al. 2019).

Natural hosts. Vector intermediate hosts are mosquitoes, primarily in the genera *Aedes* and *Mansonia*. Definitive hosts are humans (Mak 1987; Dietrich et al. 2019).

Route of infection. Introduction of L3 larvae via the bite of an infected mosquito (Mak 1987; Dietrich et al. 2019).

Site in human host. Lymphatic system (Mak 1987; Dietrich et al. 2019).

***Brugia timori* Partono et al., 1977**

Geographic distribution. Lesser Sunda Archipelago, including the islands of Timor, Sumba, Lembata, Pentar, Alor (Fischer et al. 2004).

Natural hosts. Vector intermediate hosts are mosquitoes in the genus *Anopheles*. Definitive hosts are humans (Fischer et al. 2004; Dietrich et al. 2019).

Route of infection. Introduction of L3 larvae via the bite of an infected mosquito (Fischer et al. 2004; Dietrich et al. 2019).

Site in human host. Lymphatic system (Fischer et al. 2004; Dietrich et al. 2019).

Unassigned *Brugia* species:**American *Brugia* spp.**

Geographic distribution. North, Central, and South America (Orihel and Eberhard 1998).

Natural hosts. Unknown. Vector intermediate hosts are presumed to be mosquitoes. Possible definitive hosts include raccoons, rabbits, and felids. Presumed zoonotic in humans (Orihel and Eberhard 1998).

Route of infection. Introduction of L3 larvae via the bite of an infected mosquito (Orihel and Eberhard 1998).

Site in human host. Lymphatic system (Orihel and Eberhard 1998).

Notes. Several cases of zoonotic *Brugia* spp. have reported in the Americas. The precise identification of the causal agents is not known due to limited information on the morphologic features observed on histopath. Human infection in North America has been tentatively attributed to *B. beaveri* Ash & Little, 1964 or *B. lepori* Eberhard, 1984 (Orihel and Ash 1995; Orihel and Eberhard 1998).

Genus *Dirofilaria* Railliet & Henry, 1911

Subgenus *Dirofilaria* Railliet & Henry, 1911

***Dirofilaria (D.) immitis* (Leidy, 1856)**

Filaria magalhaesi Blanchard, 1896

Dirofilaria nasuae Mazza, 1926

Dirofilaria pongoi Vogel & Vogelsang, 1930

Dirofilaria louisianensis Faust et al., 1941

Geographic distribution. Worldwide (Orihel and Eberhard 1998).

Natural hosts. Vector intermediate hosts are mosquitoes. Definitive hosts are primarily wild and domestic canids. Zoonotic in humans (Orihel and Eberhard 1998).

Route of infection. Introduction of L3 larvae via the bite of an infected mosquito (Orihel and Eberhard 1998).

Site in human host. Heart, pulmonary vessels (Orihel and Eberhard 1998).

***Dirofilaria (D.) spectans* Freitas & Lint, 1949**

Geographic distribution. South America (Orihel and Eberhard 1998).

Natural hosts. Vector intermediate hosts are presumed to be mosquitoes. Definitive hosts is the giant river otter (*Pteronura brasiliensis*). Zoonotic in humans (Orihel and Eberhard 1998).

Route of infection. Introduction of L3 larvae via the bite of an infected mosquito (Orihel and Eberhard 1998).

Site in human host. Digital artery (Orihel and Eberhard 1998).

Notes. Few features can separate *D. spectans* from *D. immitis*, thus the validity of the species and its involvement in human infections remain in question.

Subgenus *Nochtiella* Faust, 1937

***Dirofilaria (N.) repens* Railliet & Henry, 1911**

Filaria confunctivae Addario, 1885, in part

Geographic distribution. Europe, Africa, Asia (Orihel and Eberhard 1998).

Natural hosts. Vector intermediate hosts are mosquitoes. Definitive hosts are carnivores, primarily wild and domestic canids, also felids. Zoonotic in humans (Orihel and Eberhard 1998).

Route of infection. Introduction of L3 larvae via the bite of an infected mosquito (Orihel and Eberhard 1998).

Site in human host. Skin; ectopic migration to the eye (Orihel and Eberhard 1998).

Dirofilaria (N.) striata (Molin, 1858)

Geographic distribution. North, Central, and South America (Orihel and Eberhard 1998).

Natural hosts. Vector intermediate hosts are mosquitoes. Definitive hosts are wild felids. Zoonotic in humans (Orihel and Eberhard 1998).

Route of infection. Introduction of L3 larvae via the bite of an infected mosquito (Orihel and Eberhard 1998).

Site in human host. Eye (Orihel and Eberhard 1998).

Dirofilaria (N.) subdermata (Monnig, 1924)

Geographic distribution. North America (Orihel and Eberhard 1998).

Natural hosts. Vector intermediate hosts are mosquitoes. Definitive hosts are porcupines. Zoonotic in humans (Orihel and Eberhard 1998).

Route of infection. Introduction of L3 larvae via the bite of an infected mosquito (Orihel and Eberhard 1998).

Site in human host. Skin (Orihel and Eberhard 1998).

Notes. Features available in transverse histologic sections are unable to reliably distinguish *D. subdermata* from *D. ursi*; therefore, specimens recovered from human cases are generally reported as “*D. ursi*-like”.

Dirofilaria (N.) tenuis Chandler, 1942

Filaria confunctivae Addario, 1885, in part

Geographic distribution. North America (Orihel and Eberhard 1998).

Natural hosts. Vector intermediate hosts are mosquitoes. Definitive hosts are primarily raccoons. Zoonotic in humans (Orihel and Eberhard 1998).

Route of infection. Introduction of L3 larvae via the bite of an infected mosquito (Orihel and Eberhard 1998).

Site in human host. Skin; ectopic migration to the eyes (Orihel and Eberhard 1998).

Dirofilaria (N.) ursi Yamaguti, 1941

Geographic distribution. North America, Russia, Japan (Orihel and Eberhard 1998).

Natural hosts. Vector intermediate hosts are black flies in the genus *Simulium*. Definitive hosts are bears. Zoonotic in humans (Orihel and Eberhard 1998).

Route of infection. Introduction of L3 larvae via the bite of an infected black fly (Orihel and Eberhard 1998).

Site in human host. Skin (Orihel and Eberhard 1998).

Notes. Features available in transverse histologic sections are unable to reliably distinguish *D. subdermata* from *D. ursi*; therefore, specimens recovered from human cases are generally reported as “*D. ursi*-like”.

Dirofilaria, incertae sedis:

***Dirofilaria* sp. Genotype Hong Kong**

Dirofilaria hongkongensis To et al., 2012

Geographic distribution. Southeast Asia (Dantas-Torres and Otranto 2020; To et al. 2012).

Natural hosts. Vector intermediate hosts are unknown. Definitive hosts are canids. Zoonotic in humans (To et al. 2012; Dantas-Torres and Otranto 2020).

Route of infection. Presumably via the introduction of L3 larvae via the bite of an infected vector.

Site in human host. Skin (To et al. 2012; Dantas-Torres and Otranto 2020).

Notes. *Dirofilaria honkongensis* was described from dogs and humans in Hong Kong based on molecular characterization (To et al. 2012). The species was not described under the criteria of the ICZN and no type specimen was designated. To complicate matters, they described the species as ‘*Candidatus* *Dirofilaria honkongensis*’. The term *candidatus* is not used in zoological nomenclature and is usually only used for bacteria and other organisms that cannot be cultured. The species should be considered *nomen nudum* (Mathison and Pritt 2019; Dantas-Torres and Otranto 2020).

Genus *Dunnifilaria* Mullin & Balasingam, 1973

Geographic distribution. Mexico, Southeast Asia (Lam et al. 2010).

Natural hosts. Vector intermediate hosts are unknown. Definitive hosts are rodents. Zoonotic in humans (Lam et al. 2010).

Route of Infection. Presumed introduction of L3 larvae via the bite of an infected vector.

Site in human host. Eye (Lam et al. 2010).

Notes. The single human case was reported from the eye of a patient in Malaysia (Lam et al. 2010). The worm was not identified at the species level but was reported as being most consistent with *D. ramachandrani* Mullin & Balasingam, 1973, a parasite of the long-tailed giant rat in Malaysia.

Genus *Loa* Stiles, 1905***Loa loa* (Cobbold, 1864)**

Geographic distribution. West-central Africa (Whittaker et al. 2018).

Natural hosts. Vector intermediate hosts are deer flies in the genus *Chrysops*. Definitive hosts are humans (Whittaker et al. 2018).

Route of infection. Introduction of L3 larvae via the bite of an infected deer fly (Whittaker et al. 2018).

Site in human host. Skin; ectopic migration to the eyes (Whittaker et al. 2018).

Notes. Cases of *L. loa* in non-travelers outside of endemic areas should be considered as possible misidentifications. There are many case reports of *L. loa* from India, many of which appear to represent misidentifications of *Dirofilaria* (prob. *D. repens*).

Genus *Mansonella* Faust, 1929***Mansonella ozzardi* Manson, 1897**

Geographic distribution. Central and South America, Caribbean (Lima et al. 2016).

Natural hosts. Vector intermediate hosts are biting midges (genus *Culicoides*) and black flies (genus *Simulium*). Definitive hosts are humans (Lima et al. 2016).

Route of infection. Introduction of L3 larvae via the bite of an infected arthropod vector (Lima et al. 2016).

Site in human host. Skin (Lima et al. 2016).

Mansonella perstans* (Manson, 1891)**Dipetalonema berghei* Chardome & Peel, 1951**

Geographic distribution. Africa, South America, Caribbean (Mediannikov and Ranque 2018).

Natural hosts. Vector intermediate hosts are biting midges in the genus *Culicoides*. Definitive hosts are humans (Mediannikov and Ranque 2018).

Route of infection. Introduction of L3 larvae via the bite of an infected *Culicoides* biting midge (Mediannikov and Ranque 2018).

Site in human host. Peritoneal cavity, pleural cavity, pericardium, mesenteries (Mediannikov and Ranque 2018).

***Mansonella streptocerca* (MacFie & Corson, 1922)**

Geographic distribution. Sub-Saharan Africa (Mediannikov and Ranque 2018).

Natural hosts. Vector intermediate hosts are biting midges in the genus *Culicoides*. Definitive hosts are humans, non-human primates (Mediannikov and Ranque 2018).

Route of infection. Introduction of L3 larvae via the bite of an infected *Culicoides* biting midge (Mediannikov and Ranque 2018).

Site in human host. Skin (Mediannikov and Ranque 2018).

Genus *Meningonema* Orihel & Esslinger, 1973

Meningonema peruzzii Orihel & Esslinger, 1973

Geographic distribution. Africa (Orihel and Eberhard 1998).

Natural hosts. Vector intermediate hosts are unknown. Definitive hosts are monkeys. Zoonotic in humans (Orihel and Eberhard 1998).

Route of infection. Introduction of L3 larvae via the bite of an infected arthropod vector (Orihel and Eberhard 1998).

Site in human host. CNS (Orihel and Eberhard 1998).

Genus *Molinema* Freitas & Lint, 1939

Geographic distribution. North, Central, and South America (Orihel and Eberhard 1998).

Natural hosts. Vector intermediate hosts are mosquitoes. Definitive hosts are rodents. Zoonotic in humans (Orihel and Eberhard 1998).

Route of infection. Introduction of L3 larvae via the bite of an infected mosquito (Orihel and Eberhard 1998).

Site in human host. Eyes (Orihel and Eberhard 1998).

Notes. Human cases of *Molinema* are rare. Cases from the northwestern United States are believed to be attributed to either *M. arbuta* (Highby, 1943), a parasite of porcupines, or *M. sprengi* (Anderson, 1953), a parasite of beaver (Orihel and Eberhard 1998).

Genus *Onchocerca* Diesing, 1841

Onchocerca cervicalis Railliet & Henry, 1910

Geographic distribution. Worldwide (Cambra-Pellejà et al. 2020; Papini et al. 2020).

Natural hosts. Vector intermediate hosts are biting midges in the genus *Culicoides*. Definitive hosts are horses and other equids. Zoonotic in humans (Cambra-Pellejà et al. 2020; Papini et al. 2020).

Route of infection. Introduction of L3 larvae via the bite of an infected black fly (Orihel and Eberhard 1998).

Site in human host. Skin, eyes (Burr et al. 1998; Lai et al. 2014).

***Onchocerca dewittei japonica* Uni et al., 2001**

Geographic distribution. Japan (Uni et al. 2015, 2017).

Natural hosts. Vector intermediate hosts are black flies in the genus *Simulium*. Definitive hosts are wild pigs. Zoonotic in humans (Uni et al. 2015, 2017).

Route of infection. Introduction of L3 larvae via the bite of an infected black fly (Uni et al. 2015, 2017).

Site in human host. Skin (Uni et al. 2015, 2017).

***Onchocerca gutturosa* (Neumann, 1919)**

Geographic distribution. Africa, Europe, Asia, Australia (Cambra-Pellejà et al. 2020).

Natural hosts. Vector intermediate hosts are biting midges in the genus *Culicoides*. Definitive hosts are cattle. Zoonotic in humans (Cambra-Pellejà et al. 2020).

Route of infection. Introduction of L3 larvae via the bite of an infected black fly (Cambra-Pellejà et al. 2020).

Site in human host. Skin (Lai et al. 2014).

***Onchocerca jakutensis* (Gubanov, 1964)**

Geographic distribution. Europe, Asia (Cambra-Pellejà et al. 2020).

Natural hosts. Vector intermediate hosts are black flies in the genus *Simulium*. Definitive hosts are deer. Zoonotic in humans (Cambra-Pellejà et al. 2020).

Route of infection. Introduction of L3 larvae via the bite of an infected black fly (Cambra-Pellejà et al. 2020).

Site in human host. Skin, eyes (Koehsler et al. 2007).

***Onchocerca lupi* Rodonaja, 1967**

Geographic distribution. Europe, Asia, North America (Cantey et al. 2016).

Natural hosts. Putative vector intermediate hosts are black flies in the genus *Simulium*. Definitive hosts are wild and domestic canids and felids. Zoonotic in humans (Cantey et al. 2016).

Route of infection. Introduction of L3 larvae via the bite of an infected black fly (Cantey et al. 2016).

Site in human host. Skin, skeletal muscle, cervical nodules (Cantey et al. 2016).

***Onchocerca volvulus* (Leuckart, 1893)**

Geographic distribution. Sub-Saharan Africa, Yemen, Central and South America (Udall 2007).

Natural hosts. Vector intermediate hosts are black flies in the genus *Simulium*. Definitive hosts are humans (Udall 2007).

Route of infection. Introduction of L3 larvae via the bite of an infected black fly (Udall 2007).

Site in human host. Skin, subcutaneous nodules (Udall 2007).

Genus *Pelecitus* Railliet & Henry, 1910

Geographic distribution. Worldwide (Bain et al. 2011).

Natural hosts. Vector intermediate hosts are various avian live, mosquitoes, and tabanid flies. Definitive hosts are birds, lagomorphs, and marsupials. Zoonotic in humans (Bain et al. 2011).

Route of infection. Introduction of L3 larva via the bite of an infected mosquito (Bain et al. 2011).

Site in human host. Eyes (Bain et al. 2011).

Notes. The single case of human infection with *Pelecitus* was from Brazil and was not identified to the species level (Bain et al. 2011).

Genus *Wuchereria* Da Silva Araujo, 1877

Wuchereria bancrofti (Cobbold, 1877)

Wuchereria pacifica Manson-Bahr, 1941

Geographic distribution. Circumtropical, including South America, Caribbean, Africa, Asia, and Pacific Islands (Mak 1987; Dietrich et al. 2019).

Natural hosts. Vector intermediate hosts are mosquitoes, including the genera *Aedes*, *Anopheles*, and *Culex*. Definitive hosts are humans (Mak 1987; Dietrich et al. 2019).

Route of infection. Introduction of L3 larvae via the bite of an infected mosquito (Mak 1987; Dietrich et al. 2019).

Site in human host. Lymphatic system (Mak 1987; Dietrich et al. 2019).

●●●●●●●●●● Setariidae Skrjabin & Shikhobalova, 1945

Genus *Setaria* Viborg, 1795

Setaria equina (Abildgaard, 1789)

Geographic distribution. Worldwide (Nabie et al. 2017).

Natural hosts. Vector intermediate hosts are believed to be mosquitoes in the genera *Aedes* or *Culex*. Definitive hosts are horses and other equids. Zoonotic in humans (Nabie et al. 2017).

Route of infection. Introduction of L3 larvae via the bite of an infected mosquito (Nabie et al. 2017).

Site in human host. Eyes (Nabie et al. 2017).

***Setaria labiatopapillosa* (Alessandrini, 1838)**

Geographic distribution. Worldwide (Panaitescu et al. 1999).

Natural hosts. Vector intermediate hosts are believed to be mosquitoes in the genus *Aedes*. Definitive hosts are bovids. Zoonotic in humans (Panaitescu et al. 1999).

Route of infection. Introduction of L3 larvae via the bite of an infected mosquito (Panaitescu et al. 1999).

Site in human host. Eyes (Panaitescu et al. 1999).

Filaroidea, incertae sedis:

“*Microfilaria*” *bolivarensis* (Godoy et al., 1980)

Geographic distribution. Venezuela (Orihel and Eberhard 1998).

Natural hosts. Unknown; presumed to be zoonotic in humans (Orihel and Eberhard 1998).

Route of infection. Presumed introduction of L3 larvae via the bite of an infected arthropod vector.

Site in human host. Unknown, only described from microfilariae in blood (Orihel and Eberhard 1998).

“*Microfilaria*” *rodhaini* (Peel & Chardome, 1946)

Geographic distribution. Gabon (Orihel and Eberhard 1998).

Natural hosts. Vector intermediate hosts unknown. Definitive hosts presumed to be bonobos and chimpanzees. Zoonotic in humans (Orihel and Eberhard 1998).

Route of infection. Presumed introduction of L3 larvae via the bite of an infected arthropod vector

Site in human host. Unknown, only described from microfilariae in blood (Orihel and Eberhard 1998).

“*Microfilaria*” *semiclarum* (Fain, 1974)

Geographic distribution. Democratic Republic of Congo (Orihel and Eberhard 1998).

Natural hosts. Unknown; presumed to be zoonotic in humans (Orihel and Eberhard 1998).

Route of infection. Presumed introduction of L3 larvae via the bite of an infected arthropod vector.

Site in human host. Unknown, only described from microfilariae in blood (Orihel and Eberhard 1998).

●●●●●●●●●● **Spiruroidea Oerley, 1885**

●●●●●●●●●● **Gongylonematidae Sobolev, 1949**

Genus *Gongylonema* Molin, 1857

***Gongylonema pulchrum* Molin, 1857**

Geographic distribution. Worldwide (Libertin et al. 2017).

Natural hosts. Intermediate hosts are insects, primarily coprophagous beetles and cockroaches. Definitive hosts are mammals, including ruminants, carnivores, insectivores, lagomorphs. Zoonotic in humans (Libertin et al. 2017).

Route of infection. Ingestion of L3 larvae in infected insects (Libertin et al. 2017).

Site in human host. Oral mucosa (Libertin et al. 2017).

●●●●●●●●●● **Gnathostomatidae Railliet, 1895**

Genus *Gnathostoma* Owen, 1837

***Gnathostoma binucleatum* (Almeyda-Artigas, 1991)**

Geographic distribution. Central and South America (Cornaglia et al. 2016).

Natural hosts. First intermediate hosts are freshwater copepods. Second intermediate hosts are freshwater fish, amphibians, birds, rodents, and reptiles can serve as paratenic hosts. Definitive hosts are wild and domestic felids and canids. Zoonotic in humans as dead-end hosts harboring L3 larvae (Cornaglia et al. 2016).

Route of infection. Ingestion of L3 larvae in infected intermediate or paratenic hosts (Cornaglia et al. 2016).

Site in human host. Skin, heart, intestinal tract, ears, eyes, CNS (Cornaglia et al. 2016).

***Gnathostoma doloresi* (Tubangui, 1925)**

Geographic distribution. Southeast Asia, Japan (Herman and Chiodini 2009).

Natural hosts. First intermediate hosts are freshwater copepods. Second intermediate hosts are freshwater fish. Amphibians, birds, rodents, and reptiles can serve as paratenic hosts. Definitive hosts are wild and domestic pigs. Zoonotic in humans as dead-end hosts harboring L3 larvae (Herman and Chiodini 2009).

Route of infection. Ingestion of L3 larvae in infected intermediate or paratenic hosts (Herman and Chiodini 2009).

Site in human host. Skin, heart, intestinal tract, ears, eyes, CNS (Herman and Chiodini 2009).

***Gnathostoma hispidum* (Fedtschenko, 1872)**

Geographic distribution. Southeast Asia, India, Australia (Herman and Chiodini 2009).

Natural hosts. First intermediate hosts are freshwater copepods. Second intermediate hosts are freshwater fish. Amphibians, birds, rodents, and reptiles can serve as paratenic hosts. Definitive hosts are wild and domestic pigs. Zoonotic in humans as dead-end hosts harboring L3 larvae (Herman and Chiodini 2009).

Route of infection. Ingestion of L3 larvae in infected intermediate or paratenic hosts (Herman and Chiodini 2009).

Site in human host. Skin, heart, intestinal tract, ears, eyes, CNS (Herman and Chiodini 2009).

***Gnathostoma malaysiae* (Miyazaki & Dunn, 1965)**

Geographic distribution. Southeast Asia (Nomura et al. 2000).

Natural hosts. First intermediate hosts are freshwater copepods. Second intermediate hosts are freshwater fish. Amphibians, birds, rodents, and reptiles can serve as paratenic hosts. Definitive hosts are rodents, including rats. Zoonotic in humans as dead-end hosts harboring L3 larvae (Nomura et al. 2000).

Route of infection. Ingestion of L3 larvae in infected intermediate or paratenic hosts (Nomura et al. 2000).

Site in human host. Skin (Nomura et al. 2000).

Notes. The single case report of *G. malaysiae* in humans was reported from two Japanese fisherman who described eating raw freshwater shrimp in Myanmar. The species-level identification was considered presumptive (Nomura et al. 2000).

***Gnathostoma nipponicum* Yamaguti, 1941**

Geographic distribution. Japan, Korea, China (Herman and Chiodini 2009).

Natural hosts. First intermediate hosts are freshwater copepods. Second intermediate hosts are freshwater fish. Amphibians, birds, rodents, and reptiles can serve as paratenic hosts. Definitive hosts are weasels and minks. Zoonotic in humans as dead-end hosts harboring L3 larvae (Herman and Chiodini 2009).

Route of infection. Ingestion of L3 larvae in infected intermediate or paratenic hosts (Herman and Chiodini 2009).

Site in human host. Skin, heart, intestinal tract, ears, eyes, CNS (Herman and Chiodini 2009).

***Gnathostoma spinigerum* Levinsen, 1889**

Geographic distribution. Southeast Asia, southern Africa, Madagascar, Australia (Herman and Chiodini 2009).

Natural hosts. First intermediate hosts are freshwater copepods. Second intermediate hosts are freshwater fish. Amphibians, birds, rodents, and reptiles

can serve as paratenic hosts. Definitive hosts are wild and domestic canids and felids. Zoonotic in humans as dead-end hosts harboring L3 larvae (Herman and Chiodini 2009).

Route of infection. Ingestion of L3 larvae in infected intermediate or paratenic hosts (Herman and Chiodini 2009).

Site in human host. Skin, heart, intestinal tract, ears, eyes, CNS (Herman and Chiodini 2009).

●●●●●●●●●● **Camallanida Travassos, 1920**

●●●●●●●●●● **Dracunculidae Stiles, 1907**

Genus *Dracunculus* Reichard, 1759

***Dracunculus medinensis* (Linnaeus, 1758)**

Geographic distribution. Focal areas in sub-Saharan Africa, including Chad, Angola, and South Sudan (Eberhard et al. 2014b; Hopkins et al. 2020).

Natural hosts. Intermediate hosts are freshwater copepods; freshwater fish and amphibians can serve as paratenic or transport hosts. Definitive hosts are humans; canids, felids, and non-human primates can serve as reservoir hosts (Eberhard et al. 2014b; Hopkins et al. 2020).

Route of infection. Ingestion of L3 larvae in infected copepods or paratenic hosts (Eberhard et al. 2014b; Hopkins et al. 2020).

Site in human host. Subcutaneous tissues (Eberhard et al. 2014b; Hopkins et al. 2020).

Notes. Reports of *D. medinensis* infections acquired outside of historically endemic zones and/or in zones where eradication has been well-established are highly likely to represent misidentifications with other nematodes. In limited instances, these represent genuine infections with other unknown *Dracunculus* spp. (Kobayashi et al. 1986).

***Dracunculus* sp.**

Geographic distribution. Vietnam (Thach et al. 2021).

Natural hosts. Unknown; presumed zoonotic in humans (Thach et al. 2021).

Route of infection. Unknown; presumed ingestion of intermediate or reservoir host.

Site in human host. Subcutaneous abscesses (Thach et al. 2021).

Notes. There is a single report of human infection with a non-*medinensis* *Dracunculus* species from Vietnam. Genetically it is most similar to *D. insignis* (Leidy, 1858) and *D. lutrae* Crichton et Beverley-Burton, 1973, parasites of carnivores (Thach et al. 2021).

Spiruroidea, incertae sedis:**Spirurina Type X**

Geographic distribution. Japan (Goto et al. 1998; Makino et al. 2014).

Natural hosts. Intermediate or paratenic hosts are marine squid and fish. Definitive hosts unknown. Zoonotic in humans (Goto et al. 1998; Makino et al. 2014).

Route of infection. Presumed ingestion of L3 larvae in undercooked squid or fish (Goto et al. 1998; Makino et al. 2014).

Site in human host. Skin, eyes, ileum (Goto et al. 1998; Makino et al. 2014).

Notes. Around 50 infections with *Spirurina* type X larvae have been identified in humans who have consumed various seafood items, particularly small squid species. Molecular studies tentatively identified similar type X larvae as those of *Crassicauda giliakiana* Skrjabin & Andreeva, 1934, a nematode of Baird's beaked whale (*Berardius bairdii*) (Sugiyama et al. 2007).

●●●●●●●●●● **Thelazioidea Skrjabin, 1915**

●●●●●●●●●● **Thelaziidae Skrjabin, 1915**

Genus *Oxyspirura* Dräsche in Stossich, 1897

Geographic distribution. Worldwide (Dung et al. 2020).

Natural hosts. Intermediate hosts are arthropods. Definitive hosts are birds and non-human primates. Zoonotic in humans as dead-end hosts harboring L3 larvae (Dung et al. 2020).

Route of Infection. Ingestion of infected arthropod intermediate host (Dung et al. 2020).

Site in human host. Skin (Dung et al. 2020).

Notes. *Oxyspirura* has been reported as a cause of pruritic cutaneous larval migrans in a patient in Vietnam, but the specimen was not characterized at the species level (Dung et al. 2020).

Genus *Rictularia* Froelich, 1802

Geographic distribution. Worldwide (Kenney et al. 1975).

Natural hosts. Arthropods are intermediate hosts. A wide range of mammals serve as definitive hosts. Zoonotic in humans (Kenney et al. 1975).

Route of infection. Presumed ingestion of L3 larvae in infected intermediate hosts.

Site in human host. Appendix (Kenney et al. 1975).

Notes. The single human case of infection with *Rictularia* was reported from a patient in New York postmortem (Kenney et al. 1975).

Genus *Thelazia* Bosc in Blainville 1819***Thelazia californiensis* Price, 1930**

Geographic distribution. Western North America (Bradbury et al. 2018).

Natural hosts. Vector intermediate hosts are flies in the genus *Fannia*. Definitive hosts are mammals, including wild and domestic carnivores, deer, sheep, and lagomorphs. Zoonotic in humans (Bradbury et al. 2018).

Route of infection. Deposition of L3 larvae on the eye by vector fly (Bradbury et al. 2018).

Site in human host. Eye (Bradbury et al. 2018).

***Thelazia callipaeda* Railliet & Henry, 1910**

Geographic distribution. Europe, Asia (Otranto and Dutto 2008; Sah et al. 2018).

Natural hosts. Vector intermediate hosts are flies in the families Drosophilidae, primarily the genus *Phortica*. Definitive hosts are mammals, primarily carnivores and lagomorphs. Zoonotic in humans (Otranto and Dutto 2008; Sah et al. 2018).

Route of infection. Deposition of L3 larvae on the eye by vector fly (Otranto and Dutto 2008; Sah et al. 2018).

Site in human host. Eye (Otranto and Dutto 2008; Sah et al. 2018).

***Thelazia gulosa* (Railliet & Henry, 1910)**

Geographic distribution. North America, Europe, Central Asia, Australia (Bradbury et al. 2018; Bradbury et al. 2019a).

Natural hosts. Vector intermediate hosts are muscid flies in the genus *Musca*. Definitive hosts are ungulates, primarily cattle. Zoonotic in humans (Bradbury et al. 2018; Bradbury et al. 2019a).

Route of infection. Deposition of L3 larvae on the eye by vector fly (Bradbury et al. 2018; Bradbury et al. 2019a).

Site in human host. Eye (Bradbury et al. 2018; Bradbury et al. 2019a).

●●●●●●●● **Physalopteroidea Railliet, 1893**●●●●●●●● **Physalopteridae Railliet, 1893****Genus *Physaloptera* Rudolphi, 1819*****Physaloptera caucasica* (von Linstow, 1902)**

Geographic distribution. Africa, Asia (Vandepitte et al. 1964; Makki et al. 2017).

Natural hosts. Intermediate hosts are insects, especially crickets, cockroaches, and beetles. Definitive hosts are non-human primates. Zoonotic in humans (Vandepitte et al. 1964; Makki et al. 2017).

Route of infection. Presumed ingestion of L3 larvae in infected insects (Vandepitte et al. 1964; Makki et al. 2017).

Site in human host. Gastrointestinal tract (Vandepitte et al. 1964; Makki et al. 2017).

Notes. This species is often placed in the genus *Abbreviata* Travassos, 1926. Eggs consistent with *Physaloptera* spp. have been recovered from human coprolites in archaeological sites across the world, but species identity has not been established in paleoparasitological cases (Makki et al. 2017).

●●●●●●● Rhabditina Chitwood, 1933

●●●●●●● Rhabditida Orley, 1880

●●●●●●● Rhabditidae Orley, 1880

Genus *Diploscapter* Cobb, 1913

Diploscapter coronata (Cobb, 1893)

Geographic distribution. Worldwide (Morimoto et al. 2006; Chandler 2009).

Natural hosts. None (environmental); humans are incidental hosts (Morimoto et al. 2006; Chandler 2009).

Route of infection. Unknown, presumed ingestion of L3 larvae on contaminated produce (Morimoto et al. 2006; Chandler 2009).

Site in human host. Gastrointestinal tract (Morimoto et al. 2006; Chandler 2009).

Genus *Halicephalobus* Timm, 1956

Halicephalobus gingivalis (Stephanski, 1954)

Micronema deletrix Anderson & Bemrick, 1965

Geographic distribution. Worldwide (Onyiche et al. 2018).

Natural hosts. None (environmental); humans are incidental hosts (Onyiche et al. 2018).

Route of infection. Unknown, presumed inoculation of L3 larvae into wounds or mucus membranes; organ transplantation (Onyiche et al. 2018).

Site in human host. Disseminated infection with predilection for CNS (Papadi et al. 2013; Onyiche et al. 2018).

Genus *Pelodera* Schneider, 1866***Pelodera strongyloides* (Schneider, 1860)**

Geographic distribution. Worldwide (Saari and Nikander 2006).

Natural hosts. None (environmental); humans are incidental hosts (Saari and Nikander 2006).

Route of infection. Unknown, presumed direct penetration of skin by L3 larvae or possibly inoculation into wounds (Saari and Nikander 2006).

Site in human host. Skin (Orihel and Ash 1995).

Genus *Rhabditis* Dujardin, 1845

Geographic distribution. Worldwide (Fadaei Tehrani et al. 2019).

Natural hosts. None (environmental); humans are incidental hosts (Fadaei Tehrani et al. 2019).

Route of infection. Unknown, presumed inoculation of L3 larvae into wounds or mucus membranes (Fadaei Tehrani et al. 2019).

Site in human host. Small intestine, urogenital tract, ear (Teschner et al. 2014; Fadaei Tehrani et al. 2019).

Notes. Human cases of *Rhabditis* are not usually characterized at the species level. There have been cases specifically attributed to *R. axei* (Cobbold, 1884) from China (Yu et al. 2019), Iran (Meamar et al. 2007), and Zimbabwe (Goldsmid 1967) and *R. hominis* Kobayashi, 1920 from Japan and the United States (Sandground 1925). It is not always clear which cases represent true infection versus environmental contamination of clinical specimens (Sandground 1925).

●●●●●●●● Strongyloididae Chitwood & McIntosh, 1934**Genus *Strongyloides* Grassi, 1879*****Strongyloides fuellborni fuellborni* von Linstow, 1905**

Geographic distribution. Africa, Southeast Asia (Viney et al. 1991; Barratt et al. 2019).

Natural hosts. Non-human primates. Zoonotic in humans (Viney et al. 1991; Barratt et al. 2019).

Route of infection. Penetration of the skin by L3 larvae (Viney et al. 1991; Barratt et al. 2019).

Site in human host. Small intestine, disseminated infection in immunocompromised hosts, including to the lung, skin, kidneys, and brain (Viney et al. 1991; Barratt et al. 2019).

***Strongyloides fuellborni kellyi* Viney et al., 1991**

Geographic distribution. Papua New Guinea (Viney et al. 1991; Barratt et al. 2019).

Natural hosts. Non-human primates. Zoonotic in humans (Viney et al. 1991; Barratt et al. 2019).

Route of infection. Penetration of the skin by L3 larvae (Viney et al. 1991; Barratt et al. 2019).

Site in human host. Small intestine, disseminated infection in immunocompromised hosts, including to the lung, skin, kidneys, and brain (Viney et al. 1991; Barratt et al. 2019).

***Strongyloides myopotami* Artigas & Pacheco, 1933**

Geographic distribution. South America, North America, Europe, Africa, Asia (Choe et al. 2014).

Natural hosts. Nutria. Zoonotic in humans as a dead-end host (Choe et al. 2014).

Route of infection. Penetration of skin by L3 larvae (Choe et al. 2014).

Site in human host. Skin (Choe et al. 2014).

Notes. *Strongyloides myopotami* causes a short-lived dermatologic reaction in human skin, commonly referred to as ‘nutria itch’ or ‘marsh itch’. The parasite cannot survive in the human host and does not migrate beyond the dermis (Choe et al. 2014). Experimental evidence suggests the possibility of fleeting patent infection in humans (Little 1965).

***Strongyloides procyonis* Little, 1966**

Geographic distribution. North America, Asia (Choe et al. 2014).

Natural hosts. Raccoons. Zoonotic in humans as a dead-end host (Choe et al. 2014).

Route of infection. Penetration of skin by L3 larvae (Choe et al. 2014).

Site in human host. Skin (Choe et al. 2014).

Notes. *Strongyloides procyonis* causes a short-lived dermatologic reaction in human skin, commonly referred to as ‘swimmer’s itch’ or ‘marsh itch’. The parasite cannot survive in the human host and does not migrate beyond the dermis (Choe et al. 2014).

***Strongyloides stercoralis* Bavay, 1876**

Geographic distribution. Worldwide (Page et al. 2018; Barratt et al. 2019).

Natural hosts. Humans; dogs may serve as reservoir hosts (Page et al. 2018; Barratt et al. 2019).

Route of infection. Penetration of the skin by L3 larvae; organ transplantation (Abanyie et al. 2015; Page et al. 2018; Barratt et al. 2019).

Site in human host. Small intestine, disseminated infection in immunocompromised hosts, including to the lung, skin, kidneys, and brain (Page et al. 2018; Barratt et al. 2019).

- Strongylida Molin, 1861
- Chabertiidae Popova, 1952

Genus *Oesophagostomum* Molin, 1861

***Oesophagostomum aculaetum* (von Linstow, 1879)**

Geographic distribution. Southeast Asia, Japan (Polderman and Blotkamp 1995).

Natural hosts. Non-human primates. Zoonotic in humans (Polderman and Blotkamp 1995).

Route of infection. Ingestion of L3 larvae on plants (Polderman and Blotkamp 1995).

Site in human host. Large intestine (Ghai et al. 2014; Polderman and Blotkamp 1995).

***Oesophagostomum bifurcum* (Creplin, 1849)**

Geographic distribution. Sub-Saharan Africa (Polderman and Blotkamp 1995).

Natural hosts. Non-human primates. Zoonotic in humans (Polderman and Blotkamp 1995).

Route of infection. Ingestion of L3 larvae on plants (Polderman and Blotkamp 1995).

Site in human host. Large intestine (Polderman and Blotkamp 1995; Ghai et al. 2014).

***Oesophagostomum stephanostomum* Stossich, 1904**

Geographic distribution. Sub-Saharan Africa, Brazil (Polderman and Blotkamp 1995).

Natural hosts. Non-human primates. Zoonotic in humans (Glen and Brooks 1985; Polderman and Blotkamp 1995).

Route of infection. Ingestion of L3 larvae on plants (Polderman and Blotkamp 1995).

Site in human host. Large intestine (Polderman and Blotkamp 1995; Ghai et al. 2014).

- Strongylidae Baird, 1853

Genus *Ternidens* Railliet & Henry, 1909

***Ternidens deminutus* (Railliet & Henry, 1905)**

Geographic distribution. Sub-Saharan Africa (Bradbury 2019).

Natural hosts. Non-human primates. Zoonotic in humans (Bradbury 2019).

Route of infection. Unknown; presumed ingestion of L3 larvae on contaminated food or fomites, or ingestion of L3 in arthropod intermediate or paratenic hosts (Bradbury 2019).

Site in human host. Large intestine (Bradbury 2019).

●●●●●●●●●● Syngamidae Leiper, 1912

Genus *Mammomonogamus* Ryzhikovk, 1948

Mammomonogamus laryngeus Ryzhikovk, 1948

Geographic distribution. Circumtropical (Lopes-Torres et al. 2020).

Natural hosts. Mammals, including ruminants and felids. Annelids, mollusks, and arthropods may serve as paratenic hosts. Zoonotic in humans (da Costa et al. 2005; Lopes-Torres et al. 2020).

Route of infection. Unknown, presumed to be from ingestion of embryonated eggs or larvae, or the consumption of a paratenic host (da Costa et al. 2005).

Site in human host. Bronchial tree (da Costa et al. 2005).

●●●●●●●●●● Ancylostomatidae Looss, 1905

Genus *Ancylostoma* Dubini, 1843

Ancylostoma braziliense Gomes de Faria, 1910

Geographic distribution. Southern United States, Central and South America, southern Africa, Southeast Asia (Tekely et al. 2013).

Natural hosts. Wild and domestic canids and felids. Zoonotic in humans as a dead-end host (Tekely et al. 2013).

Route of infection. Penetration of skin by L3 larvae (Tekely et al. 2013).

Site in human host. Skin (Tekely et al. 2013).

Ancylostoma caninum (Ercolani, 1859)

Geographic distribution. Worldwide (Tekely et al. 2013; Shepherd et al. 2018).

Natural hosts. Wild and domestic canids and felids. Zoonotic in humans (Tekely et al. 2013; Shepherd et al. 2018).

Route of infection. Penetration of skin by L3 larvae (Tekely et al. 2013).

Site in human host. Skin, rarely small intestine (Tekely et al. 2013).

Ancylostoma ceylanicum Looss, 1911

Geographic distribution. Southeast Asia, Australia, Middle East (Traub 2013).

Natural hosts. Wild and domestic canids and felids. Zoonotic in humans (Traub 2013).

Route of infection. Penetration of skin by L3 larvae (Traub 2013).

Site in human host. Small intestine (Traub 2013).

***Ancylostoma duodenale* (Dubini, 1843)**

Geographic distribution. Worldwide in tropics and subtropics; host spots of endemicity for human infection are China, India, Egypt, northern Australia, Latin America (Brooker et al. 2004; Hotez et al. 2004).

Natural hosts. Mammals, including humans, dogs, cats (Brooker et al. 2004; Hotez et al. 2004).

Route of infection. Penetration of skin by L3 larvae (Brooker et al. 2004; Hotez et al. 2004).

Site in human host. Small intestine (Brooker et al. 2004; Hotez et al. 2004).

Genus *Bunostomum* Railliet, 1902***Bunostomum phlebotomum* (Railliet, 1900)**

Geographic distribution. Worldwide (Tekely et al. 2013).

Natural hosts. Bovids. Zoonotic in humans as a dead-end host (Tekely et al. 2013).

Route of infection. Penetration of skin by L3 larvae (Tekely et al. 2013).

Site in human host. Skin (Tekely et al. 2013).

Genus *Necator* Stiles, 1903***Necator americanus* (Stiles, 1902)**

Geographic distribution. Worldwide in tropics and subtropics; hot spots of endemicity for human infection are southern China, southern India, Southeast Asia, sub-Saharan Africa, Latin America, southeastern USA (Brooker et al. 2004; Hotez et al. 2004).

Natural hosts. Humans (Brooker et al. 2004; Hotez et al. 2004).

Route of infection. Penetration of skin by L3 larvae (Brooker et al. 2004; Hotez et al. 2004).

Site in human host. Small intestine (Brooker et al. 2004; Hotez et al. 2004).

***Necator gorillae* Noda & Yamada, 1964**

Geographic distribution. Sub-Saharan Africa (Kalousová et al. 2016).

Natural hosts. Gorillas, chimpanzees. Zoonotic in humans (Kalousová et al. 2016).

Route of infection. Penetration of skin by L3 larvae (Kalousová et al. 2016).

Site in human host. Small intestine (Kalousová et al. 2016).

Genus *Uncinaria* Frölich, 1789***Uncinaria stenocephala* (Railliet, 1884)**

Geographic distribution. Temperate and subarctic regions of the Northern Hemisphere (Tekely et al. 2013).

Natural hosts. Carnivores, including wild and domestic canids and felids. Zoonotic in humans as a dead-end host (Tekely et al. 2013).

Route of infection. Penetration of skin by L3 larvae (Tekely et al. 2013).

Site in human host. Skin (Tekely et al. 2013).

●●●●●●●● **Trichostrongylidae Leiper, 1912****Genus *Haemonchus* Cobb, 1989*****Haemonchus contortus* (Rudolphi, 1802)**

Geographic distribution. Worldwide (Ghadirian and Arfaa 1975).

Natural hosts. Many ruminants. Zoonotic in humans (Ghadirian and Arfaa 1975).

Route of infection. Ingestion of L3 larvae on contaminated plants and produce (Ghadirian and Arfaa 1975).

Site in human host. Small intestine (Ghadirian and Arfaa 1975).

Genus *Marshallagia* Orloff, 1933***Marshallagia marshalli* (Ransom, 1907)**

Geographic distribution. Worldwide (Ghadirian and Arfaa 1973).

Natural hosts. Many ruminants. Zoonotic in humans (Ghadirian and Arfaa 1973).

Route of infection. Ingestion of L3 larvae on contaminated plants and produce (Ghadirian and Arfaa 1973).

Site in human host. Small intestine (Ghadirian and Arfaa 1973).

Genus *Nematodirus* Ransom, 1907***Nematodirus abnormalis* (May, 1920)**

Geographic distribution. Worldwide (Ghadirian and Arfaa 1973; Bradbury 2006).

Natural hosts. Primarily sheep. Zoonotic in humans (Ghadirian and Arfaa 1973; Bradbury 2006).

Route of infection. Ingestion of L3 larvae on contaminated plants and produce (Ghadirian and Arfaa 1973; Bradbury 2006).

Site in human host. Small intestine (Ghadirian and Arfaa 1973; Bradbury 2006).

Genus *Ostertagia* Ransom, 1907***Ostertagia ostertagi* (Stiles, 1892)**

Geographic distribution. Worldwide (Ghadirian and Arfaa 1973; Anderson 1988).

Natural hosts. Primarily bovids; also sheep, goats, equids, and other ruminants. Zoonotic in humans (Ghadirian and Arfaa 1973; Anderson 1988).

Route of infection. Ingestion of L3 larvae on contaminated plants and produce (Ghadirian and Arfaa 1973).

Site in human host. Small intestine (Ghadirian and Arfaa 1973).

Genus *Teladorsagia* Andreeva & Satubaldin, 1953***Teladorsagia circumcincta* (Stadelman, 1894)**

Geographic distribution. Temperate climates (Ashrafi et al. 2020).

Natural hosts. Primarily sheep, also goats; zoonotic in humans (Ashrafi et al. 2020).

Route of infection. Ingestion of L3 larvae on plants and produce (Ashrafi et al. 2020).

Site in human host. Small intestine (Ashrafi et al. 2020).

Genus *Trichostrongylus* Looss, 1905***Trichostrongylus axei* (Cobbold, 1879)**

Geographic distribution. Worldwide (Sato et al. 2011).

Natural hosts. Ungulates, primarily cattle, sheep, goats, and horses. Zoonotic in humans (Sato et al. 2011).

Route of infection. Ingestion of L3 larvae on plants and produce (Cancrini et al. 1982; Sato et al. 2011).

Site in human host. Small intestine (Sato et al. 2011).

***Trichostrongylus capricola* Ransom, 1907**

Geographic distribution. Worldwide (Ghadirian and Arfaa 1973; Cancrini et al. 1982).

Natural hosts. Ungulates, primarily cattle, sheep, and goats. Zoonotic in humans (Ghadirian and Arfaa 1973; Cancrini et al. 1982).

Route of infection. Ingestion of L3 larvae on plants and produce (Ghadirian and Arfaa 1973; Cancrini et al. 1982).

Site in human host. Small intestine (Ghadirian and Arfaa 1973; Cancrini et al. 1982).

***Trichostrongylus colubriformis* (Giles, 1892)**

Geographic distribution. Worldwide; predominate in the Middle East (Sato et al. 2011).

Natural hosts. Ungulates, primarily cattle, sheep, and goats. Zoonotic in humans (Sato et al. 2011).

Route of infection. Ingestion of L3 larvae on plants and produce (Sato et al. 2011).

Site in human host. Small intestine (Sato et al. 2011).

***Trichostrongylus longispicularis* Gordon, 1933**

Trichostrongylus lerouxi Biocca et al., 1974

Geographic distribution. Worldwide (excluding Africa) (Ghadirian 1977).

Natural hosts. Ungulates, primarily cattle, sheep, and goats. Zoonotic in humans (Ghadirian 1977).

Route of infection. Ingestion of L3 larvae on plants and produce (Ghadirian 1977).

Site in human host. Small intestine (Ghadirian 1977).

***Trichostrongylus orientalis* Jimbo, 1914**

Geographic distribution. Southeast Asia, Japan (Sato et al. 2011).

Natural hosts. Ungulates, including cattle, sheep, goats, and horses. Zoonotic in humans (Sato et al. 2011).

Route of infection. Ingestion of L3 larvae on plants and produce (Sato et al. 2011).

Site in human host. Small intestine (Sato et al. 2011).

***Trichostrongylus vitrinus* Looss, 1905**

Geographic distribution. Worldwide (Ghadirian and Arfaa 1975; Cancrini et al. 1982).

Natural hosts. Ungulates, including sheep, goats. Zoonotic in humans (Ghadirian and Arfaa 1975; Cancrini et al. 1982).

Route of infection. Ingestion of L3 larvae on plants and produce (Ghadirian and Arfaa 1975; Cancrini et al. 1982).

Site in human host. Small intestine (Ghadirian and Arfaa 1975; Cancrini et al. 1982).

●●●●●●●● *Angiostrongylidae* Boehm & Gebauer, 1934**Genus *Angiostrongylus* Kamensky, 1905*****Angiostrongylus cantonensis* (Chen, 1935)**

Geographic distribution. Asia, South Pacific, Hawaii, Caribbean, Africa, southern United States, Central and South America (Cowie 2013).

Natural hosts. Intermediate hosts are terrestrial mollusks. Paratenic hosts include reptiles, amphibians, planarians. Definitive hosts are rodents, primarily rats (*Rattus*) and cotton rats (*Sigmodon*). Zoonotic in humans as a dead-end host harboring L4 larvae and young adults (Cowie 2013).

Route of infection. Ingestion of L3 larvae in infected mollusks or paratenic hosts (Cowie 2013).

Site in human host. CNS (Cowie 2013).

Angiostrongylus costaricensis Morera & Cespedes, 1971

Geographic distribution. Southern United States, Central and South America (Romero-Alegria 2014).

Natural hosts. Intermediate hosts are terrestrial mollusks. Definitive hosts are rodents. Zoonotic in humans (Romero-Alegria 2014).

Route of infection. Ingestion of L3 larvae in infected mollusks or contaminated produce (Romero-Alegria 2014).

Site in human host. Mesenteric blood vessels (Romero-Alegria 2014).

Angiostrongylus malaysiensis Bhaibulaya & Cross, 1971

Angiostrongylus cantonensis Malaysia strain

Geographic distribution. Southeast Asia (Rodpai et al. 2016).

Natural hosts. Intermediate hosts are terrestrial mollusks. Paratenic hosts include reptiles, amphibians, planarians. Definitive hosts are rodents, primarily rats. Apparently zoonotic in humans as a dead-end host harboring larvae (Rodpai et al. 2016; Watthanakulpanich et al. 2021).

Route of infection. Ingestion of L3 larvae in infected mollusks or paratenic hosts (Watthanakulpanich et al. 2021).

Site in human host. CNS (Watthanakulpanich et al. 2021).

●●●●●●●● *Metastrongylidae* Leiper, 1909

Genus *Metastrongylus* Molin, 1861

Metastrongylus elongatus Dujardin, 1845

Strongylus apri Gmelin, 1790

Geographic distribution. Worldwide (Calvopina et al. 2016).

Natural hosts. Intermediate hosts are earthworms. Definitive hosts are wild and domestic pigs. Zoonotic in humans (Calvopina et al. 2016).

Route of infection. Presumed ingestion of L3 larvae in infected earthwormsv

Site in human host. Lungs (Calvopina et al. 2016).

Metastrongylus salmi Gedoelst, 1923

Geographic distribution. Worldwide (Calvopina et al. 2016).

Natural hosts. Wild and domestic swine are definitive hosts; one incidental infection is reported in a human (Calvopina et al. 2016).

Route of infection. Presumed ingestion of L3 larvae in infected earthworms.

Site in human host. Lungs (Calvopina et al. 2016).

Excluded species

The following genera and species have been previously reported as human parasites. They are excluded from the above checklist because it is not believed they can cause parasitic infection in the human host, are based on demonstrable misidentifications, confirmation of identification is required, or because of taxonomic changes. The organisms are listed in alphabetical order.

Agamomermis spp. Members of the genus *Agamomermis* are mermithid nematodes that are free-living as adults but infect insects as larvae. Human cases are believed to represent spurious passage following accidental ingestion of worms in contaminated food or water (Leon 1946; Chabaud and Lanz 1951).

Amblyomma argentiniae Neumann, 1905. This Neotropical tick was recorded as a human parasite under its synonym *A. testudinis* (Conil, 1877) (Doss 1974), but that record is believed to be in error (Guglielmone and Robbins 2018).

Amblyomma auricularium (Conil, 1878). This Neotropical tick is a parasite of several groups of mammals and birds. Records of this species from humans are believed to represent misidentifications, primarily of *A. parvum* (Guglielmone and Robbins 2018).

Amblyomma calcaris Nakatsudi, 1942. This tick was described from a human in China (Nakatsudi 1942). Apparently the species was not described properly and the name is invalid (Guglielmone and Robbins 2018)

Amblyomma compressum (Macalister, 1872). This African tick species is a parasite of mammals, especially pangolins and rodents, birds, and reptiles. Records of this species from unknown locations in Africa (Theiler 1962) require confirmation (Guglielmone and Robbins 2018).

Amblyomma geayi Neumann, 1899. This tick is a parasite of various mammals and birds in Central and South America. A record of this species from a human (Esser et al. 2016) is based on conflicting host data in the paper and needs to be confirmed (Guglielmone and Robbins 2018).

Amblyomma helvolum Koch, 1844. This Southeast Asian tick is primarily a parasite of reptiles, and occasionally mammals. Records of this species feeding on humans

(Doss 1974) is based on an earlier publication that stated the ticks were merely crawling on humans (Audy 1960). To date, there are no records of *A. helvolum* feeding on humans (Guglielmone and Robbins 2018).

Amblyomma macfarlandi Kierans et al., 1973. This tick is a parasite of tortoises on the Galapagos Islands. The single record from a human (Guglielmone et al. 2006) was not confirmed as feeding on the human host and may have merely been an incidental finding (Guglielmone and Robbins 2018).

Amblyomma pomposum Dönitz, 1909. This African tick is a parasite of several groups of mammals and birds. Records from humans require confirmation (Guglielmone and Robbins 2018).

Amblyomma sylvaticum (De Geer, 1778). This tick is a parasite of reptiles in South Africa. Records of this species from humans are believed to represent misidentifications (Guglielmone and Robbins 2018).

Amblyomma usingeri Keirans et al., 1973. This tick is a parasite of tortoises on the Galapagos Islands. The single record from a human (Guglielmone et al. 2006) was not confirmed as feeding on the human host and may have merely been an incidental finding (Guglielmone and Robbins 2018).

Amphimermis elegans (Hagmeier, 1912). *Amphimermis elegans* is an Asian mermithid nematode parasitic on orthopteran insects. A human case (as *Mermis*) reportedly recovered from urine probably represents pseudoparasitism or contamination of the toilet by the insect host (Hasegawa et al. 1996).

Androlaelaps casalis (Berlese, 1887). This laelapid mite is a predator on other mites and insects. It was reported as a cause of human dermatitis in Israel (Rosen et al. 2002). The mouthparts of *A. casalis* are not adapted for piercing vertebrate skin and the dermatitis in that report may have been associated with *Dermanyssus gallinae*, which was also reported by the authors and is a host for *A. casalis*. More research is needed to determine if *A. casalis* can bite humans and be a cause of dermatitis.

***Anisopus* sp.** Anisopodid flies are commonly called wood gnats or window gnats. They have been implicated in intestinal and urogenital myiasis (Smith and Taylor 1966). The larvae of anisopodids breed in decaying vegetation, fermenting sap, animal manure, tree holes, mud, and sewage. Their presence in toilets, latrines, and similar should be regarded as incidental as they feed on detritus in these substrates.

Bertiella satyri Blanchard, 1891. This cestode was originally described from orangutans. Reports from humans (Chandler 1925) are believed to represent misidentifications of *B. studeri* (Sapp and Bradbury 2020).

Caccobius vulcanus (Fabricius, 1801). *Caccobius vulcanus* is a Palearctic coprophagous scarab beetles that has been implicated as a cause of scarabiasis in India (as *C. mutans* Sharp, 1875) (Iyengar 1928). Scarabiasis is a proposed phenomenon describing the colonization of the human intestinal tract with coprophagous scarab beetles, usually based on the finding of beetles in the soiled diapers of children. There is no pathology described for parasitism by scarab beetles and the phenomenon probably represents post-defecation contamination of diapers, toilets, latrines, and similar.

Clogmia albipunctata (Williston, 1893). *Clogmia albipunctata* is a psychodid fly with a nearly worldwide distribution. This species has been implicated as a cause of urogenital (El-Dib et al. 2017; Farrag et al. 2019), intestinal (Tu et al. 2007; Mokhtar et al. 2016a), and nasopharyngeal (Nevill et al. 1969; Mohammed and Smith 1976) myiasis, sometimes under the name *Telmatoscopus albipunctatus*. Psychodid flies reported as causative agents of urogenital and intestinal myiasis is usually due to the incidental finding of the flies in toilets, latrines, sinks, and bathtubs. The larvae of psychodid flies breed in biofilms, including in faucets and drains, giving the false impression they were shed in stool or urine. There is no evidence that psychodid fly larvae use human tissue as a nutritive source.

Crasodactylus punctatus Guérin-Ménéville, 1847. This carabid beetle was reported from the ear of two patients in Oman (misspelled as *Crasydactylus punctatus*) (Bhargava and Victor 1911). This beetle is predaceous on other insects and small invertebrates and its presence in the human ear canal should be regarded as incidental.

Cryptostrongylus pulmoni. The name *Cryptostrongylus pulmoni* is a provisional name given to suspect helminths associated with chronic fatigue syndrome (Klapow 1999). Images depicting this 'parasite' appear to be synthetic fibers. There is no evidence that *C. pulmoni* represents an actual animal, let alone one capable of parasitizing humans.

Cyclocephala borealis Arrow, 1911. This scarab beetle was implicated in a large-scale infestation of the ears of Boy Scouts in Pennsylvania, USA (Maddock and Fehn 1958). *Cyclocephala* species are phytophagous, and its presence in the human ear canal should be regarded as incidental.

Dermacentor cruentus Koch, 1844. This European tick was listed as a human parasite based on the original description (Doss 1974), but there is no indication in the original description it was observed feeding on humans (Guglielmone and Robbins 2018).

Dermacentor halli McIntosh, 1931. This tick is a parasite of various mammals in North and Central America. Records from humans require confirmation (Guglielmone and Robbins 2018).

Dermacentor taiwanensis Sugmito, 1935. This tick is a parasite of various mammals in China, Vietnam, and Taiwan. Records of this species from humans are believed to represent misidentifications (Guglielmone and Robbins 2018).

Dibothriocephalus alascense (Rausch & Rausch, 1956). This species was originally described from a domestic dog in the Yukon-Kuskokwim Delta of Canada. A single human case of this species was reported from an Eskimo in Alaska (Rausch et al. 1967), but the record is considered doubtful (Scholz and Kuchta 2016; Waeschenbach et al. 2017).

Diphyllobothrium cameroni Rausch, 1969. This species was originally described from the Hawaiian monk seal (*Neomonachus schauinslandi*) and has been recorded twice from humans in Japan (Kamo H. 1986), but the records are considered doubtful (Scholz and Kuchta 2016; Waeschenbach et al. 2017).

Diphyllobothrium elegans (Krabbe, 1865). This is a parasite of seals and has been described once from a human in Japan (Kamo H. 1986), but the record is considered doubtful (Scholz and Kuchta 2016; Waeschenbach et al. 2017).

Diphyllobothrium orcini Hatsushika & Shirouzu, 1990. This species was described from killer whales. There are two records from humans from Japan (Kifune 2000; Nakazawa 1992), but those records are considered doubtful (Waeschenbach et al. 2017).

Diphyllobothrium scoticum (Rennie & Reid, 1912). This species was described from the leopard seal (*Hydrurga leptonyx*) and has been recorded once from a human in Japan (Fukumoto 1988), but that record is considered doubtful (Waeschenbach et al. 2017).

Drosophila melanogaster Meigen, 1830. This common ‘fruit fly’ has been infrequently reported as a cause of nasal and ocular myiasis (Francesconi and Lupi 2012). Larvae breed in decaying vegetation and their presence in clinical specimens should be considered incidental.

Dryomyza formosa (Wiedemann, 1830). This dryomyzid fly was reported as a cause of gastrointestinal myiasis in a patient from Japan suffering from delusional parasitosis. Larvae were observed in fresh stool and it was speculated they represent spurious passage following accidental ingestion of the larvae (Chigusa et al. 2000). Dryomyzid larvae feed on decaying organic material and there is no evidence they use human intestinal tissue as a nutritive source.

Emys orbicularis (Linnaeus, 1758). *Emys orbicularis* is the Latin name of the European pond turtle. This name was used in the 23rd edition of “Manson’s Tropical Diseases” (Farrar 2014) as a species of leech that parasitizes humans. This probably represents an editorial error and may have been intended to refer to a leech that normally parasitizes the turtle.

Eristalis tenax (Linnaeus, 1758). This syrphid fly is frequently implicated in causing intestinal or urogenital myiasis (Francesconi and Lupi 2012). The larvae breed in decaying organic substrates, including manure, sewage, and contaminated water. Most reported cases are from the finding of larvae in latrines, outhouses, toilets, and similar and their presence in such locations is believed to be incidental.

***Euparyphium* spp.** Members of this genus of echinostome flukes has been recorded from humans in Laos (Toledo and Esteban 2016); however, the identity of human isolates attributed to *Euparyphium* should be confirmed (Chai et al. 2012).

Fannia scalaris (Fabricius, 1794). This fly was reported as a cause of urogenital myiasis in 1975 (Werner et al. 1975; Francesconi and Lupi 2012). The specimen was reportedly recovered in urine and likely represents contamination of a toilet, latrine, or the specimen itself.

***Gordius* spp.** A number of “gordioid worms” or “horsehair worms”, including *Gordius*, have been recovered from humans (typically in vomitus) (Kagei et al. 1966; Uchikawa et al. 1987; Herter and Nesse 1989; Lee et al. 2003). These most likely represent spurious passage following accidental ingestion of infected arthropod hosts or worms free in the environment. The finding of such worms in toilets is usually due to the drowning of the insect host by the parasite and liberation of the parasite therefrom.

- Haemaphysalis cinnabarina*** Koch, 1844. Records of this Brazilian tick from humans refer to *H. chordeilis* (Guglielmone and Robbins 2018).
- Haemaphysalis kashmirensis*** Hoogstraal & Varma, 1962. This tick is a parasite of mammals and reptiles in India and Pakistan. Records from humans require confirmation (Guglielmone and Robbins 2018).
- Haemaphysalis muhsamae*** Santos Dias, 1954. This tick is a parasite of birds in Africa. Records of this species from humans require confirmation due to the morphologic challenges in identifying this species (Guglielmone and Robbins 2018).
- Haemaphysalis proxima*** Aragão, 1911. This species has been recorded from humans in Colombia. There is no formal description for this species and the name is considered *nomen nudum* (Guglielmone and Robbins 2018). Records of this species from humans probably pertain to *H. leporispalustris* (Guglielmone and Robbins 2018).
- Haemaphysalis warburtoni*** Nuttall, 1912. This tick is a parasite of bovids and rodents in India, Nepal, and China. Records from humans may be based on misidentifications and require confirmation (Guglielmone and Robbins 2018).
- Hyalomma franchinii*** Tonelli-Rondelli, 1932. This tick is a parasite of mammals and reptiles in North Africa and the Middle East. Records from humans are based on misidentifications of *H. excavatum* and *H. marginatum* (Guglielmone and Robbins 2018).
- Hyalomma impressum*** Koch, 1844. This tick is a parasite of various mammals and birds from sub-Saharan Africa. This species has been recorded from a human in South Africa (Bedford 1927); however, *H. impressum* does not occur in South Africa and that record may represent a misidentification (Guglielmone and Robbins 2018).
- Hyalomma plubeum*** (Panzer, 1795). This name is currently considered incertae sedis, and records of *H. plumbeum* from humans currently pertain to *H. marginatum* (Guglielmone and Robbins 2018).
- Hermetia* spp.** Larvae of these soldier flies have been implicated in cases of intestinal and furuncular myiasis (Francesconi and Lupi 2012). Larvae of *Hermetia* spp. breed in decaying organic material, including outhouses. Their presence in clinical specimens is probably incidental and it is not believed they cause myiasis.
- Ixodes affinis*** Neumann, 1899. This North and Central American tick is primarily a parasite of a wide variety of mammals, including carnivores, marsupials, and ungulates. The species is difficult to identify morphologically and past records of this species from humans are believed to be misidentifications (Guglielmone and Robbins 2018).
- Ixodes humanus*** Koch, 1844. This species was described from a human in Brazil, however the identity of the species is not fully understood and it may be synonymous with a member of the genus *Amblyomma* (Guglielmone and Robbins 2018).
- Ixodes jellisoni*** Cooley & Kohls, 1938. This is a North American tick found on rodents and carnivores. Records of this species from humans are believed to be misidentifications (Guglielmone and Robbins 2018).
- Ixodes laysanensis*** Wilson, 1964. This tick is a bird parasite in Hawaii. Records of this species feed in humans (Doss 1974) are based on an earlier record that reported the tick crawling, and not feeding, on humans. To date, there is no evidence *I. laysanensis* feeds on humans (Guglielmone and Robbins 2018).

Ixodes loricatus Neumann, 1899. This tick is a parasite of marsupials and rodents in South America. Records from humans in Brazil (Serra-Freire 2011) require confirmation and are provisionally excluded as a parasite of humans (Guglielmone and Robbins 2018).

Ixodes luciae Senevet, 1940. This species is a Neotropical tick of various mammals, especially marsupials and rodents. The single record of this species on a human from Argentina (Ivancovich 1992) is believed to be based on a misidentification (Guglielmone and Robbins 2018).

Ixodes molestus James, 1923. This species was described as attacking humans in the USA, but the name is currently regarded as *nomen dubium* (Guglielmone and Robbins 2018).

Ixodes simplex Neumann, 1906. Records of this broadly-distributed Old World tick species of bats on humans from Japan cannot be confirmed and may be based on misidentifications (Guglielmone and Robbins 2018).

Ixodes trichosuri Roberts, 1960. This tick species is a parasite of marsupials and rodents in Australia. Records of this parasite infecting humans (Russell 2001) are based on unpublished records that require confirmation. Until such time, this species is removed from the list of ticks parasitizing humans (Guglielmone and Robbins 2018).

Lasioderma serricorne (Fabricius, 1792). *Lasioderma serricorne* is a cosmopolitan pest of dried, organic materials, including tobacco, cereals and other grains, dried fruit, and dried animal products. It was been reported as cause of canthariasis in infants in China (Sun et al. 2016) and Malaysia (Mokhtar et al. 2016b). Given the beetle's predilection for food products, and the age of the patients, this probably represents spurious passage following consumption of contaminated food.

***Ligula* spp.** *Ligula* species are diphyllbothriid parasites of fish-eating birds. There are two reports from humans, initially reported under the names *Diplogonophorus brauni* Leon, 1907 and *Braunia jassyensis* Leon, 1908. These records are believed to represent misidentifications (Scholz and Kuchta 2016).

Lophomonas blattarum Stein, 1860. There are many case reports in the literature of *L. blattarum* being isolated from human respiratory specimens. These all appear to be misidentifications of ciliocytophthoria, a condition whereby detached, motile epithelial cells are observed in clinical specimens (Hadziyannis et al. 2000; Li and Gao 2016). *Lophomonas blattarum* is a commensal in the gut of cockroaches and not a proven agent of human infection.

Maladera castanea (Arrow, 1913). This scarab beetle was implicated in a large-scale infestation of the ears of Boy Scouts in Pennsylvania, USA (as *Autoserica castanea*) (Maddock and Fehn 1958). *Maladera* species are phytophagous, and its presence in the human ear canal should be regarded as incidental.

Melophagus ovinus (Linnaeus, 1758). Commonly called the 'sheep ked', *M. ovinus* is a cosmopolitan parasite of domestic sheep (*Ovis aries*), as well as wild ungulates, rabbits, and wild and domestic canids. It has been reported as parasitizing humans (Zhao et al. 2018), however there is no evidence it can survive on human blood.

Musca nebulo Fabricius, 1794. This species was reported as cause of oral myiasis in India (Sharma et al. 2008). *Musca nebulo* is generally considered a synonym of *M. curviforceps* Saccà & Rivosecchi, 1956, which itself is considered a subspecies

or synonym of *M. domestica*. Also, *M. curviforceps* is considered endemic to sub-Saharan Africa (Marquez and Krafur 2002), which should preclude it from causing myiasis in India.

***Muscina* spp.** Flies in the genus *Muscina* have repeatedly been reported as causing intestinal myiasis following the recovery of fly larvae in stool (Francesconi and Lupi 2012). This probably represents post-defecation contamination of the stool specimen. There is no evidence *Muscina* uses human tissue as a nutritive source.

Onthophagus bifasciatus (Fabricius, 1781). *Onthophagus bifasciatus* is a Palearctic dung beetle that has been implicated as a causative agent of scarabiasis in India (Iyengar 1928). Scarabiasis is a proposed phenomenon describing the colonization of the human intestinal tract with coprophagous scarab beetles, usually based on the finding of beetles in the soiled diapers of children. There is no pathology described for parasitism by scarab beetles and the phenomenon probably represents post-defecation contamination of diapers, toilets, latrines, and similar.

Onthophagus unifasciatus (Schaller, 1783). *Onthophagus unifasciatus* is a Palearctic dung beetle that has been implicated as a causative agent of scarabiasis in Sri Lanka (Iyengar 1928; Gunawardena 1963). Scarabiasis is a proposed phenomenon describing the colonization of the human intestinal tract with coprophagous scarab beetles, usually based on the finding of beetles in the soiled diapers of children. There is no pathology described for parasitism by scarab beetles and the phenomenon probably represents post-defecation contamination of diapers, toilets, latrines, and similar.

Palpada scutellaris (Fabricius, 1805). This is a species of syrphid fly from Central and northern South America. It was reported as the cause of human intestinal myiasis in Costa Rica (Pérez-Bañón et al. 2020). The larvae breed in decaying organic substrates, including manure, sewage, and contaminated water. Most reported cases are from the finding of larvae in latrines, outhouses, toilets, and similar and their presence in such locations is believed to be incidental.

***Parachordodes* spp.** A number of “gordioid worms” or “horsehair worms”, including *Parachordodes*, have been recovered from humans (typically in vomitus) (Yamada et al. 2012). These most likely represent spurious passage following accidental ingestion of infected arthropod hosts or worms free in the environment. The finding of such worms in toilets is usually due to the drowning of the insect host by the parasite and liberation of the parasite therefrom.

Paragordius varius (Leidy, 1851). A number of “gordioid worms” or “horsehair worms”, including *Paragordius*, have been recovered from humans (typically in vomitus) (Ali-Khan and Ali-Khan 1977). These most likely represent spurious passage following accidental ingestion of infected arthropod hosts or worms free in the environment. The finding of such worms in toilets is usually due to the drowning of the insect host by the parasite and liberation of the parasite therefrom.

***Pericoma* spp.** *Pericoma* is a genus of Psychodidae that has been implicated as a cause of urinary myiasis in India (Singla et al. 2018). Psychodid flies reported as causative agents of urogenital and intestinal myiasis is usually due to the incidental finding of the flies in toilets, latrines, sinks, and bathtubs. The larvae of psychodid flies breed in biofilms, including in faucets and drains, giving the false impression they

were shed in stool or urine. There is no evidence psychodid fly larvae use human tissue as a nutritive source.

Piophilidae casei (Linnaeus, 1758). This species, commonly called the ‘cheese fly’, has been infrequently reported as a cause of intestinal or urogenital myiasis (Peckensneider et al. 1952; Saleh and el Sibae 1993). The larvae of this fly breeds in foodstuffs, and its presence in toilets and latrines should be considered incidental, probably following spurious passage of the larvae after eating infested food.

Psychoda albipennis Zetterstedt, 1850. This psychodid has been implicated as an agent of urogenital myiasis (Francesconi and Lupi 2012; Shimpi et al. 2018). Psychodid flies reported as causative agents of urogenital and intestinal myiasis is usually due to the incidental finding of the flies in toilets, latrines, sinks, and bathtubs. The larvae of psychodid flies breed in biofilms, including in faucets and drains, giving the false impression they were shed in stool or urine. There is no evidence psychodid fly larvae use human tissue as a nutritive source.

Psychoda alternata Say, 1824. This psychodid has been implicated as an agent of urogenital myiasis (Abul Hab 2001) and has been recorded from human sputa (Scott 1964). Psychodid flies reported as causative agents of urogenital and intestinal myiasis is usually due to the incidental finding of the flies in toilets, latrines, sinks, and bathtubs. The larvae of psychodid flies breed in biofilms, including in faucets and drains, giving the false impression they were shed in stool or urine. There is no evidence psychodid fly larvae use human tissue as a nutritive source.

Psychoda sexpunctata Curtis, 1839. This psychodid has been implicated as a source of gastrointestinal myiasis (Okada 1927). Psychodid flies reported as causative agents of urogenital and intestinal myiasis is usually due to the incidental finding of the flies in toilets, latrines, sinks, and bathtubs. The larvae of psychodid flies breed in biofilms, including in faucets and drains, giving the false impression they were shed in stool or urine. There is no evidence psychodid fly larvae use human tissue as a nutritive source.

Rhipicentor bicornis Nutall & Warburton, 1908. This tick is a parasite of various mammals in sub-Saharan Africa. This species was recorded as a parasite on humans (Doss 1974), however that record is based on a misunderstanding that a tick observed in a human dwelling meant it feeds on humans. To date, there are no records of *R. bicornis* feeding on humans (Guglielmone and Robbins 2018).

Sappinia diploidea (Hartmann & Naegler, 1908). The first case of human infection with a member of the genus *Sappinia* was reported to have been caused by *S. diploidea* based on morphologic criteria (Gelman et al. 2001). However, molecular characterization later demonstrated the species to be *S. pedata* (Qvarnstrom et al. 2009).

Scarites sulcatus Olivier, 1795. *Scarites sulcatus* is a Palearctic ground beetle that has been implicated as a cause of genital canthariasis (Paul 2007). *Scarites* species are free-living predaceous beetles. Given the habits of these beetles, this case probably represents environmental contamination or possibly urethral sounding.

***Scenopinus* spp.** Members of this genus of flies have been implicated in urogenital myiasis (Thompson et al. 1970). Larvae of *Scenopinus* are predators on the larvae of other insects. The presence of *Scenopinus* larvae in clinical specimens is probably incidental.

Trichophrya piscium Bütschli, 1899. This freshwater fish pathogen was reported as reported from sinus aspirates of a patient in Iraq (Al-Duboon and Disher 2018). Images of the suspect organism in the publication showed this to be a misidentification of ciliocytophthoria, a condition whereby detached, motile epithelial cells are observed in clinical specimens (Hadziyannis et al. 2000; Li and Gao 2016).

Tyroglyphus longior (Gervais, 1844). This grain mite has been implicated in intestinal acariasis by the finding of mites in the stool of two patients with generalized intestinal complaints (Harold Hinman and Kampmeier 1934). The finding of these mites probably represents spurious passage after incidental ingestion of the mites in contaminated foodstuffs.

Tyroglyphus putrescentiae (Schrank, 1781). This grain and mold mite has been implicated as a cause of intestinal acariasis (Khalifa et al. 2016). The finding of these mites probably represents spurious passage after incidental ingestion of the mites in contaminated foodstuffs.

Urbanorum. The name '*Urbanorum*' has been given to usual objects observed in stool specimens. Most cases have been reported from Central and South America (de Aguiar and Alves 2018). Generally referred to as protozoans, there does not appear to be any formal description of *Urbanorum* in the zoological literature and the general consensus is that these objects are nothing more than peculiar artifacts. No biochemical, ultrastructural, or genetic studies have been undertaken to confirm its status as a living organism, although the exact identity of it has not been ascertained.

Acknowledgements

The authors would like to thank Dr. Sina Adl for reviewing the manuscript prior to submission.

Disclaimer: The findings and conclusions in this report are those of the authors and do not necessarily represent the official position of the Centers for Disease Control and Prevention.

References

- Abanyie FA, Gray EB, Delli Carpini KW, Yanofsky A, McAuliffe I, Rana M, Chin-Hong PV, Barone CN, Davis JL, Montgomery SP, Huprikar S (2015) Donor-derived *Strongyloides stercoralis* infection in solid organ transplant recipients in the United States, 2009–2013. *American Journal of Transplantation* 15: 1369–1375. <https://doi.org/10.1111/ajt.13137>
- Abedkhozasteh H, Niyati M, Rahimi F, Heidari M, Farnia S, Rezaeian M (2013) First Report of *Hartmannella* keratitis in a Cosmetic Soft Contact Lens Wearer in Iran. *Iranian Journal of Parasitology* 8: 481–485.
- Abi-Akl P, Haddad G, Zaytoun G (2017) Otoacariasis: an infestation of mites in the ear. *Annals of Clinical Case Reports* 2: 1329.

- Abubakar S, Teoh BT, Sam SS, Chang LY, Johari J, Hooi PS, Lakhbeer-Singh HK, Italiano CM, Omar SF, Wong KT, Ramli N, Tan CT (2013) Outbreak of human infection with *Sarcocystis nesbitti*, Malaysia, 2012. *Emerging Infectious Diseases* 19: 1989–1991. <https://doi.org/10.3201/eid1912.120530>
- Abul Hab J (2001) A human urogenital myiasis by the larva of the moth fly *Psychoda alternata* Say (Diptera, Psychodidae). *Iraqi Journal of Medical Sciences* 1: 345–346.
- Adam RD (2001) Biology of *Giardia lamblia*. *Clinical Microbiology Reviews* 14: 447–475. <https://doi.org/10.1128/CMR.14.3.447-475.2001>
- Adl SM, Bass D, Lane CE, Lukes J, Schoch CL, Smirnov A, Agatha S, Berney C, Brown MW, Burki F, Cardenas P, Cepicka I, Chistyakova L, Del Campo J, Dunthorn M, Edvardsen B, Eglit Y, Guillou L, Hampl V, Heiss AA, Hoppenrath M, James TY, Karnkowska A, Karpov S, Kim E, Kolisko M, Kudryavtsev A, Lahr DJG, Lara E, Le Gall L, Lynn DH, Mann DG, Massana R, Mitchell EAD, Morrow C, Park JS, Pawlowski JW, Powell MJ, Richter DJ, Rueckert S, Shadwick L, Shimano S, Spiegel FW, Torruella G, Youssef N, Zlatogursky V, Zhang Q (2019) Revisions to the Classification, Nomenclature, and Diversity of Eukaryotes. *Journal of Eukaryotic Microbiology* 66: 4–119. <https://doi.org/10.1111/jeu.12691>
- Adl SM, Mathison BA (2019) Chapter 135: Taxonomy and classification of human eukaryotic parasites. In: Carroll KC, Pfäller MA, Landry ML et al. (Eds) *Manual of Clinical Microbiology*. ASM Press, Washington DC, 2620–2641.
- Aelami MH, Khoei A, Ghorbani H, Seilanian-Toosi F, Poustchi E, Hosseini-Farash BR, Moghaddas E (2019) Urinary Canthariasis Due to *Tenebrio molitor* Larva in a Ten-Year-Old Boy. *Journal of Arthropod Borne Diseases* 13: 416–419. <https://doi.org/10.18502/jad.v13i4.2239>
- Ağın H, Ayhan FY, Gülfidan G, Cevik D, Derebaşı H (2008) Severe anemia due to the pharyngeal leech *Limnatis nilotica* in a child. *Türkiye Parazitoloji Dergisi* 32: 247–248.
- Aizawa M, Morishima KJJOTJFS (2018) Distribution of *Haemadipsa japonica* in Japan before the 1980s. *Journal of the Japanese Forest Society* 100: 65–69. <https://doi.org/10.4005/jjfs.100.65>
- Ajaoud M, Es-sette N, Hamdi S, El-Idrissi AL, Riyad M, Lemrani M (2013) Detection and molecular typing of *Leishmania tropica* from *Phlebotomus sergenti* and lesions of cutaneous leishmaniasis in an emerging focus of Morocco. *Parasites & Vectors* 6: e217. <https://doi.org/10.1186/1756-3305-6-217>
- Akiyoshi DE, Dilo J, Pearson C, Chapman S, Tumwine J, Tzipori S (2003) Characterization of *Cryptosporidium meleagridis* of human origin passaged through different host species. *Infection and Immunity* 71: 1828–1832. <https://doi.org/10.1128/IAI.71.4.1828-1832.2003>
- Al-Arfaj AM, Mullen GR, Rashad R, Abdel-Hameed A, OConnor BM, Alkalife IS, Dute RR (2007) A human case of otoacariasis involving a histiostomatid mite (Acari: Histiostomatidae). *The American Journal of Tropical Medicine and Hygiene* 76: 967–971. <https://doi.org/10.4269/ajtmh.2007.76.967>
- Al-Bajalan MMM, Al-Jaf SMA, Niranji SS, Abdulkareem DR, Al-Kayali KK, Kato H (2018) An outbreak of *Leishmania major* from an endemic to a non-endemic region posed a public health threat in Iraq from 2014–2017: Epidemiological, molecular and phylogenetic studies. *PLoS Neglected Tropical Diseases* 12: e0006255. <https://doi.org/10.1371/journal.pntd.0006255>

- Al-Duboon A-H, Disher MA (2018) First Incidence of the Ciliophoran Freshwater Fish Pathogen *Trichophrya piscium* Bütschli, 1899 as a Human Pathogen from Basrah, Iraq. *Biological and Applied Environmental Research* 2: 154–161.
- Ale A, Victor B, Praet N, Gabriel S, Speybroeck N, Dorny P, Devleeschauwer B (2014) Epidemiology and genetic diversity of *Taenia asiatica*: a systematic review. *Parasites & Vectors* 7: e45. <https://doi.org/10.1186/1756-3305-7-45>
- Ali-Khan FE, Ali-Khan Z (1977) *Paragordius varius* (Leidy) (Nematomorpha) infection in man: a case report from Quebec (Canada). *Journal of Parasitology* 63: 174–176. <https://doi.org/10.2307/3280141>
- Ali IK (2015) Intestinal amebae. *Clinics in Laboratory Medicine* 35: 393–422. <https://doi.org/10.1016/j.cll.2015.02.009>
- Almallah Z (1968) Internal hirudiniasis in man with *Limnatis nilotica*, in Iraq. *Journal of Parasitology* 54: 637–638. <https://doi.org/10.2307/3277105>
- Almeria S, Cinar HN, Dubey JP (2019) *Cyclospora cayetanensis* and Cyclosporiasis: An Update. *Microorganisms* 7(9): e317. <https://doi.org/10.3390/microorganisms7090317>
- Aloto D, Eticha E (2018) Leeches: A review on their pathogenic and beneficial effects. *Journal of Veterinary Science & Technology* 9: 1000511. <https://doi.org/10.4172/2157-7579.1000511>
- Amanatfard E, Youssefi MR, Barimani A (2014) Human Dermatitis Caused by *Ophionyssus natricis*, a Snake Mite. *Iranian Journal of Parasitology* 9: 594–596.
- Amin O, Margolis A (1998) Redescription of *Bolbosoma capitatum* (Acanthocephala: Polymorphidae) from False Killer Whale off Vancouver Island, with Taxonomic Reconsideration of the Species and Synonymy of *B. physeteris*. *Journal of the Helminthological Society of Washington* 65: 179–188.
- Ammerman NC, Swanson KI, Anderson JM, Schwartz TR, Seaberg EC, Glass GE, Norris DE (2004) Spotted-fever group *Rickettsia* in *Dermacentor variabilis*, Maryland. *Emerging Infectious Diseases* 10: 1478–1481. <https://doi.org/10.3201/eid1008.030882>
- Anderson AL, Chaney E (2009) Pubic lice (*Phthirus pubis*): history, biology and treatment vs. knowledge and beliefs of US college students. *International Journal of Environmental Research and Public Health* 6: 592–600. <https://doi.org/10.3390/ijerph6020592>
- Anderson N (1988) Aspects of the biology of *Ostertagia ostertagi* in relation to the genesis of ostertagiasis. *Veterinary Parasitology* 27: 13–21. [https://doi.org/10.1016/0304-4017\(88\)90057-X](https://doi.org/10.1016/0304-4017(88)90057-X)
- Anderson RC (2000) *Nematode Parasites of Vertebrates: Their Development and Transmission*, 2nd edn. CABI Publishing, New York, 672 pp. <https://doi.org/10.1079/9780851994215.0001>
- Andreassen J, Ito A, Ito M, Nakao M, Nakaya K (2004) *Hymenolepis microstoma*: direct life cycle in immunodeficient mice. *Journal of helminthology* 78: 1–5. <https://doi.org/10.1079/JOH2003207>
- Angelou A, Tsakou K, Mpranditsas K, Sioutas G, Moores DA, Papadopoulos E (2020) Giant kidney worm: novel report of *Diocotophyma renale* in the kidney of a dog in Greece. *Helminthologia* 57: 43–48. <https://doi.org/10.2478/helm-2020-0008>
- Apanaskevich D, Apanaskevich M (2016) Description of two new species of *Dermacentor* Koch, 1844 (Acari: Ixodidae) from Oriental Asia. *Systematic Parasitology* 93: 159–171. <https://doi.org/10.1007/s11230-015-9614-8>

- Apanaskevich D, Horak I, Camicas J (2007) Redescription of *Haemaphysalis (Rhipistoma) elliptica* (Koch, 1844), an old taxon of the *Haemaphysalis* [*Rhipistoma leachi* group from East and southern Africa, and of *Haemaphysalis (Rhipistoma) leachi* (Audouin, 1826) (Ixodida, Ixodidae). The Onderstepoort journal of veterinary research 74: 181–208. <https://doi.org/10.4102/ojvr.v74i3.122>
- Apanaskevich DA, Horak IG (2008) The genus *Hyalomma*. VI. Systematics of *H. (Euhyalomma) truncatum* and the closely related species, *H. (E.) albiparmatum* and *H. (E.) nitidum* (Acari: Ixodidae). Experimental and Applied Acarology 44: 115–136. <https://doi.org/10.1007/s10493-008-9136-z>
- Apanaskevich MA, Apanaskevich DA (2015) Description of New *Dermacentor* (Acari: Ixodidae) Species from Malaysia and Vietnam. Journal of Medical Entomology 52: 156–162. <https://doi.org/10.1093/jme/tjv001>
- Arai HP (1980) Biology of the tapeworm *Hymenolepis diminuta*. Academic Press, Cambridge, 746 pp.
- Arfuso F, Gaglio G, Ferrara MC, Abbate F, Giannetto S, Brianti E (2019) First record of infestation by nasal leeches, *Limnatis nilotica* (Hirudinida, Praobdellidae), from cattle in Italy. Journal of Veterinary Medical Science 81: 1419–1423. <https://doi.org/10.1292/jvms.19-0247>
- Ariyaratne S, Dilrukshi PRMP, Amarasinghe PH, Rajakaruna RS (2011) Intra-aural ecdysis of *Dermacentor auratus* Supino, 1897, in a human host. Ceylon Medical Journal 56: 133–134. <https://doi.org/10.4038/cmj.v56i3.3612>
- Arizono N, Kuramochi T, Kagei N (2012) Molecular and histological identification of the acanthocephalan *Bolbosoma cf. capitatum* from the human small intestine. Parasitology International 61: 715–718. <https://doi.org/10.1016/j.parint.2012.05.011>
- Arlian LG, Morgan MS (2017) A review of *Sarcoptes scabiei*: past, present and future. Parasites & Vectors 10: e297. <https://doi.org/10.1186/s13071-017-2234-1>
- Ash LR, Orihel TC (2007) Atlas of human parasitology. ASCP Press, Chicago, 540 pp.
- Ashford R, Crewe W (2003) Parasites of Homo sapiens: An Annotated Checklist of the Protozoa, Helminths and Arthropods for which we are Home, 2nd edn. Taylor & Francis, London and New York, 150 pp. <https://doi.org/10.5962/bhl.title.61568>
- Ashrafi K, Sharifdini M, Heidari Z, Rahmati B, Kia EB (2020) Zoonotic transmission of *Teladorsagia circumcincta* and *Trichostrongylus* species in Guilan province, northern Iran: molecular and morphological characterizations. BMC Infectious Diseases 20: e28. <https://doi.org/10.1186/s12879-020-4762-0>
- Attia RA, Tolba ME, Yones DA, Bakir HY, Eldeek HE, Kamel S (2012) *Capillaria philippinensis* in Upper Egypt: has it become endemic? American Journal of Tropical Medicine & Hygiene 86: 126–133. <https://doi.org/10.4269/ajtmh.2012.11-0321>
- Attias M, Teixeira DE, Benchimol M, Vommaro RC, Crepaldi PH, De Souza W (2020) The life-cycle of *Toxoplasma gondii* reviewed using animations. Parasites & Vectors 13: e588. <https://doi.org/10.1186/s13071-020-04445-z>
- Audicana MT, Kennedy MW (2008) *Anisakis simplex*: from obscure infectious worm to inducer of immune hypersensitivity. Clinical Microbiology Reviews 21: 360–379. [table of contents] <https://doi.org/10.1128/CMR.00012-07>

- Audy JRN, Nadchatram M, Boo-Liat L (1960) Malaysian parasites. XLIX. Host distribution of Malayan ticks (Ixodoidea). Studies from the Institute for Medical Research, Federation of Malaya 29: 225–246.
- Aznar FJ, Perez-Ponce de Leon G, Raga JA (2006) Status of *Corynosoma* (Acanthocephala: Polymorphidae) based on anatomical, ecological, and phylogenetic evidence, with the erection of *Pseudocorynosoma* n. gen. Journal of Parasitology 92: 548–564. <https://doi.org/10.1645/GE-715R.1>
- Baer JG, Sandars DF (1956) The first record of *Railletina* (*Railletina*) *celebensis* (Janicki, 1902), (Cestoda) in man from Australia, with a critical survey of previous cases. Journal of Helminthology 30: 173–182. <https://doi.org/10.1017/S0022149X00033137>
- Bain O, Otranto D, Diniz DG, dos Santos JN, de Oliveira NP, Frota de Almeida IN, Frota de Almeida RN, Frota de Almeida LN, Dantas-Torres F, de Almeida Sobrinho EF (2011) Human intraocular filariasis caused by *Pelecitus* sp. nematode, Brazil. Emerging Infectious Diseases 17: 867–869. <https://doi.org/10.3201/eid1705.101309>
- Barkati S, Gottstein B, Müller N, Sheitoyan-Pesant C, Metrakos P, Chen T, Garceau R, Libman MD, Ndao M, Yansouni CP (2018) First Human Case of Metacestode Infection Caused by *Versteria* sp. in a Kidney Transplant Recipient. Clinical Infectious Diseases 68: 680–683. <https://doi.org/10.1093/cid/ciy602>
- Barker HS, Snyder JsW, Hicks AB, Yanoviak SP, Southern P, Dhakal BK, Ghimire GR, Couturier MR (2014) First Case Reports of *Ignatzschineria* (*Schineria*) *indica* Associated with Myiasis. Journal of Clinical Microbiology 52: 4432–4434. <https://doi.org/10.1128/JCM.02183-14>
- Barratt JLN, Lane M, Talundzic E, Richins T, Robertson G, Formenti F, Pritt B, Verocai G, Nascimento de Souza J, Mato Soares N, Traub R, Buonfrate D, Bradbury RS (2019) A global genotyping survey of *Strongyloides stercoralis* and *Strongyloides fuelleborni* using deep amplicon sequencing. PLoS Neglected Tropical Diseases 13: e0007609. <https://doi.org/10.1371/journal.pntd.0007609>
- Barton D, Pichelin S (1999) *Acanthocephalus bufonis* (Acanthocephala) from *Bufo marinus* (Bufonidae: Amphibia) in Hawaii. Parasite (Paris, France) 6: 269–272. <https://doi.org/10.1051/parasite/1999063269>
- Batu N, Wang Y, Liu Z, Huang T, Bao W, He H, Geri L (2020) Molecular epidemiology of *Rickettsia* sp. and *Coxiella burnetii* collected from *Hyalomma asiaticum* in Bactrian camels (*Camelus bactrianus*) in inner Mongolia of China. Ticks and Tick-borne Diseases 11: 101548. <https://doi.org/10.1016/j.ttbdis.2020.101548>
- Beard CB, Occi J, Bonilla DL, Egizi AM, Fonseca DM, Mertins JW, Backenson BP, Bajwa WI, Barbarin AM, Bertone MA, Brown J, Connally NP, Connell ND, Eisen RJ, Falco RC, James AM, Krell RK, Lahmers K, Lewis N, Little SE, Neault M, Pérez de León AA, Randall AR, Ruder MG, Saleh MN, Schappach BL, Schroeder BA, Seraphin LL, Wehtje M, Wormser GP, Yabsley MJ, Halperin W (2018) Multistate Infestation with the Exotic Disease-Vector Tick *Haemaphysalis longicornis* – United States, August 2017–September 2018. MMWR Morbidity and Mortality Weekly Report 67: 1310–1313. <https://doi.org/10.15585/mmwr.mm6747a3>

- Beaver PC, Duron RA, Little MD (1977) Trematode eggs in the peritoneal cavity of man in Honduras. *American Journal of Tropical Medicine and Hygiene* 26: 684–687. <https://doi.org/10.4269/ajtmh.1977.26.684>
- Beck W, Pfister K (2006) Humanpathogene Milben als Zoonoseerreger. *Wiener klinische Wochenschrift* 118: 27–32. <https://doi.org/10.1007/s00508-006-0678-y>
- Bedford GAH (1927) Check-List and Host-List of the External Parasites found on South African Mammalia, Aves, and Reptilia. In: 11th and 12th Reports of the Director of Veterinary Education and Research, Department of Agriculture, Union of South Africa, Praetoria, 705–817.
- Behr MA, Gyorkos TW, Kokoskin E, Ward BJ, MacLean JD (1998) North American liver fluke (*Metorchis conjunctus*) in a Canadian aboriginal population: a submerging human pathogen? *Canadian Journal of Public Health* 89: 258–259. <https://doi.org/10.1007/BF03403931>
- Bellanger AP, Cabaret O, Costa JM, Foulet F, Bretagne S, Botterel F (2008) Two unusual occurrences of trichomoniasis: rapid species identification by PCR. *Journal of Clinical Microbiology* 46: 3159–3161. <https://doi.org/10.1128/JCM.00322-08>
- Bermudez CSE, Troyo A (2018) A review of the genus *Rickettsia* in Central America. *Research and Reports in Tropical Medicine* 9: 103–112. <https://doi.org/10.2147/RRTM.S160951>
- Bern C, Kjos S, Yabsley MJ, Montgomery SP (2011) *Trypanosoma cruzi* and Chagas Disease in the United States. *Clinical Microbiology Reviews* 24: 655–681. <https://doi.org/10.1128/CMR.00005-11>
- Berzunza-Cruz M, Rodriguez-Moreno A, Gutierrez-Granados G, Gonzalez-Salazar C, Stephens CR, Hidalgo-Mihart M, Marina CF, Rebollar-Tellez EA, Bailon-Martinez D, Balcells CD, Ibarra-Cerdena CN, Sanchez-Cordero V, Becker I (2015) *Leishmania (L.) mexicana* infected bats in Mexico: novel potential reservoirs. *PLoS Neglected Tropical Diseases* 9: e0003438. <https://doi.org/10.1371/journal.pntd.0003438>
- Beser J, Bujila I, Wittesjö B, Lebbad M (2020) From mice to men: Three cases of human infection with *Cryptosporidium ditrichi*. *Infection, Genetics and Evolution* 78: 104120. <https://doi.org/10.1016/j.meegid.2019.104120>
- Betson M, Nejsum P, Bendall RP, Deb RM, Stothard JR (2014) Molecular epidemiology of ascariasis: a global perspective on the transmission dynamics of *Ascaris* in people and pigs. *Journal of Infectious Diseases* 210: 932–941. <https://doi.org/10.1093/infdis/jiu193>
- Betson M, Søre MJ, Nejsum P (2015) Human Trichuriasis: Whipworm Genetics, Phylogeny, Transmission and Future Research Directions. *Current Tropical Medicine Reports* 2: 209–217. <https://doi.org/10.1007/s40475-015-0062-y>
- Bhargava D, Victor R (1911) Carabid beetle invasion of the ear in Oman. *Wilderness and Environmental Medicine* 10: 157–160. [https://doi.org/10.1580/1080-6032\(1999\)010\[0157:CBIOTE\]2.3.CO;2](https://doi.org/10.1580/1080-6032(1999)010[0157:CBIOTE]2.3.CO;2)
- Bitam I, Dittmar K, Parola P, Whiting MF, Raoult D (2010) Fleas and flea-borne diseases. *International Journal of Infectious Diseases* 14: E667–E676. <https://doi.org/10.1016/j.ijid.2009.11.011>
- Blair D (2019) Paragonimiasis. In: Toledo R, Fried B (Eds) *Digenetic Trematodes*. *Advances in Experimental Medicine and Biology*, vol. 1154. Springer, Cham, 105–138. https://doi.org/10.1007/978-3-030-18616-6_5

- Blair D, Xu Z-B, Agatsuma T (1999) Paragonimiasis and the Genus *Paragonimus*. In: Baker JR, Muller R, Rollinson D (Eds) *Advances in Parasitology*. Academic Press, 113–222. [https://doi.org/10.1016/S0065-308X\(08\)60149-9](https://doi.org/10.1016/S0065-308X(08)60149-9)
- Boinas F, Ribeiro R, Madeira S, Palma M, de Carvalho IL, Nuncio S, Wilson AJ (2014) The medical and veterinary role of *Ornithodoros erraticus* complex ticks (Acari: Ixodida) on the Iberian Peninsula. *Journal of Vector Ecology* 39: 238–248. <https://doi.org/10.1111/jvec.12098>
- Bonilla DL, Durden LA, Eremeeva ME, Dasch GA (2013) The biology and taxonomy of head and body lice – implications for louse-borne disease prevention. *PLoS Pathogens* 9: e1003724. <https://doi.org/10.1371/journal.ppat.1003724>
- Bonner M, Fresno M, Girones N, Guillen N, Santi-Rocca J (2018) Reassessing the Role of *Entamoeba gingivalis* in Periodontitis. *Frontiers in Cellular and Infection Microbiology* 8: e379. <https://doi.org/10.3389/fcimb.2018.00379>
- Booton GC, Visvesvara GS, Byers TJ, Kelly DJ, Fuerst PA (2005) Identification and distribution of *Acanthamoeba* species genotypes associated with nonkeratitis infections. *Journal of Clinical Microbiology* 43: 1689–1693. <https://doi.org/10.1128/JCM.43.4.1689-1693.2005>
- Bousquet Y (1990) Beetles associated with stored products in Canada: an identification guide. Canadian Government Pub Centre, Ottawa, 214 pp.
- Bouzig M, Hunter PR, Chalmers RM, Tyler KM (2013) *Cryptosporidium* Pathogenicity and Virulence. *Clinical Microbiology Reviews* 26: 115–134. <https://doi.org/10.1128/CMR.00076-12>
- Braae UC, Thomas LF, Robertson LJ, Dermauw V, Dorny P, Willingham AL, Saratsis A, Devleeschauwer B (2018) Epidemiology of *Taenia saginata* taeniosis/cysticercosis: a systematic review of the distribution in the Americas. *Parasites & Vectors* 11: e518. <https://doi.org/10.1186/s13071-018-3079-y>
- Bradbury R (2006) An imported case of trichostrongylid infection in Tasmania and a review of human trichostrongylidiosis. *Annals of the Australasian College of Tropical Medicine* 7: 25–28.
- Bradbury RS (2014) Free-living amoebae recovered from human stool samples in *Strongyloides* agar culture. *Journal of Clinical Microbiology* 52: 699–700. <https://doi.org/10.1128/JCM.02738-13>
- Bradbury RS (2019) *Ternidens deminutus* Revisited: A Review of Human Infections with the False Hookworm. *Tropical Medicine and Infectious Disease* 4(3): e106. <https://doi.org/10.3390/tropicalmed4030106>
- Bradbury RS, Breen KV, Bonura EM, Hoyt JW, Bishop HS (2018) Case Report: Conjunctival Infestation with *Thelazia gulosa*: A Novel Agent of Human Thelaziasis in the United States. *American Journal of Tropical Medicine & Hygiene* 98: 1171–1174. <https://doi.org/10.4269/ajtmh.17-0870>
- Bradbury RS, Gustafson DT, Sapp SGH, Fox M, de Almeida M, Boyce M, Iwen P, Herrera V, Ndubuisi M, Bishop HS (2019a) A Second Case of Human Conjunctival Infestation With *Thelazia gulosa* and a Review of *T. gulosa* in North America. *Clinical Infectious Diseases* 70(3): 518–520. <https://doi.org/10.1093/cid/ciz469>
- Bradbury RS, Roy S, Ali IK, Morrison JR, Waldner D, Hebbeln K, Aldous W, Jepson R, Delavan HR, Ndubuisi M, Bishop HS (2019b) Case Report: Cervicovaginal Co-Colonization with *Entamoeba gingivalis* and *Entamoeba polecki* in Association with an Intrauterine Device. *American Journal of Tropical Medicine and Hygiene* 100: 311–313. <https://doi.org/10.4269/ajtmh.18-0522>

- Brant SV, Jouet D, Ferte H, Loker ES (2013) *Anserobilharzia* gen. n. (Digenea, Schistosomatidae) and redescription of *A. brantae* (Farr & Blankemeyer, 1956) comb. n. (syn. *Trichobilharzia brantae*), a parasite of geese (Anseriformes). *Zootaxa* 3670: 193–206. <https://doi.org/10.11646/zootaxa.3670.2.5>
- Brasil P, Zalis MG, de Pina-Costa A, Siqueira AM, Junior CB, Silva S, Areas ALL, Pelajo-Machado M, de Alvarenga DAM, da Silva Santelli ACF, Albuquerque HG, Cravo P, Santos de Abreu FV, Peterka CL, Zanini GM, Suarez Mutis MC, Pissinatti A, Lourenco-de-Oliveira R, de Brito CFA, de Fatima Ferreira-da-Cruz M, Culleton R, Daniel-Ribeiro CT (2017) Outbreak of human malaria caused by *Plasmodium simium* in the Atlantic Forest in Rio de Janeiro: a molecular epidemiological investigation. *The Lancet Global Health* 5: e1038–e1046. [https://doi.org/10.1016/S2214-109X\(17\)30333-9](https://doi.org/10.1016/S2214-109X(17)30333-9)
- Braverman I, Dano I, Saah D, Gapany B (1994) Aural myiasis caused by flesh fly larva, *Sarcophaga haemorrhoidalis*. *Journal of Otolaryngology* 23: 204–205.
- Bray RS, Ashford RW, Bray MA (1973) The parasite causing cutaneous leishmaniasis in Ethiopia. *Transactions of The Royal Society of Tropical Medicine and Hygiene* 67: 345–348. [https://doi.org/10.1016/0035-9203\(73\)90111-9](https://doi.org/10.1016/0035-9203(73)90111-9)
- Broce AB, Zurek L, Kalisch JA, Brown R, Keith DL, Gordon D, Goedeke J, Welbourn C, Moser J, Ochoa R, Azziz-Baumgartner E, Yip F, Weber J (2006) *Pyemotes herfsi* (Acari: Pyemotidae), a Mite New to North America as the Cause of Bite Outbreaks. *Journal of Medical Entomology* 43: 610–613. <https://doi.org/10.1093/jmedent/43.3.610>
- Brooker S, Bethony J, Hotez PJ (2004) Human hookworm infection in the 21st century. *Advances in Parasitology* 58: 197–288. [https://doi.org/10.1016/S0065-308X\(04\)58004-1](https://doi.org/10.1016/S0065-308X(04)58004-1)
- Brown S, De Jonckheere JF (1999) A reevaluation of the amoeba genus *Vahlkampfia* based on SSUrDNA sequences. *European Journal of Protistology* 35: 49–54. [https://doi.org/10.1016/S0932-4739\(99\)80021-2](https://doi.org/10.1016/S0932-4739(99)80021-2)
- Brun R, Blum J, Chappuis F, Burri C (2010) Human African trypanosomiasis. *Lancet* 375: 148–159. [https://doi.org/10.1016/S0140-6736\(09\)60829-1](https://doi.org/10.1016/S0140-6736(09)60829-1)
- Brunet J, Benoild A, Kremer S, Dalvit C, Lefebvre N, Hansmann Y, Chenard MP, Mathieu B, Grimm F, Deplazes P, Pfaff AW, Abou-Bacar A, Marescaux C, Candolfi E (2015) First Case of Human Cerebral *Taenia martis* Cysticercosis. *Journal of Clinical Microbiology* 53: 2756–2759. <https://doi.org/10.1128/JCM.01033-15>
- Burgess MJ, Rosenbaum ER, Pritt BS, Haselow DT, Ferren KM, Alzghoul BN, Rico JC, Sloan LM, Ramanan P, Purushothaman R, Bradsher RW (2017) Possible Transfusion-Transmitted *Babesia divergens*-like/MO-1 Infection in an Arkansas Patient. *Clinical Infectious Diseases* 64: 1622–1625. <https://doi.org/10.1093/cid/cix216>
- Burr Jr WE, Brown MF, Eberhard ML (1998) Zoonotic *Onchocerca* (Nematoda:Filarioidea) in the cornea of a Colorado resident. *Ophthalmology* 105: 1494–1497. [https://doi.org/10.1016/S0161-6420\(98\)98035-6](https://doi.org/10.1016/S0161-6420(98)98035-6)
- Bush SE, Duszynski DW, Nickol BB (2009) Acanthocephala from amphibians in China with the description of a new species of *Pseudoacanthocephalus* (Echinorhynchida). *Journal of Parasitology* 95(6): 1440–1445. <https://doi.org/10.1645/GE-2101.1>
- Cabello RR, Ruiz AC, Feregrino RR, Romero LC, Feregrino RR, Zavala JT (2011) *Dipylidium caninum* infection. *BMJ Case Reports* 2011. <https://doi.org/10.1136/bcr.07.2011.4510>

- Caccio S, Pinter E, Fantini R, Mezzaroma I, Pozio E (2002) Human infection with *Cryptosporidium felis*: case report and literature review. *Emerging Infectious Diseases* 8: 85–86. <https://doi.org/10.3201/eid0801.010269>
- Cafiero MA, Barlaam A, Camarda A, Radeski M, Mul M, Sparagano O, Giangaspero A (2019) *Dermanysus gallinae* attacks humans. Mind the gap! *Avian Pathology* 48: S22–S34. <https://doi.org/10.1080/03079457.2019.1633010>
- Calvopina M, Caballero H, Morita T, Korenaga M (2016) Human Pulmonary Infection by the Zoonotic *Metastrongylus salmi* Nematode. The First Reported Case in the Americas. *The American Journal of Tropical Medicine and Hygiene* 95: 871–873. <https://doi.org/10.4269/ajtmh.16-0247>
- Calvopiña M, Cevallos W, Atherton R, Saunders M, Small A, Kumazawa H, Sugiyama H (2015) High prevalence of the liver fluke *Amphimerus* sp. in domestic cats and dogs in an area for human amphimeriasis in Ecuador. *PLoS Neglected Tropical Diseases* 9: e0003526. <https://doi.org/10.1371/journal.pntd.0003526>
- Calvopiña M, Cevallos W, Kumazawa H, Eisenberg J (2011) High prevalence of human liver infection by *Amphimerus* spp. flukes, Ecuador. *Emerging Infectious Diseases* 17: 2331–2334. <https://doi.org/10.3201/eid1712.110373>
- Cama VA, Mathison BA (2015) Infections by Intestinal Coccidia and *Giardia duodenalis*. *Clinics in Laboratory Medicine* 35: 423–444. <https://doi.org/10.1016/j.cl.2015.02.010>
- Cambra-Pellejà M, Gandasegui J, Balaña-Fouce R, Muñoz J, Martínez-Valladares M (2020) Zoonotic Implications of *Onchocerca* Species on Human Health. *Pathogens* 9: e761. <https://doi.org/10.3390/pathogens9090761>
- Campos DMB, Barbosa AP, Oliveira JA, Tavares GG, Cravo PVL, Ostermayer AL (2017) Human lagochilascariasis—A rare helminthic disease. *PLoS Neglected Tropical Diseases* 11: e0005510. <https://doi.org/10.1371/journal.pntd.0005510>
- Cancrini G, Boemi G, Iori A, Corselli A (1982) [Human infestations by *Trichostrongylus axei*, *T. capricola* and *T. vitrinus*: 1st report in Italy]. *Parassitologia* 24: 145–149.
- Cantanhêde LM, Mattos CB, de Souza Ronconi C, Filgueira CPB, da Silva Júnior CF, Limeira C, de Jesus Silva HP, Ferreira GEM, Porrozzi R, Ferreira RGM, Cupolillo E (2019) First report of *Leishmania (Viannia) lindenbergi* causing tegumentary leishmaniasis in the Brazilian western Amazon region. *Parasite* 26: e30. <https://doi.org/10.1051/parasite/2019030>
- Cantey PT, Weeks J, Edwards M, Rao S, Ostovar GA, Dehority W, Alzona M, Swoboda S, Christiaens B, Ballan W, Hartley J, Terranella A, Weatherhead J, Dunn JJ, Marx DP, Hicks MJ, Rauch RA, Smith C, Dishop MK, Handler MH, Dudley RW, Chundu K, Hobohm D, Feiz-Erfan I, Hakes J, Berry RS, Stepensaski S, Greenfield B, Shroeder L, Bishop H, de Almeida M, Mathison B, Eberhard M (2016) The Emergence of Zoonotic *Onchocerca lupi* Infection in the United States – A Case-Series. *Clinical Infectious Diseases* 62: 778–783. <https://doi.org/10.1093/cid/civ983>
- Cao W-C, Zhan L, De Vlas SJ, Wen B-H, Yang H, Richardus JH, Habbema JDF (2008) Molecular Detection of Spotted Fever Group *Rickettsia* in *Dermacentor silvarum* from a Forest Area of Northeastern China. *Journal of Medical Entomology* 45: 741–744. <https://doi.org/10.1093/jmedent/45.4.741>

- Carlson JC, Fox MS (2009) A sticktight flea removed from the cheek of a two-year-old boy from Los Angeles. *Dermatology Online Journal* 15(1): 4. <https://doi.org/10.5070/D36VB8P1B1>
- Carroll KC, Pfaller MA, Landry ML, McAdam AJ, Patel R, Richter SS, Warnock DW (2019) *Manual of Clinical Microbiology*. ASM Press, Washington DC, 2 volumes. <https://doi.org/10.1128/9781555819842>
- Castro EA, Thomaz-Soccol V, Augur C, Luz E (2007) *Leishmania (Viannia) braziliensis*: epidemiology of canine cutaneous leishmaniasis in the State of Parana (Brazil). *Experimental Parasitology* 117: 13–21. <https://doi.org/10.1016/j.exppara.2007.03.003>
- Cembranelli SB, Souto FO, Ferreira-Paim K, Richinho TT, Nunes PL, Nascentes GA, Ferreira TB, Correia D, Lages-Silva E (2013) First evidence of genetic intraspecific variability and occurrence of *Entamoeba gingivalis* in HIV(+)/AIDS. *PLoS ONE* 8: e82864. <https://doi.org/10.1371/journal.pone.0082864>
- Cengiz ZT, Yilmaz H, Dulger AC, Cicek M (2010) Human infection with *Dicrocoelium dendriticum* in Turkey. *Annals of Saudi Medicine* 30: 159–161. <https://doi.org/10.4103/0256-4947.60525>
- Chabaud AG, Lanz P (1951) Pseudo-parasitism of Man by *Agamomermis* sp. *Annales de Parasitologie Humaine et Comparée* 26: 376–378. <https://doi.org/10.1051/parasite/1951264376>
- Chai JY, Jung BK (2020) Foodborne intestinal flukes: A brief review of epidemiology and geographical distribution. *Acta Tropica* 201: 105210. <https://doi.org/10.1016/j.actatropica.2019.105210>
- Chai JY, Sohn WM, Yong TS, Eom KS, Min DY, Hoang EH, Phammasack B, Insiengmay B, Rim HJ (2012) Echinostome flukes recovered from humans in Khammouane Province, Lao PDR. *The Korean Journal of Parasitology* 50: 269–272. <https://doi.org/10.3347/kjp.2012.50.3.269>
- Chaiwong T, Tem-Eiam N, Limpavithayakul M, Boongunha N, Poolphol W, Sukontason KL (2014) Aural myiasis caused by *Parasarcophaga (Liosarcophaga) dux* (Thomson) in Thailand. *Tropical Biomedicine* 31: 496–498.
- Chako CZ, Tyler JW, Schultz LG, Chiguma L, Beerntsen BT (2010) Cryptosporidiosis in people: it's not just about the cows. *Journal of Veterinary Internal Medicine* 24: 37–43. <https://doi.org/10.1111/j.1939-1676.2009.0431.x>
- Chakrabarti A (1986) Human notoedric scabies from contact with cats infested with *Notoedres cati*. *International Journal of Dermatology* 25: 646–648. <https://doi.org/10.1111/j.1365-4362.1986.tb04527.x>
- Champour M, Chinikar S, Mohammadi G, Razmi G, Shah-Hosseini N, Khakifrouz S, Mostafavi E, Jalali T (2016) Molecular epidemiology of Crimean-Congo hemorrhagic fever virus detected from ticks of one humped camels (*Camelus dromedarius*) population in northeastern Iran. *Journal of Parasitic Diseases* 40: 110–115. <https://doi.org/10.1007/s12639-014-0458-y>
- Chancey RJ, Sapp SGH, Fox M, Bishop HS, Ndubuisi M, de Almeida M, Montgomery SP, Congeni B (2021) Patent *Macracanthorhynchus ingens* infection in a 17-month-old child, Ohio. *Open Forum Infectious Diseases* 8(2): ofaa641. <https://doi.org/10.1093/ofid/ofaa641>
- Chandler AC (1925) New records of *Bertiella satyri* (Cestoda) in man and apes. *Parasitology* 17: 421–425. <https://doi.org/10.1017/S0031182000004844>

- Chandler AC (2009) *Diploscapter coronata* as a facultative parasite of man, with a general review of vertebrate parasitism by rhabditoid worms. *Parasitology* 30: 44–55. <https://doi.org/10.1017/S0031182000010817>
- Chandler DJ, Fuller LC (2019) A review of scabies: An infestation more than skin deep. *Dermatology* 235: 79–90. <https://doi.org/10.1159/000495290>
- Chappell CL, Okhuysen PC, Langer-Curry RC, Lupo PJ, Widmer G, Tzipori S (2015) *Cryptosporidium muris*: infectivity and illness in healthy adult volunteers. *American Journal of Tropical Medicine and Hygiene* 92: 50–55. <https://doi.org/10.4269/ajtmh.14-0525>
- Charruau P, Ocegüera-Figueroa A, Cedeño-Vázquez JR, Pérez-Rivera SD (2020) Record of *Haementeria acuecuyetzin* (Ocegüera-Figueroa, 2008) in Morelet's Crocodiles from Quintana Roo, Mexico. *Comparative Parasitology* 87: 89–92[, 84]. <https://doi.org/10.1654/1525-2647-87.1.89>
- Chavatte J-M, Karadjian G, Landau I (2018) Half a century after its discovery, new insights on *Anthemiosoma garnhami* (Sporozoa, Piroplasmida): morphology, molecular characterisation and phylogenetic position. *Parasitology Research* 117: 3917–3925. <https://doi.org/10.1007/s00436-018-6101-6>
- Chen L, Hu S, Jiang W, Zhao J, Li N, Guo Y, Liao C, Han Q, Feng Y, Xiao L (2019) *Cryptosporidium parvum* and *Cryptosporidium hominis* subtypes in crab-eating macaques. *Parasites & Vectors* 12: e350. <https://doi.org/10.1186/s13071-019-3604-7>
- Chen SH, Liu Q, Zhang YN, Chen JX, Li H, Chen Y, Steinmann P, Zhou XN (2010) Multi-host model-based identification of *Armillifer agkistrodantis* (Pentastomida), a new zoonotic parasite from China. *PLoS Neglected Tropical Diseases* 4: e647. <https://doi.org/10.1371/journal.pntd.0000647>
- Chervy L (2002) The terminology of larval cestodes or metacestodes. *Systematic Parasitology* 52: 1–33. <https://doi.org/10.1023/A:1015086301717>
- Chigusa Y, Kawakami K, Shimada M, Kurahashi H, Matsuda H (2006) Hospital-acquired oral myiasis due to *Boettcherisca septentrionalis* (Diptera: Sarcophagidae) in Shimane Prefecture, Japan. *Medical Entomology and Zoology* 57: 139–143. <https://doi.org/10.7601/mez.57.139>
- Chigusa Y, Kirinoki M, Matsuda H (2005) Nosocomial myiasis due to *Sarcophaga peregrina* in an intensive care unit (ICU) in Japan. *Medical Entomology and Zoology* 56: 355–358. <https://doi.org/10.7601/mez.56.355>
- Chigusa Y, Shinonaga S, Koyama Y, Terano A, Kirinoki M, Matsuda H (2000) Suspected intestinal myiasis due to *Dryomyza formosa* in a Japanese schizophrenic patient with symptoms of delusional parasitosis. *Medical and Veterinary Entomology* 14: 453–457. <https://doi.org/10.1046/j.1365-2915.2000.00266.x>
- Cho JH, Kim JB, Cho CS, Huh S, Ree HI (1999) An infestation of the mite *Sancassania berlesei* (Acari: Acaridae) in the external auditory canal of a Korean man. *The Journal of Parasitology* 85: 133–134. <https://doi.org/10.2307/3285717>
- Choe S, Lee D, Park H, Oh M, Jeon HK, Eom KS (2014) *Strongyloides myopotami* (Secernentea: Strongyloididae) from the intestine of feral nutrias (*Myocastor coypus*) in Korea. *The Korean Journal of Parasitology* 52: 531–535. <https://doi.org/10.3347/kjp.2014.52.5.531>

- Choi JY, Cho BK, Lee YB, Yu DS, Jun BC, Lee IY, Kim JW (2018) An uncommon presentation of human otoacariasis by *Haemaphysalis longicornis*. *Annals of Dermatology* 30: 348–350. <https://doi.org/10.5021/ad.2018.30.3.348>
- Christensen NO, Mutani A, Frandsen F (1983) A review of the biology and transmission ecology of African bovine species of the genus *Schistosoma*. *Zeitschrift für Parasitenkunde* 69: 551–570. <https://doi.org/10.1007/BF00926667>
- Christoffersen ML, (2013) A systematic monograph of the recent Pentastomida with a compilation of their hosts. *Zoologische Mededelingen* 87: 1–206.
- Christoffersen ML, De Assis JE (2015) Pentastomida. *Revista IDE@ – SEA, Ibero Diversidad Entomológica @ccesible* 98B: 1–10.
- Chunge R, Katsivo M, Kok P, Wamwea M, Kinoti S (1986) *Schistosoma bovis* in human stools in Kenya. *Transactions of The Royal Society of Tropical Medicine and Hygiene* 80(5): 849–849. [https://doi.org/10.1016/0035-9203\(86\)90404-9](https://doi.org/10.1016/0035-9203(86)90404-9)
- Cnops L, Huyse T, Maniewski U, Soentjens P, Bottieau E, Van Esbroeck M, Clerinx J (2020) Acute schistosomiasis with a *Schistosoma mattheei* × *Schistosoma haematobium* hybrid species in a cluster of 34 travelers infected in South Africa. *Clinical Infectious Diseases* 72(10): 1693–1698. <https://doi.org/10.1093/cid/ciaa312>
- Coatney GR, Collins WE, Warren M, Contacos PG (1971) The primate malarias. U.S. National Institute of Allergy and Infectious Diseases; for sale by the Supt. of Docs., U.S. Govt. Print. Off., Washington, Bethesda, Md., 366 pp.
- Colley DG, Bustinduy AL, Secor WE, King CH (2014) Human schistosomiasis. *Lancet* 383: 2253–2264. [https://doi.org/10.1016/S0140-6736\(13\)61949-2](https://doi.org/10.1016/S0140-6736(13)61949-2)
- Collins WE, Jeffery GM (2005) *Plasmodium ovale*: parasite and disease. *Clinical Microbiology Reviews* 18: 570–581. <https://doi.org/10.1128/CMR.18.3.570-581.2005>
- Collins WE, Jeffery GM (2007) *Plasmodium malariae*: parasite and disease. *Clinical Microbiology Reviews* 20: 579–592. <https://doi.org/10.1128/CMR.00027-07>
- Condlova S, Horcickova M, Sak B, Kvetonova D, Hlaskova L, Konecny R, Stanko M, McEvoy J, Kvac M (2018) *Cryptosporidium apodemi* sp. n. and *Cryptosporidium ditrichi* sp. n. (Apicomplexa: <https://doi.org/10.1016/j.ejop.2017.12.006> Cryptosporidiidae) in *Apodemus* spp. *European Journal of Protistology* 63: 1–12.
- Connors EE, Vinetz JM, Weeks JR, Brouwer KC (2016) A global systematic review of Chagas disease prevalence among migrants. *Acta Tropica* 156: 68–78. <https://doi.org/10.1016/j.actatropica.2016.01.002>
- Conrad PA, Kjemtrup AM, Carreno RA, Thomford J, Wainwright K, Eberhard M, Quick R, Telford III SR, Herwaldt BL (2006) Description of *Babesia duncani* n.sp. (Apicomplexa: Babesiidae) from humans and its differentiation from other piroplasms. *International Journal of Parasitology* 36: 779–789. <https://doi.org/10.1016/j.ijpara.2006.03.008>
- Conraths FJ, Deplazes P (2015) *Echinococcus multilocularis*: Epidemiology, surveillance and state-of-the-art diagnostics from a veterinary public health perspective. *Veterinary Parasitology* 213: 149–161. <https://doi.org/10.1016/j.vetpar.2015.07.027>
- Contacos PG, Coatney GR, Orihel TC, Collins WE, Chin W, Jeter MH (1970) Transmission of *Plasmodium schwetzi* from the Chimpanzee to Man by Mosquito Bite. *The American Journal of Tropical Medicine and Hygiene* 19: 190–195. <https://doi.org/10.4269/ajtmh.1970.19.190>

- Cooley RA, Kohls, G. M. (1944) The genus *Amblyomma* (Ixodidae) in the United States. *Journal of Parasitology* 30: 77–111. <https://doi.org/10.2307/3272571>
- Cooley RA, Kohls GM (1945) The Argasidae of North America, Central America and Cuba. *Annals of the Entomological Society of America* 38(1): 13–13. <https://doi.org/10.1093/aesa/38.1.13>
- Cope JR, Landa J, Nethercut H, Collier SA, Glaser C, Moser M, Puttagunta R, Yoder JS, Ali IK, Roy SL (2019) The Epidemiology and Clinical Features of *Balamuthia mandrillaris* Disease in the United States, 1974–2016. *Clinical Infectious Diseases* 68: 1815–1822. <https://doi.org/10.1093/cid/ciy813>
- Coral-Almeida M, Gabriel S, Abatih EN, Praet N, Benitez W, Dorny P (2015) *Taenia solium* human cysticercosis: A systematic review of sero-epidemiological data from endemic zones around the world. *PLoS Neglected Tropical Diseases* 9: e0003919. <https://doi.org/10.1371/journal.pntd.0003919>
- Cornaglia J, Jean M, Bertrand K, Aumaitre H, Roy M, Nickel B (2016) Gnathostomiasis in Brazil: an emerging disease with a challenging diagnosis. *Journal of Travel Medicine* 24(1): taw074. <https://doi.org/10.1093/jtm/taw074>
- Cornet M, Florent M, Lefebvre A, Wertheimer C, Perez-Eid C, Bangs MJ, Bouvet A (2003) Tracheopulmonary myiasis caused by a mature third-instar *Cuterebra* larva: case report and review. *Journal of Clinical Microbiology* 41: 5810–5812. <https://doi.org/10.1128/JCM.41.12.5810-5812.2003>
- Coura JR (2015) The main sceneries of Chagas disease transmission. The vectors, blood and oral transmissions – a comprehensive review. *Memórias do Instituto Oswaldo Cruz* 110: 277–282. <https://doi.org/10.1590/0074-0276140362>
- Cowie RH (2013) Biology, systematics, life cycle, and distribution of *Angiostrongylus cantonensis*, the cause of rat lungworm disease. *Hawai'i Journal of Medicine & Public Health* 72: 6–9.
- Crocco L, Catalá S (1997) Host preferences of *Triatoma sordida*. *Annals of Tropical Medicine and Parasitology* 91: 927–930. <https://doi.org/10.1080/00034983.1997.11813220>
- Cross JH (1992) Intestinal capillariasis. *Clinical Microbiology Reviews* 5: 120–129. <https://doi.org/10.1128/CMR.5.2.120>
- Cumberlidge N, Rollinson D, Vercruyse J, Tchuem Tchuenté LA, Webster B, Clark PF (2018) *Paragonimus* and paragonimiasis in West and Central Africa: unresolved questions. *Parasitology* 145: 1748–1757. <https://doi.org/10.1017/S0031182018001439>
- Cundall DB, Whitehead SM, Hechtel FO (1986) Severe anaemia and death due to the pharyngeal leech *Myxobdella africana*. *Transactions of The Royal Society of Tropical Medicine and Hygiene* 80: 940–944. [https://doi.org/10.1016/0035-9203\(86\)90265-8](https://doi.org/10.1016/0035-9203(86)90265-8)
- Cunningham LJ, Olson PD (2010) Description of *Hymenolepis microstoma* (Nottingham strain): a classical tapeworm model for research in the genomic era. *Parasites & Vectors* 3: e123. <https://doi.org/10.1186/1756-3305-3-123>
- D'Alessandro A, Rausch RL (2008) New aspects of neotropical polycystic (*Echinococcus vogeli*) and unicyclic (*Echinococcus oligarthrus*) echinococcosis. *Clinical Microbiology Reviews* 21: 380–401. [table of contents] <https://doi.org/10.1128/CMR.00050-07>
- da Costa JC, Delgado ML, Vieira P, Afonso A, Conde B, Cross JH (2005) Syngamioniasis in tourist. *Emerging Infectious Diseases* 11: 1976–1977. <https://doi.org/10.3201/eid1112.050713>
- Dantas-Torres F (2010) Biology and ecology of the brown dog tick, *Rhipicephalus sanguineus*. *Parasites & Vectors* 3: e26. <https://doi.org/10.1186/1756-3305-3-26>

- Dantas-Torres F, Chomel BB, Otranto D (2012) Ticks and tick-borne diseases: a One Health perspective. *Trends in Parasitology* 28: 437–446. <https://doi.org/10.1016/j.pt.2012.07.003>
- Dantas-Torres F, Otranto D (2020) On the validity of “*Candidatus* *Dirofilaria hongkongensis*” and on the use of the provisional status *Candidatus* in zoological nomenclature. *Parasites & Vectors* 13: e287. <https://doi.org/10.1186/s13071-020-04158-3>
- Daron J, Boissière A, Ngoubangoye B, Boundenga L, Houze S, Arnathau C, Sidobre C, Trape J-F, Durant P, Renaud F, Fontaine M, Prugnonne F, Rougeron V (2020) Population genomic evidence of a Southeast Asian origin of *Plasmodium vivax*. *bioRxiv* 2020.04.29.067439. <https://doi.org/10.1101/2020.04.29.067439>
- Das K, Nair LV, Ghosal A, Sardar SK, Dutta S, Ganguly S (2019) Genetic characterization reveals evidence for an association between water contamination and zoonotic transmission of a *Cryptosporidium* sp. from dairy cattle in West Bengal, India. *Food and Waterborne Parasitology* 17: e00064. <https://doi.org/10.1016/j.fawpar.2019.e00064>
- Davidson MM, Williams H, Macleod JAJ (1991) Louping ill in man: A forgotten disease. *Journal of Infection* 23: 241–249. [https://doi.org/10.1016/0163-4453\(91\)92756-U](https://doi.org/10.1016/0163-4453(91)92756-U)
- de Aguiar RPS, Alves LL (2018) *Urbanorum* spp: First Report in Brazil. *American Journal of Case Reports* 19: 486–490. <https://doi.org/10.12659/AJCR.908653>
- de Sousa MA (2014) On opportunist infections by *Trypanosoma lewisi* in humans and its differential diagnosis from *T. cruzi* and *T. rangeli*. *Parasitology Research* 113: 4471–4475. <https://doi.org/10.1007/s00436-014-4132-1>
- Delshad E, Rubin AI, Almeida L, Niedt GW (2008) Cuterebra cutaneous myiasis: case report and world literature review. *International Journal of Dermatology* 47: 363–366. <https://doi.org/10.1111/j.1365-4632.2008.03532.x>
- Deplazes P, Eichenberger RM, Grimm F (2019) Wildlife-transmitted *Taenia* and *Versteria* cysticercosis and coenurosis in humans and other primates. *International Journal for Parasitology: Parasites and Wildlife* 9: 342–358. <https://doi.org/10.1016/j.ijppaw.2019.03.013>
- Dergousoff SJ, Chilton NB (2012) Association of different genetic types of *Francisella*-like organisms with the Rocky Mountain wood tick (*Dermacentor andersoni*) and the American dog tick (*Dermacentor variabilis*) in localities near their northern distributional limits. *Applied Environmental Microbiology* 78: 965–971. <https://doi.org/10.1128/AEM.05762-11>
- Desbois N, Pratlong F, Quist D, Dedet JP (2014) *Leishmania (Leishmania) martiniquensis* n. sp. (Kinetoplastida: Trypanosomatidae), description of the parasite responsible for cutaneous leishmaniasis in Martinique Island (French West Indies). *Parasite* 21: e12. <https://doi.org/10.1051/parasite/2014011>
- Despommier D (2003) Toxocariasis: clinical aspects, epidemiology, medical ecology, and molecular aspects. *Clinical Microbiology Reviews* 16: 265–272. <https://doi.org/10.1128/CMR.16.2.265-272.2003>
- Desquesnes M, Ravel S, Cuny G (2002) PCR identification of *Trypanosoma lewisi*, a common parasite of laboratory rats. *Kinetoplastid Biology and Disease* 1: 2–2. <https://doi.org/10.1186/1475-9292-1-2>
- Devkota R, Brant SV, Thapa A, Loker ES (2014) Sharing schistosomes: the elephant schistosome *Bivitellobilharzia nairi* also infects the greater one-horned rhinoceros (*Rhinoceros unicornis*) in Chitwan National Park, Nepal. *Journal of Helminthology* 88: 32–40. <https://doi.org/10.1017/S0022149X12000697>

- Di Cave D, Monno R, Bottalico P, Guerriero S, D'Amelio S, D'Orazi C, Berrilli F (2009) *Acanthamoeba* T4 and T15 genotypes associated with keratitis infections in Italy. *European Journal of Clinical Microbiology & Infectious Diseases* 28: 607–612. <https://doi.org/10.1007/s10096-008-0682-4>
- Diaz JH (2009) Mite-transmitted dermatoses and infectious diseases in returning travelers. *Journal of Travel Medicine* 17: 21–31. <https://doi.org/10.1111/j.1708-8305.2009.00352.x>
- Dietrich CF, Chaubal N, Hoerauf A, Kling K, Piontek MS, Steffgen L, Mand S, Dong Y (2019) Review of dancing parasites in lymphatic filariasis. *Ultrasound International Open* 5: E65–E74. <https://doi.org/10.1055/a-0918-3678>
- Dilrukshi PRMP, Yasawardene ADKSN, Amerasinghe PH, Amerasinghe FP (2004) Human otoparasitosis: a retrospective study from an area of Sri Lanka. *Transactions of The Royal Society of Tropical Medicine and Hygiene* 98: 489–495. <https://doi.org/10.1016/j.trstmh.2003.12.008>
- Dimasuay KGB, Lavilla OJY, Rivera WL (2013) New hosts of *Simplicimonas similis* and *Trichomitus batrachorum* identified by 18S ribosomal RNA gene sequences. *Journal of Parasitology Research* 2013: 1–5. <https://doi.org/10.1155/2013/831947>
- Dini LA, Frean JA (2005) Clinical Significance of Mites in Urine. *Journal of Clinical Microbiology* 43: 6200–6201. <https://doi.org/10.1128/JCM.43.12.6200-6201.2005>
- Disney RH (2008) Natural history of the scuttle fly, *Megaselia scalaris*. *Annual Review of Entomology* 53: 39–60. <https://doi.org/10.1146/annurev.ento.53.103106.093415>
- Domrow R, Derrick E (1964) *Ixodes holocyclus* the man-biting tick in SE Queensland. *Australian Journal of Science* 27: 234–236.
- Dorresteijn GM, Van Bronswijk JEMH (1979) *Trixacarus caviae* Fain, Howell & Hyatt 1972 (Acari: Sarcoptidae) as a cause of mange in guinea-pigs and papular urticaria in man. *Veterinary Parasitology* 5: 389–398. [https://doi.org/10.1016/0304-4017\(79\)90029-3](https://doi.org/10.1016/0304-4017(79)90029-3)
- Doss MA, Farr MM, Roach KF, Anastos G (1974) II. Hosts. Part 2. In: United States Department of Agriculture USGPO (Ed.) *Index-Catalogue of Medical and Veterinary Zoology*. Special Publication 3. Ticks and Tick-borne Diseases. United States Government Print Office, Washington DC, 489–976.
- Dua HS, Azuara-Blanco A, Hossain M, Lloyd J (1998) Non-*Acanthamoeba* amebic keratitis. *Cornea* 17: 675–677. <https://doi.org/10.1097/00003226-199811000-00018>
- Dubar M, Zaffino ML, Remen T, Thilly N, Cunat L, Machouart MC, Bisson C (2020) Protozoans in subgingival biofilm: clinical and bacterial associated factors and impact of scaling and root planing treatment. *Journal of Oral Microbiology* 12: 1693222. <https://doi.org/10.1080/20002297.2019.1693222>
- Dubey JP (2015) Foodborne and waterborne zoonotic sarcocystosis. *Food and Waterborne Parasitology* 1: 2–11. <https://doi.org/10.1016/j.fawpar.2015.09.001>
- Dubey JP, Almeria S (2019) *Cystoisospora belli* infections in humans: the past 100 years. *Parasitology* 146: 1490–1527. <https://doi.org/10.1017/S0031182019000957>
- Dubey JP, van Wilpe E, Calero-Bernal R, Verma SK, Fayer R (2015) *Sarcocystis heydorni*, n. sp. (Apicomplexa: Sarcocystidae) with cattle (*Bos taurus*) and human (*Homo sapiens*) cycle. *Parasitology Research* 114: 4143–4147. <https://doi.org/10.1007/s00436-015-4645-2>
- Duboucher C, Caby S, Dufernez F, Chabé M, Gantois N, Delgado-Viscogliosi P, Billy C, Barré E, Torabi E, Capron M, Pierce RJ, Dei-Cas E, Viscogliosi E (2006) Molecular Identifica-

- tion of *Tritrichomonas foetus*-Like Organisms as Coinfecting Agents of Human *Pneumocystis* Pneumonia. *Journal of Clinical Microbiology* 44: 1165–1168. <https://doi.org/10.1128/JCM.44.3.1165-1168.2006>
- Dung DT, Hop NT, Tho TH, Nawa Y, Doanh PN (2020) Pruritic cutaneous nematodiasis caused by avian eyeworm *Oxyspirura* larvae, Vietnam. *Emerging Infectious Diseases* 26: 786–788. <https://doi.org/10.3201/eid2604.191592>
- Dunn JJ, Columbus ST, Aldeen WE, Davis M, Carroll KC (2002) *Trichuris vulpis* recovered from a patient with chronic diarrhea and five dogs. *Journal of Clinical Microbiology* 40: 2703–2704. <https://doi.org/10.1128/JCM.40.7.2703-2704.2002>
- Eberhard ML, Hellstein JW, Lanzel EA (2014a) Zoonotic anatrachosomiasis in a mother and daughter. *Journal of Clinical Microbiology* 52: 3127–3129. <https://doi.org/10.1128/JCM.01236-14>
- Eberhard ML, Mathison B, Bishop H, Handoo NQ, Hellstein JW (2010) Zoonotic anatrachosomiasis in an Illinois resident. *American Journal of Tropical Medicine and Hygiene* 83: 342–344. <https://doi.org/10.4269/ajtmh.2010.10-0144>
- Eberhard ML, Ruiz-Tiben E (2014) Cutaneous emergence of *Eustrongylides* in two persons from South Sudan. *American Journal of Tropical Medicine and Hygiene* 90: 315–317. <https://doi.org/10.4269/ajtmh.13-0638>
- Eberhard ML, Ruiz-Tiben E, Hopkins DR, Farrell C, Toe F, Weiss A, Withers PC, Jenks MH, Thiele EA, Cotton JA, Hance Z, Holroyd N, Cama VA, Tahir MA, Mounda T (2014b) The peculiar epidemiology of dracunculiasis in Chad. *American Journal of Tropical Medicine and Hygiene* 90: 61–70. <https://doi.org/10.4269/ajtmh.13-0554>
- Eberhard ML, Thiele EA, Yembo GE, Yibi MS, Cama VA, Ruiz-Tiben E (2015) Thirty-seven human cases of sparganosis from Ethiopia and South Sudan caused by *Spirometra* spp. *American Journal of Tropical Medicine and Hygiene* 93: 350–355. <https://doi.org/10.4269/ajtmh.15-0236>
- Eberwein P, Haeupler A, Kuepper F, Wagner D, Kern WV, Muntau B, Racz P, Agostini H, Popper S (2013) Human infection with marten tapeworm. *Emerging Infectious Diseases* 19: 1152–1154. <https://doi.org/10.3201/eid1907.121114>
- Eckert J, Deplazes P (2004) Biological, epidemiological, and clinical aspects of echinococcosis, a zoonosis of increasing concern. *Clinical Microbiology Reviews* 17: 107–135. <https://doi.org/10.1128/CMR.17.1.107-135.2004>
- Eiras JC, Pavanelli GC, Takemoto RM, Nawa Y (2018) An overview of fish-borne nematodiasis among returned travelers for recent 25 years – unexpected diseases sometimes far away from the origin. *The Korean Journal of Parasitology* 56: 215–227. <https://doi.org/10.3347/kjp.2018.56.3.215>
- Eisen L (2007) Seasonal pattern of host-seeking activity by the human-biting adult life stage of *Dermacentor andersoni* (Acari: Ixodidae). *Journal of Medical Entomology* 44: 359–366.
- El-Dib NA, El Wahab WMA, Hamdy DA, Ali MI (2017) Case report of human urinary myiasis caused by *Clogmia albipunctata* (Diptera: Psychodidae) with morphological description of larva and pupa. *Journal of Arthropod Borne Diseases* 11: 533–538.
- el-Shazly AM, Morsy TA, Dawoud HA (2004) Human Monieziasis expansa: the first Egyptian parasitic zoonosis. *Journal of the Egyptian Society of Parasitology* 34: 515–518.

- Elelu N (2018) Tick-borne relapsing fever as a potential veterinary medical problem. *Veterinary Medicine and Science* 4: 271–279. <https://doi.org/10.1002/vms3.108>
- Elkins CA, Nickol BB (1983) The epizootiology of *Macracanthorhynchus ingens* in Louisiana. *Journal of Parasitology* 69: 951–956. <https://doi.org/10.2307/3281063>
- Elliott DC (1986) Tapeworm (*Moniezia expansa*) and its effect on sheep production: the evidence reviewed. *New Zealand Veterinary Journal* 34(5): 61–65. <https://doi.org/10.1080/00480169.1986.35289>
- Elliott I, Pearson I, Dahal P, Thomas NV, Roberts T, Newton PN (2019) Scrub typhus ecology: a systematic review of *Orientia* in vectors and hosts. *Parasites & Vectors* 12: e513. <https://doi.org/10.1186/s13071-019-3751-x>
- Enzien M, McKhann HI, Margulis L (1989) Ecology and life history of an amoebomastigote, *Paratetramitus jugosus*, from a microbial mat: new evidence for multiple fission. *Biology Bulletin* 177: 110–129. <https://doi.org/10.2307/1541839>
- Eom KS, Rim HJ (1993) Morphologic descriptions of *Taenia asiatica* sp. n. *The Korean Journal of Parasitology* 31: 1–6. <https://doi.org/10.3347/kjp.1993.31.1.1>
- Escobedo AA, Almirall P, Alfonso M, Cimerman S, Chacin-Bonilla L (2014) Sexual transmission of giardiasis: a neglected route of spread? *Acta Tropica* 132: 106–111. <https://doi.org/10.1016/j.actatropica.2013.12.025>
- Esser HJ, Foley JE, Bongers F, Herre EA, Miller MJ, Prins HH, Jansen PA (2016) Host body size and the diversity of tick assemblages on Neotropical vertebrates. *International Journal for Parasitology: Parasites and Wildlife* 5: 295–304. <https://doi.org/10.1016/j.ijppaw.2016.10.001>
- Esslinger JH (1962) Development of *Porocephalus crotali* (Humboldt, 1808) (Pentastomida) in experimental intermediate hosts. *Journal of Parasitology* 48: 452–456. <https://doi.org/10.2307/3275214>
- Estambale BB, Knight R, Chungue R (1992) Haematemesia and severe anaemia due to a pharyngeal leech (*Myxobdella africana*) in a Kenyan child: a case report. *Transactions of The Royal Society of Tropical Medicine and Hygiene* 86(4): 458–458. [https://doi.org/10.1016/0035-9203\(92\)90271-D](https://doi.org/10.1016/0035-9203(92)90271-D)
- Estrada-Pena A, Jongejan F (1999) Ticks feeding on humans: a review of records on human-biting Ixodoidea with special reference to pathogen transmission. *Experimental and Applied Acarology* 23: 685–715. <https://doi.org/10.1023/A:1006241108739>
- Estrada-Peña A, Sánchez N, Estrada-Sánchez A (2012) An assessment of the distribution and spread of the tick *Hyalomma marginatum* in the western Palearctic under different climate scenarios. *Vector Borne Zoonotic Diseases* 12: 758–768. <https://doi.org/10.1089/vbz.2011.0771>
- Faccini-Martínez ÁA, Botero-García CA (2016) Regarding Tick-Borne Relapsing Fever in the Americas; Some Historical Aspects of a Forgotten Disease in Colombia. *Veterinary Sciences* 3(4): e33. <https://doi.org/10.3390/vetsci3040033>
- Fadaei Tehrani M, Sharifdini M, Zahabiun F, Latifi R, Kia EB (2019) Molecular characterization of human isolates of *Strongyloides stercoralis* and *Rhabditis* spp. based on mitochondrial cytochrome c oxidase subunit 1 (cox1). *BMC Infectious Diseases* 19: e776. <https://doi.org/10.1186/s12879-019-4407-3>

- Fagundes-Silva GA, Romero GAS, Cupolillo E, Yamashita EPG, Gomes-Silva A, Guerra JAdO, Da-Cruz AM (2015) *Leishmania (Viannia) naiffi*: rare enough to be neglected? Memórias do Instituto Oswaldo Cruz 110: 797–800. <https://doi.org/10.1590/0074-02760150128>
- Fain A (1960) La pentastomose chez l’homme. Bulletin de l’Academie Royale de Médecine de Belgique 25: 516–532.
- Fain A (1961) Les pentastomides de l’Afrique Centrale. Annales Musée Royale de l’Afrique Centrale (8) Sciences Zoologiques 92: 1–115.
- Fain A, Vandepitte J (1957) A new trematode, *Poikilorchis congolensis*, n.g., n.sp., living in subcutaneous retroauricular cysts in man from the Belgian Congo. Nature 179: 740–740. <https://doi.org/10.1038/179740a0>
- Fan Q-H, Zhang Z-Q (2004) Revision of *Rhizoglyphus* Claparède (Acari: Acaridae) of Australasia and Oceania. Systematic & Applied Acarology Society, London, 374 pp.
- Farley J (1971) A review of the family Schistosomatidae: excluding the genus *Schistosoma* from mammals. Journal of helminthology 45: 289–320. <https://doi.org/10.1017/S0022149X00000572>
- Farrag HMM, Huseein EAM, Almatary AM, Othman RA (2019) Morphological and initial molecular characterization of *Clogmia albipunctatus* larvae (Diptera: Psychodidae) causing urinary myiasis in Egypt. PLoS Neglected Tropical Diseases 13: e0007887. <https://doi.org/10.1371/journal.pntd.0007887>
- Farrar J, Hotez PJ, Junghans T, Kang G, Lalloo D, White NJ (2014) Manson’s Tropical Diseases. Saunders Elsevier, Philadelphia, 1360 pp.
- Fayer R (2004) *Sarcocystis* spp. in human infections. Clin Microbiol Rev 17: 894–902. [table of contents] <https://doi.org/10.1128/CMR.17.4.894-902.2004>
- Fayer R, Esposito DH, Dubey JP (2015) Human infections with *Sarcocystis* species. Clinical Microbiology Reviews 28: 295–311. <https://doi.org/10.1128/CMR.00113-14>
- Fayer R, Santín M, Trout JM (2008) *Cryptosporidium ryanae* n. sp. (Apicomplexa: Cryptosporidiidae) in cattle (*Bos taurus*). Veterinary Parasitology 156: 191–198. <https://doi.org/10.1016/j.vetpar.2008.05.024>
- Fayer R, Trout JM, Xiao L, Morgan UM, Lai AA, Dubey JP (2001) *Cryptosporidium canis* n. sp. from domestic dogs. Journal of Parasitology 87: 1415–1422. [https://doi.org/10.1645/0022-3395\(2001\)087\[1415:Ccnsfd\]2.0.Co;2](https://doi.org/10.1645/0022-3395(2001)087[1415:Ccnsfd]2.0.Co;2)
- Feldmeier H, Heukelbach J, Ugbomoiko US, Sentongo E, Mbabazi P, von Samson-Himmelstjerna G, Krantz I (2014) Tungiasis – a neglected disease with many challenges for global public health. PLoS Neglected Tropical Diseases 8: e3133. <https://doi.org/10.1371/journal.pntd.0003133>
- Feng Y, Ortega Y, He G, Das P, Xu M, Zhang X, Fayer R, Gatei W, Cama V, Xiao L (2007) Wide geographic distribution of *Cryptosporidium bovis* and the deer-like genotype in bovines. Veterinary Parasitology 144: 1–9. <https://doi.org/10.1016/j.vetpar.2006.10.001>
- Feng Y, Yang W, Ryan U, Zhang L, Kvac M, Koudela B, Modry D, Li N, Fayer R, Xiao L (2011) Development of a multilocus sequence tool for typing *Cryptosporidium muris* and *Cryptosporidium andersoni*. Journal of Clinical Microbiology 49: 34–41. <https://doi.org/10.1128/JCM.01329-10>
- Fernandes LF, Pimenta FC, Fernandes FF (2009) First report of human myiasis in Goiás state, Brazil: frequency of different types of myiasis, their various etiological agents, and associated factors. The Journal of Parasitology 95: 32–38. <https://doi.org/10.1645/GE-1103.1>

- Ferreira L de L, Pereira MH, Guarneri AA (2015) Revisiting *Trypanosoma rangeli* transmission involving susceptible and non-susceptible hosts. PLoS ONE 10: e0140575. <https://doi.org/10.1371/journal.pone.0140575>
- Fischer P, Supali T, Maizels RM (2004) Lymphatic filariasis and *Brugia timori*: prospects for elimination. Trends in Parasitology 20: 351–355. <https://doi.org/10.1016/j.pt.2004.06.001>
- Fogden SCL, Proctor J (1985) Notes on the Feeding of Land Leeches (*Haemadipsa zeylanica* Moore and *H. picta* Moore) in Gunung Mulu National Park, Sarawak. Biotropica 17: 172–174. <https://doi.org/10.2307/2388511>
- Földvári G, Široký P, Szekeres S, Majoros G, Sprong H (2016) *Dermacentor reticulatus*: a vector on the rise. Parasites & Vectors 9: e314. <https://doi.org/10.1186/s13071-016-1599-x>
- Foxx TS, Ewing SA (1969) Morphologic features, behavior, and life history of *Cheyletiella yasguri*. American Journal of Veterinary Research 30: 269–285.
- Francesconi F, Lupi O (2012) Myiasis. Clinical Microbiology Reviews 25: 79–105. <https://doi.org/10.1128/CMR.00010-11>
- Franco JR, Simarro PP, Diarra A, Jannin JG (2014) Epidemiology of human African trypanosomiasis. Clinical Epidemiology 6: 257–275. <https://doi.org/10.2147/CLEP.S39728>
- Franzini-Junior E, Mendes MT, Anê A, da Costa TA, Silva MV, Hernandez CG, Pelli A, Sales-Campos H, Oliveira CJF (2018) Biology of *Meccus pallidipennis* (Hemiptera: Reduviidae) to Other Conditions Than That Encountered in Their Native Habitat. Journal of Arthropod Borne Diseases 12: 262–268.
- Freeman R (2011) Studies on the biology of *Taenia crassiceps* (Zeder, 1800) Rudolphi, 1810 (Cestoda). Canadian Journal of Zoology 40: 969–990. <https://doi.org/10.1139/z62-086>
- Freeman RS, Stuart PF, Cullen SJ, Ritchie AC, Mildon A, Fernandes BJ, Bonin R (1976) Fatal human infection with mesocercariae of the trematode *Alaria americana*. American Journal of Tropical Medicine and Hygiene 25: 803–807. <https://doi.org/10.4269/ajtmh.1976.25.803>
- Fujita T, Waga E, Kitaoka K, Imagawa T, Komatsu Y, Takanashi K, Anbo F, Anbo T, Katuki S, Ichihara S, Fujimori S, Yamasaki H, Morishima Y, Sugiyama H, Katahira H (2016) Human infection by acanthocephalan parasites belonging to the genus *Corynosoma* found from small bowel endoscopy. Parasitology International 65: 491–493. <https://doi.org/10.1016/j.parint.2016.07.002>
- Fukumoto S, Yazaki S, Maejima J, Kamo H, Takao Y, Tsutsumi H (1988) The first report of human infection with *Diphyllobothrium scoticum* (Rennie et Reid, 1912). Japanese Journal of Parasitology 37: 84–90.
- Galan-Puchades MT (2019) Diagnosis and treatment of human sparganosis. Lancet Infectious Diseases 19: 465–465. [https://doi.org/10.1016/S1473-3099\(19\)30166-5](https://doi.org/10.1016/S1473-3099(19)30166-5)
- Garcia-Varela M, Perez-Ponce de Leon G, de la Torre P, Cummings MP, Sarma SS, Lacleite JP (2000) Phylogenetic relationships of Acanthocephala based on analysis of 18S ribosomal RNA gene sequences. Journal of Molecular Evolution 50: 532–540. <https://doi.org/10.1007/s002390010056>
- Garcia HH, Gonzalez AE, Evans CA, Gilman RH, Cysticercosis Working Group in P (2003) *Taenia solium* cysticercosis. Lancet 362: 547–556. [https://doi.org/10.1016/S0140-6736\(03\)14117-7](https://doi.org/10.1016/S0140-6736(03)14117-7)
- Gargili A, Thangamani S, Bente D (2013) Influence of laboratory animal hosts on the life cycle of *Hyalomma marginatum* and implications for an in vivo transmission model for

- Crimean-Congo hemorrhagic fever virus. *Frontiers in Cellular and Infection Microbiology* 3: e39. <https://doi.org/10.3389/fcimb.2013.00039>
- Gautret P, Mockenhaupt FP, von Sonnenburg F, Rothe C, Libman M, Van De Winkel K, Bouteau E, Grobusch MP, Hamer DH, Esposito DH, Parola P, Schlagenhauf P, GeoSentinel Surveillance N (2015) Local and international implications of schistosomiasis acquired in Corsica, France. *Emerging Infectious Diseases* 21: 1865–1868. <https://doi.org/10.3201/eid2110.150881>
- Gavin PJ, Kazacos KR, Shulman ST (2005) Baylisascariasis. *Clinical Microbiology Reviews* 18: 703–718. <https://doi.org/10.1128/CMR.18.4.703-718.2005>
- Gelman BB, Rauf SJ, Nader R, Popov V, Borkowski J, Chaljub G, Nauta HW, Visvesvara GS (2001) Amoebic encephalitis due to *Sappinia diploidea*. *JAMA* 285: 2450–2451. <https://doi.org/10.1001/jama.285.19.2450>
- Gern L (2005) [The biology of the Ixodes ricinus tick]. *Therapeutische Umschau* 62(11): 707–712. <https://doi.org/10.1024/0040-5930.62.11.707>
- Getty TA, Peterson JW, Fujioka H, Walsh AM, Sam-Yellowe TY (2021) *Colpodella* sp. (ATCC 50594) Life Cycle: Myzocytosis and Possible Links to the Origin of Intracellular Parasitism. *Tropical Medicine and Infectious Disease* 6(3): e127.
- Ghadirian E (1977) Human Infection with *Trichostrongylus lerouxi* (Biocca, Chabaud, and Ghadirian, 1974) in Iran. *The American Journal of Tropical Medicine and Hygiene* 26: 1212–1213. <https://doi.org/10.4269/ajtmh.1977.26.1212>
- Ghadirian E, Arfaa F (1973) First report of human infection with *Haemonchus contortus*, *Ostertagia ostertagi*, and *Marshallagia marshalli* (Family Trichostrongylidae) in Iran. *The Journal of Parasitology* 59: 1144–1145. <https://doi.org/10.2307/3278661>
- Ghadirian E, Arfaa F (1975) Present status of trichostrongyliasis in Iran. *American Journal of Tropical Medicine and Hygiene* 24: 935–941. <https://doi.org/10.4269/ajtmh.1975.24.935>
- Ghai RR, Chapman CA, Omeja PA, Davies TJ, Goldberg TL (2014) Nodule worm infection in humans and wild primates in Uganda: cryptic species in a newly identified region of human transmission. *PLoS Neglected Tropical Diseases* 8: e2641. <https://doi.org/10.1371/journal.pntd.0002641>
- Gharpure R, Bliton J, Goodman A, Ali IKM, Yoder J, Cope JR (2020) Epidemiology and clinical characteristics of primary amebic meningoencephalitis caused by *Naegleria fowleri*: A global review. *Clinical Infectious Diseases* 73(1): e19–e27. <https://doi.org/10.1093/cid/ciaa520>
- Ghavami MB, Djalilvand A (2015) First Record of Urogenital Myiasis Induced by *Megaselia scalaris* (Diptera: Phoridae) from Iran. *Journal of Arthropod Borne Diseases* 9: 274–280.
- Ghobakhloo N, Motazedian MH, Naderi S, Ebrahimi S (2019) Isolation of *Crithidia* spp. from lesions of immunocompetent patients with suspected cutaneous leishmaniasis in Iran. *Tropical Medicine & International Health* 24: 116–126. <https://doi.org/10.1111/tmi.13042>
- Ghosh S, Banerjee P, Sarkar A, Datta S, Chatterjee M (2012) Coinfection of *Leptomonas seymouri* and *Leishmania donovani* in Indian leishmaniasis. *Journal of Clinical Microbiology* 50: 2774–2778. <https://doi.org/10.1128/JCM.00966-12>
- Glen DR, Brooks DR (1985) Phylogenetic relationships of some strongylate nematodes of primates. *Proceedings of the Helminthological Society of Washington* 52: 227–236.

- Goddard J, Baker GT, Ferrari FG, Ferrari C (2012) Bed bugs (*Cimex lectularius*) and bat bugs (several *Cimex* species): a confusing issue. *Outlooks on Pest Management* 23: 125–127. <https://doi.org/10.1564/23jun09>
- Goldsmid JM (1967) *Rhabditis (Rhabditella) axei* in the urine of an African in Rhodesia. *Journal of Helminthology* 41: 305–308. <https://doi.org/10.1017/S0022149X00021842>
- Goldsmid JM, Muir M (1972) *Inermicapsifer madagascariensis* (Davaine, 1870), Baer, 1956 (platyhelminthes: cestoda) as a parasite of man in Rhodesia. *Central African Journal of Medicine* 18: 205–207.
- Goto Y, Tamura A, Ishikawa O, Miyachi Y, Ishii T, Akao N (1998) Creeping eruption caused by a larva of the suborder Spirurina type X. *British Journal of Dermatology* 139: 315–318. <https://doi.org/10.1046/j.1365-2133.1998.02375.x>
- Gottstein B, Pozio E, Nockler K (2009) Epidemiology, diagnosis, treatment, and control of trichinellosis. *Clinical Microbiology Reviews* 22: 127–145. <https://doi.org/10.1128/CMR.00026-08>
- Grady CA, McDonald DR, Poppen CF, Pritt BS (2011) *Dermacentor* tick attached to tympanic membrane. *Lancet* 378(9788): 347–347. [https://doi.org/10.1016/S0140-6736\(11\)60981-1](https://doi.org/10.1016/S0140-6736(11)60981-1)
- Graf J (1999) Symbiosis of *Aeromonas veronii* biovar *sobria* and *Hirudo medicinalis*, the medicinal leech: a novel model for digestive tract associations. *Infection and Immunity* 67: 1–7. <https://doi.org/10.1128/IAI.67.1.1-7.1999>
- Graham AM, Davenport A, Moshnikova VS, Gilmour LJ, Fabiani M, Bishop MA, Cook AK (2021) *Heterobilharzia americana* infection in dogs: A retrospective study of 60 cases (2010–2019). *Journal of Veterinary Internal Medicine* 35: 1361–1367. <https://doi.org/10.1111/jvim.16127>
- Gray A, Capewell P, Loney C, Katzer F, Shiels BR, Weir W (2019) Sheep as host species for zoonotic *Babesia venatorum*, United Kingdom. *Emerging Infectious Diseases* 25: 2257–2260. <https://doi.org/10.3201/eid2512.190459>
- Greigert V, Abou-Bacar A, Brunet J, Nourrisson C, Pfaff AW, Benarbia L, Pereira B, Randrianarivojosia M, Razafindrakoto JL, Solotiana Rakotomalala R, Morel E, Candolfi E, Poirier P (2018) Human intestinal parasites in Mahajanga, Madagascar: The kingdom of the Protozoa. *PLoS ONE* 13: e0204576. <https://doi.org/10.1371/journal.pone.0204576>
- Grimaldi Junior G, Kreutzer RD, Hashiguchi Y, Gomez EA, Mimory T, Tesh RB (1992) Description of *Leishmania equatorensis* sp. n (Kinetoplastida: Trypanosomatidae), a new parasite infecting arboreal mammals in Ecuador. *Memórias do Instituto Oswaldo Cruz* 87: 221–228. <https://doi.org/10.1590/S0074-02761992000200009>
- Grün AL, Stemplewitz B, Scheid P (2014) First report of an *Acanthamoeba* genotype T13 isolate as etiological agent of a keratitis in humans. *Parasitology Research* 113: 2395–2400. <https://doi.org/10.1007/s00436-014-3918-5>
- Guerra JA, Prestes SR, Silveira H, Coelho LI, Gama P, Moura A, Amato V, Barbosa M, Ferreira LC (2011) Mucosal *Leishmaniasis* caused by *Leishmania (Viannia) braziliensis* and *Leishmania (Viannia) guyanensis* in the Brazilian Amazon. *PLoS Neglected Tropical Diseases* 5: e980. <https://doi.org/10.1371/journal.pntd.0000980>

- Guerrant RL, Walker DH, Weller PF (2006) Tropical infectious diseases : principles, pathogens & practice. Churchill Livingstone, Philadelphia.
- Guglielmono AA, Beati L, Barros-Battesti DM, Labruna MB, Nava S, Venzal JM, Mangold AJ, Szabo MP, Martins JR, Gonzalez-Acuna D, Estrada-Pena A (2006) Ticks (Ixodidae) on humans in South America. *Experimental and Applied Acarology* 40: 83–100. <https://doi.org/10.1007/s10493-006-9027-0>
- Guglielmono AA, Robbins RG (2018) Hard ticks (Acari: Ixodida: Ixodidae) parasitizing humans: A global overview. Springer International Publishing, Cham, 668 pp.
- Guhl F, Vallejo GA (2003) *Trypanosoma (Herpetosoma) rangeli* Tejera, 1920: an updated review. *Memórias do Instituto Oswaldo Cruz* 98: 435–442. <https://doi.org/10.1590/S0074-02762003000400001>
- Gui Z, Wu L, Cai H, Mu L, Yu J-F, Fu S-Y, Si X-Y (2021) Genetic diversity analysis of *Dermacentor nuttalli* within Inner Mongolia, China. *Parasites & Vectors* 14: e131. <https://doi.org/10.1186/s13071-021-04625-5>
- Guimaraes LO, Bajay MM, Wunderlich G, Bueno MG, Rohe F, Catao-Dias JL, Neves A, Malafronte RS, Curado I, Kirchgatter K (2012) The genetic diversity of *Plasmodium malariae* and *Plasmodium brasilianum* from human, simian and mosquito hosts in Brazil. *Acta Tropica* 124: 27–32. <https://doi.org/10.1016/j.actatropica.2012.05.016>
- Gunawardena K (1963) A Study of *Onthophagus unifasciatus* Schall (Scarabaeidae-Coprinii) and Scarabiasis in Ceylon. *Indian Journal of Medical Research* 51: 654–660.
- Güner ES, Watanabe M, Hashimoto N, Kadosaka T, Kawamura Y, Ezaki T, Kawabata H, Imai Y, Kaneda K, Masuzawa T (2004) *Borrelia turcica* sp. nov., isolated from the hard tick *Hyalomma aegyptium* in Turkey. *International Journal of Systematic and Evolutionary Microbiology* 54: 1649–1652. <https://doi.org/10.1099/ijs.0.03050-0>
- Guo X (2020) Proteomics Analysis of *Hydatigera taeniaeformis* Metacestode Stage. *Frontiers in Veterinary Science* 7: e474. <https://doi.org/10.3389/fvets.2020.00474>
- Hadziyannis E, Yen-Lieberman B, Hall G, Procop GW (2000) Ciliocytophthoria in clinical virology. *Archives of Pathology & Laboratory Medicine* 124: 1220–1223.
- Hale AJ, Mathison B, Pritt B, Collins K (2019) Endemic bot fly larvae infection in Northern New York State. *IDCases* 16: e00531. <https://doi.org/10.1016/j.idcr.2019.e00531>
- Hall-Mendelin S, Craig SB, Hall RA, O'Donoghue P, Atwell RB, Tulsiani SM, Graham GC (2011) Tick paralysis in Australia caused by *Ixodes holocyclus* Neumann. *Annals of Tropical Medicine and Parasitology* 105: 95–106. <https://doi.org/10.1179/136485911X12899838413628>
- Hall RL, Lindsay A, Hammond C, Montgomery SP, Wilkins PP, da Silva AJ, McAuliffe I, de Almeida M, Bishop H, Mathison B, Sun B, Largusa R, Jones JL (2012) Outbreak of human trichinellosis in Northern California caused by *Trichinella murrelli*. *American Journal of Tropical Medicine and Hygiene* 87: 297–302. <https://doi.org/10.4269/ajtmh.2012.12-0075>
- Hamilton H, Pontiff KL, Bolton M, Bradbury RS, Mathison BA, Bishop H, de Almeida M, Ogden BW, Barnett E, Rastanis D, Klar AL, Uzodi AS (2018) *Trichomonas vaginalis* Brain Abscess in a Neonate. *Clinical Infectious Diseases* 66: 604–607. <https://doi.org/10.1093/cid/cix908>

- Hanya G, Morishima K, Koide T, Otani Y, Hongo S, Honda T, Okamura H, Higo Y, Hattori M, Kondo Y, Kurihara Y, Jin S, Otake A, Shiroishi I, Takakuwa T, Yamamoto H, Suzuki H, Kajimura H, Hayakawa T, Suzuki-Hashido N, Nakano T (2019) Host selection of hematophagous leeches (*Haemadipsa japonica*): Implications for iDNA studies. *Ecological Research* 34: 842–855. <https://doi.org/10.1111/1440-1703.12059>
- Harcourt-Brown F (2002) Infectious diseases of domestic rabbits. *Textbook of Rabbit Medicine*: 361–385. <https://doi.org/10.1016/B978-075064002-2.50019-9>
- Harold Hinman E, Kampmeier RH (1934) Intestinal Acariasis Due to *Tyroglyphus longior* Gervais. *American Journal of Tropical Medicine and Hygiene* s1–14(4): 355–362. <https://doi.org/10.4269/ajtmh.1934.s1-14.355>
- Hart BJ, Fain A (1988) Morphological and biological studies of medically important household mites. *Acarologia* 29: 285–295.
- Hartmeyer GN, Stensvold CR, Fabricius T, Marmolin ES, Hoegh SV, Nielsen HV, Kemp M, Vestergaard LS (2019) *Plasmodium cynomolgi* as cause of malaria in tourist to Southeast Asia, 2018. *Emerging Infectious Diseases* 25: 1936–1939. <https://doi.org/10.3201/eid2510.190448>
- Hasegawa H, Korenaga M, Kumazawa H, Imamura K (1996) *Mermis* sp. found from human urine in Kochi Prefecture, Japan. (Nematoda: Mermithidae). *Japanese Journal of Parasitology* 45: 43–46.
- Heisch RB, Harvey AE (1953) The biology of *Ornithodoros graingeri* Heisch & Guggisberg, 1952. *Parasitology* 43: 131–132. <https://doi.org/10.1017/S0031182000018382>
- Herc E, Pritt B, Huizenga T, Douce R, Hysell M, Newton D, Sidge J, Losman E, Sherbeck J, Kaul DR (2018) Probable Locally Acquired *Babesia divergens*-Like Infection in Woman, Michigan, USA. *Emerging Infectious Diseases* 24(8): 1558–1560. <https://doi.org/10.3201/eid2408.180309>
- Herman JS, Chiodini PL (2009) Gnathostomiasis, another emerging imported disease. *Clinical Microbiology Reviews* 22: 484–492. <https://doi.org/10.1128/CMR.00003-09>
- Hernandez-Orts JS, Scholz T, Brabec J, Kuzmina T, Kuchta R (2015) High morphological plasticity and global geographical distribution of the Pacific broad tapeworm *Adenocephalus pacificus* (syn. *Diphyllobothrium pacificum*): molecular and morphological survey. *Acta Tropica* 149: 168–178. <https://doi.org/10.1016/j.actatropica.2015.05.017>
- Herter CD, Nesse RE (1989) Pseudoparasitism with *Gordius robustus*. *American Family Physician* 39: 139–142.
- Herwaldt BL, Caccio S, Gherlinzoni F, Aspöck H, Slemenda SB, Piccaluga P, Martinelli G, Edelhofer R, Hollenstein U, Poletti G, Pampiglione S, Loschenberger K, Tura S, Pieni-azek NJ (2003) Molecular characterization of a non-*Babesia divergens* organism causing zoonotic babesiosis in Europe. *Emerging Infectious Diseases* 9: 942–948. <https://doi.org/10.3201/eid0908.020748>
- Heyworth MF (2016) *Giardia duodenalis* genetic assemblages and hosts. *Parasite* 23: e13. [5 pp.] <https://doi.org/10.1051/parasite/2016013>
- Hii JL, Kan SK, Au Yong KS (1978) A record of *Limnatis maculosa* (Blanchard) (Hirudinea: Arynchobdellida) taken from the nasal cavity of man in Sabah, Malaysia. *Medical Journal of Malaysia* 32: 247–248.

- Hinnebusch BJ, Bland DM, Bosio CF, Jarrett CO (2017) Comparative ability of *Oropsylla montana* and *Xenopsylla cheopis* fleas to transmit *Yersinia pestis* by two different mechanisms. *PLoS Neglected Tropical Diseases* 11: e0005276. <https://doi.org/10.1371/journal.pntd.0005276>
- Hira PR, Assad RM, Okasha G, Al-Ali FM, Iqbal J, Mutawali KE, Disney RH, Hall MJ (2004) Myiasis in Kuwait: nosocomial infections caused by *Lucilia sericata* and *Megaselia scalaris*. *American Journal of Tropical Medicine and Hygiene* 70: 386–389. <https://doi.org/10.4269/ajtmh.2004.70.386>
- Hoberg EP (2002) Taenia tapeworms: their biology, evolution and socioeconomic significance. *Microbes and Infections* 4: 859–866. [https://doi.org/10.1016/S1286-4579\(02\)01606-4](https://doi.org/10.1016/S1286-4579(02)01606-4)
- Hoffman GL (1955) Research notes: notes on the life cycle of *Fibricola cratera* (Trematoda: Strigeida). *Journal of Parasitology* 41: 327–327. <https://doi.org/10.2307/3274227>
- Hong Q, Feng J, Liu H, Li X, Gong L, Yang Z, Yang W, Liang X, Zheng R, Cui Z, Wang W, Chen D (2016) Prevalence of *Spirometra mansoni* in dogs, cats, and frogs and its medical relevance in Guangzhou, China. *International Journal of Infectious Diseases* 53: 41–45. <https://doi.org/10.1016/j.ijid.2016.10.013>
- Hong S-H, Kim S-Y, Song BG, Roh JY, Cho CR, Kim C-N, Um T-H, Kwak YG, Cho S-H, Lee S-E (2019) Detection and characterization of an emerging type of *Babesia* sp. similar to *Babesia motasi* for the first case of human babesiosis and ticks in Korea. *Emerging Microbes & Infections* 8: 869–878. <https://doi.org/10.1080/22221751.2019.1622997>
- Hoogstraal H (1979) The epidemiology of tick-borne Crimean-Congo hemorrhagic fever in Asia, Europe, and Africa. *Journal of Medical Entomology* 15: 307–417. <https://doi.org/10.1093/jmedent/15.4.307>
- Hoogstraal H (1985) Argasid and Nuttalliellid ticks as parasites and vectors. *Advances in Parasitology* 24: 135–238. [https://doi.org/10.1016/S0065-308X\(08\)60563-1](https://doi.org/10.1016/S0065-308X(08)60563-1)
- Hooshyar H, Rostamkhani P, Rezaeian M (2015) An annotated checklist of the human and animal Entamoeba (Amoebida: Endamoebidae) species – A review article. *Iranian Journal of Parasitology* 10: 146–156.
- Hopkins DR, Weiss AJ, Roy SL, Yerian S, Sapp SGH (2020) Progress toward global eradication of Dracunculiasis, January 2019–June 2020. *MMWR Morbidity Mortality Weekly Report* 69: 1563–1568. <https://doi.org/10.15585/mmwr.mm6943a2>
- Horák P, Kolárová L, Adema CM (2002) Biology of the schistosome genus *Trichobilharzia*. *Advances in Parasitology* 52: 155–233. [https://doi.org/10.1016/S0065-308X\(02\)52012-1](https://doi.org/10.1016/S0065-308X(02)52012-1)
- Horák P, Mikeš L, Lichtenbergová L, Skála V, Soldánová M, Brant SV (2015) Avian schistosomes and outbreaks of cercarial dermatitis. *Clinical Microbiology Reviews* 28: 165–190. <https://doi.org/10.1128/CMR.00043-14>
- Hotez PJ, Brooker S, Bethony JM, Bottazzi ME, Loukas A, Xiao S (2004) Hookworm infection. *New England Journal of Medicine* 351: 799–807. <https://doi.org/10.1056/NEJMra032492>
- Howes RE, Battle KE, Mendis KN, Smith DL, Cibulskis RE, Baird JK, Hay SI (2016) Global Epidemiology of *Plasmodium vivax*. *American Journal of Tropical Medicine and Hygiene* 95: 15–34. <https://doi.org/10.4269/ajtmh.16-0141>

- Hua R, Xie Y, Song H, Shi Y, Zhan J, Wu M, Gu X, Peng X, Yang G (2019) Echinococcus canadensis G8 tapeworm infection in a sheep, China, 2018. *Emerging Infectious Diseases* 25: 1420–1422. <https://doi.org/10.3201/eid2507.181585>
- Huang Y-L, Huang D-N, Wu W-H, Yang F, Zhang X-M, Wang M, Tang Y-J, Zhang Q, Peng L-F, Zhang R-L (2018) Identification and characterization of the causative triatomine bugs of anaphylactic shock in Zhanjiang, China. *Infectious Diseases of Poverty* 7: e127. <https://doi.org/10.1186/s40249-018-0509-1>
- Huynh T, Thean J, Maini R (2001) *Dipetalonema reconditum* in the human eye. *British Journal of Ophthalmology* 85: 1384–1384. <https://doi.org/10.1136/bjo.85.11.1384i>
- Idro R, Marsh K, John CC, Newton CR (2010) Cerebral malaria: mechanisms of brain injury and strategies for improved neurocognitive outcome. *Pediatric Research* 68: 267–274. <https://doi.org/10.1203/PDR.0b013e3181eee738>
- Ikuno H, Akao S, Yamasaki H (2018) Epidemiology of *Diphyllobothrium nihonkaiense* Diphyllobothriasis, Japan, 2001–2016. *Emerging Infectious Diseases* 24(8): 1428–1434. <https://doi.org/10.3201/eid2408.171454>
- Ioannou P, Vamvoukaki R (2019) *Armillifer* infections in humans: A systematic review. *Tropical Medicine and Infectious Disease* 4(2): e80. <https://doi.org/10.3390/tropicalmed4020080>
- Ionita M, Verale MG, Lyons ET, Spraker TR (2008) Hookworms (*Uncinaria lucasi*) and acanthocephalans (*Corynosoma* spp. and *Bolbosoma* spp.) found in dead northern fur seals (*Callorhinus ursinus*) on St. Paul Island, Alaska in 2007. *Parasitology Research* 103: 1025–1029. <https://doi.org/10.1007/s00436-008-1087-0>
- Ivancovich JC, Luciana CA (1992) Las garrapatas de Argentina. Monografía de la Asociación Argentina de Parasitología Veterinaria, Buenos Aires, 95 pp.
- Iwasaki S, Takebayashi S, Watanabe T (2007) Tick bites in the external auditory canal. *Auris Nasus Larynx* 34: 375–377. <https://doi.org/10.1016/j.anl.2006.09.013>
- Iyengar M (1928) Infestation of the human intestine by corpid beetles in Bengal. *The Indian Medical Gazette* 63(7): 365–369.
- Jaenson TGT, Väriv K, Fröjdman I, Jääskeläinen A, Rundgren K, Versteirt V, Estrada-Peña A, Medlock JM, Golovljova I (2016) First evidence of established populations of the taiga tick *Ixodes persulcatus* (Acari: Ixodidae) in Sweden. *Parasites & Vectors* 9: e377. <https://doi.org/10.1186/s13071-016-1658-3>
- Jariyapan N, Daroontum T, Jaiwong K, Chanmol W, Intakhan N, Sor-Suwan S, Siriyasatien P, Somboon P, Bates MD, Bates PA (2018) *Leishmania (Mundinia) orientalis* n. sp. (Trypanosomatidae), a parasite from Thailand responsible for localised cutaneous leishmaniasis. *Parasites & Vectors* 11: e351. <https://doi.org/10.1186/s13071-018-2908-3>
- Jeon H-K, Park H, Lee D, Choe S, Kang Y, Bia MM, Lee S-H, Sohn W-M, Hong S-J, Chai J-Y, Eom KS (2018a) Genetic and morphologic identification of *Spirometra ranarum* in Myanmar. *The Korean Journal of Parasitology* 56: 275–280. <https://doi.org/10.3347/kjp.2018.56.3.275>
- Jeon H-K, Park H, Lee D, Choe S, Kim K-H, Huh S, Sohn W-M, Chai J-Y, Eom KS (2015) Human infections with *Spirometra decipiens* Plerocercoids identified by morphologic and genetic analyses in Korea. *The Korean Journal of Parasitology* 53: 299–305. <https://doi.org/10.3347/kjp.2015.53.3.299>

- Jeon HK, Kim KH, Sohn WM, Eom KS (2018b) Differential diagnosis of human sparganosis using multiplex PCR. *The Korean Journal of Parasitology* 56: 295–300. <https://doi.org/10.3347/kjp.2018.56.3.295>
- Jeratthitikul E, Jiranunskul P, Nakano T, Sucharit C, Panha S (2020) A new species of buffalo leech in the genus *Hirudinaria* Whitman, 1886 (Arhynchobdellida, Hirudinidae) from Thailand. *Zookeys* 933: 1–14. <https://doi.org/10.3897/zookeys.933.49314>
- Ji-Tuan X (1997) Four misdiagnosed cases of visceral bleeding caused by *Haemadipha japonica*. *Southeast Asian Journal of Tropical Medicine and Public Health* 28: 673–674.
- Jia N, Zheng YC, Jiang JF, Jiang RR, Jiang BG, Wei R, Liu HB, Huo QB, Sun Y, Chu YL, Fan H, Chang QC, Yao NN, Zhang WH, Wang H, Guo DH, Fu X, Wang YW, Krause PJ, Song JL, Cao WC (2018) Human Babesiosis caused by a *Babesia crassa*-like pathogen: A case series. *Clinical Infectious Diseases* 67: 1110–1119. <https://doi.org/10.1093/cid/ciy212>
- Jiang JF, Jiang RR, Chang QC, Zheng YC, Jiang BG, Sun Y, Jia N, Wei R, Liu HB, Huo QB, Wang H, von Fricken ME, Cao WC (2018) Potential novel tick-borne *Colpodella* species parasite infection in patient with neurological symptoms. *PLoS Neglected Tropical Diseases* 12: e0006546. <https://doi.org/10.1371/journal.pntd.0006546>
- Jiang W, Roellig DM, Lebbad M, Beser J, Troell K, Guo Y, Li N, Xiao L, Feng Y (2020) Subtype distribution of zoonotic pathogen *Cryptosporidium felis* in humans and animals in several countries. *Emerging Microbes & Infections* 9: 2446–2454. <https://doi.org/10.1080/22221751.2020.1840312>
- Jiang Y, Ren J, Yuan Z, Liu A, Zhao H, Liu H, Chu L, Pan W, Cao J, Lin Y, Shen Y (2014) *Cryptosporidium andersoni* as a novel predominant *Cryptosporidium* species in outpatients with diarrhea in Jiangsu Province, China. *BMC Infectious Diseases* 14: e555. <https://doi.org/10.1186/s12879-014-0555-7>
- Jiménez PA, Jaimes JE, Ramírez JD (2019) A summary of *Blastocystis* subtypes in North and South America. *Parasites & Vectors* 12: e376. <https://doi.org/10.1186/s13071-019-3641-2>
- Jongejan F, Berger L, Busser S, Deetman I, Jochems M, Leenders T, de Sitter B, van der Steen F, Wentzel J, Stoltz H (2020) *Amblyomma hebraeum* is the predominant tick species on goats in the Mnisi Community Area of Mpumalanga Province South Africa and is co-infected with *Ehrlichia ruminantium* and *Rickettsia africae*. *Parasites & Vectors* 13: e172. <https://doi.org/10.1186/s13071-020-04059-5>
- Joshi PP, Shegokar VR, Powar RM, Herder S, Katti R, Salkar HR, Dani VS, Bhargava A, Jannin J, Truc P (2005) Human trypanosomiasis caused by *Trypanosoma evansi* in India: the first case report. *American Journal of Tropical Medicine and Hygiene* 73: 491–495. <https://doi.org/10.4269/ajtmh.2005.73.491>
- Jurberg J, Galvão C (2006) Biology, ecology, and systematics of Triatominae (Heteroptera, Reduviidae), vectors of Chagas disease, and implications for human health. *Landesmuseum Neue Serie* 50: 1096–1116.
- Kaewpitoon N, Kaewpitoon SJ, Pengsaa P, Sripa B (2008) *Opisthorchis viverrini*: the carcinogenic human liver fluke. *World Journal of Gastroenterology* 14: 666–674. <https://doi.org/10.3748/wjg.14.666>
- Kagei N, Oshima T, Inoue I, Kumasaki T (1966) First human case of gordiid worm in Japan (Nematomorpha: Gordiidae). *Japanese Journal of Parasitology* 15: 79–81.

- Kalousová B, Hasegawa H, Petrželková KJ, Sakamaki T, Kooriyma T, Modrý D (2016) Adult hookworms (*Necator* spp.) collected from researchers working with wild western lowland gorillas. *Parasites & Vectors* 9: e75.
- Kamo HYS, Fukumoto S, Maejima J, Sakagucki Y (1986) Two unknown marine species of the genus *Diphyllbothrium* from human cases. *Japanese Journal of Tropical Medicine and Hygiene* 14: 79–86. <https://doi.org/10.2149/tmh1973.14.79>
- Kannangara WW (1971) *Paratelfusa rugosa* as the second intermediate host of *Achillurbainia*, a trematode transmissible to man. *Journal of Parasitology* 57: 683–684. <https://doi.org/10.2307/3277944>
- Kapoor AK, Agarwal S, Singh A, Pandey R, Khare V (2019) Diarrhoea due to intestinal acariasis. *Journal of Clinical and Diagnostic Research* 13: EL01–EL02. <https://doi.org/10.7860/JCDR/2019/40291.12676>
- Kassari H, Dehghani R, Kasiri M, Dehghani M, Kasiri R (2020) A review on the reappearance of Crimean-Congo hemorrhagic fever, a tick-borne nairovirus. *Entomology and Applied Science Letters* 7(1): 81–90.
- Kato H, Bone AE, Mimori T, Hashiguchi K, Shiguango GF, Gonzales SV, Velez LN, Guevara AG, Gomez EA, Hashiguchi Y (2016) First human cases of *Leishmania* (*Viannia*) *lainsoni* infection and a search for the vector sand flies in Ecuador. *PLoS Neglected Tropical Diseases* 10: e0004728. <https://doi.org/10.1371/journal.pntd.0004728>
- Kato H, Seki C, Kubo M, Gonzales-Cornejo L, Caceres AG (2021) Natural infections of *Pintomyia verrucarum* and *Pintomyia maranonensis* by *Leishmania* (*Viannia*) *peruviana* in the Eastern Andes of northern Peru. *PLoS Neglected Tropical Diseases* 15: e0009352. <https://doi.org/10.1371/journal.pntd.0009352>
- Keh B, Lane RS (1987) *Cheyletiella blakei*, an ectoparasite of cats, as cause of cryptic arthropod infestations affecting humans. *Western Journal of Medicine* 146: 192–194.
- Ken KM, Shockman SC, Sirichotiratana M, Lent MP, Wilson ML (2014) Dermatoses associated with mites other than *Sarcoptes*. *Seminars in Cutaneous Medicine and Surgery* 33: 110–115. <https://doi.org/10.12788/j.sder.0104>
- Kenney M, Eveland LK, Yermakov V, Kassouny DY (1975) A case of *Rictularia* infection of man in New York. *The American Journal of Tropical Medicine and Hygiene* 24: 596–599. <https://doi.org/10.4269/ajtmh.1975.24.596>
- Khalifa RMA, Abdellatif MZM, Ahmed AK, Yones DA, El-Mazary AM, Aly LH, El-Seify MA, Haridi MA (2016) First case of intestinal acariasis from Egypt. *SpringerPlus* 5: e28. <https://doi.org/10.1186/s40064-015-1584-4>
- Khan M, Beckham JD, Piquet AL, Tyler KL, Pastula DM (2019) An overview of Powassan virus disease. *Neurohospitalist* 9: 181–182. <https://doi.org/10.1177/1941874419844888>
- Khan NA (2006) *Acanthamoeba*: biology and increasing importance in human health. *FEMS Microbiology Reviews* 30: 564–595. <https://doi.org/10.1111/j.1574-6976.2006.00023.x>
- Kiakojouri K, Omran S, Rajabnia R, Pournajaf A, Armaki M, Karami M (2018) A case of human acute otoacariasis caused by *Rhizoglyphus* sp, the first report from Iran. *Journal of Acute Disease* 7: 220–222. <https://doi.org/10.4103/2221-6189.244175>
- Kiewra D, Czułowska A, Dyczko D, Zieliński R, Plewa-Tutaj K (2019) First record of *Haemaphysalis concinna* (Acari: Ixodidae) in Lower Silesia, SW Poland. *Experimental and Applied Acarology* 77: 449–454. <https://doi.org/10.1007/s10493-019-00344-w>

- Kifune T, Hatsushika R, Ushirogawa H, Takeda S, Kono K, Shimizu T (2000) A case study of human infection with diphyllbothriid tapeworm (Cestoda: Pseudophyllidea) found from a man in Fukuoka prefecture. *Japanese Medical Bulletin of Fukuoka University* 27: 93–100.
- Kim HJ, Yong TS, Shin MH, Lee KJ, Park GM, Suvonkulov U, Kovalenko D, Yu HS (2020) Phylogenetic characteristics of *Echinococcus granulosus* sensu lato in Uzbekistan. *The Korean Journal of Parasitology* 58: 205–210. <https://doi.org/10.3347/kjp.2020.58.2.205>
- Kim JY, Cho SH, Joo HN, Tsuji M, Cho SR, Park IJ, Chung GT, Ju JW, Cheun HI, Lee HW, Lee YH, Kim TS (2007) First case of human babesiosis in Korea: detection and characterization of a novel type of *Babesia* sp. (KO1) similar to ovine babesia. *Journal of Clinical Microbiology* 45: 2084–2087. <https://doi.org/10.1128/JCM.01334-06>
- Kim JY, Lim HY, Shin SE, Cha HK, Seo J-H, Kim S-K, Park SH, Son GH (2018) Comprehensive transcriptome analysis of *Sarcophaga peregrina*, a forensically important fly species. *Scientific Data* 5: e180220. <https://doi.org/10.1038/sdata.2018.220>
- Kirchhoff LV (2011) Epidemiology of American trypanosomiasis (Chagas disease). *Advances in Parasitology* 75: 1–18. <https://doi.org/10.1016/B978-0-12-385863-4.00001-0>
- Kissinger P (2015) *Trichomonas vaginalis*: a review of epidemiologic, clinical and treatment issues. *BMC Infectious Diseases* 15: e307. <https://doi.org/10.1186/s12879-015-1055-0>
- Klapow LA (1999) Roundworm-like specimens in the sputum of chronic fatigue syndrome patients and controls in open and blind analyses. In: Patarca-Montero R (Ed.) *Chronic Fatigue Syndrome: Advances in Epidemiologic, Clinical, and Basic Science Research*. Hawthorn Medical Press, Binghamton, 247–248.
- Klatt S, Simpson L, Maslov DA, Konthur Z (2019) *Leishmania tarentolae*: Taxonomic classification and its application as a promising biotechnological expression host. *PLoS Neglected Tropical Diseases* 13: e0007424. [29 pp.] <https://doi.org/10.1371/journal.pntd.0007424>
- Kobayashi A, Katakura K, Hamada A, Suzuki T, Hataba Y, Tashiro N, Yoshida A (1986) Human case of Dracunculiasis in Japan. *The American Journal of Tropical Medicine and Hygiene* 35: 159–161. <https://doi.org/10.4269/ajtmh.1986.35.159>
- Koehler AV, Robson JMB, Spratt DM, Hann J, Beveridge I, Walsh M, McDougall R, Bromley M, Hume A, Sheorey H, Gasser RB (2021) Ocular Filariasis in Human Caused by *Breinvilia (Johnstonema) annulipapillata* Nematode, Australia. *Emerging Infectious Diseases* 27: 297–300. <https://doi.org/10.3201/eid2701.203585>
- Koehler AV, Wang T, Haydon SR, Gasser RB (2018) *Cryptosporidium viatorum* from the native Australian swamp rat *Rattus lutreolus* – An emerging zoonotic pathogen? *International Journal for Parasitology: Parasites and Wildlife* 7: 18–26. <https://doi.org/10.1016/j.ijppaw.2018.01.004>
- Koehler AV, Whipp MJ, Haydon SR, Gasser RB (2014) *Cryptosporidium cuniculus* – new records in human and kangaroo in Australia. *Parasites & Vectors* 7: e492. <https://doi.org/10.1186/s13071-014-0492-8>
- Koehler M, Soleiman A, Aspöck H, Auer H, Walochnik J (2007) *Onchocerca jakutensis* filariasis in humans. *Emerging Infectious Diseases* 13: 1749–1752. <https://doi.org/10.3201/eid1311.070017>
- Kogoashiwa Y, Moro Y, Kono N (2009) A case of tick bite (*Ixodes persulcatus*) in the external auditory canal. *Otolaryngol Head Neck Surg (Tokyo)* 81: 670–672.
- Kopacz Ž, Kváč M, Karpiński P, Hendrich AB, Sasiadek MM, Leszczyński P, Sak B, McEvoy J, Kicia M (2019) The first evidence of *Cryptosporidium meleagridis* infection in a colon

- Adenocarcinoma from an immunocompetent patient. *Frontiers in Cellular and Infection Microbiology* 9: 1–5. <https://doi.org/10.3389/fcimb.2019.00035>
- Kraeva N, Butenko A, Hlaváčová J, Kostygov A, Myšková J, Grybchuk D, Leštinová T, Votýpka J, Volf P, Opperdoes F, Flegontov P, Lukeš J, Yurchenko V (2015) *Leptomonas seymouri*: Adaptations to the Dixenous life cycle analyzed by genome sequencing, transcriptome profiling and co-infection with *Leishmania donovani*. *PLoS Pathogens* 11: e1005127. [23 pp.] <https://doi.org/10.1371/journal.ppat.1005127>
- Kreppel KS, Telfer S, Rajerison M, Morse A, Baylis M (2016) Effect of temperature and relative humidity on the development times and survival of *Synopsyllus fonquerniei* and *Xenopsylla cheopis*, the flea vectors of plague in Madagascar. *Parasites & Vectors* 9: e82. <https://doi.org/10.1186/s13071-016-1366-z>
- Kreutzer RD, Corredor A, Grimaldi Jr G, Grogl M, Rowton ED, Young DG, Morales A, McMahon-Pratt D, Guzman H, Tesh RB (1991) Characterization of *Leishmania colombiensis* sp. n. (Kinetoplastida: Trypanosomatidae), a new parasite infecting humans, animals, and phlebotomine sand flies in Colombia and Panama. *American Journal of Tropical Medicine and Hygiene* 44: 662–675. <https://doi.org/10.4269/ajtmh.1991.44.662>
- Kruger FJ, Evans AC (2009) Do all human urinary infections with *Schistosoma mattheei* Represent hybridization between *S. haematobium* and *S. mattheei*? *Journal of Helminthology* 64: 330–332. <https://doi.org/10.1017/S0022149X00012384>
- Kuchta R, Brabec J, Kubackova P, Scholz T (2013) Tapeworm *Diphyllobothrium dendriticum* (Cestoda) – neglected or emerging human parasite? *PLoS Neglected Tropical Diseases* 7: e2535.
- Kuchta R, Kolodziej-Sobocinska M, Brabec J, Mlocicki D, Salamatin R, Scholz T (2021) Sparganosis (*Spirometra*) in Europe in the Molecular Era. *Clinical Infectious Diseases* 72: 882–890. <https://doi.org/10.1093/cid/ciaa1036>
- Kuchta R, Scholz T, Brabec J, Bray RA (2008) Suppression of the tapeworm order Pseudophylloidea (Platyhelminthes: Eucestoda) and the proposal of two new orders, Bothriocephalidea and Diphyllobothriidea. *International Journal of Parasitology* 38: 49–55. <https://doi.org/10.1016/j.ijpara.2007.08.005>
- Kuchta R, Serrano-Martinez ME, Scholz T (2015) Pacific broad tapeworm *Adenocephalus pacificus* as a causative agent of globally reemerging Diphyllobothriosis. *Emerging Infectious Diseases* 21: 1697–1703. <https://doi.org/10.3201/eid2110.150516>
- Kukina IV, Guzeeva TM, Zelya OP, Ganushkina LA (2018) Fatal human babesiosis caused by *Babesia divergens* in an asplenic host. *IDCases* 13: e00414. <https://doi.org/10.1016/j.idcr.2018.e00414>
- Kukina IV, Zelya OP, Guzeeva TM, Karan LS, Perkovskaya IA, Tymoshenko NI, Guzeeva MV (2019) Severe babesiosis caused by *Babesia divergens* in a host with intact spleen, Russia, 2018. *Ticks and Tick-borne Diseases* 10(6): 101262. <https://doi.org/10.1016/j.ttbdis.2019.07.006>
- Kulakova NV, Khasnatinov MA, Sidorova EA, Adel'shin RV, Belikov SI (2014) Molecular identification and phylogeny of *Dermacentor nuttalli* (Acari: Ixodidae). *Parasitology Research* 113: 1787–1793. <https://doi.org/10.1007/s00436-014-3824-x>
- Kulkarni GK, Hanumante MM, Nagabhushanam R (1977) Some aspects of osmotic biology of the freshwater leech, *Poecilobdella viridis* (Blanchard). *Hydrobiologia* 56: 103–108. <https://doi.org/10.1007/BF00023346>

- Kutschera U, Elliott J (2014) The European medicinal leech *Hirudo medicinalis* L.: Morphology and occurrence of an endangered species. *Zoosystematics and Evolution* 90(2): 271–280. <https://doi.org/10.3897/zse.90.8715>
- Kváč M, Hofmannová L, Hlásková L, Květoňová D, Vítovec J, McEvoy J, Sak B (2014) *Cryptosporidium erinacei* n. sp. (Apicomplexa: Cryptosporidiidae) in hedgehogs. *Veterinary Parasitology* 201: 9–17. <https://doi.org/10.1016/j.vetpar.2014.01.014>
- Kváč M, Kestřánová M, Pinková M, Květoňová D, Kalinová J, Wagneřová P, Kotková M, Vítovec J, Ditrich O, McEvoy J, Stenger B, Sak B (2013) *Cryptosporidium scrofarum* n. sp. (Apicomplexa: Cryptosporidiidae) in domestic pigs (*Sus scrofa*). *Veterinary Parasitology* 191: 218–227. <https://doi.org/10.1016/j.vetpar.2012.09.005>
- Kvac M, Vlnata G, Jezkova J, Horcickova M, Konecny R, Hlaskova L, McEvoy J, Sak B (2018) *Cryptosporidium occultus* sp. n. (Apicomplexa: Cryptosporidiidae) in rats. *European Journal of Protistology* 63: 96–104. <https://doi.org/10.1016/j.ejop.2018.02.001>
- Kwak ML (2018) The first records of human infestation by the hard tick *Ixodes (Endopalpiger) australiensis* (Acari: Ixodidae), with a review of human infestation by ticks in Australia. *Experimental and Applied Acarology* 74: 185–190. <https://doi.org/10.1007/s1007/s10493-018-0217-3>
- Kwo EH, Lim BL (1968) A new record for a *Paragonimus*-like trematode. *Achillurbainia nouveli* Dollfus, 1939 in West Malaysia. *Medical Journal of Malaysia* 22: 231.
- Kyany'a C, Eyase F, Odundo E, Kipkirui E, Kipkemoi N, Kirera R, Philip C, Ndonge J, Kirui M, Ombogo A, Koech M, Bulimo W, Hulseberg CE (2019) First report of *Entamoeba moshkovskii* in human stool samples from symptomatic and asymptomatic participants in Kenya. *Tropical Diseases, Travel Medicine and Vaccines* 5: 23. <https://doi.org/10.1186/s40794-019-0098-4>
- Lado P, Glon MG, Klompen H (2021) Integrative taxonomy of *Dermacentor variabilis* (Ixodida: Ixodidae) with description of a new species, *Dermacentor similis* n. sp. *Journal of Medical Entomology* 2021: tjab134. <https://doi.org/10.1093/jme/tjab134>
- Lado P, Nava S, Mendoza-Uribe L, Caceres AG, Delgado-de la Mora J, Licona-Enriquez JD, Delgado-de la Mora D, Labruna MB, Durden LA, Allerdice MEJ, Paddock CD, Szabó MPJ, Venzal JM, Guglielmone AA, Beati L (2018) The *Amblyomma maculatum* Koch, 1844 (Acari: Ixodidae) group of ticks: phenotypic plasticity or incipient speciation? *Parasites & Vectors* 11: e610. <https://doi.org/10.1186/s13071-018-3186-9>
- Lai JH, Walsh NM, Pritt BS, Sloan L, Gibson LE, Desormeau L, Haldane DJ (2014) Cutaneous manifestations of a zoonotic *Onchocerca* species in an adult male, acquired in Nova Scotia, Canada. *Journal of Clinical Microbiology* 52: 1768–1770. <https://doi.org/10.1128/JCM.03358-13>
- Lai Y-T, Nakano T, Chen J-H (2011) Three species of land leeches from Taiwan, *Haemadipsa rjukjuana* comb. n., a new record for *Haemadipsa picta* Moore, and an updated description of *Tritetrabdella taiwana* (Oka). *Zookeys* 139: 1–12. <https://doi.org/10.3897/zookeys.139.1711>
- Lai YT (2019) Beyond the epistaxis: voluntary nasal leech (*Dinobdella ferox*) infestation revealed the leech behaviours and the host symptoms through the parasitic period. *Parasitology* 146: 1477–1485. <https://doi.org/10.1017/S0031182019000751>
- Lainson R, Braga RR, De Souza AA, Povoia MM, Ishikawa EA, Silveira FT (1989) *Leishmania (Viannia) shawi* sp. n., a parasite of monkeys, sloths and procyonids in Amazonian Brazil.

- Annales de Parasitologie Humaine et Comparée 64: 200–207. <https://doi.org/10.1051/parasite/1989643200>
- Lainson R, Shaw JJ (1989) *Leishmania (Viannia) naiffi* sp. n., a parasite of the armadillo, *Dasypus novemcinctus* (L.) in Amazonian Brazil. Annales de Parasitologie Humaine et Comparée 64: 3–9. <https://doi.org/10.1051/parasite/19896413>
- Lalremruata A, Magris M, Vivas-Martinez S, Koehler M, Esen M, Kempaiah P, Jeyaraj S, Perkins DJ, Mordmuller B, Metzger WG (2015) Natural infection of *Plasmodium brasilianum* in humans: Man and monkey share quartan malaria parasites in the Venezuelan Amazon. EBioMedicine 2: 1186–1192. <https://doi.org/10.1016/j.ebiom.2015.07.033>
- Lam HH, Subrayan V, Khaw G (2010) Intraocular filarial infection. Eye (London) 24: 1825–1826. <https://doi.org/10.1038/eye.2010.141>
- Lamon C, Greer GJ (1986) Human infection with an anoplocephalid tapeworm of the genus *Mathevotaenia*. American Journal of Tropical Medicine and Hygiene 35: 824–826. <https://doi.org/10.4269/ajtmh.1986.35.824>
- Landero A, Hernandez F, Abasolo MA, Rechy DA, Nunez P (1991) Cerebral sparganosis caused by *Spirometra mansonoides*. Case report. Journal of Neurosurgery 75: 472–474. <https://doi.org/10.3171/jns.1991.75.3.0472>
- Lass-Flörl C, Mayr A (2007) Human Protothecosis. Clinical Microbiology Reviews 20: 230–242. <https://doi.org/10.1128/CMR.00032-06>
- Latif B, Omar E, Heo CC, Othman N, Tappe D (2011) Human pentastomiasis caused by *Armillifer moniliformis* in Malaysian Borneo. American Journal of Tropical Medicine and Hygiene 85: 878–881. <https://doi.org/10.4269/ajtmh.2011.11-0404>
- Laufer AS, Siddall ME, Graf J (2008) Characterization of the digestive-tract microbiota of *Hirudo orientalis*, a european medicinal leech. Applied Environmental Microbiology 74: 6151–6154. <https://doi.org/10.1128/AEM.00795-08>
- Ledee DR, Iovieno A, Miller D, Mandal N, Diaz M, Fell J, Fini ME, Alfonso EC (2009) Molecular identification of t4 and t5 genotypes in isolates from *Acanthamoeba* keratitis patients. Journal of Clinical Microbiology 47: 1458–1462. <https://doi.org/10.1128/JCM.02365-08>
- Lee JK, Moraru GM, Stokes JV, Benton AN, Wills RW, Nabors HP, Smith CL, Lawrence AM, Willeford BV, Varela-Stokes AS (2019) *Amblyomma maculatum*-associated rickettsiae in vector tissues and vertebrate hosts during tick feeding. Experimental and Applied Acarology 77: 187–205. <https://doi.org/10.1007/s10493-019-00343-x>
- Lee KJ, Bae YT, Kim DH, Deung YK, Ryang YS, Im KI, Yong TS (2003) Gordius worm found in a three year old girl's vomitus. YMJ Yonsei Medical Journal 44(3): 557–560. <https://doi.org/10.3349/ymj.2003.44.3.557>
- Lee LM, Wallace RS, Clyde VL, Gendron-Fitzpatrick A, Sibley SD, Stuchin M, Lauck M, O'Connor DH, Nakao M, Lavikainen A, Hoberg EP, Goldberg TL (2016) Definitive hosts of *Versteria* tapeworms (Cestoda: Taeniidae) causing fatal infection in North America. Emerging Infectious Diseases 22: 707–710. <https://doi.org/10.3201/eid2204.151446>
- Lee SH, Chai JY (2001) A review of *Gymnophalloides seoi* (Digenea: Gymnophallidae) and human infections in the Republic of Korea. The Korean Journal of Parasitology 39: 85–118. <https://doi.org/10.3347/kjp.2001.39.2.85>
- Lee SH, Chai JY, Seo BS (1984) Two cases of human infection by adult of *Spirometra erinacei*. The Korean Journal of Parasitology 22: 66–71. <https://doi.org/10.3347/kjp.1984.22.1.66>

- Leelayoova S, Siripattanapipong S, Manomat J, Piyaraj P, Tan-Ariya P, Bualert L, Mungthin M (2017) Leishmaniasis in Thailand: A review of causative agents and situations. *American Journal of Tropical Medicine and Hygiene* 96: 534–542. <https://doi.org/10.4269/ajtmh.16-0604>
- Lehman B, Leal Jr SM, Procop GW, O’Connell E, Shaik J, Nash TE, Nutman TB, Jones S, Braunthal S, Shah SN, Cruise MW, Mukhopadhyay S, Banzon J (2019) Disseminated Metacystode *Versteria* Species Infection in Woman, Pennsylvania, USA(1). *Emerging Infectious Diseases* 25: 1429–1431. <https://doi.org/10.3201/eid2507.190223>
- Leidenberger S, Bostrom S, Wayland MT (2020) Host records and geographical distribution of *Corynosoma magdalenii*, *C. semerme* and *C. strumosum* (Acanthocephala: Polymorphidae). *Biodiversity Data Journal* 8: e50500. <https://doi.org/10.3897/BDJ.8.e50500>
- Leles D, Gardner SL, Reinhard K, Iniguez A, Araujo A (2012) Are *Ascaris lumbricoides* and *Ascaris suum* a single species? *Parasites & Vectors* 5: e42. <https://doi.org/10.1186/1756-3305-5-42>
- Leon LA (1946) Fourth case of human infection with *Agamomermis*. *Revista de Medicina Tropical y Parasitologia* 12: 25–26.
- Lescano AG, Zunt J (2013) Other cestodes: sparganosis, coenurosis and *Taenia crassiceps* cysticercosis. *Handbook of Clinical Neurology* 114: 335–345. <https://doi.org/10.1016/B978-0-444-53490-3.00027-3>
- Levecke B, Dorny P, Vercammen F, Visser LG, Van Esbroeck M, Vercruysse J, Verweij JJ (2015) Transmission of *Entamoeba nuttalli* and *Trichuris trichiura* from Nonhuman Primates to Humans. *Emerging Infectious Diseases* 21: 1871–1872. <https://doi.org/10.3201/eid2110.141456>
- Li CD, Yang HL, Wang Y (2010) *Capillaria hepatica* in China. *World Journal of Gastroenterology* 16: 698–702. <https://doi.org/10.3748/wjg.v16.i6.698>
- Li H, Zhang PH, Du J, Yang ZD, Cui N, Xing B, Zhang XA, Liu W (2019) *Rickettsia japonica* Infections in Humans, Xinyang, China, 2014–2017. *Emerging Infectious Diseases* 25: 1719–1722. <https://doi.org/10.3201/eid2509.171421>
- Li N, Xiao L, Alderisio K, Elwin K, Cebelinski E, Chalmers R, Santin M, Fayer R, Kvac M, Ryan U, Sak B, Stanko M, Guo Y, Wang L, Zhang L, Cai J, Roellig D, Feng Y (2014) Subtyping *Cryptosporidium ubiquitum*, a zoonotic pathogen emerging in humans. *Emerging Infectious Diseases* 20: 217–224. <https://doi.org/10.3201/eid2002.121797>
- Li R, Gao ZC (2016) *Lophomonas blattarum* infection or just the movement of ciliated epithelial cells? *Chinese Medical Journal (Engl)* 129(6): 739–742. <https://doi.org/10.4103/0366-6999.178025>
- Li WC, Wang K, Gu Y (2018) Occurrence of *Blastocystis* sp. and *Pentatrichomonas hominis* in sheep and goats in China. *Parasites & Vectors* 11: e93. <https://doi.org/10.1186/s13071-018-2671-5>
- Li WC, Ying M, Gong PT, Li JH, Yang J, Li H, Zhang XC (2016) *Pentatrichomonas hominis*: prevalence and molecular characterization in humans, dogs, and monkeys in Northern China. *Parasitology Research* 115: 569–574. <https://doi.org/10.1007/s00436-015-4773-8>
- Li X, Li J, Zhang X, Yang Z, Yang J, Gong P (2017) Prevalence of *Pentatrichomonas hominis* infections in six farmed wildlife species in Jilin, China. *Veterinary Parasitology* 244: 160–163. <https://doi.org/10.1016/j.vetpar.2017.07.032>

- Liao EC, Chang KC (2012) Images in clinical medicine. Mites in the external auditory canal. *New England Journal of Medicine* 367: e19. <https://doi.org/10.1056/NEJMicm1010983>
- Liautaud B, Vignier N, Miossec C, Plumelle Y, Kone M, Delta D, Ravel C, Cabié A, Desbois N (2015) First case of visceral leishmaniasis caused by *Leishmania martiniquensis*. *The American Journal of Tropical Medicine and Hygiene* 92: 317–319. <https://doi.org/10.4269/ajtmh.14-0205>
- Libertin CR, Reza M, Peterson JH, Lewis J, Hata DJ (2017) Human *Gongylonema pulchrum* infection: esophageal symptoms and need for prolonged albendazole therapy. *American Journal of Tropical Medicine and Hygiene* 96: 873–875. <https://doi.org/10.4269/ajtmh.16-0852>
- Lidani KCF, Andrade FA, Bavia L, Damasceno FS, Beltrame MH, Messias-Reason IJ, Sandri TL (2019) Chagas Disease: From Discovery to a Worldwide Health Problem. *Frontiers in Public Health* 7: e166. [13 pp.] <https://doi.org/10.3389/fpubh.2019.00166>
- Liew JWK, Bukhari FDM, Jeyaprakasam NK, Phang WK, Vythilingam I, Lau YL (2021) Natural *Plasmodium inui* Infections in Humans and *Anopheles cracens* Mosquito, Malaysia. *Emerging Infectious Diseases* 27: 2700–2703. <https://doi.org/10.3201/eid2710.210412>
- Lima NF, Veggiani Aybar CA, Dantur Juri MJ, Ferreira MU (2016) *Mansonella ozzardi*: a neglected New World filarial nematode. *Pathogens and Global Health* 110: 97–107. <https://doi.org/10.1080/20477724.2016.1190544>
- Little MD (1965) Dermatitis in a human volunteer infected with *Strongyloides* of nutria and raccoon. *American Journal of Tropical Medicine and Hygiene* 14: 1007–1009. <https://doi.org/10.4269/ajtmh.1965.14.1007>
- Liu A, Zhang J, Zhao J, Zhao W, Wang R, Zhang L (2015) The first report of *Cryptosporidium andersoni* in horses with diarrhea and multilocus subtype analysis. *Parasites & Vectors* 8: e483. <https://doi.org/10.1186/s13071-015-1102-0>
- Liu GH, Wu CY, Song HQ, Wei SJ, Xu MJ, Lin RQ, Zhao GH, Huang SY, Zhu XQ (2012) Comparative analyses of the complete mitochondrial genomes of *Ascaris lumbricoides* and *Ascaris suum* from humans and pigs. *Gene* 492: 110–116. <https://doi.org/10.1016/j.gene.2011.10.043>
- Liyanaarachchi DR, Rajakaruna RS, Dikkumbura AW, De Silva A, Rajapakse RP (2015) Ticks (Acarina: Ixodida) infesting five reptile species in Sri Lanka with sixteen new host records. *Zootaxa* 3964: 146–148. <https://doi.org/10.11646/zootaxa.3964.1.11>
- Lombert HAPM, Lukoschus FS, O'Connor BM (1982) The Life-cycle of *Cosmoglyphus inaequalis* Fain & Caceres, 1973, with Comments on the Systematic Position of the Genus: Results of the Namaqualand-Namibia Expedition of the King Léopold III Foundation for the Exploration and Protection of Nature (1980). Institut Royal des Sciences Naturelles de Belgique, Brussels, 17 pp.
- Lopes-Torres EJ, da Silva Pinheiro RH, Rodrigues RAR, Francez LdC, Gonçalves EC, Giese EG (2020) Additional characterization of the adult worm *Mammomonogamus laryngeus* (Railliet, 1899) and the tissue lesions caused by the infection in buffaloes. *Veterinary Parasitology* 283: 109164. <https://doi.org/10.1016/j.vetpar.2020.109164>
- Lopez-Escamilla E, Sanchez-Aguillon F, Alatorre-Fernandez CP, Aguilar-Zapata D, Arroyo-Escalante S, Arellano T, Moncada-Barron D, Romero-Valdovinos M, Martinez-Hernandez

- F, Rodriguez-Zulueta P, Maravilla P (2013) New *Tetratrichomonas* species in two patients with pleural empyema. *Journal of Clinical Microbiology* 51: 3143–3146. <https://doi.org/10.1128/JCM.01056-13>
- Lopez J, Krishnavajhala A, Garcia M, Bermúdez S (2016) Tick-borne relapsing fever spirochetes in the Americas. *Veterinary Sciences* 3(3): e16. [18 pp.] <https://doi.org/10.3390/vetsci3030016>
- Lopez MC, Leon CM, Fonseca J, Reyes P, Moncada L, Olivera MJ, Ramirez JD (2015) Molecular epidemiology of *Entamoeba*: First description of *Entamoeba moshkovskii* in a rural area from Central Colombia. *PLoS ONE* 10: e0140302. <https://doi.org/10.1371/journal.pone.0140302>
- Luce-Fedrow A, Lehman ML, Kelly DJ, Mullins K, Maina AN, Stewart RL, Ge H, John HS, Jiang J, Richards AL (2018) A review of scrub Typhus (*Orientia tsutsugamushi* and related organisms): Then, now, and tomorrow. *Tropical Medicine and Infectious Disease* 3(1): e8. [30 pp.] <https://doi.org/10.3390/tropicalmed3010008>
- Lukes J, Mauricio IL, Schonian G, Dujardin JC, Soteriadou K, Dedet JP, Kuhls K, Tintaya KW, Jirku M, Chocholova E, Haralambous C, Pratlong F, Obornik M, Horak A, Ayala FJ, Miles MA (2007) Evolutionary and geographical history of the *Leishmania donovani* complex with a revision of current taxonomy. *Proceedings of the National Academy of Sciences of the United States of America* 104: 9375–9380. <https://doi.org/10.1073/pnas.0703678104>
- Lv Y, Guo X-G, Jin D-C (2018) Research Progress on *Leptotrombidium deliense*. *The Korean Journal of Parasitology* 56: 313–324. <https://doi.org/10.3347/kjp.2018.56.4.313>
- Lysaght TB, Wooster ME, Jenkins PC, Koniaris LG (2018) Myiasis-induced sepsis: a rare case report of *Wohlfahrtiimonas chitiniclastica* and *Ignatzschineria indica* bacteremia in the continental United States. *Medicine (Baltimore)* 97: e13627. <https://doi.org/10.1097/MD.00000000000013627>
- Ma J, Sun MM, He JJ, Liu GH, Ai L, Chen MX, Zhu XQ (2017) *Fasciolopsis buski* (Digenea: Fasciolidae) from China and India may represent distinct taxa based on mitochondrial and nuclear ribosomal DNA sequences. *Parasit Vectors* 10: e101. <https://doi.org/10.1186/s13071-017-2039-2>
- MacLean JD, Arthur JR, Ward BJ, Gyorkos TW, Curtis MA, Kokoskin E (1996) Common-source outbreak of acute infection due to the North American liver fluke *Metorchis conjunctus*. *Lancet* 347: 154–158. [https://doi.org/10.1016/S0140-6736\(96\)90342-6](https://doi.org/10.1016/S0140-6736(96)90342-6)
- Macnish MG, Ryan UM, Behnke JM, Thompson RCA (2003) Detection of the rodent tapeworm *Rodentolepis* (= *Hymenolepis*) *microstoma* in humans. A new zoonosis? *International Journal of Parasitology* 33: 1079–1085. [https://doi.org/10.1016/S0020-7519\(03\)00137-1](https://doi.org/10.1016/S0020-7519(03)00137-1)
- Maddock DR, Fehn CF (1958) Human ear invasions by adult scarabaeid beetles. *Journal of Economic Entomology* 51: 546–547. <https://doi.org/10.1093/jee/51.4.546>
- Madison-Antenucci S, Kramer LD, Gebhardt LL, Kauffman E (2020) Emerging tick-borne diseases. *Clinical Microbiology Reviews* 33(2): e00083-18. <https://doi.org/10.1128/CMR.00083-18>
- Mairena H, Solano M, Venegas W (1989) Human dermatitis caused by a nymph of *Sebekia*. *American Journal of Tropical Medicine and Hygiene* 41: 352–354. <https://doi.org/10.4269/ajtmh.1989.41.352>
- Mak JW (1987) Epidemiology of lymphatic filariasis. *Ciba Foundation Symposium* 127: 5–14. <https://doi.org/10.1002/9780470513446.ch2>

- Makino T, Mori N, Sugiyama H, Mizawa M, Seki Y, Kagoyama K, Shimizu T (2014) Creeping eruption due to *Spirurina* type X larva. *Lancet* 384(9959): 2082–2082. [https://doi.org/10.1016/S0140-6736\(14\)61644-5](https://doi.org/10.1016/S0140-6736(14)61644-5)
- Makiya K, Tsukamoto M, Horio M, Kuroda Y (1988) [A case report of nasal infestation by the leech, *Dinobdella ferox*]. *Journal of UOEH* 10: 203–209 <https://doi.org/10.7888/juoe.10.203>.
- Makki M, Dupouy-Camet J, Seyed Sajjadi SM, Moravec F, Reza Naddaf S, Mobedi I, Malekafzali H, Rezaeian M, Mohebbali M, Kargar F, Mowlavi G (2017) Human spiruridiasis due to *Physaloptera* spp. (Nematoda: Physalopteridae) in a grave of the Shahr-e Sukhteh archeological site of the Bronze Age (2800–2500 BC) in Iran. *Parasite (Paris, France)* 24: 18–18. <https://doi.org/10.1051/parasite/2017019>
- Mans DRA, Kent AD, Hu RVPF, Schalling HDFH (2017) Epidemiological, biological and clinical aspects of leishmaniasis with special emphasis on Busi Yasi in Suriname. *Journal of Clinical & Experimental Dermatology Research* 8(2): e1000388. [16 pp.] <https://doi.org/10.4172/2155-9554.1000388>
- Mantini C, Souppart L, Noël C, Duong TH, Mornet M, Carroger G, Dupont P, Masseret E, Goustille J, Capron M, Duboucher C, Dei-Cas E, Viscogliosi E (2009) Molecular Characterization of a New *Tetratrichomonas* Species in a Patient with Empyema. *Journal of Clinical Microbiology* 47: 2336–2339. <https://doi.org/10.1128/JCM.00353-09>
- Mariana A, Srinovianti N, Ho TM, Halimaton I, Hatikah A, Shaharudin MH, Rosmaliza I, Wan Ishlah L, Sathananthar KS (2008) Intra-aural ticks (Acari: Mesastigmata: Ixodidae) from human otoacariasis cases in Pahang, Malaysia. *Asian Pacific Journal of Tropical Medicine* 1: 20–24.
- Maritz JM, Land KM, Carlton JM, Hirt RP (2014) What is the importance of zoonotic trichomonads for human health? *Trends in Parasitology* 30: 333–341. <https://doi.org/10.1016/j.pt.2014.05.005>
- Marquez JG, Krafusur ES (2002) Gene flow among geographically diverse housefly populations (*Musca domestica* L.): A worldwide survey of mitochondrial diversity. *Journal of Heredity* 93: 254–259. <https://doi.org/10.1093/jhered/93.4.254>
- Marrone F, Canale DE (2019) Occurrence, distribution and bibliography of the medicinal leech *Hirudo verbana* Carena, 1820 (Hirudinea, Hirudinidae) in Sicily (Italy). *Biogeographia – The Journal of Integrative Biogeography* 34: 33–38. <https://doi.org/10.21426/B634143807>
- Martins TF, Barbieri ARM, Costa FB, Terassini FA, Camargo LMA, Peterka CRL, de C Pacheco R, Dias RA, Nunes PH, Marcili A, Scofield A, Campos AK, Horta MC, Guilloux AGA, Benatti HR, Ramirez DG, Barros-Battesti DM, Labruna MB (2016) Geographical distribution of *Amblyomma cajennense* (sensu lato) ticks (Parasitiformes: Ixodidae) in Brazil, with description of the nymph of *A. cajennense* (sensu stricto). *Parasites & Vectors* 9: e186. <https://doi.org/10.1186/s13071-016-1460-2>
- Marty M, Lemaitre M, Kémoun P, Morrier JJ, Monsarrat P (2017) *Trichomonas tenax* and periodontal diseases: a concise review. *Parasitology* 144: 1417–1425. <https://doi.org/10.1017/S0031182017000701>
- Maruyama SR, de Santana AKM, Takamiya NT, Takahashi TY, Rogerio LA, Oliveira CAB, Milanezi CM, Trombela VA, Cruz AK, Jesus AR, Barreto AS, da Silva AM, Almeida RP, Ribeiro

- JM, Silva JS (2019) Non-*Leishmania* parasite in fatal visceral leishmaniasis-like disease, Brazil. *Emerging Infectious Diseases* 25: 2088–2092. <https://doi.org/10.3201/eid2511.181548>
- Mas-Coma S, Bargues MD, Valero MA (2006) Gastrodiscoidiasis, a plant-borne zoonotic disease caused by the amphistome fluke *Gastrodiscoides hominis* (Trematoda: Gastrodiscidae). *Revista Iberica de Parasitologia* 66: 75–81.
- Mathison BA, Bishop HS, Sanborn CR, Dos Santos Souza S, Bradbury R (2016) *Macracanthorhynchus ingens* infection in an 18-month-old child in Florida: A case report and review of acanthocephaliasis in humans. *Clinical Infectious Diseases* 63: 1357–1359. <https://doi.org/10.1093/cid/ciw543>
- Mathison BA, Bradbury RS, Pritt BS (2021) Medical parasitology taxonomy update, January 2018 to May 2020. *Journal of Clinical Microbiology* 59: e01308–01320. <https://doi.org/10.1128/JCM.01308-20>
- Mathison BA, da Silva AJ (2018) Anisakiasis. In: Ortega Y, Sterling CR (Eds) *Foodborne Parasites*. Springer Nature, New York, 159–174. https://doi.org/10.1007/978-3-319-67664-7_8
- Mathison BA, Pritt BS (2014) Laboratory identification of arthropod ectoparasites. *Clinical Microbiology Reviews* 27: 48–67. <https://doi.org/10.1128/CMR.00008-13>
- Mathison BA, Pritt BS (2015) *Arthropod benchtop reference guide*. College of American Pathologists, Northfield, 82 pp.
- Mathison BA, Pritt BS (2018) Chapter 5: Parasitology. In: Pritt BS (Ed.) *Atlas of Fundamental Infectious Diseases Histopathology: A Guide for Daily Practice*. College of American Pathologists, Northfield, 193–288.
- Mathison BA, Pritt BS (2019) Medical Parasitology Taxonomy Update, 2016–2017. *Journal of Clinical Microbiology* 57(2): e01067–18. <https://doi.org/10.1128/JCM.01067-18>
- Mathison BA, Pritt BS (2020) Correction for Mathison and Pritt, “Medical Parasitology Taxonomy Update, 2016–2017”. *Journal of Clinical Microbiology* 58(7): e00822–20. <https://doi.org/10.1128/JCM.00822-20>
- Mathison BA, Pritt BS (2021) Sleeping with the enemy: Everything you need to know about the biology, clinical significance, and laboratory identification of bed bugs. *Clinical Microbiology Newsletter* 43: 1–7. <https://doi.org/10.1016/j.clinmicnews.2020.12.004>
- Mazlumoglu MR (2018) Ticks in the external auditory canal: A common situation in rural areas. *Journal of Head Neck & Spine Surgery* 2(2): e555581. <https://doi.org/10.19080/JHNSS.2018.02.555581>
- Mazyad SA, Sanad EM, Morsy TA (2001) Two types of scab mites infesting man and sheep in North Sinai. *Journal of the Egyptian Society of Parasitology* 31: 213–222.
- McDonald ME (1981) *Key to Trematodes Reported in Waterfowl*. Vol. 142. Department of the Interior, Fish and Wildlife Service, Resource Publication, Washington DC, 156 pp.
- McGhee RB, Cosgrove WB (1980) Biology and physiology of the lower Trypanosomatidae. *Microbiology Reviews* 44: 140–173. <https://doi.org/10.1128/mr.44.1.140-173.1980>
- McHale B, Callahan RT, Paras KL, Weber M, Kimbrell L, Velázquez-Jiménez Y, McManamon R, Howerth EW, Verocai GG (2020) Sparganosis due to *Spirometra* sp. (Cestoda; Diphyllbothriidae) in captive meerkats (*Suricata suricatta*). *International Journal for Parasitology: Parasites and Wildlife* 13: 186–190. <https://doi.org/10.1016/j.ijpaw.2020.10.005>

- McKelvie P, Reardon K, Bond K, Spratt DM, Gangell A, Zochling J, Daffy J (2013) A further patient with parasitic myositis due to *Haycocknema perplexum*, a rare entity. *Journal of Clinical Neuroscience* 20: 1019–1022. <https://doi.org/10.1016/j.jocn.2012.08.009>
- Meamar AR, Kia EB, Zahabiun F, Jafari-Mehr A, Moghadam A, Sadjjadi SM (2007) The occurrence of severe infections with *Rhabditis axei* in AIDS patients in Iran. *Journal of Helminthology* 81: 351–352. <https://doi.org/10.1017/S0022149X07792301>
- Mediannikov O, Ranque S (2018) Mansonellosis, the most neglected human filariasis. *New Microbes and New Infections* 26(Supplement 1): S19–S22. <https://doi.org/10.1016/j.nmni.2018.08.016>
- Meister L, Ochsendorf F (2016) Head Lice: Epidemiology, Biology, Diagnosis, and Treatment. *Deutsches Ärzteblatt International* 113: 763–772. <https://doi.org/10.3238/arztebl.2016.0763>
- Melo CM, Cruz ACFG, Lima AFVA, Silva LR, Madi RR, Jeraldo VdLS, Mercado R (2018) Triatomine fauna and recent epidemiological dynamics of chagas disease in an endemic area of Northeast Brazil. *Canadian Journal of Infectious Diseases and Medical Microbiology* 2018: e7020541. <https://doi.org/10.1155/2018/7020541>
- Mendez O, Szmulewicz G, Gatta C, Clementel V, Menghi C (2002) *Retortamonas intestinalis*: A little-known parasite. *Acta Bioquímica Clínica Latinoamericana* 36: 357–363.
- Miarinjara A, Rogier C, Harimalala M, Ramihangihajason TR, Boyer S (2016) *Xenopsylla brasiliensis* fleas in plague focus areas, Madagascar. *Emerging Infectious Diseases* 22: 2207–2208. <https://doi.org/10.3201/eid2212.160318>
- Miles MA, Arias JR, Valente SA, Naiff RD, de Souza AA, Pova MM, Lima JA, Cedillos RA (1983) Vertebrate hosts and vectors of *Trypanosoma rangeli* in the Amazon Basin of Brazil. *American Journal of Tropical Medicine and Hygiene* 32: 1251–1259. <https://doi.org/10.4269/ajtmh.1983.32.1251>
- Millan J, Ferroglio E, Solano-Gallego L (2014) Role of wildlife in the epidemiology of *Leishmania infantum* infection in Europe. *Parasitology Research* 113: 2005–2014. <https://doi.org/10.1007/s00436-014-3929-2>
- Miller D (2008) Bed Bugs (Hemiptera: Cimicidae: *Cimex* spp.). In: Capinera JL (Ed.) *Encyclopedia of Entomology*. Springer Netherlands, Dordrecht, 405–417.
- Milman O, Dik B (2017) First report of cheyletiellosis due to the skin mite *Cheyletiella parastivorax* Megnin, 1878 in a human in Turkey. *Elyns Journal of Microbes* 2: 105.
- Miura M, Hayasaka S, Yamada T, Hayasaka Y, Kamimura K (2005) Ophthalmomyiasis caused by larvae of *Boettcherisca peregrina*. *Japanese Journal of Ophthalmology* 49: 177–179. <https://doi.org/10.1007/s10384-004-0155-y>
- Mohammadi S, Lutermann H, Hoffmann S, Emami-Khoyi A, Webster HJ, Fagir D, Bennett NC, van Vuuren BJ (2021) Morphological and Molecular Characterization of the Plague Vector *Xenopsylla brasiliensis*. *Journal of Parasitology* 107: 289–294. <https://doi.org/10.1645/20-44>
- Mohammed N, Smith KGV (1976) Nasopharyngeal myiasis in man caused by larvae of *Clogmia* (= *Telmetoscopus*) *albipunctatus* Williston (Psychodidae, Dipt.). *Transactions of The Royal Society of Tropical Medicine and Hygiene* 70: 91–91. [https://doi.org/10.1016/0035-9203\(76\)90022-5](https://doi.org/10.1016/0035-9203(76)90022-5)

- Möhl K, Große K, Hamedy A, Wüste T, Kabelitz P, Lückner E (2009) Biology of *Alaria* spp. and human exposition risk to *Alaria* mesocercariae – a review. *Parasitology Research* 105: e1. <https://doi.org/10.1007/s00436-009-1444-7>
- Mokhtar AS, Braima KA, Peng Chin H, Jeffery J, Mohd Zain SN, Rohela M, Lau YL, Jamaiah I, Wilson JJ, Abdul-Aziz NM (2016a) Intestinal myiasis in a Malaysian patient caused by larvae of *Clogmia albipunctatus* (Diptera: Psychodidae). *Journal of Medical Entomology* 53: 957–960. <https://doi.org/10.1093/jme/tjw014>
- Mokhtar AS, Sridhar GS, Mahmud R, Jeffery J, Lau YL, Wilson J-J, Abdul-Aziz NM (2016b) First case report of canthariasis in an infant caused by the larvae of *Lasioderma serricorne* (Coleoptera: Anobiidae). *Journal of Medical Entomology* 53: 1234–1237. <https://doi.org/10.1093/jme/tjw071>
- Moles A (1982) Parasite-host records of Alaskan fishes. NOAA Technical Report NMFS SSRF-760. In: Commerce USDo (Ed.) National Oceanic and Atmospheric Administration, Rockdale, 41 pp.
- Montoya JG, Remington JS (2008) Management of *Toxoplasma gondii* infection during pregnancy. *Clinical Infectious Diseases* 47: 554–566. <https://doi.org/10.1086/590149>
- Moore JP (1930) Leeches (Hirudinea) from China with descriptions of new species. *Proceedings of the Academy of Natural Science of Philadelphia* 82: 169–192.
- Moraru GM, Goddard J (2019) The Goddard guide to arthropods of medical importance. CRC Press, Boca Raton, [xx,] 515 pp. <https://doi.org/10.1201/b22250>
- Mordvinov VA, Yurlova NI, Ogorodova LM, Katokhin AV (2012) *Opisthorchis felineus* and *Metorchis bilis* are the main agents of liver fluke infection of humans in Russia. *Parasitology International* 61: 25–31. <https://doi.org/10.1016/j.parint.2011.07.021>
- Morimoto N, Korenaga M, Komatsu C, Sugihara S, Nishida M, Yasuoka M, Kumazawa H, Sasaki M, Hashiguchi Y (1996) A case report of an overseas-traveler's diarrhea probably caused by *Chilomastix mesnili* infection. *Japanese Journal of Tropical Medicine and Hygiene* 24(3): 177–180. <https://doi.org/10.2149/tmh1973.24.177>
- Morimoto N, Korenaga M, Yagyū K, Kagei N, Fujieda M, Bain O, Wakiguchi H, Hashiguchi Y, Sugiura T (2006) Morphological observations and the effects of artificial digestive fluids on the survival of *Diploscapter coronata* from a Japanese patient. *Journal of Helminthology* 80: 341–348. <https://doi.org/10.1017/JOH2006361>
- Moser JC (1975) Biosystematics of the straw itch mite with special reference to nomenclature and dermatology. *Transactions of the Royal Entomological Society of London* 127: 185–191. <https://doi.org/10.1111/j.1365-2311.1975.tb00568.x>
- Mouchet F, Develoux M, Magasa MB (1988) *Schistosoma bovis* in human stools in Republic of Niger. *Transactions of The Royal Society of Tropical Medicine and Hygiene* 82(2): 257–257. [https://doi.org/10.1016/0035-9203\(88\)90438-5](https://doi.org/10.1016/0035-9203(88)90438-5)
- Mueller JF, Strano AJ (1974) *Sparganium proliferum*, a sparganium infected with a virus? *Journal of Parasitology* 60: 15–19. <https://doi.org/10.2307/3278671>
- Mukhtar MM, Eisawi OA, Amanfo SA, Elamin EM, Imam ZS, Osman FM, Hamed ME (2019) *Plasmodium vivax* cerebral malaria in an adult patient in Sudan. *Malaria Journal* 18: e316. <https://doi.org/10.1186/s12936-019-2961-1>
- Mullen GR, Durden LA (2019) *Medical and veterinary entomology*. Elsevier, Cambridge, 792 pp.

- Munasinghe VS, Vella NG, Ellis JT, Windsor PA, Stark D (2013) Cyst formation and faecal-oral transmission of *Dientamoeba fragilis* – the missing link in the life cycle of an emerging pathogen. *International Journal for Parasitology* 43: 879–883. <https://doi.org/10.1016/j.ijpara.2013.06.003>
- Murillo AC, Mullens BA (2017) A review of the biology, ecology, and control of the northern fowl mite, *Ornithonyssus sylviarum* (Acari: Macronyssidae). *Veterinary Parasitology* 246: 30–37. <https://doi.org/10.1016/j.vetpar.2017.09.002>
- Na L, Gao JF, Liu GH, Fu X, Su X, Yue DM, Gao Y, Zhang Y, Wang CR (2016) The complete mitochondrial genome of *Metorchis orientalis* (Trematoda: Opisthorchiidae): Comparison with other closely related species and phylogenetic implications. *Infection, Genetics and Evolution* 39: 45–50. <https://doi.org/10.1016/j.meegid.2016.01.010>
- Nabie R, Spotin A, Rouhani S (2017) Subconjunctival setariasis due to *Setaria equina* infection; a case report and a literature review. *Parasitology International* 66: 930–932. <https://doi.org/10.1016/j.parint.2016.10.017>
- Nakao M, Lavikainen A, Iwaki T, Haukisalme V, Konyaev S, Oku Y, Okamoto M, Ito A (2013) Molecular phylogeny of the genus *Taenia* (Cestoda: Taeniidae): Proposals for the resurrection of *Hydatigera* Lamarck, 1816 and the creation of a new genus *Versteria*. *International Journal of Parasitology* 43: 427–437. <https://doi.org/10.1016/j.ijpara.2012.11.014>
- Nakao M, Okamoto M, Sako Y, Yamasaki H, Nakaya K, Ito A (2002) A phylogenetic hypothesis for the distribution of two genotypes of the pig tapeworm *Taenia solium* worldwide. *Parasitology* 124: 657–662. <https://doi.org/10.1017/S0031182002001725>
- Nakao Y, Tanigawa T, Shibata R (2017) Human otoacariasis caused by *Amblyomma testudinarium*: Diagnosis and management: Case report. *Medicine* 96: e7394. <https://doi.org/10.1097/MD.0000000000007394>
- Nakatsudi K (1942) [One new tick from North Manchukou]. *Journal of the Agricultural Chemical Society of Japan (Tokyo)* 1: 329–332.
- Nakazawa M, Amano T, Oshima T (1992) The first record of human infection with *Diphyllobothrium orcinii* Hatsushika and Shirouzu, 1990. *Japanese Journal of Parasitology* 41: 306–313.
- Nasr M (1941) The occurrence of *Prohemistomum vivax* (Sonsino, 1892) Azim, 1933. *Infection in man, with a redescription of the Parasite. Laboratory and Medical Progress* 2: 135–149.
- Naudé TW, Heyne H, Merwe IR, Benic MJ (2001) Spinose ear tick, *Otobius megnini* (Dugès, 1884) as the cause of an incident of painful otitis externa in humans. *Journal of the South African Veterinary Association* 72: 118–119. <https://doi.org/10.4102/jsava.v72i3.633>
- Neimanis AS, Moraes C, Bergman A, Bignert A, Höglund J, Lundström K, Strömberg A, Bäcklin BM (2016) Emergence of the Zoonotic Biliary Trematode *Pseudamphistomum truncatum* in Grey Seals (*Halichoerus grypus*) in the Baltic Sea. *PLoS ONE* 11: e0164782. <https://doi.org/10.1371/journal.pone.0164782>
- Nelder MP, Russell CB, Sheehan NJ, Sander B, Moore S, Li Y, Johnson S, Patel SN, Sider D (2016) Human pathogens associated with the blacklegged tick *Ixodes scapularis*: a systematic review. *Parasites & Vectors* 9: e265. <https://doi.org/10.1186/s13071-016-1529-y>
- Nemejc K, Sak B, Kvetonova D, Kernerova N, Rost M, Cama VA, Kvac M (2013) Occurrence of *Cryptosporidium suis* and *Cryptosporidium scrofarum* on commercial swine farms in the

- Czech Republic and its associations with age and husbandry practices. *Parasitology Research* 112: 1143–1154. <https://doi.org/10.1007/s00436-012-3244-8>
- Nesemann H, Sharma S (2001) Leeches of the suborder Hirudiniiformes (Hirudinea: Haemopidae, Hirudinidae, Haemadipsidae) from the Ganga watershed (Nepal, India: Bihar). *Annalen des Naturhistorischen Museums in Wien* 103: 77–88.
- Nevill EM, Basson PA, Schoonraad JH, Swanepoel K (1969) A case of nasal myiasis caused by the larvae of *Telmatoscopus albipunctatus* (Williston) 1893 (Diptera: Psychodidae). *South African Medical Journal* 43: 512–514.
- Ng JS, Eastwood K, Walker B, Durrheim DN, Massey PD, Porigneaux P, Kemp R, McKinnon B, Laurie K, Miller D, Bramley E, Ryan U (2012) Evidence of *Cryptosporidium* transmission between cattle and humans in northern New South Wales. *Experimental Parasitology* 130: 437–441. <https://doi.org/10.1016/j.exppara.2012.01.014>
- Ngamprasertwong T, Thirakhupt K, Panha S (2007) Two new species of land leeches from Thailand (Hirudiniiformes: Haemadipsidae). *Tropical Natural History* 7: 155–159.
- Ngobeni R, Samie A, Moonah S, Watanabe K, Petri Jr WA, Gilchrist C (2017) *Entamoeba* species in South Africa: Correlations with the host microbiome, parasite burdens, and first description of *Entamoeba bangladeshi* outside of Asia. *Journal of Infectious Diseases* 216: 1592–1600. <https://doi.org/10.1093/infdis/jix535>
- Nickol BB, Helle E, Valtonen ET (2002) *Corynosoma magdaleni* in gray seals from the Gulf of Bothnia, with emended descriptions of *Corynosoma strumosum* and *Corynosoma magdaleni*. *Journal of Parasitology* 88: 1222–1229. [https://doi.org/10.1645/0022-3395\(2002\)088\[1222:CMIGSF\]2.0.CO;2](https://doi.org/10.1645/0022-3395(2002)088[1222:CMIGSF]2.0.CO;2)
- Nieuwenhuys Ev, Gatti F (1968) A case of chronic mastoid infestation by a rare parasite: (*Poikilorchis congolensis*). *Acta Oto-Laryngologica* 66: 444–448. <https://doi.org/10.3109/00016486809126309>
- Nkouawa A, Haukisalmi V, Li T, Nakao M, Lavikainen A, Chen X, Henttonen H, Ito A (2016) Cryptic diversity in hymenolepidid tapeworms infecting humans. *Parasitology international* 65: 83–86. <https://doi.org/10.1016/j.parint.2015.10.009>
- Nollen PM, Kanev I (1995) The Taxonomy and Biology of Philophthalmid Eye-flukes. In: Baker JR, Muller R, Rollinson D (Eds) *Advances in Parasitology*. Academic Press, 205–269. [https://doi.org/10.1016/S0065-308X\(08\)60492-3](https://doi.org/10.1016/S0065-308X(08)60492-3)
- Nomura Y, Nagakura K, Kagei N, Tsutsumi Y, Araki K, Sugawara M (2000) Gnathostomiasis possibly caused by *Gnathostoma malaysiae*. *Tokai Journal of Experimental and Clinical Medicine* 25: 1–6.
- Ntoukas V, Tappe D, Pfütz D, Simon M, Holzmann T (2013) Cerebellar cysticercosis caused by larval *Taenia crassiceps* tapeworm in immunocompetent woman, Germany. *Emerging Infectious Diseases* 19: 2008–2011. <https://doi.org/10.3201/eid1912.130284>
- Nuprasert W, Putaporntip C, Pariyakanok L, Jongwutiwes S (2010) Identification of a novel t17 genotype of *Acanthamoeba* from environmental isolates and t10 genotype causing keratitis in Thailand. *Journal of Clinical Microbiology* 48: 4636–4640. <https://doi.org/10.1128/JCM.01090-10>
- Oceguera-Figueroa A (2008) A new glossiphoniid leech from Catemaco Lake, Veracruz, Mexico. *Journal of Parasitology* 94: 375–380. <https://doi.org/10.1645/GE-1240.1>

- Oceguera-Figueroa A, León-Règagnon V (2014) [Biodiversity of leeches (Annelida: Euhirudinea) in Mexico]. *Mexican Journal of Biodiversity* 85: 183–189.
- Oda T, Imajima M, Mori A, Fujita K, Tsukidate S, Murata M, Mori S (1984) Infestation of a blood-sucking leech, *Hirudo nipponia* Whitman, in human's eye. *Medical Entomology and Zoology* 35: 319–321. <https://doi.org/10.7601/mez.35.319>
- Odei MA (1966) A note on dicrocoeliasis and *Fasciola gigantica* infection in livestock in Northern Ghana, with a record of spurious and of genuine *Dicrocoelium hospes* infections in man. *Annals of Tropical Medicine and Parasitology* 60: 215–218. <https://doi.org/10.1080/00034983.1966.11686408>
- Oguike MC, Betson M, Burke M, Nolder D, Stothard JR, Kleinschmidt I, Proietti C, Bousema T, Ndounga M, Tanabe K, Ntege E, Culleton R, Sutherland CJ (2011) *Plasmodium ovale curtisi* and *Plasmodium ovale wallikeri* circulate simultaneously in African communities. *International Journal of Parasitology* 41: 677–683. <https://doi.org/10.1016/j.ijpara.2011.01.004>
- Okada JK (1927) A new case of digestive tube myiasis caused by the larva of *Psychoda 6-punctata* Curt. *Annales de Parasitologie Humaine et Comparee (Paris)* 5: 105–106. <https://doi.org/10.1051/parasite/1927052105>
- Okono T, Ushirogawa H, Matoba K, Hatsushika R (2010) Bibliographical studies on human cases 487 of hard tick (Acarina: Ixodidae) bites in Japan (7) cases of unidentified tick infestation. *Kawasaki Medical Journal* 39: 127–141.
- Oleaga A, Rey O, Polack B, Grech-Angelini S, Quilichini Y, Perez-Sanchez R, Boireau P, Mulero S, Brunet A, Rognon A, Vallee I, Kincaid-Smith J, Allienne JF, Boissier J (2019) Epidemiological surveillance of schistosomiasis outbreak in Corsica (France): Are animal reservoir hosts implicated in local transmission? *PLoS Neglected Tropical Diseases* 13: e0007543. <https://doi.org/10.1371/journal.pntd.0007543>
- Onyiche TE, Okute TO, Oseni OS, Okoro DO, Biu AA, Mbaya AW (2018) Parasitic and zoonotic meningoencephalitis in humans and equids: Current knowledge and the role of *Halicephalobus gingivalis*. *Parasite Epidemiology and Control* 3: 36–42. <https://doi.org/10.1016/j.parepi.2017.12.002>
- Ord RL, Lobo CA (2015) Human babesiosis: pathogens, prevalence, diagnosis and treatment. *Current Clinical Microbiology Reports* 2: 173–181. <https://doi.org/10.1007/s40588-015-0025-z>
- Orihel TC, Ash LR (1995) Parasites in human tissues. ASCP Press, Chicago, [xxi,] 386 pp.
- Orihel TC, Eberhard ML (1998) Zoonotic filariasis. *Clinical Microbiology Reviews* 11: 366–381. <https://doi.org/10.1128/CMR.11.2.366>
- Orosz E, Szentmary N, Kiss HJ, Farkas A, Kucsera I, Nagy ZZ (2018) First report of *Acanthamoeba* genotype T8 human keratitis. *Acta Microbiologica et Immunologica Hungarica* 65: 73–79. <https://doi.org/10.1556/030.65.2018.007>
- Ortega YR, Sanchez R (2010) Update on *Cyclospora cayetanensis*, a food-borne and waterborne parasite. *Clinical Microbiology Reviews* 23: 218–234. <https://doi.org/10.1128/CMR.00026-09>
- Osava CF, Ramos VDN, Rodrigues AC, Dos Reis Neto HV, Martins MM, Pascoal JO, Yokosawa J, Szabo MPJ (2016) *Amblyomma sculptum* (*Amblyomma cajennense* complex) tick population maintained solely by domestic pigs. *Veterinary Parasitology: Regional Studies and Reports* 6: 9–13. <https://doi.org/10.1016/j.vprsr.2016.11.002>

- Otranto D, Dutto M (2008) Human thelaziasis, Europe. *Emerging Infectious Diseases* 14: 647–649. <https://doi.org/10.3201/eid1404.071205>
- Otranto D, Eberhard ML (2011) Zoonotic helminths affecting the human eye. *Parasites & Vectors* 4: e41. <https://doi.org/10.1186/1756-3305-4-41>
- Oxford University Press (2020) Oxford English and Spanish Dictionary, Thesaurus, and Spanish to English Translator. <https://www.lexico.com/> [accessed July 10, 2020]
- Padgett KA, Bonilla D, Eremeeva ME, Glaser C, Lane RS, Porse CC, Castro MB, Messenger S, Espinosa A, Hacker J, Kjemtrup A, Ryan B, Scott JJ, Hu R, Yoshimizu MH, Dasch GA, Kramer V (2016) The Eco-epidemiology of Pacific Coast Tick Fever in California. *PLoS Neglected Tropical Diseases* 10: e0005020. [17 pp.] <https://doi.org/10.1371/journal.pntd.0005020>
- Padgett KA, Boyce WM (2004) Life-history studies on two molecular strains of mesocestoides (Cestoda: Mesocestoididae): identification of sylvatic hosts and infectivity of immature life stages. *Journal of Parasitology* 90(1): 108–113. <https://doi.org/10.1645/GE-100R1>
- Padgett KA, Lane RS (2001) Life cycle of *Ixodes pacificus* (Acari: Ixodidae): Timing of developmental processes under field and laboratory conditions. *Journal of Medical Entomology* 38: 684–693. <https://doi.org/10.1603/0022-2585-38.5.684>
- Page W, Judd JA, Bradbury RS (2018) The unique life cycle of *Strongyloides stercoralis* and Implications for public health action. *Tropical Medicine and Infectious Disease* 3(2): e53. <https://doi.org/10.3390/tropicalmed3020053>
- Pakharukova MY, Mordvinov VA (2016) The liver fluke *Opisthorchis felineus*: biology, epidemiology and carcinogenic potential. *Transactions of The Royal Society of Tropical Medicine and Hygiene* 110: 28–36. <https://doi.org/10.1093/trstmh/trv085>
- Pal S, Negi V, Bisht R, Juyal D (2018) Bite of a mite: A case of human otoacariasis caused by *Cosmoglyphus* species (Acari: Acaridae). *Journal of Clinical and Diagnostic Research* 12(3): DD03–DD05. <https://doi.org/10.7860/JCDR/2018/34277.11274>
- Paleri V, Ruckley RW (2001) Recurrent infestation of the mastoid cavity with *Caloglyphus berlesei*: an occupational hazard. *The Journal of Laryngology and Otology* 115: 652–653. <https://doi.org/10.1258/0022215011908513>
- Palmer ED (1946) Intestinal canthariasis due to *Tenebrio molitor*. *Journal of Parasitology* 32(1): 54–55. <https://doi.org/10.2307/3272703>
- Panaitescu D, Preda A, Bain O, Vasile-Bugarin AC (1999) Four cases of human filariasis due to *Setaria labiatopapillosa* found in Bucharest, Romania. *Roumanian Archives of Microbiology and Immunology* 58: 203–207.
- Papadi B, Boudreaux C, Tucker JA, Mathison B, Bishop H, Eberhard ME (2013) *Halicephalobus gingivalis*: a rare cause of fatal meningoencephalomyelitis in humans. *American Journal of Tropical Medicine and Hygiene* 88: 1062–1064. <https://doi.org/10.4269/ajtmh.12-0730>
- Papini RA, Lubas G, Sgorbini M (2020) Incidental detection of *Onchocerca* Microfilariae in donkeys (*Equus asinus*) in Italy: Report of four cases. *Frontiers in Veterinary Science* 7: e569916. <https://doi.org/10.3389/fvets.2020.569916>
- Parker ID, Lopez RR, Silvy NJ, Pierce BL, Watts KG, Myers EP, Gibbs SEJ, Davis DS, Beaver JT, Lund AA (2020) Florida key deer abundance and recovery following new world screwworm infestation. *Southeastern Naturalist* 19(2): 179–191. <https://doi.org/10.1656/058.019.0201>

- Pattullo KM, Wobeser G, Lockerbie BP, Burgess HJ (2013) *Babesia odocoilei* infection in a Saskatchewan elk (*Cervus elaphus canadensis*) herd. *Journal of Veterinary Diagnostic Investigation* 25: 535–540. <https://doi.org/10.1177/1040638713491746>
- Paul J (2007) *Scarites sulcatus* Olivier (Carabidae) as an agent of genital canthariasis. *The Coleopterist* 16: 34–34.
- Peckensneider LE, Pokorny C, Hellwig CA (1952) Intestinal infestation with maggots of the cheese fly (*Piophilha casei*). *Journal of the American Medical Association* 149: 262–263. <https://doi.org/10.1001/jama.1952.72930200005011b>
- Pérez-Bañón C, Rojas C, Vargas M, Mengual X, Rojo S (2020) A world review of reported myiasis caused by flower flies (Diptera: Syrphidae), including the first case of human myiasis from *Palpada scutellaris* (Fabricius, 1805). *Parasitology Research* 119: 815–840. <https://doi.org/10.1007/s00436-020-06616-4>
- Perry BD, Lessard P, Norval RA, Kundert K, Kruska R (1990) Climate, vegetation and the distribution of *Rhipicephalus appendiculatus* in Africa. *Parasitology Today* 6: 100–104. [https://doi.org/10.1016/0169-4758\(90\)90224-R](https://doi.org/10.1016/0169-4758(90)90224-R)
- Perry RD, Fetherston JD (1997) *Yersinia pestis* – etiologic agent of plague. *Clinical Microbiology Reviews* 10: 35–66. <https://doi.org/10.1128/CMR.10.1.35>
- Pezzi M, Cultrera R, Chicca M, Leis M (2015) Furuncular myiasis caused by *Cordylobia rodhaini* (Diptera: Calliphoridae): A case report and a literature review. *Journal of Medical Entomology* 52: 151–155. <https://doi.org/10.1093/jme/tju027>
- Phalee A, Wongsawad C, Rojanapaibul A, Chai JY (2015) Experimental life history and biological characteristics of *Fasciola gigantica* (Digenea: Fasciolidae). *The Korean Journal of Parasitology* 53: 59–64. <https://doi.org/10.3347/kjp.2015.53.1.59>
- Phillips AJ, Arauco-Brown R, Ocegüera-Figueroa A, Gomez GP, Beltrán M, Lai Y-T, Siddall ME (2010) *Tyrannobdella rex* g. gen. n. sp. and the evolutionary origins of mucosal leech infestations. *PLoS ONE* 5: e10057. <https://doi.org/10.1371/journal.pone.0010057>
- Phillips AJ, Salas-Montiel R, Kvist S, Ocegüera-Figueroa A (2019) Phylogenetic position and description of a new species of medicinal leech from the Eastern United States. *Journal of Parasitology* 105: 587–597. <https://doi.org/10.1645/18-119>
- Phillips AJ, Salas-Montiel R, Ocegüera-Figueroa A (2016) Distribution of the New England medicinal leech, *Macrobdella sestertia* Whitman, 1886 and redeterminations of specimens of *Macrobdella* (Annelida: Clitellata: Macrobdellidae) at the National Museum of Natural History, Smithsonian Institution. *Proceedings of the Biological Society of Washington* 129: 103–113[111]. <http://dx.doi.org/10.2988/0006-324X-129.Q2.103>
- Polderman AM, Blotkamp J (1995) *Oesophagostomum* infections in humans. *Parasitology Today* 11: 451–456. [https://doi.org/10.1016/0169-4758\(95\)80058-1](https://doi.org/10.1016/0169-4758(95)80058-1)
- Pombi M, Giacomi A, Barlozzari G, Mendoza-Roldan J, Macri G, Otranto D, Gabrielli S (2020) Molecular detection of *Leishmania (Sauroleishmania) tarentolae* in human blood and *Leishmania (Leishmania) infantum* in *Sergentomyia minuta*: unexpected host-parasite contacts. *Medical and Veterinary Entomology* 34: 470–475. <https://doi.org/10.1111/mve.12464>
- Pothirat T, Tantiworawit A, Chaiwarith R, Jariyapan N, Wannasan A, Siriyasatien P, Supparatpinyo K, Bates MD, Kwakye-Nuako G, Bates PA (2014) First isolation of *Leishmania* from Northern Thailand: case report, identification as *Leishmania martiniquensis* and phyloge-

- netic position within the *Leishmania enriettii* complex. PLoS Neglected Tropical Diseases 8: e3339. <https://doi.org/10.1371/journal.pntd.0003339>
- Poulsen CS, Stensvold CR (2016) Systematic review on *Endolimax nana*: A less well studied intestinal ameba. Tropical Parasitology 6: 8–29. <https://doi.org/10.4103/2229-5070.175077>
- Powar RM, Shegokar VR, Joshi PP, Dani VS, Tankhiwale NS, Truc P, Jannin J, Bhargava A (2006) A rare case of human trypanosomiasis caused by *Trypanosoma evansi*. Indian Journal of Medical Microbiology 24: 72–74. <https://doi.org/10.4103/0255-0857.19904>
- Powell RF, Palmer SM, Palmer CH, Smith EB (1977) *Cheyletiella* dermatitis. International Journal of Dermatology 16: 679–682. <https://doi.org/10.1111/j.1365-4362.1977.tb01882.x>
- Pratlong F, Lami P, Ravel C, Balard Y, Dereure J, Serres G, Baidouri FE, Dedet JP (2013) Geographical distribution and epidemiological features of Old World *Leishmania infantum* and *Leishmania donovani* foci, based on the isoenzyme analysis of 2277 strains. Parasitology 140: 423–434. <https://doi.org/10.1017/S0031182012001825>
- Procop GW (2009) North American paragonimiasis (Caused by *Paragonimus kellicotti*) in the context of global paragonimiasis. Clinical Microbiology Reviews 22: 415–446. <https://doi.org/10.1128/CMR.00005-08>
- Prüter H, Sitko J, Krone O (2017) Having bird schistosomes in mind—the first detection of *Bilharziella polonica* (Kowalewski 1895) in the bird neural system. Parasitology Research 116: 865–870. <https://doi.org/10.1007/s00436-016-5359-9>
- Puleston RL, Mallaghan CM, Modha DE, Hunter PR, Nguyen-Van-Tam JS, Regan CM, Nichols GL, Chalmers RM (2014) The first recorded outbreak of cryptosporidiosis due to *Cryptosporidium cuniculus* (formerly rabbit genotype), following a water quality incident. Journal of Water and Health 12: 41–50. <https://doi.org/10.2166/wh.2013.097>
- Qian M-B, Chen Y-D, Liang S, Yang G-J, Zhou X-N (2012) The global epidemiology of clonorchiasis and its relation with cholangiocarcinoma. Infectious Diseases of Poverty 1: e4. <https://doi.org/10.1186/2049-9957-1-4>
- Qiu MH, Ma KC, Fan PC, Lu SS (2005) [Discovery of a new species of the pentastomid genus *Porocephalus* (Humboldt, 1811) from Taiwan, China and its pathogenic features]. Zhongguo Ji Sheng Chong Xue Yu Ji Sheng Chong Bing Za Zhi 23: 69–72.
- Qvarnstrom Y, da Silva AJ, Schuster FL, Gelman BB, Visvesvara GS (2009) Molecular confirmation of *Sappinia pedata* as a causative agent of amoebic encephalitis. Journal of Infectious Diseases 199: 1139–1142. <https://doi.org/10.1086/597473>
- Ramírez-Hernández A, Uchoa F, Serpa MCdA, Binder LC, Souza CE, Labruna MB (2020) Capybaras (*Hydrochoerus hydrochaeris*) as amplifying hosts of *Rickettsia rickettsii* to *Amblyomma sculptum* ticks: Evaluation during primary and subsequent exposures to *R. rickettsii* infection. Ticks and Tick-borne Diseases 11(5): 101463. <https://doi.org/10.1016/j.ttbdis.2020.101463>
- Ramos JM, Jado I, Padilla S, Masia M, Anda P, Gutierrez F (2013) Human infection with *Rickettsia sibirica mongolitimonae*, Spain, 2007–2011. Emerging Infectious Diseases 19: 267–269. <https://doi.org/10.3201/eid1902.111706>
- Raskova V, Kvetonova D, Sak B, McEvoy J, Edwinston A, Stenger B, Kvac M (2013) Human cryptosporidiosis caused by *Cryptosporidium tyzzeri* and *C. parvum* isolates presumably transmitted from wild mice. Journal of Clinical Microbiology 51: 360–362. <https://doi.org/10.1128/JCM.02346-12>

- Rather PA, Hassan I (2014) Human demodex mite: the versatile mite of dermatological importance. *Indian Journal of Dermatology* 59: 60–66. <https://doi.org/10.4103/0019-5154.123498>
- Rausch RL, Scott EM, Rausch VR (1967) Helminths in eskimos in Western Alaska, with particular reference to *Diphyllobothrium* infection and anaemia. *Transactions of The Royal Society of Tropical Medicine and Hygiene* 61: 351–357. [https://doi.org/10.1016/0035-9203\(67\)90008-9](https://doi.org/10.1016/0035-9203(67)90008-9)
- Ready PD (2014) Epidemiology of visceral leishmaniasis. *Clinical Epidemiology* 6: 147–154. <https://doi.org/10.2147/CLEP.S44267>
- Reisenman CE, Lawrence G, Guerenstein PG, Gregory T, Dotson E, Hildebrand JG (2010) Infection of kissing bugs with *Trypanosoma cruzi*, Tucson, Arizona, USA. *Emerging Infectious Diseases* 16: 400–405. <https://doi.org/10.3201/eid1603.090648>
- Ribeiro G, dos Santos CGS, Lanza F, Reis J, Vaccarezza F, Diniz C, Miranda DLP, de Araújo RF, Cunha GM, de Carvalho CMM, Fonseca EOL, dos Santos RF, de Sousa OMF, Reis RB, de Araújo WN, Gurgel-Gonçalves R, dos Reis MG (2019) Wide distribution of *Trypanosoma cruzi*-infected triatomines in the State of Bahia, Brazil. *Parasites & Vectors* 12: e604. <https://doi.org/10.1186/s13071-019-3849-1>
- Richardson DJ, Leveille A, Belsare AV, Al-Warid HS, Gompper ME (2017) Geographic Distribution Records of *Macracanthorhynchus ingens* (Archiacanthocephala: Oligacanthorhynchidae) from the Raccoon, *Procyon lotor* in North America. *Journal of the Arkansas Academy of Science* 71: 203–205.
- Riley WA (1919) A mouse oxyurid, *Syphacia obvelata*, as a parasite of man. *Journal of Parasitology* 6: 89–93. <https://doi.org/10.2307/3270899>
- Rivas AK, Alcover M, Martínez-Orellana P, Montserrat-Sangrà S, Nachum-Biala Y, Bardagí M, Fisa R, Riera C, Baneth G, Solano-Gallego L (2018) Clinical and diagnostic aspects of feline cutaneous leishmaniasis in Venezuela. *Parasites & Vectors* 11: e141. <https://doi.org/10.1186/s13071-018-2747-2>
- Rizzoli A, Silaghi C, Obiegala A, Rudolf I, Hubalek Z, Foldvari G, Plantard O, Vayssier-Tausat M, Bonnet S, Spitalska E, Kazimirova M (2014) *Ixodes ricinus* and its transmitted pathogens in urban and peri-urban areas in Europe: New hazards and relevance for public health. *Frontiers in Public Health* 2: e251. <https://doi.org/10.3389/fpubh.2014.00251>
- Robert-Gangneux F, Darde ML (2012) Epidemiology of and diagnostic strategies for toxoplasmosis. *Clinical Microbiology Reviews* 25: 264–296. <https://doi.org/10.1128/CMR.05013-11>
- Roberts FHS (1970) *Australian Ticks*. CSIRO, Melbourne, 267 pp.
- Rodpai R, Intapan PM, Thanchomngang T, Sanpool O, Sadaow L, Laymanivong S, Aung WP, Phosuk I, Laummaunwai P, Maleewong W (2016) *Angiostrongylus cantonensis* and *A. malaysiensis* broadly overlap in Thailand, Lao PDR, Cambodia and Myanmar: A molecular survey of larvae in land snails. *PLoS ONE* 11: e0161128. <https://doi.org/10.1371/journal.pone.0161128>
- Rodriguez-Morales AJ, Vera-Ospina JJ, Berthel-Vergara JM, Silvera-Arenas LA, Villamil-Gomez WE (2018) Accidental ulcer infestation due to *Tenebrio molitor* in an AIDS patient: canthariasis. *International Journal of Infectious Diseases* 73: 253–254. <https://doi.org/10.1016/j.ijid.2018.04.3993>
- Romero-Alegria A, Behllassen-Garcia M, Velasco-Tirado V, Garcia-Kingo A, Alvela-Suarez L, Pardo-Lledias J, Sanchez MC (2014) *Angiostrongylus costaricensis*: systematic review of case reports. *Advances in Infectious Diseases* 4: 36–41. <https://doi.org/10.4236/aid.2014.41007>

- Romig T, Ebi D, Wassermann M (2015) Taxonomy and molecular epidemiology of *Echinococcus granulosus* sensu lato. *Veterinary Parasitology* 213: 76–84. <https://doi.org/10.1016/j.vetpar.2015.07.035>
- Rosen S, Yeruham I, Braverman Y (2002) Dermatitis in humans associated with the mites *Pyemotes tritici*, *Dermanyssus gallinae*, *Ornithonyssus bacoti* and *Androlaelaps casalis* in Israel. *Medical and Veterinary Entomology* 16: 442–444. <https://doi.org/10.1046/j.1365-2915.2002.00386.x>
- Rossiter A (1997) Occupational otitis externa in chicken catchers. *The Journal of laryngology and Otology* 111: 366–367. <https://doi.org/10.1017/S0022215100137338>
- Rowan DJ, Said S, Schuetz AN, Pritt BS (2020) A case of *Cystoisospora (Isospora) belli* infection with multiple life stages identified on endoscopic small bowel biopsies. *International Journal of Surgical Pathology* 28: 884–886. <https://doi.org/10.1177/1066896920901589>
- Rubel F, Brugger K, Pfeffer M, Chitimia-Dobler L, Didyk YM, Leverenz S, Dautel H, Kahl O (2016) Geographical distribution of *Dermacentor marginatus* and *Dermacentor reticulatus* in Europe. *Ticks and Tick-borne Diseases* 7: 224–233. <https://doi.org/10.1016/j.ttbdis.2015.10.015>
- Russell RC (2001) The medical significance of Acari in Australia. In: Halliday RB, Walter DE, Proctor HC, Norton RA, Colloff MJ (Eds) *Acarology Proceedings of the 10th International Congress*. CSIRO, Melbourne, 535–546.
- Ryan U, Caccio SM (2013) Zoonotic potential of *Giardia*. *International Journal of Parasitology* 43: 943–956. <https://doi.org/10.1016/j.ijpara.2013.06.001>
- Ryan U, Zahedi A, Papparini A (2016) *Cryptosporidium* in humans and animals—a one health approach to prophylaxis. *Parasite Immunology* 38: 535–547. <https://doi.org/10.1111/pim.12350>
- Ryan UM, Monis P, Enemark HL, Sulaiman I, Samarasinghe B, Read C, Buddle R, Robertson I, Zhou L, Thompson RC, Xiao L (2004) *Cryptosporidium suis* n. sp. (Apicomplexa: Cryptosporidiidae) in pigs (*Sus scrofa*). *Journal of Parasitology* 90: 769–773. <https://doi.org/10.1645/GE-202R1>
- Ryan UM, Power M, Xiao L (2008) *Cryptosporidium fayeri* n. sp. (Apicomplexa: Cryptosporidiidae) from the Red Kangaroo (*Macropus rufus*). *Journal of Eukaryotic Microbiology* 55: 22–26. <https://doi.org/10.1111/j.1550-7408.2007.00299.x>
- Saari SA, Nikander SE (2006) *Pelodera* (syn. *Rhabditis*) *strongyloides* as a cause of dermatitis – a report of 11 dogs from Finland. *Acta Veterinaria Scandinavica* 48: e18. <https://doi.org/10.1186/1751-0147-48-18>
- Sage KM, Johnson TL, Teglas MB, Nieto NC, Schwan TG (2017) Ecological niche modeling and distribution of *Ornithodoros hermsi* associated with tick-borne relapsing fever in western North America. *PLoS Neglected Tropical Diseases* 11: e0006047. <https://doi.org/10.1371/journal.pntd.0006047>
- Sah R, Calatri M, Toledo R (2019) An autochthonous human case of fasciolopsiasis in Nepal. *The Korean Journal of Parasitology* 57: 295–298. <https://doi.org/10.3347/kjp.2019.57.3.295>
- Sah R, Khadka S, Adhikari M, Niraula R, Shah A, Khatri A, Donovan S (2018) Human thelaziasis: emerging ocular pathogen in Nepal. *Open Forum Infectious Diseases* 5(10): ofy237. <https://doi.org/10.1093/ofid/ofy237>

- Saini R, Pui JC, Burgin S (2004) Rickettsialpox: report of three cases and a review. *Journal of the American Academy of Dermatology* 51: S137–S142. <https://doi.org/10.1016/j.jaad.2004.03.036>
- Saksirisampant W, Eamudomkarn C, Jeon H-K, Eom KS, Assavapongpaiboon B, Sintuwong S, Tulvatana W (2020) Ocular sparganosis: The first report of *Spirometra ranarum* in Thailand. *The Korean Journal of Parasitology* 58: 577–581. <https://doi.org/10.3347/kjp.2020.58.5.577>
- Saleh MS, el Sibae MM (1993) Urino-genital myiasis due to *Piophilha casei*. *Journal of the Egyptian Society of Parasitology* 23: 737–739.
- Sambon LW (1922) A synopsis of the family Linguatulidae [part I]. *Journal of Tropical Medicine and Hygiene* 25: 188–206.
- Sánchez-Montes S, Fernández-Figueroa E, González-Guzmán S, Cervantes VP, Ballados-González GG, Rangel-Escareño C, Cárdenas-Ovando RA, Becker I (2020) New records of *Haemaphysalis leporispalustris* in the Transmexican Volcanic Belt province of Mexico with detection of rickettsial infection. *Parasitology Research* 119: 1969–1973. <https://doi.org/10.1007/s00436-020-06684-6>
- Sandground JH (1925) Observations of *Rhabditis hominis* Kobayashi in the United States. *Journal of Parasitology* 11: 140–148. <https://doi.org/10.2307/3270989>
- Sapp SGH, Bradbury RS (2020) The forgotten exotic tapeworms: a review of uncommon zoonotic Cyclophyllidae. *Parasitology* 147: 533–558. <https://doi.org/10.1017/S003118202000013X>
- Saracho-Bottero MN, Tarragona EL, Sebastian PS, Venzal JM, Mangold AJ, Guglielmo AA, Nava S (2018) Ticks infesting cattle and humans in the Yungas Biogeographic Province of Argentina, with notes on the presence of tick-borne bacteria. *Experimental and Applied Acarology* 74: 107–116. <https://doi.org/10.1007/s10493-018-0208-4>
- Sargeant PG, Patrick S, O’Keeffe D (1992) Human infections of *Entamoeba chattoni* masquerade as *Entamoeba histolytica*. *Transactions of the Royal Society of Tropical Medicine and Hygiene* 86: 633–634. [https://doi.org/10.1016/0035-9203\(92\)90162-6](https://doi.org/10.1016/0035-9203(92)90162-6)
- Sargison N, Herman J, Pilkington J, Buckland P, Watt K, Chambers A, Chaudhry U (2018) Molecular confirmation of *Hymenolepis hibernia* in field mice (*Apodemus sylvaticus*) from St Kilda has potential to resolve a host-parasite relationship. *International Journal for Parasitology Parasites and Wildlife* 7: 364–368. <https://doi.org/10.1016/j.ijppaw.2018.09.007>
- Sarkar S, Bhattacharya P (2008) Cerebral malaria caused by *Plasmodium vivax* in adult subjects. *Indian Journal of Critical Care Medicine* 12: 204–205. <https://doi.org/10.4103/0972-5229.45084>
- Sataeva TP, Kutya SA, Smirnova SN, Kakakova VV (2018) A historical review of the study on biology of the dwarf tapeworm *Hymenolepis nana*. *Rossiiskii Parazitologicheskii Shurnal* 2018: 18–26. <https://doi.org/10.31016/1998-8435-2018-12-1-18-26>
- Sato M, Yoonuan T, Sanguankiat S, Nuamtanong S, Pongvongsa T, Phimmayoi I, Phanhanan V, Boupna B, Moji K, Waikagul J (2011) Short report: Human *Trichostrongylus colubri-formis* infection in a rural village in Laos. *American Journal of Tropical Medicine and Hygiene* 84: 52–54. <https://doi.org/10.4269/ajtmh.2011.10-0385>
- Sato Y, Mekata H, Sudaryatma PE, Kirino Y, Yamamoto S, Ando S, Sugimoto T, Okabayashi T (2021) Isolation of severe fever with thrombocytopenia syndrome virus from various tick

- species in area with human severe fever with thrombocytopenia syndrome cases. *Vector-Borne Zoonotic Diseases* 21(5): 378–384. <https://doi.org/10.1089/vbz.2020.2720>
- Sawyer R, Lepont F, Stuart D, Kramer A (1981) Growth and reproduction of the giant glosiphoniid Leech *Haementeria ghilianii*. *Biological Bulletin* 160: 322–331. <https://doi.org/10.2307/1540892>
- Scheid PL (2019) *Vermamoeba vermiformis* – a free-living amoeba with public health and environmental health significance. *The Open Parasitology Journal* 7: 40–47. <https://doi.org/10.2174/1874421401907010040>
- Scheid PL, Lam TT, Sinsch U, Balczun C (2019) *Vermamoeba vermiformis* as etiological agent of a painful ulcer close to the eye. *Parasitology Research* 118: 1999–2004. <https://doi.org/10.1007/s00436-019-06312-y>
- Schmidt GD (1969) *Paracanthocephalus rauschi* sp. n. (Acanthocephala, Paracanthocephalidae) from grayling, *Thymallus arcticus* (Pallas), in Alaska. *Canadian Journal of Zoology* 47: 383–385. <https://doi.org/10.1139/z69-071>
- Schmidt GD (1971) Acanthocephalan infections of man, with two new records. *Journal of Parasitology* 57: 582–584. <https://doi.org/10.2307/3277920>
- Schneider-Crease I, Griffin RH, Gomery MA, Dorny P, Noh JC, Handali S, Chastain HM, Wilkins PP, Nunn CL, Snyder-Mackler N, Beehner JC, Bergman TJ (2017) Identifying wildlife reservoirs of neglected taeniid tapeworms: Non-invasive diagnosis of endemic *Taenia serialis* infection in a wild primate population. *PLoS Neglected Tropical Diseases* 11: e0005709. [18 pp.] <https://doi.org/10.1371/journal.pntd.0005709>
- Schöler A, Maier WA, Kampen H (2006) Multiple environmental factor analysis in habitats of the harvest mite *Neotrombicula autumnalis* (Acari: Trombiculidae) suggests extraordinarily high euryoecious biology. *Experimental and Applied Acarology* 39: 41–62. <https://doi.org/10.1007/s10493-006-0025-z>
- Scholz T, Garcia HH, Kuchta R, Wicht B (2009) Update on the human broad tapeworm (genus *Diphyllobothrium*), including clinical relevance. *Clinical Microbiology Reviews* 22: 146–160. [Table of Contents] <https://doi.org/10.1128/CMR.00033-08>
- Scholz T, Kuchta R (2016) Fish-borne, zoonotic cestodes (*Diphyllobothrium* and relatives) in cold climates: A never-ending story of neglected and (re)-emergent parasites. *Food and Waterborne Parasitology* 4: 23–28. <https://doi.org/10.1016/j.fawpar.2016.07.002>
- Schuster FL, Ramirez-Avila L (2008) Current world status of *Balantidium coli*. *Clinical Microbiology Reviews* 21: 626–638. <https://doi.org/10.1128/CMR.00021-08>
- Schultz H (1975) Human infestation by *Ophionyssus natricis* snake mite. *British Journal of Dermatology* 93: 695–697. <https://doi.org/10.1111/j.1365-2133.1975.tb05120.x>
- Schwan TG, Corwin MD, Brown SJ (1992) *Argas (Argas) monolakensis*, new species (Acari: Ixodoidea: Argasidae), a parasite of California gulls on islands in Mono Lake, California: description, biology, and life cycle. *Journal of Medical Entomology* 29: 78–97. <https://doi.org/10.1093/jmedent/29.1.78>
- Schwebke JR, Burgess D (2004) Trichomoniasis. *Clinical Microbiology Reviews* 17: 794–803. <https://doi.org/10.1128/CMR.17.4.794-803.2004>
- Scott HG (1964) Filter fly larva (*Psychoda alternata*) from human sputa. *Florida Entomologist* 47: 53–53. <https://doi.org/10.2307/3493767>

- Scott JD, Sajid MS, Pascoe EL, Foley JE (2021) Detection of *Babesia odocoilei* in Humans with Babesiosis Symptoms. *Diagnostics* 11(6): e947. <https://dx.doi.org/10.3390%2Fdiagnostics11060947>
- Scott MJ, Concha C, Welch JB, Phillips PL, Skoda SR (2017) Review of research advances in the screwworm eradication program over the past 25 years. *Entomologia Experimentalis et Applicata* 164: 226–236. <https://doi.org/10.1111/eea.12607>
- Senanayake SN, Papparini A, Latimer M, Andriolo K, Dasilva AJ, Wilson H, Xayavong MV, Collignon PJ, Jeans P, Irwin PJ (2012) First report of human babesiosis in Australia. *Medical Journal of Australia* 196: 350–352. <https://doi.org/10.5694/mja11.11378>
- Serra-Freire NM, Sena LMM, Borsoi ABP (2011) Parasitismo humano por carrapatos na Mata Atlântica, Rio de Janeiro, Brasil. *EntomoBrasilis* 4(2): 67–72. <https://doi.org/10.12741/ebrasilis.v4i2.101>
- Sharma J, Mamatha GP, Acharya R (2008) Primary oral myiasis: a case report. *Medicina Oral, Patologia Oral, Cirugia Bucal* 13(11): E714–E716.
- Sharma SN, Kumawat R, Singh SK (2019) Kyasanur Forest disease: vector surveillance and its control. *Journal of Communicable Diseases* 51: 38–44. <https://doi.org/10.24321/0019.5138.201915>
- Shaw J, Pratlong F, Floeter-Winter L, Ishikawa E, El Baidouri F, Ravel C, Dedet JP (2015) Characterization of *Leishmania (Leishmania) waltoni* n. sp. (Kinetoplastida: Trypanosomatidae), the Parasite Responsible for Diffuse Cutaneous Leishmaniasis in the Dominican Republic. *American Journal of Tropical Medicine and Hygiene* 93: 552–558. <https://doi.org/10.4269/ajtmh.14-0774>
- Shepherd C, Wangchuk P, Loukas A (2018) Of dogs and hookworms: man's best friend and his parasites as a model for translational biomedical research. *Parasites & Vectors* 11: e59. <https://doi.org/10.1186/s13071-018-2621-2>
- Shimpi R, Patel D, Raval K (2018) Human urinary myiasis by *Psychoda albipennis*: A case report and review of literature. *Urology case reports* 21: 122–123. <https://doi.org/10.1016/j.eucr.2018.08.015>
- Shirley DT, Farr L, Watanabe K, Moonah S (2018) A review of the global burden, new diagnostics, and current therapeutics for amebiasis. *Open Forum Infectious Diseases* 5: ofy161. <https://doi.org/10.1093/ofid/ofy161>
- Siddiqui R, Makhlof Z, Khan NA (2021) The increasing importance of *Vermamoeba vermiformis*. *Journal of Eukaryotic Microbiology* 68(5): e12857. <https://doi.org/10.1111/jeu.12857>
- Siebers R (2014) Dust mites in urine. *New Zealand Journal of Medical Laboratory Science* 68: 00.
- Silveira I, Pacheco RC, Szabo MP, Ramos HG, Labruna MB (2007) *Rickettsia parkeri* in Brazil. *Emerging Infectious Diseases* 13: 1111–1113. <https://doi.org/10.3201/eid1307.061397>
- Simner PJ (2017) Medical parasitology taxonomy update: January 2012 to December 2015. *Journal of Clinical Microbiology* 55: 43–47. <https://doi.org/10.1128/JCM.01020-16>
- Singh B, Daneshvar C (2013) Human infections and detection of *Plasmodium knowlesi*. *Clinical Microbiology Reviews* 26: 165–184. <https://doi.org/10.1128/CMR.00079-12>
- Singla N, Chander J, Lehl SS (2018) Urinary myiasis by *Pericoma* species: a case report. *Journal of Medical College Chandigarh* 8: 36–38.

- Sivajothi S, Sudhakara Reddy B, Rayulu VC, Sreedevi C (2015) *Notoedres cati* in cats and its management. *Journal of Parasitic Diseases* 39: 303–305. <https://doi.org/10.1007/s12639-013-0357-7>
- Smadi Re, Amr Z, Katbeh-Bader A, Obaidat N, Tawarah M, Hasan H (2014) Facial myiasis and canthariasis associated with systemic lupus panniculitis: a case report. *International Journal of Dermatology* 53(11): 1365–1369. <https://doi.org/10.1111/ijd.12500>
- Smith KGV, Taylor E (1966) *Anisopus* Larvae (Diptera) in cases of intestinal and urino-genital myiasis. *Nature* 210: 852–852. <https://doi.org/10.1038/210852a0>
- Snow RW, Guerra CA, Noor AM, Myint HY, Hay SI (2005) The global distribution of clinical episodes of *Plasmodium falciparum* malaria. *Nature* 434: 214–217. <https://doi.org/10.1038/nature03342>
- Soares RP, das Graças Evangelista L, Laranja LS, Diotaiuti L (2000) Population dynamics and feeding behavior of *Triatoma brasiliensis* and *Triatoma pseudomaculata*, main vectors of Chagas disease in Northeastern Brazil. *Memórias do Instituto Oswaldo Cruz* 95: 151–155. <https://doi.org/10.1590/S0074-02762000000200003>
- Spencer GJ (1963) Attacks on humans by *Ixodes angustus* Neumann, the coastal squirrel tick, and *I. soricis* Gregson, the shrew tick. *Proceedings of the Entomological Society of British Columbia* 60: 40.
- Spiegel JR, Saag KG, Tsang TK (1989) Infectious diarrhea secondary to *Enteromonas hominis*. *American Journal of Gastroenterology* 84: 1313–1314.
- Springer YP, Casillas S, Helfrich K, Mocan D, Smith M, Arriaga G, Mixson L, Castrodale L, McLaughlin J (2017) Two Outbreaks of Trichinellosis Linked to Consumption of Walrus Meat – Alaska, 2016–2017. *MMWR Morbidity Mortality Weekly Report* 66: 692–696. <https://doi.org/10.15585/mmwr.mm6626a3>
- Stark D, Barratt J, Chan D, Ellis JT (2016) *Dientamoeba fragilis*, the Neglected Trichomonad of the Human Bowel. *Clinical Microbiology Reviews* 29: 553–580. <https://doi.org/10.1128/CMR.00076-15>
- Stark D, Garcia LS, Barratt JL, Phillips O, Roberts T, Marriott D, Harkness J, Ellis JT (2014) Description of *Dientamoeba fragilis* cyst and precystic forms from human samples. *Journal of Clinical Microbiology* 52: 2680–2683. <https://doi.org/10.1128/JCM.00813-14>
- Stead D, du Plessis D, Sun LM, Freaun J (2021) *Anthemosoma garnhami* in an HIV-Infected Man from Zimbabwe Living in South Africa. *Emerging Infectious Disease* 27(7): 1991–1993. <https://doi.org/10.3201/eid2707.204759>
- Stensvold CR, Suresh GK, Tan KS, Thompson RC, Traub RJ, Viscogliosi E, Yoshikawa H, Clark CG (2007) Terminology for *Blastocystis* subtypes – a consensus. *Trends in Parasitology* 23: 93–96. <https://doi.org/10.1016/j.pt.2007.01.004>
- Stensvold CR, Winiacka-Krusnell J, Lier T, Lebbad M (2018) Evaluation of a PCR method for detection of *Entamoeba polecki*, with an overview of its molecular epidemiology. *Journal of Clinical Microbiology* 56: e00154-18. <https://doi.org/10.1128/JCM.00154-18>
- Stěrba J, Barus V (1976) First record of *Strobilocercus fasciolaris* (Taenidae-larvae) in man. *Folia Parasitologica* 23: 221–226.
- Steverding D (2017) The history of leishmaniasis. *Parasites & Vectors* 10: e82. <https://doi.org/10.1186/s13071-017-2028-5>

- Stewart A, Armstrong M, Graves S, Hajkovicz K (2017) *Rickettsia australis* and queensland tick typhus: A rickettsial spotted fever group infection in Australia. *American Journal of Tropical Medicine and Hygiene* 97: 24–29. <https://doi.org/10.4269/ajtmh.16-0915>
- Strasek-Smrđel K, Korva M, Pal E, Rajter M, Skvarc M, Avsic-Zupanc T (2020) Case of *Babesia crassa*-like infection, Slovenia, 2014. *Emerging Infectious Diseases* 26: 1038–1040. <https://doi.org/10.3201/eid2605.191201>
- Sugiyama H, Morishima Y, Arakawa K, Kishiro T, Kawanaka M (2007) Recent advances in the studies on larval spirurid nematode. *Japanese Society of Systematic Parasitology Circulation* 25: 4–7.
- Sukmee T, Siripattanapipong S, Mungthin M, Worapong J, Rangsin R, Samung Y, Kongkaew W, Bumrungsana K, Chanachai K, Apiwathanasorn C, Rujirojindakul P, Wattanasri S, Ungchusak K, Leelayoova S (2008) A suspected new species of *Leishmania*, the causative agent of visceral leishmaniasis in a Thai patient. *International Journal of Parasitology* 38: 617–622. <https://doi.org/10.1016/j.ijpara.2007.12.003>
- Sukontason KL, Sanit S, Klong-Klaew T, Tomberlin JK, Sukontason K (2014) *Sarcophaga (Liosarcophaga) dux* (Diptera: Sarcophagidae): A flesh fly species of medical importance. *Biological Research* 47: 14–14. <https://doi.org/10.1186/0717-6287-47-14>
- Sumner JW, Durden LA, Goddard J, Stromdahl EY, Clark KL, Reeves WK, Paddock CD (2007) Gulf Coast ticks (*Amblyomma maculatum*) and *Rickettsia parkeri*, United States. *Emerging Infectious Diseases* 13: 751–753. <https://doi.org/10.3201/eid1305.061468>
- Sun X, Wang L-F, Feng Y, Xie H, Zheng X-Y, He A, Karim MR, Lv Z-Y, Wu Z-D (2016) A case report: A rare case of infant gastrointestinal canthariasis caused by larvae of *Lasioderma serricornis* (Fabricius, 1792) (Coleoptera: Anobiidae). *Infectious Diseases of Poverty* 5: e34. <https://doi.org/10.1186/s40249-016-0129-6>
- Suwannatrai A, Saichua P, Haswell M (2018) Epidemiology of *Opisthorchis viverrini* Infection. *Advances in Parasitology* 101: 41–67. <https://doi.org/10.1016/bs.apar.2018.05.002>
- Suwannayod S, Sanit S, Sukontason K, Sukontason KL (2013) *Parasarcophaga (Liopygia) ruficornis* (Diptera: Sarcophagidae): a flesh fly species of medical importance. *Tropical Biomedicine* 30: 174–180.
- Suzuki J, Kobayashi S, Osuka H, Kawahata D, Oishi T, Sekiguchi K, Hamada A, Iwata S (2016) Characterization of a human isolate of *Tritrichomonas foetus* (cattle/swine genotype) infected by a zoonotic opportunistic infection. *Journal of Veterinary Medical Science* 78: 633–640. <https://doi.org/10.1292/jvms.15-0644>
- Sweatman GK (1958) Biology of *Otodectes cynotis*, the ear canker mite of carnivores. *Canadian Journal of Zoology* 36: 849–862. <https://doi.org/10.1139/z58-072>
- Swei A, O'Connor KE, Couper LI, Thekkiniath J, Conrad PA, Padgett KA, Burns J, Yoshimizu MH, Gonzales B, Munk B, Shirkey N, Konde L, Ben Mamoun C, Lane RS, Kjemtrup A (2019) Evidence for transmission of the zoonotic apicomplexan parasite *Babesia duncani* by the tick *Dermacentor albipictus*. *International Journal of Parasitology* 49: 95–103. <https://doi.org/10.1016/j.ijpara.2018.07.002>
- Szabó M, Pinter A, Labruna M (2013) Ecology, biology and distribution of spotted-fever tick vectors in Brazil. *Frontiers in Cellular and Infection Microbiology* 3: e27. <https://doi.org/10.3389/fcimb.2013.00027>

- Takada N, Fujita H (1992) Description of *Ixodes columnae* sp. nov., associated with a nymphal sp. N2 and a larval sp. L1 (Acarina: Ixodidae). Journal of the Acarological Society of Japan 1: 37–44. <https://doi.org/10.2300/acari.1.37>
- Takahashi K, Ito T, Sato T, Goto M, Kawamoto T, Fujinaga A, Yanagawa N, Saito Y, Nakao M, Hasegawa H, Fujiya M (2016) Infection with fully mature *Corynosoma* cf. *validum* causes ulcers in the human small intestine. Clinical Journal of Gastroenterology 9: 114–117. <https://doi.org/10.1007/s12328-016-0646-7>
- Talagrand-Reboul E, Boyer PH, Bergström S, Vial L, Boulanger N (2018) Relapsing fevers: neglected tick-borne diseases. Frontiers in Cellular and Infection Microbiology 8: e98. <https://doi.org/10.3389/fcimb.2018.00098>
- Tan KS (2008) New insights on classification, identification, and clinical relevance of *Blastocystis* spp. Clinical Microbiology Reviews 21: 639–665. <https://doi.org/10.1128/CMR.00022-08>
- Tanaka M, Makiuchi T, Komiyama T, Shiina T, Osaki K, Tachibana H (2019) Whole genome sequencing of *Entamoeba nuttalli* reveals mammalian host-related molecular signatures and a novel octapeptide-repeat surface protein. PLoS Neglected Tropical Diseases 13: e0007923. <https://doi.org/10.1371/journal.pntd.0007923>
- Tang TH, Wong SS, Lai CK, Poon RW, Chan HS, Wu TC, Cheung YF, Poon TL, Tsang YP, Tang WL, Wu AK (2017) Molecular Identification of *Spirometra erinaceieuropaei* tapeworm in cases of human sparganosis, Hong Kong. Emerging Infectious Diseases 23: 665–668. <https://doi.org/10.3201/eid2304.160791>
- Tappe D, Berkholz J, Mahlke U, Lobeck H, Nagel T, Haeupler A, Muntau B, Racz P, Poppert S (2016a) Molecular identification of zoonotic tissue-invasive tapeworm larvae other than *Taenia solium* in suspected human cysticercosis cases. Journal of Clinical Microbiology 54: 172–174. <https://doi.org/10.1128/JCM.02171-15>
- Tappe D, Büttner DW (2009) Diagnosis of human visceral pentastomiasis. PLoS Neglected Tropical Diseases 3: e320. <https://doi.org/10.1371/journal.pntd.0000320>
- Tappe D, Sulyok M, Riu T, Rozsa L, Bodo I, Schoen C, Muntau B, Babocsay G, Hardi R (2016b) Co-infections in visceral pentastomiasis, Democratic Republic of the Congo. Emerging Infectious Diseases 22: 1333–1339. <https://doi.org/10.3201/eid2208.151895>
- Tarragona EL, Soares JF, Costa FB, Labruna MB, Nava S (2016) Vectorial competence of *Amblyomma tonelliae* to transmit *Rickettsia rickettsii*. Medical and Veterinary Entomology 30: 410–415. <https://doi.org/10.1111/mve.12189>
- Tekely E, Szostakiewicz B, Wawrzycki B, Kądziela-Wypyska G, Juszkiewicz-Borowiec M, Pietrzak A, Chodorowska G (2013) Cutaneous larva migrans syndrome: a case report. Postępy Dermatol Alergol 30: 119–121. <https://doi.org/10.5114/pdia.2013.34164>
- Teschner M, Würfel W, Sedlacek L, Suerbaum S, Tappe D, Hornef MW (2014) Outer ear canal infection with *Rhabditis* sp. nematodes in a human. Journal of Clinical Microbiology 52: 1793–1795. <https://doi.org/10.1128/JCM.00115-14>
- Tessler M, Weiskopf SR, Berniker L, Hersch R, McCarthy KP, Yu DW, Siddall ME (2018) Bloodlines: mammals, leeches, and conservation in southern Asia. Systematics and Biodiversity 16: 488–496. <https://doi.org/10.1080/14772000.2018.1433729>
- Thach PN, van Doorn HR, Bishop HS, Fox MS, Sapp SGH, Cama VA, Van Duyet L (2021) Human infection with an unknown species of *Dracunculus* in Vietnam. International Journal of Infectious Diseases 105: 739–742. <https://doi.org/10.1016/j.ijid.2021.02.018>

- Thakur L, Kushwaha HR, Negi A, Jain A, Jain M (2020) *Leptomonas seymouri* co-infection in cutaneous leishmaniasis cases caused by *Leishmania donovani* from Himachal Pradesh, India. *Frontiers in Cellular and Infection Microbiology* 10: e345. <https://doi.org/10.3389/fcimb.2020.00345>
- Theiler G (1962) The Ixodoidea Parasites of Vertebrates in Africa South of the Sahara (Ethiopian Region). Report to the Director of Veterinary Services, Onderstepoort, South Africa, 255 pp.
- Thompson Jr JH, Knutson LV, Culp OS (1970) Larva of *Scenopinus* sp. (Diptera: Scenopinidae) causing human urogenital myiasis? *Mayo Clinic Proceedings* 45: 597–601.
- Thompson RC (2015) Neglected zoonotic helminths: *Hymenolepis nana*, *Echinococcus canadensis* and *Ancylostoma ceylanicum*. *Clinical Microbiology and Infection* 21: 426–432. <https://doi.org/10.1016/j.cmi.2015.01.004>
- Tilak R, Kunwar R, Wankhade UB, Tilak VW (2011) Emergence of *Schoengastiella ligula* as the vector of scrub typhus outbreak in Darjeeling: has *Leptotrombidium deliense* been replaced? *Indian Journal of Public Health* 55: 92–99. <https://doi.org/10.4103/0019-557X.85239>
- Timms S, Ferro DN, Emberson RM (1981) General biology and nomenclature of *Sancassania berlesei* (Michael). *Acarologia* 22: 385–390.
- Tiwari S, Karuna T, Rautaraya B (2014) *Hymenolepis diminuta* infection in a child from a rural area: A rare case report. *Journal of Laboratory Physicians* 6: 58–59. <https://doi.org/10.4103/0974-2727.129096>
- To KK, Wong SS, Poon RW, Trendell-Smith NJ, Ngan AH, Lam JW, Tang TH, AhChong AK, Kan JC, Chan KH, Yuen KY (2012) A novel *Dirofilaria* species causing human and canine infections in Hong Kong. *Journal of Clinical Microbiology* 50: 3534–3541. <https://doi.org/10.1128/JCM.01590-12>
- Tolan Jr RW (2011) Fascioliasis due to *Fasciola hepatica* and *Fasciola gigantica* infection: An update on this ‘Neglected’ neglected tropical disease. *Laboratory Medicine* 42: 107–116. <https://doi.org/10.1309/LMLFBB8PW4SA0YJI>
- Tolba ME, Husein EA, Farrag HM, Mohamed Hel D, Kobayashi S, Suzuki J, Ali TA, Sugano S (2016) *Allovalbikampfia spelaea* causing keratitis in humans. *PLoS Neglected Tropical Diseases* 10: e0004841. <https://doi.org/10.1371/journal.pntd.0004841>
- Toledo R, Esteban JG (2016) An update on human echinostomiasis. *Transactions of The Royal Society of Tropical Medicine and Hygiene* 110: 37–45. <https://doi.org/10.1093/trstmh/trv099>
- Torres-Guerrero E, Quintanilla-Cedillo MR, Ruiz-Esmenjaud J, Arenas R (2017) Leishmaniasis: A review. *F1000Research* 6: 750. <https://doi.org/10.12688/f1000research.11120.1>
- Traub RJ (2013) *Ancylostoma ceylanicum*, a re-emerging but neglected parasitic zoonosis. *International Journal of Parasitology* 43: 1009–1015. <https://doi.org/10.1016/j.ijpara.2013.07.006>
- Traversa D, Di Cesare A, Lia RP, Castagna G, Meloni S, Heine J, Strube K, Milillo P, Otranto D, Meckes O, Schaper R (2011) New insights into morphological and biological features of *Capillaria aerophila* (Trichocephalida, Trichuridae). *Parasitology Research* 109: 97–104. <https://doi.org/10.1007/s00436-011-2406-4>
- Trontelj P, Utevsky SY (2005) Celebrity with a neglected taxonomy: molecular systematics of the medicinal leech (genus *Hirudo*). *Molecular Phylogenetics and Evolution* 34: 616–624. <https://doi.org/10.1016/j.ympev.2004.10.012>

- Tu WC, Chen HC, Chen KM, Tang LC, Lai SC (2007) Intestinal myiasis caused by larvae of *Telmatoscopus albipunctatus* in a Taiwanese man. *Journal of Clinical Gastroenterology* 41: 400–402. <https://doi.org/10.1097/01.mcg.0000212615.66713.ba>
- Uchikawa R, Akune K, Inoue I, Kagei N, Sato A (1987) A human case of hair worm (*Gordius* sp.) infection in Kagoshima, Japan. *Kiseichugaku Zasshi* 36: 358–360.
- Udall DN (2007) Recent updates on onchocerciasis: Diagnosis and treatment. *Clinical Infectious Diseases* 44: 53–60. <https://doi.org/10.1086/509325>
- Udonsom R, Prasertbun R, Mahittikorn A, Mori H, Changbunjong T, Komalamisra C, Pintong AR, Sukthana Y, Popruk S (2018) *Blastocystis* infection and subtype distribution in humans, cattle, goats, and pigs in central and western Thailand. *Infection, Genetics and Evolution* 65: 107–111. <https://doi.org/10.1016/j.meegid.2018.07.007>
- Umehara A, Kawakami Y, Araki J, Uchida A (2007) Molecular identification of the etiological agent of the human anisakiasis in Japan. *Parasitology International* 56: 211–215. <https://doi.org/10.1016/j.parint.2007.02.005>
- Uni S, Fukuda M, Ogawa K, Lim YA-L, Agatsuma T, Bunchom N, Saijuntha W, Otsuka Y, Bhasu SAP, Udin ASM, Zainuri NA, Omar H, Nakatani J, Matsubayashi M, Maruyama H, Ramli R, Azirun MSB, Takaoka H (2017) Zoonotic infection with *Onchocerca dewittei japonica* in an 11-year-old boy in Kansai Region, Western Honshu, Japan. *Parasitology International* 66(5): 593–595. <https://doi.org/10.1016/j.parint.2017.06.006>
- Uni S, Fukuda M, Otsuka Y, Hiramatsu N, Yokobayashi K, Takahashi H, Murata S, Kusatake K, Morita E, Maruyama H, Hasegawa H, Shiwaku K, Ramli R, Azirun MS, Takaoka H (2015) New zoonotic cases of *Onchocerca dewittei japonica* (Nematoda: Onchocercidae) in Honshu, Japan. *Parasites & Vectors* 8: e59. <https://doi.org/10.1186/s13071-015-0655-2>
- Usinger RL (1966) Monograph of the Cimicidae (Hemiptera – Heteroptera). Entomological Society of America, College Park, Maryland, 572 pp.
- Utevsky SY, Trontelj P (2005) A new species of the medicinal leech (Oligochaeta, Hirudinida, *Hirudo*) from Transcaucasia and an identification key for the genus *Hirudo*. *Parasitology Research* 98: e61. <https://doi.org/10.1007/s00436-005-0017-7>
- Valdivia HO, Almeida LV, Roatt BM, Reis-Cunha JL, Pereira AA, Gontijo C, Fujiwara RT, Reis AB, Sanders MJ, Cotton JA, Bartholomeu DC (2017) Comparative genomics of canine-isolated *Leishmania (Leishmania) amazonensis* from an endemic focus of visceral leishmaniasis in Governador Valadares, southeastern Brazil. *Scientific Reports* 7: e40804. <https://doi.org/10.1038/srep40804>
- Van de Heyning J, Thienpont D (1977) Otitis externa in man caused by the mite *Otodectes cynotis*. *Laryngoscope* 87: 1938–1941. <https://doi.org/10.1002/lary.1977.87.11.1938>
- van Henten S, Adriaansen W, Fikre H, Akuffo H, Diro E, Hailu A, Van der Auwera G, van Griensven J (2018) Cutaneous Leishmaniasis Due to *Leishmania aethiopica*. *EClinicalMedicine* 6: 69–81. <https://doi.org/10.1016/j.eclinm.2018.12.009>
- Van Vinh Chau N, Buu Chau L, Desquesnes M, Herder S, Phu Huong Lan N, Campbell JJ, Van Cuong N, Yimming B, Chalermwong P, Jittapalpong S, Ramon Franco J, Tri Tue N, Rabaa MA, Carrique-Mas J, Pham Thi Thanh T, Tran Vu Thieu N, Berto A, Thi Hoa N, Van Minh Hoang N, Canh Tu N, Khac Chuyen N, Wills B, Tinh Hien T, Thwaites GE, Yacoub S, Baker S (2016) A clinical and epidemiological investigation of the first reported

- human infection with the zoonotic parasite *Trypanosoma evansi* in Southeast Asia. *Clinical Infectious Diseases* 62: 1002–1008. <https://doi.org/10.1093/cid/ciw052>
- Vandepitte J, Michaux JL, Fain A, Gatti F (1964) First records of *Physaloptera* infection in man in the Congo. *Annales de la Societe Belge de Medecine Tropicale* 44: 1067–1076.
- Vanderick F, Fain A, Langi S, Vanbalen H (1964) [2 new cases of human coenurosis caused by *Taenia brauni* in Rwanda, with orbital localization of the coenurus]. *Annales des Sociétés Belges de Médecine Tropicale, de Parasitologie, et de Mycologie* 44: 1077–1079.
- Varcasia A, Tamponi C, Toscirri G, Pipia AP, Dore F, Schuster RK, Kandil OM, Manunta ML, Scala A (2015) Is the red fox (*Vulpes vulpes*) a competent definitive host for *Taenia multiceps*? *Parasites & Vectors* 8: e491. <https://doi.org/10.1186/s13071-015-1096-7>
- Varma MGR, Webb HE, Pavri KM (1960) Studies on the transmission of Kyasanur Forest disease virus by *Haemaphysalis spinigera* newman. *Transactions of The Royal Society of Tropical Medicine and Hygiene* 54: 509–516. [https://doi.org/10.1016/0035-9203\(60\)90024-9](https://doi.org/10.1016/0035-9203(60)90024-9)
- Vatansever Z (2017a) *Hyalomma anatolicum* Koch, 1844 (Figs 158–160). In: Estrada-Peña A, Mihalca AD, Petney TN (Eds) *Ticks of Europe and North Africa: A Guide to Species Identification*. Springer International Publishing, Cham, 391–395. https://doi.org/10.1007/978-3-319-63760-0_74
- Vatansever Z (2017b) *Hyalomma asiaticum* Schülze and Schlottke, 1929. In: Estrada-Peña A, Mihalca AD, Petney TN (Eds) *Ticks of Europe and North Africa: A Guide to Species Identification*. Springer International Publishing, Cham, 403–404. https://doi.org/10.1007/978-3-319-63760-0_76
- Veracx A, Raoult D (2012) Biology and genetics of human head and body lice. *Trends in Parasitology* 28: 563–571. <https://doi.org/10.1016/j.pt.2012.09.003>
- Verma A, Manchanda S, Kumar N, Sharma A, Goel M, Banerjee PS, Garg R, Singh BP, Balharbi F, Lejon V, Deborggraeve S, Singh Rana UV, Puliyl J (2011) *Trypanosoma lewisi* or *T. lewisi*-like infection in a 37-day-old Indian infant. *American Journal of Tropical Medicine and Hygiene* 85: 221–224. <https://doi.org/10.4269/ajtmh.2011.11-0002>
- Verweij JJ, Polderman AM, Clark CG (2001) Genetic Variation among human isolates of uninucleated cyst-producing *Entamoeba* species. *Journal of Clinical Microbiology* 39: 1644–1646. <https://doi.org/10.1128/JCM.39.4.1644-1646.2001>
- Vial L (2009) Biological and ecological characteristics of soft ticks (Ixodida: Argasidae) and their impact for predicting tick and associated disease distribution. *Parasite* 16: 191–202. <https://doi.org/10.1051/parasite/2009163191>
- Viney ME, Ashford RW, Barnish G (1991) A taxonomic study of *Strongyloides* Grassi, 1879 (Nematoda) with special reference to *Strongyloides fuelleborni* von linstow, 1905 in man in Papua New Guinea and the description of a new subspecies. *Systematic Parasitology* 18: 95–109. <https://doi.org/10.1007/BF00017661>
- Visvesvara GS (2013) Infections with free-living amebae. *Handbook of Clinical Neurology* 114: 153–168. <https://doi.org/10.1016/B978-0-444-53490-3.00010-8>
- Visvesvara GS, Sriram R, Qvarnstrom Y, Bandyopadhyay K, Da Silva AJ, Pieniazek NJ, Cabral GA (2009) *Paravahlkampfia francinae* n. sp. masquerading as an agent of primary amoebic meningoencephalitis. *Journal of Eukaryotic Microbiology* 56: 357–366. <https://doi.org/10.1111/j.1550-7408.2009.00410.x>

- Voronova A, Chelomina GN (2018) Genetic diversity and phylogenetic relations of salmon trematode *Nanophyetus japonensis*. *Parasitology International* 67: 267–276. <https://doi.org/10.1016/j.parint.2018.01.002>
- Voronova AN, Chelomina GN, Besprozvannykh VV, Tkach VV (2017) Genetic divergence of human pathogens *Nanophyetus* spp. (Trematoda: Troglotrematidae) on the opposite sides of the Pacific Rim. *Parasitology* 144: 601–612. <https://doi.org/10.1017/S0031182016002171>
- Vos LJ, Robertson T, Binotto E (2016) *Haycocknema perplexum*: an emerging cause of parasitic myositis in Australia. *Communicable Diseases Intelligence Quarterly Report* 40: E496–E499.
- Waeschenbach A, Brabec J, Scholz T, Littlewood DTJ, Kuchta R (2017) The catholic taste of broad tapeworms – multiple routes to human infection. *International Journal of Parasitology* 47: 831–843. <https://doi.org/10.1016/j.ijpara.2017.06.004>
- Wagner R, Stallmeister N (2000) *Cheyletiella dermatitis* in humans, dogs and cats. *British Journal of Dermatology* 143: 1110–1112. <https://doi.org/10.1046/j.1365-2133.2000.03869x>
- Wagnerová P, Sak B, McEvoy J, Rost M, Matysiak AP, Ježková J, Kváč M (2015) Genetic diversity of *Cryptosporidium* spp. including novel identification of the *Cryptosporidium muris* and *Cryptosporidium tyzzeri* in horses in the Czech Republic and Poland. *Parasitology Research* 114: 1619–1624. <https://doi.org/10.1007/s00436-015-4353-y>
- Waldron LS, Cheung-Kwok-Sang C, Power ML (2010) Wildlife-associated *Cryptosporidium fayeri* in human, Australia. *Emerging Infectious Diseases* 16: 2006–2007. <https://doi.org/10.3201/eid1612.100715>
- Walker ED, Stobierski MG, Poplar ML, Smith TW, Murphy AJ, Smith PC, Schmitt SM, Cooley TM, Kramer CM (1998) Geographic distribution of ticks (Acari: Ixodidae) in Michigan, with emphasis on *Ixodes scapularis* and *Borrelia burgdorferi*. *Journal of Medical Entomology* 35: 872–882. <https://doi.org/10.1093/jmedent/35.5.872>
- Walochnik J, Haller-Schober E, Kolli H, Picher O, Obwaller A, Aspöck H (2000) Discrimination between clinically relevant and nonrelevant *Acanthamoeba* strains isolated from contact lens-wearing keratitis patients in Austria. *Journal of Clinical Microbiology* 38: 3932–3936. <https://doi.org/10.1128/JCM.38.11.3932-3936.2000>
- Walochnik J, Mulec J (2009) Free-living amoebae in carbonate precipitating microhabitats of Karst Caves and a new vahlkampfiid amoeba. *Acta Protozoologica* 48: 25–33.
- Wang J, Gao S, Zhang S, He X, Liu J, Liu A, Li Y, Liu G, Luo J, Guan G, Yin H (2020) Rapid detection of *Babesia motasi* responsible for human babesiosis by cross-priming amplification combined with a vertical flow. *Parasites & Vectors* 13: e377. <https://doi.org/10.1186/s13071-020-04246-4>
- Watson J (2008) New building, old parasite: Mesostigmatid mites – an ever-present threat to barrier facilities. *ILAR Journal* 49: 303–309. <https://doi.org/10.1093/ilar.49.3.303>
- Wattanakuppanich D, Jakkul W, Chanapromma C, Ketboonlue T, Dekumyoy P, Lv Z, Chan AHE, Thaenkham U, Chaisiri K (2021) Co-occurrence of *Angiostrongylus malaysiensis* and *Angiostrongylus cantonensis* DNA in cerebrospinal fluid: Evidence from human eosinophilic meningitis after ingestion of raw snail dish in Thailand. *Food and Waterborne Parasitology* 24: e00128. <https://doi.org/10.1016/j.fawpar.2021.e00128>

- Webster BL, Southgate VR, Littlewood DTJ (2006) A revision of the interrelationships of *Schistosoma* including the recently described *Schistosoma guineensis*. *International Journal of Parasitology* 36: 947–955. <https://doi.org/10.1016/j.ijpara.2006.03.005>
- Weiss DJ, Lucas TCD, Nguyen M, Nandi AK, Bisanzio D, Battle KE, Cameron E, Twohig KA, Pfeffer DA, Rozier JA, Gibson HS, Rao PC, Casey D, Bertozzi-Villa A, Collins EL, Dalrymple U, Gray N, Harris JR, Howes RE, Kang SY, Keddie SH, May D, Rumisha S, Thorn MP, Barber R, Fullman N, Huynh CK, Kulikoff X, Kutz MJ, Lopez AD, Mokdad AH, Naghavi M, Nguyen G, Shackelford KA, Vos T, Wang H, Smith DL, Lim SS, Murray CJL, Bhatt S, Hay SI, Gething PW (2019) Mapping the global prevalence, incidence, and mortality of *Plasmodium falciparum*, 2000–17: a spatial and temporal modelling study. *Lancet* 394: 322–331. [https://doi.org/10.1016/S0140-6736\(19\)31097-9](https://doi.org/10.1016/S0140-6736(19)31097-9)
- Wendt S, Trawinski H, Schubert S, Rodloff AC, Mössner J, Lübbert C (2019) The Diagnosis and Treatment of Pinworm Infection. *Deutsches Ärzteblatt International* 116: 213–219. <https://doi.org/10.3238/arztebl.2019.0213>
- Werner H, Rall E, Hendrischk A (1975) [Urogenital myiasis with *Fannia scalaris*]. *Deutsche Medizinische Wochenschrift* 100: 1397–1398. <https://doi.org/10.1055/s-0028-1106394>
- Westblade LF, Simon MS, Mathison BA, Kirkman LA (2017) *Babesia microti*: from mice to ticks to an increasing number of highly susceptible humans. *Journal of Clinical Microbiology* 55: 2903–2912. <https://doi.org/10.1128/JCM.00504-17>
- Whittaker C, Walker M, Pion SDS, Chesnais CB, Boussinesq M, Basáñez MG (2018) The Population Biology and Transmission Dynamics of *Loa loa*. *Trends in Parasitology* 34: 335–350. <https://doi.org/10.1016/j.pt.2017.12.003>
- Wikswø ME, Hu R, Dasch GA, Krueger L, Arugay A, Jones K, Hess B, Bennett S, Kramer V, Ereemeeva ME (2014) Detection and identification of spotted fever group rickettsiae in *Dermacentor* species from Southern California. *Journal of Medical Entomology* 45: 509–516. <https://doi.org/10.1093/jmedent/45.3.509>
- Wilkialis J (1973) The biology of nutrition of *Haemanteria costata*. *Zoologica Poloniae* 23: 213–225.
- Wolf MJ, Watkins HR, Schwan WR (2020) *Ixodes scapularis*: Vector to an increasing diversity of human pathogens in the Upper Midwest. *Wisconsin Medical Journal* 119: 16–21.
- Wolfe MS (2007) *Dicrocoelium dendriticum* or *Dicrocoelium hospes*. *Clinical Infectious Diseases* 44: 1522–1522. <https://doi.org/10.1086/517835>
- Wolmarans CT, de Kock KN, van der Walt MP (1990) An experimental *Schistosoma mattheei* infection in man. *Onderstepoort Journal of Veterinary Research* 57: 211–214.
- Won S, Park BK, Kim BJ, Kim HW, Kang JG, Park TS, Seo HY, Eun Y, Kim KG, Chae JS (2014) Molecular identification of *Haemadipsa rjukjuana* (Hirudiniformes: Haemadipsidae) in Gageo Island, Korea. *The Korean Journal of Parasitology* 52: 169–175. <https://doi.org/10.3347/kjp.2014.52.2.169>
- Wormser GP, McKenna D, Piedmonte N, Vinci V, Egizi AM, Backenson B, Falco RC (2020) First recognized human bite in the United States by the Asian longhorned tick, *Haemaphysalis longicornis*. *Clinical Infectious Diseases* 70: 314–316. <https://doi.org/10.1093/cid/ciz449>

- Wozniak E, DeNardo D (2000) The biology, clinical significance and control of the common snake mite, *Ophionyssus natricis*, in captive reptiles. *Journal of Herpetological Medicine and Surgery* 10: 4–10. <https://doi.org/10.5818/1529-9651-10.3.4>
- Wozniak EJ, Lawrence G, Gorchakov R, Alamgir H, Dotson E, Sissel B, Sarkar S, Murray KO (2015) The biology of the triatomine bugs native to south central Texas and assessment of the risk they pose for autochthonous Chagas disease exposure. *The Journal of Parasitology* 101: 520–528. <https://doi.org/10.1645/15-748>
- Xiao L, Bern C, Arrowood M, Sulaiman I, Zhou L, Kawai V, Vivar A, Lal AA, Gilman RH (2002) Identification of the *Cryptosporidium* pig genotype in a human patient. *Journal of Infectious Diseases* 185: 1846–1848. <https://doi.org/10.1086/340841>
- Xu N, Liu H, Jiang Y, Yin J, Yuan Z, Shen Y, Cao J (2020) First report of *Cryptosporidium viatorum* and *Cryptosporidium occultus* in humans in China, and of the unique novel *C. viatorum* subtype XVaA3h. *BMC Infectious Diseases* 20: e16. <https://doi.org/10.1186/s12879-019-4693-9>
- Yamada M, Tegoshi T, Abe N, Urabe M (2012) Two human cases infected by the horsehair worm, *Parachordodes* sp. (Nematomorpha: Chordodidae), in Japan. *The Korean Journal of Parasitology* 50: 263–267. <https://doi.org/10.3347/kjp.2012.50.3.263>
- Yamaguti S (1963) *Systema Helminthum*. Vol. V. Acanthocephala. Interscience Publishers, New York, London, [vii +] 423 pp.
- Yamamoto S, Yamada M, Matsumura K (2018) Mimicking small intestinal anisakiasis. *Gastroenterology* 154(6): e9–e10. <https://doi.org/10.1053/j.gastro.2017.08.020>
- Yamauchi T, Yoshigou H, Itoh T (2013) Occurrence of *Parabdelia quadrioculata* (Annelida: Hirudinida: Glossiphoniidae) in Japan, with a first case of human infestation by the leech. *Comparative Parasitology* 80: 134–135[132]. <https://doi.org/10.1654/4593.1>
- Yang T (1996) *Fauna Sinica Annelida Hirudinea*. Science Press, Bei Jing, 260 pp.
- Yao C, Köster LS (2015) *Tritrichomonas foetus* infection, a cause of chronic diarrhea in the domestic cat. *Veterinary Research* 46: e35. <https://doi.org/10.1186/s13567-015-0169-0>
- Yao MH, Wu F, Tang LF (2008) Human pentastomiasis in China: case report and literature review. *Journal of Parasitology* 94: 1295–1298. <https://doi.org/10.1645/GE-1597.1>
- Yap NJ, Hossain H, Nada-Raja T, Ngui R, Muslim A, Hoh BP, Khaw LT, Kadir KA, Simon Divis PC, Vythilingam I, Singh B, Lim YA (2021) Natural Human Infections with *Plasmodium cynomolgi*, *P. inui*, and 4 other Simian Malaria Parasites, Malaysia. *Emerging Infectious Disease* 27: 2187–2191. <https://doi.org/10.3201/eid2708.204502>
- Yoder JS, Straif-Bourgeois S, Roy SL, Moore TA, Visvesvara GS, Ratard RC, Hill VR, Wilson JD, Linscott AJ, Crager R, Kozak NA, Sriram R, Narayanan J, Mull B, Kahler AM, Schneeberger C, da Silva AJ, Poudel M, Baumgarten KL, Xiao L, Beach MJ (2012) Primary amebic meningoencephalitis deaths associated with sinus irrigation using contaminated tap water. *Clinical Infectious Diseases* 55: e79–e85. <https://doi.org/10.1093/cid/cis626>
- Yoshida A, Doanh PN, Maruyama H (2019) *Paragonimus* and paragonimiasis in Asia: An update. *Acta Tropica* 199: 105074. <https://doi.org/10.1016/j.actatropica.2019.105074>
- Yu L, Zhang ZQ, He L (2010) Two new species of *Pyemotes* closely related to *P. tritici* (Acari: Pyemotidae). *Zootaxa* 2723: 1–40. <https://doi.org/10.11646/zootaxa.2723.1.1>

- Yu PX, Zhao Q, Meng ZM, Ji YJ (2019) [Rhabditis axei found in urine routine examination: a case report]. Zhongguo Xue Xi Chong Bing Fang Zhi Za Zhi 31: 565–566. <https://doi.org/10.16250/j.32.1374.2018181> [in chinese]
- Yuan CL, Keeling PJ, Krause PJ, Horak A, Bent S, Rollend L, Hua XG (2012) *Colpodella* spp.-like parasite infection in woman, China. Emerging Infectious Diseases 18: 125–127. <https://doi.org/10.3201/eid1801.110716>
- Zahedi A, Paparini A, Jian F, Robertson I, Ryan U (2016) Public health significance of zoonotic *Cryptosporidium* species in wildlife: Critical insights into better drinking water management. International Journal for Parasitology: Parasites and Wildlife 5: 88–109. <https://doi.org/10.1016/j.ijppaw.2015.12.001>
- Zainuddin N, Nair P, Razali F (2016) Leech in the Nose – an unusual cause of epistaxis. Malaysian Family Physician 11: 33–34.
- Zaloni CA, Pillay A, Secor W, Humburg B, Aber R (2011) Rare case of trichomonal peritonitis. Emerging Infectious Diseases 17: 1312–1313. <https://doi.org/10.3201/eid1707.100892>
- Zhan XD, Li CP, Yang BH, Zhu YX, Tian Y, Shen J, Zhao JH (2017) Investigation on the zoonotic trematode species and their natural infection status in Huainan areas of China. Nutrition Hospitalaria 34: 175–179. <https://doi.org/10.20960/nh.994>
- Zhao L, He B, Li K-R, Li F, Zhang L-Y, Li X-Q, Liu Y-H (2018) First report of *Anaplasma ovis* in pupal and adult *Melophagus ovinus* (sheep ked) collected in South Xinjiang, China. Parasites & Vectors 11: e258. <https://doi.org/10.1186/s13071-018-2788-6>
- Zumpt F (1965) Myiasis in Man and Animals in the Old World: A Textbook for Physicians, Veterinarians, and Zoologists. Butterworths, Oxford, 267 pp.