

Four new species of the genus *Luzonomyza* Malloch (Diptera, Lauxaniidae) from China

Wenliang Li¹, Xulong Chen¹, Keli Feng¹, Shengjuan Zhao², Ding Yang³

1 College of Horticulture and Plant Protection, Henan University of Science and Technology, Luoyang 471023, China **2** College of Food & Bioengineering, Henan University of Science and Technology, Luoyang 471023, China **3** Department of Entomology, China Agricultural University, Beijing 100193, China

Corresponding author: Wenliang Li (wenliangli@haust.edu.cn; laughonce@126.com)

Academic editor: Marc De Meyer | Received 7 May 2021 | Accepted 14 November 2021 | Published 1 December 2021

<http://zoobank.org/2494F10A-DACB-4C0F-8C01-AB7A8A4E63BB>

Citation: Li W, Chen X, Feng K, Zhao S, Yang D (2021) Four new species of the genus *Luzonomyza* Malloch (Diptera, Lauxaniidae) from China. ZooKeys 1074: 43–59. <https://doi.org/10.3897/zookeys.1074.68392>

Abstract

Four species of the genus *Luzonomyza* Malloch, 1929 from southwest China are described as new to science: *Luzonomyza vittifacies* Li & Yang, **sp. nov.**, *L. serrata* Li & Yang, **sp. nov.**, *L. honghensis* Li & Yang, **sp. nov.**, and *L. brevis* Li & Yang, **sp. nov.** A key to *Luzonomyza* species is also presented.

Keywords

Lauxaniinae, Lauxanioidea, new species, Oriental region, Schizophora

Introduction

The genus *Luzonomyza* Malloch, 1929 was erected as a subgenus of *Trigonometopus* Macquart 1835 on the basis of a single species from the Philippines, *Trigonometopus bakeri* Bezzi, 1913. From the 20th to the 21st century, considerable contributions were made to the genus by several taxonomists. In his key, Stuckenberg (1971) first recognized *Luzonomyza* (= *Luzomyia*) as a valid subgenus. Shatalkin (1997) described a new species *Trigonometopus (Tetroxyrhina) nigripalpis* from Thailand (at that time, *Tetroxyrhina* Hendel, 1938 was also considered a subgenus of *Trigonometopus*), and

later, he described another new species, *L. sinica* Shatalkin, 1998, from China and Thailand. A new species (*L. japonica*) was described from Japan in *Trigonometopus*, and the male genitalia was figured (Sasakawa 2002). Papp (2007) described three new species, *L. pseudoforicula*, *L. sasakawai*, and *L. vietnamensis*, providing several figures (the epandrium of *L. bakeri*, the wing and male genitalia of *L. sinica* and *L. pseudoforicula*, and the head of *L. sasakawai* and *L. vietnamensis*) and presented a key to the Old World genera of Trigonometopini, including *Luzonomyza* and a key to the species of the genus *Luzonomyza*. Shi and Yang (2015) also described two new species and presented a key to species of the genus. To date, 13 species of *Luzonomyza* are known in the world, and all are distributed in the Oriental Region. Seven species occur in China, including four new species from southwest China described in this paper.

Luzonomyza can be recognized by the combination of the following features: head triangular, frons projecting beyond eye, usually with brown longitudinal band and with short setulae on anterior half; ocellar setae short. Fronto-facial angle acute. Face long, inclinate, some species with distinct genual spine, parafacial with a row of short setulae on lower half along inner margin. Gena with a row of long setae extending from middle to lower half of parafacial, but not in line with setulae on inner margin. Mesonotum with wide brown band; three post-sutural dorsocentral setae; katepisternum with a seta. Wing hyaline, anteriorly usually pale brown to dark brown, cross-veins r-m and dm-cu with brown spots. Abdomen yellow except posterior margin of tergites usually brown or black. (Malloch 1929; Stuckenberg 1971; Papp 2007; Shi and Yang 2015).

Materials and methods

Material

All specimens were collected in Yunnan Province, China. The specimens of *L. brevis*, *L. honghensis*, and *L. serrata* were captured alive and fixed in 75% ethanol. The specimen of *L. vittifacies* was killed with ethyl acetate and dried for morphological examination. All specimens are deposited at the China Agricultural University, Beijing, China (CAUC).

Morphological descriptions

General terminology follows Cumming and Wood (2009) and Gaimari and Silva (2010). Genitalia preparations were made by removing and macerating the apical portion of the abdomen in cold saturated NaOH for 6 h, after which they were rinsed and neutralized for dissection and study. After examination in glycerin, they were transferred to fresh glycerin and stored in a microvial pinned below the specimen or moved to ethanol in a tube together with the wet specimens.

Taxonomy

Key to the known species of the genus *Luzonomyza* Malloch

(modified from Papp 2007 and Shi and Yang 2015)

- 1 All tarsi entirely yellow..... *L. vietnamensis* Papp
- At least a pair of tarsi apices brown to black..... 2
- 2 Male surstylus with broad, slightly bifid apex, which is concave in lateral view *L. bakeri* (Bezzi)
- Male surstylus not bifid apex in lateral view 3
- 3 Wing unicolorous 4
- Wing patterned..... 5
- 4 Wing hyaline, very faintly yellow-tinged along anterior margin; haltere yellow with faintly brown-tinged knob; male epandrium without dorsoapical processes; postgonites simple and long
..... *L. japonica* (Sasakawa)
- Wing unicolorous brown; knob of haltere rather dark; male epandrium with extremely long dorsoapical processes, postgonites short with characteristic apex *L. pseudoforicula* Papp
- 5 Palpus black *L. nigripalpis* (Shatalkin)
- Palpus yellow 6
- 6 Male genitalia with epandrium lacking dorsoapical processes.....
..... *L. sasakawai* Papp
- Male genitalia with epandrium having strong dorsoapical processes 7
- 7 Mesonotum with 3 brown stripes *L. sinica* Shatalkin
- Mesonotum with 4 brown stripes 8
- 8 Frons with 3 brown longitudinal stripes, median longitudinal stripe wide extending from anterior margin to ocellar triangle, lateral longitudinal stripes narrow *L. brevis* sp. nov.
- Frons with a brown or blackish brown longitudinal stripe, extending from anterior margin to ocellar triangle..... 9
- 9 Acrostichal setulae in 2 rows 10
- Acrostichal setulae in 4 rows 11
- 10 Hind femur without long anteroventral seta; epandrium with a pair of long black coniform dorsal processes, surstylus tapering distally and incurved
..... *L. gaimarii* Shi & Yang
- Hind femur with a long anteroventral seta; epandrium with a pair of short black coniform dorsal processes, surstylus bluntly rounded, not incurved distally *L. hirsute* Shi & Yang
- 11 Face without brown stripes *L. honghensis* sp. nov.
- Face with 2 brown stripes..... 12

- 12 Legs yellow, fore tarsomeres 4 and 5 brown; abdomen yellow, tergites 2–6 black on posterior margin, but tergite 6 yellow posteromedially; hypandrium broad, membranous; pregonite confluent with hypandrium; phallus with a pair of lateral acute processes near base in ventral view, concavity V-shaped, with a dorsal acute process subapically in lateral view, apex acute and curved dorsally *L. vittifacies* sp. nov.
- Legs pale yellow, fore femur and basal half of tibia brown, tarsomeres 4 and 5 pale brown, mid and hind femora and tibiae yellow but pale yellow at apex; abdomen yellow, tergites 2–6 blackish brown on posterior margin; hypandrium V-shaped, with 2 pairs of lateral acute processes; pregonite indistinct, hypandrium, pregonite and phallus confluent together; both sides of phallus serrated, with a dorsal process at apex in lateral view *L. serrata* sp. nov.

Species descriptions

Luzonomyza brevis Li & Yang, sp. nov.

<http://zoobank.org/55D839C9-A2BD-4545-976B-C7C0B3F989F2>

Figures 1–8

Type material. *Holotype*. ♂ (CAUC), CHINA, Yunnan Province, Longling County, Xiaoheshan Reserve; 24°33'51"N, 98°49'20"E; 1840 m; 26 Apr 2012; Wenliang Li leg.


Etymology. Latin, *brevis*, referring to the epandrium with short dorsoapical processes. It is an adjective in apposition.

Diagnosis. Frons with 3 brown longitudinal stripes, median longitudinal stripe wide, extending from anterior margin to ocellar triangle, lateral longitudinal stripes narrow; face with a diamond-shaped brown marking in the middle, four angles of marking extending to facial margins. Mesonotum with 4 brown stripes. Epandrium with paired dorsoapical processes; phallopodeme claviform, shorter than phallus.

Description. Male. Body length 4.8 mm, wing length 4.3 mm.


Head yellow. Face with an angular hump on middle of basal half, lateral margin brown on apical half, with a diamond-shaped brown marking in the middle, 4 angles of marking extending to facial margins; parafacial with sparse short hairs, with a black spot between eye and antennal bases, and with 5 long setae extending to gena. Frons ~1.2× longer than wide and parallel-sided, with 3 brown longitudinal stripes, a broader brown median longitudinal stripe extending from anterior margin to ocellar triangle, 2 lateral longitudinal stripes narrower and pale brown, anterior half with short setulae; ocellar triangle grayish black, ocellar setae very small, hair-like, anterior fronto-orbital seta reclinate, shorter than the posterior one. Gena with broad brown stripe, ~1/3 height of eye. Antenna yellow, rounded apically, ~1.2× longer than high; arista brown except for yellow base, pubescent. Proboscis yellow with white and black setulae, and with a pair of irregular blackish-brown lateral spots apically; palpus yellow with black setulae.

Thorax (Fig. 3) brownish yellow, with grayish-white pruinescence. Mesonotum with 4 brown stripes extending to tip of scutellum, occupying most of scutellum. 0+3



Figures 1–4. *Luzonomyza brevis* sp. nov. Male **1** wing **2** habitus, lateral view **3** thorax, dorsal view **4** abdomen, dorsal view.

dorsocentral setae, anteriormost dorsocentral seta away from scutal suture; acrostichal setulae in 4 rows; a pair of prescutellar setae, shorter than anteriormost dorsocentral seta. Dorsal and posterior margin of anepisternum and dorsal margin of katepisternum pale yellow. One anepisternal seta, 1 katepisternal seta. **Legs** mostly yellow; fore tar-



Figures 5–8. *Luzonomyza brevis* sp. nov. Male **5** syntergosternite and epandrium, lateral view **6** epandrial complex, posterior view **7** phallic complex, ventral view **8** phallic complex, lateral view. Scale bar: 0.5 mm.

someres 2–5 pale brown, mid-legs differently colored: only right mid femur blackish brown but brownish yellow at base. Fore femur with 7 posterodorsal setae and 6 posteroventral setae, fore tibia with a long dorsal preapical seta and a short apicoventral seta. Mid tibia with a strong dorsal preapical seta and an apicoventral seta. Hind femur with a weak preapical anterodorsal seta; hind tibia with a long dorsal preapical seta and a short apicoventral seta. **Wing** pale brown along costal margin, extending to M_1 , a brown spot on each of the crossveins r-m and dm-cu; subcostal cell brown; costa with 2nd (between R_1 and R_{2+3}), 3rd (between R_{2+3} and R_{4+5}) and 4th (between R_{4+5} and M_1) sections in ratios of 9.7: 1.5: 1.4; r-m on middle of discal cell; ultimate and penultimate sections of M_1 in ratios of 6.6: 3.5; ultimate section of CuA_1 ~1/5 of penultimate. Haltere pale yellow.

Abdomen (Fig. 4) yellow, tergites 2–6 blackish brown on posterior margin. **Male genitalia** (Figs 5–8): syntergosternite confluent with epandrium, near triangular. Epandrium with a pair of long conical dorsoapical processes in lateral view, a pair of lateral processes broad apically and narrow basally on anterior margin. Surstylus

situated in ventral angle and small; hypandrium V-shaped, with 2 pairs of inner apical processes. Gonopod absent, hypandrium confluent with phallus. Phallus with a pair of lateral preapical acute processes in ventral view; dorsal process broader in lateral view, vertical to phallus, anterior apical angle acute. Phallus deeply concave apically, phallapodeme claviform, shorter than phallus.

Female. Unknown.

Remarks. This new species is very similar to *Luzonomyza sasakawai* from Thailand and Vietnam by the body markings and wing type, but it can be separated from the latter by the 3 narrow brown stripes on the frons; by the brown fore tarsomeres 2–5; by the epandrium with dorsoapical processes. In *Luzonomyza sasakawai*, the frons has a narrow brown stripe; the fore tarsomeres 3–5 are black; the epandrium is without dorsoapical process.

Distribution. China (Yunnan).

***Luzonomyza honghensis* Li & Yang, sp. nov.**

<http://zoobank.org/D8D043A7-C691-4830-9AFA-8BDC6625CAB3>

Figures 9–16

Type material. Holotype. ♂ (CAUC), CHINA, Yunnan Province, Honghe Hani and Yi Autonomous Prefecture, Qimaba township; 22°48'33"N, 102°14'31"E; 1010 m; 11 Jun 2013; Jinying Yang leg.


Etymology. The specific name refers to the holotype locality, Honghe Prefecture. It is a noun in genitive case.

Diagnosis. Frons with a brown median stripe extending from anterior margin to ocellar triangle; gena with broad brown stripe. Acrostichal setulae in 4 rows. Epandrium with a pair of dorsoapical processes in lateral view and a pair of lateral processes on anterior margin; hypandrium, gonopod, and phallus confluent together. Phallus with 2 pairs of lateral processes and a pair of median processes apically. Phallapodeme longer than phallus.

Description. Male. Body length 3.7 mm.


Head (Fig. 9) yellow. Face with an angular hump on middle of basal half; parafacial with sparse short hairs, with a black spot between eye and antennal bases, and with 5 long setae extending to gena. Frons ~1.3× longer than wide and parallel-sided, with a brown median stripe extending from anterior margin to ocellar triangle, and frons with short setulae on anterior half; ocellar triangle grayish black, ocellar setae very small, hair-like; fronto-orbital setae missing. Gena with broad brown stripe, ~1/2 height of eye. Antenna yellow, first flagellomere and arista missing. Proboscis yellow with white and black setulae, and with a pair of irregular lateral spots apically; palpus yellow with black setulae.

Thorax (Fig. 11) brownish yellow, with grayish-white pruinescence. Mesonotum with 4 brown stripes extending to tip of scutellum. 0+3 dorsocentral setae, anteriormost dorsocentral seta away from scutal suture; acrostichal setulae in 4 rows; a pair of



Figures 9–12. *Luzonomyza honghensis* sp. nov. Male **9** head, anterior view **10** habitus, lateral view **11** thorax, dorsal view **12** abdomen, dorsal view.

prescutellar setae, all setae on thorax missing. Dorsal margin of anepisternum and katepisternum pale yellow. One anepisternal seta, 1 katepisternal seta. **Legs** yellow, fore tarsomeres and hind legs missing, mid tarsomeres 4 and 5 brown. Fore femur with 7 posterodorsal setae and 7 posteroventral setae, fore tibia with a long dorsal preapical



Figures 13–16. *Luzonomyza honghensis* sp. nov. Male **13** syntergosternite and epandrium, lateral view **14** epandrial complex, posterior view **15** phallic complex, ventral view **16** phallic complex, lateral view. Scale bar: 0.5 mm.

seta and a short apicoventral seta. Mid tibia with a strong dorsal preapical seta and an apicoventral seta. Hind femur with a weak preapical anterodorsal seta; hind tibia with a long dorsal preapical seta and a short apicoventral seta. **Wing** pale brown along anterior margin, a brown spot on each of the crossveins r-m and dm-cu; subcostal cell brown but pale brown apically. Haltere pale yellow.

Abdomen (Fig. 12) yellow, tergites 2–6 blackish brown on posterior margin but yellow laterally. **Male genitalia** (Figs 13–16): syntergosternite confluent with epandrium, broad dorsally and narrow ventrally. Epandrium with a pair of long black conical dorsoapical processes in lateral view, with a pair of lateral processes on anterior margin. Surstylus situated in ventral angle and small, ventral margin with setae, hypandrium V-shaped, disconnected in the middle and with 2 pairs of inner processes apically. Gonopod short and thick, extending to both sides; hypandrium, gonopod and phallus confluent together. Phallus with 2 pairs of different lateral processes and a pair of median processes apically; median processes claviform in lateral view. Phallus deeply concave apically, phallopodeme claviform, longer than phallus.

Female. Unknown.

Remarks. This new species is very similar to *Luzonomyza sinica* from China (Hainan) and Thailand in the body markings, wing type, and surstylus, but it can be separated from the latter by the brown mid tarsomeres 4 and 5 and the 2 pairs of apical processes of the phallus. In *Luzonomyza sinica*, the mid tarsi are yellow and the phallus is not bifurcated apically.

Distribution. China (Yunnan).

***Luzonomyza serrata* Li & Yang, sp. nov.**

<http://zoobank.org/2820D475-A55B-49B3-8C3E-1D6318626B66>

Figures 17–25

Type material. Holotype. ♂ (CAUC), CHINA, Yunnan Province, Xishuangbanna Dai Autonomous Prefecture, Menglun Zhiwuyuan; 21°55'12"N, 101°16'20"E; 580 m; 22 Apr 2007; Wenliang Li leg. **Paratype.** 1 ♂ (CAUC), CHINA, Yunnan Province, Xishuangbanna Dai Autonomous Prefecture, Menglun No. 55 area; 21°56'5"N, 101°14'48"E; 540 m; 23 Apr 2007; Hui Dong leg.


Etymology. Latin, *serrata*, referring to the serrated sides of the phallus.

Diagnosis. Face with 2 brown longitudinal stripes, gena with apical half of inner margin and ventral margin brown. Proboscis apically with a pair of Y-shaped brown spots. Mesonotum with 4 brown stripes extending to tip of scutellum. *Legs* pale yellow, fore femur with 7 posteroventral setae. Gonopod indistinct; both sides of phallus serrated. Phallus indistinct concave apically.

Description. Male. Body length 3.4–4.0 mm, wing length 3.5–3.8 mm.


Head (Fig. 17) yellow. Face with an angular hump on middle of basal half and 2 brown longitudinal stripes; parafacial with sparse short hairs and with 4 long setae extending to gena, with a black spot between eye and antennal bases, gena with apical half of inner margin and ventral margin brown. Frons ~1.4× longer than wide and parallel-sided, with a brown median stripe extending from anterior margin to ocellar triangle and with short setulae on anterior half; ocellar triangle grayish black, ocellar setae very small, hair-like; anterior fronto-orbital seta reclinate, shorter than the posterior one. Gena ~half height of eye and with broad brown stripe. Antenna yellow, first flagellomere and arista missing. Proboscis yellow with white and black setulae, and apically with a pair of Y-shaped brown spots; palpus yellow with black setulae.

Thorax (Fig. 20) brownish yellow, with grayish-white pruinescence. Mesonotum with 4 brown stripes extending to tip of scutellum, occupying most of scutellum. 0+3 dorsocentral setae, anteriormost dorsocentral seta away from scutal suture; acrostichal setulae in 4 rows; a pair of prescutellar setae, all setae on thorax missing. Dorsal margin of anepisternum and katapisternum yellow. One anepisternal seta, 1 katapisternal seta. **Legs** pale yellow, fore femur and basal half of tibia brown, tarsomeres 4 and 5 pale brown, mid and hind femora and tibiae yellow but pale yellow apically. Fore femur with 7 posterodorsal setae and 7 posteroventral setae, fore tibia with a long dorsal



Figures 17–21. *Luzonomyza serrata* sp. nov. Male **17** head, anterior view **18** wing **19** habitus, lateral view **20** thorax, dorsal view **21** abdomen, dorsal view.

preapical seta and a short apicoventral seta. Mid tibia with a strong dorsal preapical seta and an apicoventral seta. Hind femur with a weak preapical anterodorsal seta; hind tibia with a long dorsal preapical seta and a short apicoventral seta. **Wing** brown along costal margin, extending to M_1 , a brown spot on each of the crossveins r-m and dm-cu; subcostal cell brown; costa with 2nd (between R_1 and R_{2+3}), 3rd (between R_{2+3} and R_{4+5}),



Figures 22–25. *Luzonomyza serrata* sp. nov. Male **22** sytergosternite and epandrium, lateral view **23** epandrial complex, posterior view **24** phallic complex, ventral view **25** phallic complex, lateral view. Scale bar: 0.5 mm.

and 4th (between R_{4+5} and M_1) sections in ratios of 9.0: 1.6: 1.2; r-m beyond middle of discal cell; ultimate and penultimate sections of M_1 in ratios of 5.9: 3.0; ultimate section of CuA_1 ~1/5 of penultimate. Haltere pale yellow.

Abdomen (Fig. 21) yellow, tergites 2–6 blackish brown on posterior margin. **Male genitalia** (Figs 22–25): sytergosternite confluent with epandrium, broad dorsally and narrow ventrally. Epandrium with a pair of long black conical dorsoapical processes in lateral view, with a pair of lateral processes on anterior margin. Surstylus absent, ventral margin with setae, hypandrium V-shaped, with a pair of lateral acute processes. Gonopod indistinct; hypandrium, gonopod and phallus confluent together. Both sides of phallus serrated, with a dorsal process apically in lateral view. Phallus indistinct concave apically. Phallapodeme claviform, shorter than phallus.

Female. Unknown.

Remarks. This new species is very similar to *Luzonomyza pseudoforicula* from Thailand in the body markings, wing type, and leg color, but it can be separated from the latter by the yellow antenna and the phallus being serrated on both sides. In *Luzonomyza pseudoforicula*, the antenna is black, and the phallus is not serrated in ventral view.

Distribution. China (Yunnan).

***Luzonomyza vittifacies* Li & Yang, sp. nov.**

<http://zoobank.org/E092BDD5-2FD7-4705-A2B6-E57085A69696>

Figures 26–34

Type material. Holotype. ♂ (CAUC), CHINA, Yunnan Province, Xishuangbanna Dai Autonomous Prefecture, Mengla County, Yaoqu township; 21°43'32"N, 101°32'37"E; 780 m; 26 Apr 2007; Wenliang Li leg.


Etymology. This epithet is an adjectival combination of the Latin adjective *vitti* (vittate) and noun *facies* (face), referring to the face with brown longitudinal stripes.

Diagnosis. Head yellow; face with a pair of brown stripes extending from antennal base and confluent on ventral margin. Thorax brownish yellow with grayish-white pruinescence; legs yellow, fore tarsomeres 4 and 5 brown. Abdomen yellow, tergites 2–6 black on posterior margin but tergite 6 yellow posteromedially. *Male genitalia:* hypandrium broad, membranous; pregonite confluent with hypandrium; phallus with a pair of lateral acute processes near base in ventral view, apical concave V-shaped, with a dorsal acute subapical process in lateral view, apex acute and curved dorsally.

Description. Male. Body length 3.2 mm, wing length 3.4 mm.


Head (Fig. 26) yellow. Face brownish yellow, with a pair of brown longitudinal stripes extending from antennal bases and confluent on ventral margin, face with an angular hump on middle of basal half; inner margin of parafacial with a brown stripe, broadening towards gena, with a black round spot between eye and antennal bases, parafacial with sparse short setulae, and 6 short black setae extending from parafacial ventral corner to gena. Frons as long as wide and parallel-sided, with a blackish-brown median line extending from anterior margin to ocellar triangle, and with short setulae on anterior half, denser on anterior margin; ocellar triangle grayish black; ocellar setae very small, hair-like, anterior fronto-orbital seta (situated at middle of fronto-orbital plate) reclinate, shorter than the posterior one. Gena $\sim 1/3$ height of eye and with broad brown stripe. Antenna yellow, first flagellomere brownish yellow and rounded apically, $\sim 1.5\times$ longer than high; arista brown, pubescent. Proboscis mostly yellow except blackish brown at tip, with white and black setulae; palpus yellow with black setulae.

Thorax (Fig. 29) brownish yellow, with grayish-white pruinescence. Mesonotum with 4 broad brown stripes, 2 brown median stripes extending to tip of scutellum; 0+3 dorsocentral setae, anteriormost dorsocentral seta away from scutal suture; acrostichal setulae in 4 rows; a pair of prescutellar setae, missing. One anepisternal seta,



Figures 26–30. *Luzonomyza vittifacies* sp. nov. Male **26** head, anterior view **27** wing **28** habitus, lateral view **29** thorax, dorsal view **30** abdomen, dorsal view.

1 katepisternal seta. **Legs** mostly yellow, fore tarsomeres 4 and 5 brown. Fore femur with 8 posterodorsal setae and 6 posteroventral setae, fore tibia with a long dorsal preapical seta and a short apicoventral seta. Mid tibia with a strong dorsal preapical seta and an apicoventral seta. Hind tibia with a long dorsal preapical seta and a short apicoventral seta. **Wing** brown along costal margin extending to M_1 and with hyaline



Figures 31–34. *Luzonomyza vittifacies* sp. nov. Male **31** sytergosternite and epandrium, lateral view **32** epandrial complex, posterior view **33** phallic complex, ventral view **34** phallic complex, lateral view. Scale bar: 0.5 mm.

spots, a brown spot on each of the crossveins r-m and dm-cu; subcostal cell brown; costa with 2nd (between R₁ and R₂₊₃), 3rd (between R₂₊₃ and R₄₊₅) and 4th (between R₄₊₅ and M₁) sections in ratios of 7.1: 1.4: 1.2; r-m beyond middle of discal cell; ultimate and penultimate sections of M₁ in ratios of 4.9: 2.5; ultimate section of CuA₁ ~1/5 of penultimate. Haltere yellow.

Abdomen (Fig. 30) yellow, tergites 2–6 black on posterior margin, but tergite 6 yellow posteromedially. **Male genitalia** (Figs 31–34): sytergosternite confluent with

epandrium. Epandrium with a pair of black long conical dorsoapical processes in lateral view. Surstylus broken, hypandrium broad and membranous. Gonopod confluent with hypandrium, phallus consisting of a pair of sclerites in ventral view and a pair of lateral acute processes near base, with V-shaped apical concavity, broad basally and narrow apically in lateral view, with a dorsal acute process subapically, apex acute and curved dorsally. Phallapodeme claviform in ventral view. Ejaculatory apodeme bent.

Female. Unknown.

Remarks. This new species is very similar to *Luzonomyza gaimarii* from China (Yunnan) in the wing spots and the thoracic and abdominal spots, but it can be separated from the latter by the pair of brown stripes on the face that extend down from antennal base and are confluent on ventral margin; by the 4 rows of acrostichal setulae; by r-m extending beyond middle of discal cell; by the absence of a lateral concavity in the hypandrium; and by the phallus being acute and curved dorsally in lateral view. In *Luzonomyza gaimarii*, the face has no spot; the acrostichal setulae are arranged in 2 rows; crossvein r-m is in the middle of the discal cell; the hypandrium has a lateral concavity; and the phallus is not curved dorsally in lateral view.

Distribution. China (Yunnan).

Discussion

This study increases the number of known *Luzonomyza* species in the world to 13. Seven species occur in China, accounting for more than half of the total species worldwide. Except for *L. sinica* in Hainan Province, the other six species are all found in Yunnan Province. There may be several reasons for the high number of species in Yunnan. One reason may be the location in the southwest of China, which has diverse climate types and a unique geographical location with suitable climate conditions. It has been considered to be a hotspot of global biodiversity for many years, and its fauna has attracted much attention. Another important reason may be that the seemingly more limited distribution of this genus may be the result of insufficient sampling outside Yunnan. There are still many areas in China with rich biological biodiversity that remain poorly investigated. We believe that the species diversity of the genus *Luzonomyza* in China may be underestimated, and there may be new species in other areas, so further investigations are needed.

Acknowledgements

The research was funded by the National Natural Science Foundation of China (32070477, 31301903), the National Science & Technology Fundamental Resources Investigation Program of China (2018FY100400, 2019FY100400), and the Biodiversity Survey and Assessment Project of the Ministry of Ecology and Environment, China (2019HJ2096001006).

References

- Bezzi M (1913) Studies in Philippine Diptera. 1. The Philippine Journal Science, Section D 8: 305–332.
- Cumming JM, Wood DM (2009) Adult Morphology and Terminology. In: Brown BV, Borkent A, Cumming JM, Wood DM, Woodley NE, Zumbado MA (Eds) Manual of Central American Diptera. Volume 1. NRC Research Press, Ottawa, 9–50.
- Gaimari SD, Silva VC (2010) Lauxaniidae (Lauxaniid flies). In: Brown BV, Borkent A, Cumming JM, Wood DM, Woodley NE (Coords) Manual of Central American Diptera (Vol. 2). NRC Research Press, Ottawa, 971–995.
- Malloch JR (1929) Notes on some Oriental Sapromyzid flies (Diptera), with particular reference to the Philippine species. Proceedings of the United States National Museum 74(6): 1–97. <https://doi.org/10.5479/si.00963801.74-2751.1>
- Papp L (2007) A review of the Old World Trigonometopini Becker (Diptera: Lauxaniidae). Annales Historico-Naturales Musei Nationalis Hungarici 99: 55–95.
- Sasakawa M (2002) Oriental Lauxaniidae (Diptera) Part 3. Fauna of the Lauxaniidae in Japan (Ryukyus) and Formosa. Scientific Reports of Kyoto Prefectural University Human Environment and Agriculture 54: 33–61.
- Shatalkin AI (1997) East-Asian species of Lauxaniidae (Diptera). Genera *Trigonometopus* Macquart, *Protrigonometopus* Hendel. Dipterological Research 8(3): 163–168.
- Shatalkin AI (1998) New and little-known Lauxaniidae (Diptera) from Asia. Russian Entomological Journal 7(3–4): 209–218.
- Shi L, Yang D (2015) Two new species of the genus *Luzonomyza* from China (Diptera, Lauxaniidae). Zootaxa 3964(1): 87–94. <http://doi.org/10.11646/zootaxa.3964.1.5>
- Stuckenberg BR (1971) A review of the Old World genera of Lauxaniidae (Diptera). Annals of the Natal Museum 20: 499–610.