

New species of the genus *Inversidens* Haas, 1911 (Unionoida, Unionidae, Gonideinae) from Jiangxi Province, China

Ruiwen Wu¹, Xiongjun Liu², Takaki Kondo³, Shan Ouyang⁴, Xiaoping Wu⁴

1 School of Life Science, Shanxi Normal University, Linfen 041000, China **2** School of Life Science, Jiaying University, Meizhou 514015, China **3** Division of Natural Science, Osaka Kyoiku University, Osaka 582-8582, Japan **4** School of Life Sciences, Nanchang University, Nanchang 330031, China

Corresponding authors: Xiongjun Liu (609449126@qq.com); Xiaoping Wu (xpwu@ncu.edu.cn)

Academic editor: Graham Oliver | Received 24 May 2021 | Accepted 9 July 2021 | Published 3 August 2021

<http://zoobank.org/BA743F4E-2452-432A-9796-E1DEBC1D4CAD>

Citation: Wu R, Liu X, Kondo T, Ouyang S, Wu X (2021) New species of the genus *Inversidens* Haas, 1911 (Unionoida, Unionidae, Gonideinae) from Jiangxi Province, China. ZooKeys 1054: 85–93. <https://doi.org/10.3897/zookeys.1054.69075>

Abstract

We diagnose and describe a new freshwater mussel species of the genus *Inversidens*, *I. rentianensis* **sp. nov.** from Jiangxi Province, China based on morphological characters and molecular data. This paper includes a morphological description and photograph of the holotype, and partial sequences of mitochondrial COI as DNA barcode data.

Keywords

COI, freshwater mussel, genetic distances, morphology, taxonomy

Introduction

The genus *Inversidens* Haas, 1911 belongs to the subfamily Gonideinae in the family Unionidae. The genus was first depicted by Haas (1911) as a subgenus of *Nodularia* with two species, i.e., *Unio brandtii* Kobelt, 1879 and *Nodularia parcedentata* Haas, 1911, both restricted to Japan. Later, Haas (1969) further classified *Unio reinianus* Kobelt, 1879, *Unio haconensis* Ihering, 1893, *Unio japanensis* Lea, 1859, *Unio pantoensis* Neumayr, 1899 within *Inversidens*. All species were restricted to Japan, except for

U. pantoensis, which was distributed in China. By comparing the conchological characters, Kondo (1982) believed that *U. brandtii* was not morphologically distinct from *N. parcedentata*, and regarded *N. parcedentata* as a variety of *U. brandtii*. Based on the morphology of the glochidium, Habe (1991) removed *U. reinianus* from *Inversidens*, and established a new genus *Inversiunio*. Based on morphological characteristics of the shell, Kondo (1998) moved *Unio haconensis*, which was regarded as a synonym of *Unio jokohamensis* (Ihering, 1893), into *Inversiunio*. Furthermore, Starobogatov (1970) used *Unio japonensis* as the type species for his newly established genus *Pronodularia*.

Currently, only two species are recognized within *Inversidens*, the Japanese endemic *I. brandtii* and *I. pantoensis* in China (Fig. 1A, B; Kondo 2008; He and Zhuang 2013; Lopes-Lima et al. 2020; Graf and Cummings 2021a, b; MolluscaBase eds. 2021).

In this study, we diagnose and describe a new *Inversidens* species from Jiangxi Province, China. In addition, we provide estimations of the intraspecific and interspecific genetic distances within *Inversidens* based on the mitochondrial COI barcode to examine species validity.

Materials and methods

Specimen collection and identification

In March 2018, four samples were collected from the Mianshui River, Rentian Town, Ganzhou City, Jiangxi Province, China (25.989557°N, 116.131333°E). All type and voucher specimens are deposited at the Biological Museum of Nanchang University, China (NCFM180325–NCFM180328).

DNA extraction and COI amplification

Of the four individuals, only two samples had tissues. Total genomic DNA was extracted from dissected somatic tissue using TIANamp Marine Animals DNA Kit (Tiangen Biotech, Beijing, China) according to the manufacturer's instructions.

Mitochondrial cytochrome oxidase subunit I (COI) gene sequences have been widely used for species delimitation of freshwater mussels based on genetic distance and the criteria of monophyly (Elderkin et al. 2016; Lopes-Lima et al. 2019; Smith et al. 2019). Polymerase chain reaction (PCR) amplification of the COI gene with a 680-base pair fragment was performed using a primer pair consisting of LCO1490 and HCO2198 (Folmer et al. 1994). Thermal cycling conditions were 98 °C for 10 s, followed by 35 cycles of 94 °C for 1 min, 50 °C for 1 min, 72 °C for 1–2 min, and a final extension of 72 °C for 7 min, following the TaKaRa Ex manufacturer's protocol. The amplified PCR products were purified and sequenced by Sangon Biotech (Shanghai). The PCR product size for the COI amplicon was 680 bp. The sequences obtained in this study have been uploaded to GenBank.

DNA barcode dataset construction

We constructed a mitochondrial COI dataset with the newly obtained sequences from this study and the available *Inversidens brandtii* sequences from GenBank.

Previously published sequences were downloaded from GenBank and added to the dataset, i.e., 17 species of the subfamily Gonideinae and four species of the subfamily Unioninae for the ingroup, and one species of the subfamily Parreysiinae as the outgroup.

As a result, a total of 29 COI sequences were used for this study. Sequence details and GenBank accession numbers are shown in Table 1.

All COI nucleotide sequences were translated to amino acid sequences using MEGA 5.0 (Tamura et al. 2011) and aligned based on the amino acid sequences using the program MUSCLE (Edgar 2004) with default settings. We calculated and compared inter- and intraspecific distances with MEGA 5.0 using the uncorrected *p*-distance. Standard error was assessed using 1000 bootstrap replicates.

Table 1. List of sequences used in this study. (*) Sequenced from this study.

Taxa	GenBank accession number
UNIONIDAE Rafinesque, 1820	
Parreysiinae Henderson, 1935	
<i>Indonaia andersoniana</i> (Nevill, 1877)	KX865835
Unioninae Rafinesque, 1820	
<i>Acuticosta chinensis</i> (Lea, 1868)	MG462919
<i>Inversiunio jokohamensis</i> (Ihering, 1893)	LC518985
<i>Inversiunio reinianus</i> (Kobelt, 1879)	LC518976
<i>Nodularia douglasiae</i> (Griffith & Pidgeon, 1833)	NC_026111
Gonideinae Ortmann, 1916	
<i>Pseudodon bogani</i> Bolotov, Kondakov & Konopleva in Bolotov et al. 2017	MF352216
<i>Pseudodon manueli</i> Konopleva, Kondakov & Vikrev in Bolotov et al. 2017	MF352228
<i>Monodontina cambodjensis</i> (Petit de la Saussaye, 1865)	KP795028
<i>Pilsbryconcha exilis</i> (Lea, 1838)	KP795024
<i>Chamberlainia hainesiana</i> (Lea, 1856)	KX822635
<i>Sinohyriopsis cumingii</i> (Lea, 1852)	NC_011763
<i>Sinohyriopsis schlegelii</i> (Martens, 1861)	NC_015110
<i>Lamprotula caveata</i> (Heude, 1877)	KX822646
<i>Lamprotula leaii</i> (Griffith & Pidgeon, 1833)	NC_023346
<i>Potomida littoralis</i> (Cuvier, 1798)	JN243905
<i>Pronodularia japonensis</i> (Lea, 1859)	KX822659
<i>Gonidea angulata</i> (Lea, 1838)	DQ272371
<i>Leguminaia wheatleyi</i> (Lea, 1862)	KX822651
<i>Microcondylaea bonellii</i> (Férussac, 1827)	KX822652
<i>Sinosolenaia carinata</i> (Heude, 1877)	KX822669
<i>Ptychorhynchus pfisteri</i> (Heude, 1874)	KY067440
<i>Parvasolenaia rivularis</i> (Heude, 1877)	KX966393
<i>Inversidens brandtii</i> (Kobelt, 1879)	AB040827
<i>Inversidens brandtii</i> (Kobelt, 1879)	MT020598
<i>Inversidens brandtii</i> (Kobelt, 1879)	MT020597
<i>Inversidens brandtii</i> (Kobelt, 1879)	LC519005
<i>Inversidens brandtii</i> (Kobelt, 1879)	LC519004
<i>Inversidens rentianensis</i> sp. nov. 1*	MZ073336
<i>Inversidens rentianensis</i> sp. nov. 2*	MZ073337

Phylogenetic analysis

Bayesian inference (BI) analyses were inferred in MrBayes Version 2.01 (Ronquist et al. 2012), using GTRGAMMAI model of nucleotide substitution. Four chains were run simultaneously for 10 million generations and trees were sampled every 1000 generations. The first 25% of these trees were discarded as burn-in when computing the consensus tree (50% Majority Rule). Sufficient mixing of the chains was considered to have been reached when the average standard deviation of split frequencies was below 0.01. Additionally, IQ-TREE was run for Maximum Likelihood (ML) tree reconstruction, using partition models with 1000 ultrafast bootstraps (Minh et al. 2013).

Taxonomy

Inversidens rentianensis Wu & Wu, sp. nov.

<http://zoobank.org/62424717-9514-4C7D-9C0E-240F1D95F03E>

Fig. 1C

Type specimens. *Holotype*. CHINA • Jiangxi Province, Ganzhou City, Rentian Town (壬田镇), Mianshui River (25.989557°N, 116.131333°E), 13 March 2018, coll. Xiongjun Liu (NCFM180325). *Paratypes*. Same data as holotype (NCFM180326–NCFM180328).

Diagnosis. *Inversidens rentianensis* sp. nov. is morphologically distinct from the other two recognized species within the genus by shell shape, beak position and nacre colour (Table 2). Diagnostic characteristics: shell irregularly subtriangular; curvature of the ventral margin slight, nearly straight; umbo situated 1/2 of shell length; nacre reddish.

Description. Shell irregularly subtriangular, medium thickness, and quite inflated. Anterior margin regularly rounded; ventral margin nearly straight; posterior margin obliquely arc-shaped. Umbo prominent and slightly eroded. Umbo sculptured with feebly wavy wrinkles. Posterior slope formed by the ventral margin and posterior margin low, triangular. Epidermis shining black or with brownish-yellow hue. Only one cardinal tooth in each valve, shape triangular. Laterals thick, a little curved, 2 in each valve. Nacre reddish-bronze in colour.

Length 43–52 mm, height 29–36 mm.

Etymology. The specific epithet is derived from the type locality, Rentian Town.

Distribution. The species is known only from Mianshui River, Rentian Town, Ganzhou City, Jiangxi Province, China (present study) (Fig. 2).

GenBank accession number. Holotype, NCFM180325: MZ073336; paratypes, NCFM180326: MZ073337.

Molecular analyses. Pairwise COI sequence divergences from *Inversidens brandtii* and *Inversidens rentianensis* sp. nov. were conducted using MEGA 5.0. Based on the uncorrected *p*-distance model, the intraspecific divergences of *I. brandtii* and *I. rentianensis* sp. nov. were both 0.00%. The interspecific divergence of *I.*




Figure 1. Photographs of *Inversidens* taxa **A** *I. brandtii* **B** *I. pantoensis* **C** *I. rentianensis* sp. nov. Photos: **[A, B]** from the MUSSEL Project, **[C]** from this study, NCFM180325 (holotype), scale is 2 cm.

brandtii and *I. rentianensis* sp. nov. was 10.1%. Both BI and ML trees obtained a completely consistent topology. Consistent topology relationships are shown in Figure 3. In the phylogenetic trees, *I. rentianensis* sp. nov. formed a well-supported

Table 2. Conchological characters of *Inversidens rentianensis* sp. nov., *Inversidens brandtii*, *Inversidens pantoensis*. Characteristic descriptions of *I. brandtii* and *I. pantoensis* are referenced from Kondo (1982, 2008) and He and Zhuang (2013).

	<i>I. rentianensis</i> sp. nov.	<i>I. brandtii</i>	<i>I. pantoensis</i>
Shell shape	Irregularly subtriangular	Ovate	Inequilateral, quadrate
Umbo position	1/2 of shell length	1/4 of shell length	1/3 of shell length
Umbo sculpture	Feebly wavy wrinkles	Rippled	Angularly wrinkled
Surface sculpture	Concentric ridges	Concentric ridges	Irregular growth lines
Nacre colour	Reddish	Milk-white	Bluish
Posterior slope	Sharp	Blunt	Blunt
Ventral margin	Nearly straight	Arc-shaped	Long and straight



Figure 2. Photograph of sampling site of *Inversidens rentianensis* sp. nov. in China.

sister-group relationship with *Inversidens brandtii* (PP = 1.00, BS = 100; Fig. 3). The genera *Pronodularia* and *Inversiunio* belong to different clades well-separated from *Inversidens* (Fig. 3).

Remarks. Species delineation can be problematic in the presence of morphological ambiguities due to phenotypic plasticity and convergence (e.g., cryptic species), especially in mollusks (Zieritz et al. 2010; Inoue et al. 2013). The use of molecular genetics can aid species delineation in the case of phenotypic plasticity and/or convergence (Pieri et al. 2018; Wu et al. 2018). *Inversidens rentianensis* sp. nov. can be




Figure 3. Phylogenetic tree of freshwater mussels inferred from Bayesian Inference (BI) and Maximum Likelihood (ML) analyses of COI barcode. Support values above the branches are posterior probabilities (PP)/bootstrap support (BS). Red font indicates the new species from this study.

distinguished from congeneric species based on diagnostic characteristics of the shell. In this study, we also analyzed the interspecific divergence between *Inversidens brandtii* and *Inversidens rentianensis* sp. nov. based on the COI barcode. The results showed that the average interspecific divergence between the two species was 10.1%, which was much higher than intraspecific divergences. Genetic analysis conducted in this study supports *I. rentianensis* sp. nov. as a valid species, which can be easily distinguished by the COI barcode.

Acknowledgments

We thank the reviewer Dr Manuel Lopes-Lima, the other anonymous reviewer and the Subject Editor Dr Graham Oliver for valuable comments that have greatly improved this manuscript. This work was supported by the National Natural Science Foundation of China under Grant No.31772412.

References

- Doucet-Beaupré H, Blier PU, Chapman EG, Piontkivska H, Dufresne F, Sietman BE, Mulcrone RS, Hoeh WR (2012) *Pyganodon* (Bivalvia: Unionoida: Unionidae) phylogenetics: a male- and female-transmitted mitochondrial DNA perspective. *Molecular Phylogenetics and Evolution* 63: 430–444. <https://doi.org/10.1016/j.ympev.2012.01.017>
- Edgar RC (2004) Muscle: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Research* 32: 1792–1797. <https://doi.org/10.1093/nar/gkh340>
- Elderkin CL, Clewing C, Wembo Ndeo O, Albrecht C (2016) Molecular phylogeny and DNA barcoding confirm cryptic species in the African freshwater oyster *Etheria elliptica* Lamarck, 1807 (Bivalvia: Etheriidae). *Biological Journal of the Linnean Society* 118(2): 369–381. <https://doi.org/10.1111/bij.12734>
- Folmer O, Black M, Hoeh W, Lutz R, Vrijenhoek R (1994) DNA primers for amplification of mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates. *Molecular Marine Biology & Biotechnology* 3: 294–299.
- Graf DL, Cummings KS (2021a) A ‘big data’ approach to global freshwater mussel diversity (Bivalvia: Unionoida), with an updated checklist of genera and species. *Journal of Molluscan Studies* 87(1): 1–36. <https://doi.org/10.1093/mollus/eyaa034>
- Graf DL, Cummings KS (2021b) The Freshwater Mussels (Unionoida) of the World (and other less consequential bivalves). MUSSEL Project Web Site. <http://www.mussel-project.net/>
- Haas F (1969) Superfamilia Unionacea. Berlin: Walter de Gruyter.
- Haas F (1911) Die Unioniden. [in] H.C. Küster, Systematisches Conchylien-Cabinet von Martini und Chemnitz 9 (pt. 2, h. 45): 89–112.
- Haas F (1920) Die Neumayrschen Najaden aus der Ausbeute des Grafen Széchenyi. *Senckenbergiana* 2: 146–151.
- Habe T (1991) Catalogue of non-marine molluscs in Japan. *Hitachiobi* 56: 3–7.
- He J, Zhuang Z (2013) The Freshwater Bivalves of China. ConchBooks.
- Inoue K, Harris JL, Robertson CR, Johnson NA, Randklev CR (2019) A comprehensive approach uncovers hidden diversity in freshwater mussels (Bivalvia: Unionidae) with the description of a novel species. *Cladistics* 36(1): 88–113. <https://doi.org/10.1111/cla.12386>
- Inoue K, Hayes DM, Harris JL, Christian AD (2013) Phylogenetic and morphometric analyses reveal ecophenotypic plasticity in freshwater mussels *Obovaria jacksoniana* and *Villosa arkansasensis* (Bivalvia: Unionidae). *Ecology and Evolution* 3(8): 2670–2683. <https://doi.org/10.1002/ece3.649>
- Jones JW, Neves RJ, Ahlstedt SA, Hallerman EM (2006) A holistic approach to taxonomic evaluation of two closely related endangered freshwater mussel species, the Oyster mussel *Epioblasma capsaeformis* and Tan Riffleshell *Epioblasma florentina* walker (Bivalvia: Unionidae). *Journal of Molluscan Studies* 72(3): 267–283. <https://doi.org/10.1093/mollus/eyl004>
- Kondo T (1982) Taxonomic revision of *Inversidens* (Bivalvia: Unionidae). *Venus* 41(3): 181–198.
- Kondo T (1998) Revision of the genus *Inversiunio* (Bivalvia: Unionidae). *Venus* 57(2): 85–93.
- Kondo T (2008) Monograph of Unionoida in Japan (Mollusca: Bivalvia). Special Publication of the Malacological Society of Japan (3): 32–34.

- Lopes-Lima M, Burlakova L, Karatayev A, Gomes-dos-Santos A, Zieritz A, Froufe E, Bogan AE (2019) Revisiting the North American freshwater mussel genus *Quadrula sensu lato* (Bivalvia: Unionidae): Phylogeny, taxonomy and species delineation. *Zoologica Scripta* 48(3): 313–336. <https://doi.org/10.1111/zsc.12344>
- Lopes-Lima M, Hattori A, Kondo T, Hee Lee J, Ki Kim S, Shirai A, Hayashi H, Usui T, Sakuma K, Toriya T, Sunamura Y, Ishikawa H, Hoshino N, Kusano Y, Kumaki H, Utsugi Y, Yabe S, Yoshinari Y, Hiruma H, Tanaka A, Sao K, Ueda T, Sano I, Miyazaki J-I, Gonçalves DV, Klishko OK, Konopleva ES, Vikhrev IV, Kondakov AV, Gofarov MY, Bolotov IN, Sayenko EM, Soroka M, Zieritz A, Bogan AE, Froufe E (2020) Freshwater mussels (Bivalvia: Unionidae) from the rising sun (Far East Asia): phylogeny, systematics, and distribution. *Molecular Phylogenetics and Evolution* 146: e106755. <https://doi.org/10.1016/j.ympev.2020.106755>
- Minh BQ, Nguyen MAT, von Haeseler A (2013) Ultrafast approximation for phylogenetic bootstrap. *Molecular biology and evolution* 30(5): 1188–1195. <https://doi.org/10.1093/molbev/mst024>
- MolluscaBase eds (2021) MolluscaBase. <http://www.molluscabase.org> [on 2021-06-21.]
- Pieri AM, Inoue K, Johnson NA, Smith CH, Harris JL, Robertson C, Randklev CR (2018) Molecular and morphometric analyses reveal cryptic diversity within freshwater mussels (Bivalvia: Unionidae) of the western Gulf coastal drainages of the USA. *Biological Journal of the Linnean Society* 124(2): 261–277. <https://doi.org/10.1093/biolinnean/bly046>
- Ronquist F, Teslenko M, Mark PV, Ayres DL, Darling A, Höhna S, Larget B, Liu L, Marc A, Suchard MA, Huelsenbeck JP (2012) Mrbayes 3.2: Efficient Bayesian phylogenetic inference and model choice across a large model space. *Systematic Biology* 61: 539–542. <https://doi.org/10.1093/molbev/mst024>
- Simpson CT (1914) A descriptive catalogue of the naiades, or pearly fresh-water mussels. Parts I–III. Bryant Walker, Detroit, Michigan. <https://www.biodiversitylibrary.org/page/11344898>
- Smith CH, Johnson NA, Inoue K, Doyle RD, Randklev CR (2019) Integrative taxonomy reveals a new species of freshwater mussel, *Potamilus streckersoni* sp. nov. (Bivalvia: Unionidae): implications for conservation and management. *Systematics and Biodiversity* 17(4): 331–348. <https://doi.org/10.1080/14772000.2019.1607615>
- Starobogatov Y (1970) Fauna of Molluscs and Zoogeographic Division of Continental Waterbodies of the Globe. Leningard Nauka.
- Tamura K, Peterson D, Peterson N, Stecher G, Nei M, Kumar S (2011) Mega5: Molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Molecular Biology and Evolution* 28: 2731–2739. <https://doi.org/10.1093/molbev/msr121>
- Wu RW, Liu YT, Wang S, Liu XJ, Zanatta DT, Roe KJ, Song XL, An CT, Wu XP (2018) Testing the utility of DNA barcodes and a preliminary phylogenetic framework for Chinese freshwater mussels (Bivalvia: Unionidae) from the middle and lower Yangtze River. *PLoS ONE* 13: e0200956. <https://doi.org/10.1371/journal.pone.0200956>
- Zieritz A, Hoffman JL, Amos W, Aldridge DC (2010) Phenotypic plasticity and genetic isolation-by-distance in the freshwater mussel *Unio pictorum* (Mollusca: Unionoida). *Evolutionary Ecology* 24(4): 923–938. <https://doi.org/10.1007/s10682-009-9350-0>