

Additions to the Limoniidae and Pediciidae fauna of Morocco, with an updated checklist (Diptera, Tipuloidea)

Ouafaa Driauach¹, Boutaina Belqat¹

¹ Laboratory «Ecology, Biodiversity and Environment», Department of Biology, Faculty of Sciences, University Abdelmalek Essaâdi, Tétouan, 93030, Morocco

Corresponding author: Ouafaa Driauach (driauach@gmail.com)

Academic editor: J. Salmela | Received 2 December 2015 | Accepted 20 January 2016 | Published 15 February 2016

<http://zoobank.org/F408D3D4-0FBD-4841-9178-5A21EBDF6770>

Citation: Driauach O, Belqat B (2016) Additions to the Limoniidae and Pediciidae fauna of Morocco, with an updated checklist (Diptera, Tipuloidea). ZooKeys 563: 129–146. doi: 10.3897/zookeys.563.7384

Abstract

Eighteen species of Limoniidae and two species of Pediciidae are recorded for the first time in Morocco, of which 15 species are new to North Africa. An updated checklist of Moroccan short-palped craneflies (Limoniidae and Pediciidae) is appended, containing 73 species in 25 genera.

Keywords

Morocco, new records, North Africa, Short-palped crane flies, updated checklist

Introduction

Limoniidae and Pediciidae belong to the Tipuloidea together with Cylindrotomidae and Tipulidae. The Limoniidae comprise more than 741 species already known in the Westpalaearctic region, from which only 102 species are known to occur in Morocco (49), Algeria (32), Tunisia (9), Libye (1) and Egypt (11). The Pediciidae are the second smallest family of the Tipuloidea, with 77 recognized taxa in the Westpalaearctic of which only 4 were recorded from North Africa, in Morocco (Driauach et al. 2013, Oosterbroek 2015).

Adults of Limoniidae and Pediciidae are associated with moist environments. They are usually collected from damp vegetation along the borders of various water bodies and from damp forests (Starý et al. 2005, De Jong et al. 2008). Their immature stages are usually found in a wide range of aquatic and semi-aquatic habitats (Reusch and Oosterbroek 1997).

Limoniidae and Pediciidae are poorly known in Morocco. They only received sporadic faunistic records made by European entomologists (Pierre 1922a, b, Pierre 1924, Séguay 1941a, b, Vaillant 1956, Starý 1971, Starý 1974, Savchenko et al. 1992, Eiroa 2000, Pârvu et al. 2006, Starý 2006, Starý and Oosterbroek 2008, Starý 2009, Starý 2011).

Only a single attempt was made to summarize the knowledge of the Moroccan fauna in the first checklist of the short-palped crane flies of Morocco (Driauach et al. 2013) and very recent research also revealed a new species, *Baeoura staryi* from Morocco including an identification key of the genus *Baeoura* in the West Palaearctic species (Driauach and Belqat 2015).

Prior to the present study, there were 53 species recorded from Morocco (Oosterbroek 2015). New findings increase the number of Moroccan Limoniidae and Pediciidae to 73. Altogether 44 species of Limoniidae and Pediciidae are collected and recorded in this work. Of these, 18 species of Limoniidae and two species of Pediciidae are recorded for the first time in Morocco of which 15 species are new to North Africa.

Material and methods

The material presented in this work come from field campaigns carried out by the authors between 2011 and 2015 in 77 sites distributed over a mountainous area (Rif, High Atlas, Middle Atlas, Beni Snassen) and arid area (Sahara) of Morocco. A total of 564 specimens of short-palped crane flies (515 specimens of Limoniidae and 49 specimens of Pediciidae) were identified. Adults were mostly collected by sweeping vegetation with entomological hand net whereas other specimens were reared in the laboratory from the substratum taken from several zones of water bodies margins and kept under the laboratory conditions. Fresh material was stored in 70% alcohol. For identification, the male terminalia, if necessary, were prepared by boiling in a solution of 10% KOH and preserved in glycerine. All the specimens were collected by the authors and are deposited in the Diptera Collection of the Faculty of Sciences, University Abdelmalek Essââdi, Tétouan, Morocco. The Distribution as it is given for individual species is based on Oosterbroek (2015).

The checklist of Moroccan short-palped crane flies is updated. New records are marked with an asterisk (*) and bold set text. A brief description of the material examined is given and the distribution of the identified species is provided, together with notes on the breeding sites (Table 1).

Table I. Sampling sites (in alphabetic order) for the adults of Limoniidae and Pediciidae in Moroccan regions, 2011–2015, with localities, geographical coordinates and altitudes.

Site Rif Mts	Locality	Province	Altitude (m)	Geographical coordinates	Type of habitat
Tributary of Oued El Fondak	Aïn Lahcen, Khandek Melouka	Tétouan	291	35°33.544'N, 05°34.995'W	River of the middle course
Tributary of Oued El Kébir	Route de Beni Yedder	Chefchaouen	145	35°25.864'N, 05°27.610'W	River of the middle course
Tributary of Oued Hachef	Dar Chaoui	Tanger-Assila	47	35°31.458'N, 05°43.771'W	Small stream
Tributary of Oued Majiou	Nord Majiou	Chefchaouen	786	35°06.730'N, 05°11.309'W	Small stream
Tributary of Oued Ouara	Khandek Romana, Jbel Khizana	Chefchaouen	1003	35°02.579'N 05°12.872'W	River of the middle course
Tributary of Oued Taïda	Sidi Brahim Ben Arrif	Larache	860	35°20.398'N, 05°32.471'W	River of the higher course
Tributary of Oued Zarka	Zarka	Chefchaouen	128	35°31.220'N, 05°20.461'W	Stream
Aïn Afersiw	Mezine	Chefchaouen	746	35°05.945'N, 05°20.439'W	Spring
Aïn Bab Tariouente	Ânasar	Chefchaouen	1440	35°01.110'N, 05°00.586'W	Spring and stream
Aïn Boughaba	Jbel Bou Bessouï, Cedar Forest	Chefchaouen	1526	34°58.779'N, 04°46.366'W	Spring
Aïn Chouk	Aïn Chouk	Larache	5	35°09.803'N, 06°06.473'W	Marsh
Aïn El Ma Bared	Bouzquate	Chefchaouen	1246	35°00.333'N, 05°12.105'W	Spring and irrigation canal
Aïn El Maouzil	Parc National de Tâlassemteane	Chefchaouen	1106	35°04.577'N, 05°10.406'W	Spring and brook
Aïn El Malkab	Parc National de Tâlassemteane	Chefchaouen	1278	35°05.509'N, 05°09.443'W	Spring and brook
Aïn Quanquiben	Jbel Bou Bessouï	Chefchaouen	1600	34°57.750'N, 04°40.783'W	Spring
Aïn Mâze	Jbel Bouhachem	Larache	1294	35°14.381'N, 05226.316'W	Spring
Aïn Ras El Ma	Majiou	Chefchaouen	856	35°06.873'N, 5°11.388'W	Spring
Aïn Sidi Brahim Ben Arrif (Site 1)	Route moulay Abdessalam	Larache	882	35°20.415'N, 05°32.535'W	Small stream
Aïn Sidi Brahim Ben Arrif (Site 2)	Route moulay Abdessalam	Larache	897	35°20.398'N, 05°32.712'W	Spring and brook
Aïn Takhmijoute	Bab Rouida, Parc National de Tâlassemteane	Chefchaouen	1512	35°06.881'N, 05°08.270'W	Spring
Âounsar Aheramen	Majiou	Chefchaouen	855	35°06.162'N, 05°10.739'W	Spring
Barrage Ajras	Aïn Lahcen	Tétouan	49	35°23.833'N, 05°29.902'W	Barrage
Barrage Moulay Bouchra	Larbaa Beni Hassan	Tétouan	364	35°15.864'N, 0521.221'W	Barrage
Cascade Chrafate	Chrafate	Chefchaouen	820	35°03.997'N, 05°06.434'W	Waterfall
Cascade Zarka	Zarka	Tétouan	147	35°31.211'N, 05°20.477'W	Waterfall
Daya Aïn Jdioui	Aïn Jdioui	Tanger-Assilah	5	35°34.074'N, 05°55.499'W	Pond

Site	Locality	Province	Altitude (m)	Geographical coordinates	Type of habitat
Daya Amsmil	Jbel Bouhachem	Tétonan	1059	35°15.596'N, 05°25.917'W	Bog
Daya El Ânassar	Bab Berred	Chefchaouen	1183	35°00.788'N, 04°57.419'W	Bog
Daya Fifi	Fifi	Chefchaouen	1202	35°1.367'N, 05°12.335'W	Bog
Daya Jbel Zemzem	Jbel Zemzem	Tétonan	233	35°45.553'N, 05°22.340'W	Pond
Daya Mezine	Mezine	Chefchaouen	778	35°06.104'N, 5°21.177'W	Pond
Daya Mgħara	Dar Chaoui	Tétonan	44	35°31.147'N, 05°42.963'W	Pond
Daya Mtahen	Jbel Bouhachem	Larache	966	35°16.195'N, 05°26.158'W	Pond
Daya near Ain Afersiw	Mezine	Chefchaouen	716	35°06.069'N, 5°20.337'W	Pond
Daya Tazia	Route Moulay Abdessalam	Larache	721	35°20.814'N, 05°33.139'W	Pond
Daya Rmali	El Malāab, Parc National de Talassentane	Chefchaouen	1276	35°05.563'N, 05°09.488'W	Pond
Daya Afraie	Tanaqoub	Chefchaouen	600	35°05.634'N, 05°26.028'W	Bog
Lac Badriouen	Badriouen	Tanger	34	35°42.083'N, 05°52.116'W	Lack
Maison forestière	Parc National de Talassentane	Chefchaouen	1674	35°08.076'N, 05°08.262'W	Spring and brook
Mari El Kheyel	Jbel Bouhachem	Tétonan	989	35°15.985'N, 05°26.209'W	Marshland
Oued Aārate	Dardara	Chefchaouen	269	35°07.381'N, 05°17.456'W	River of the middle course
Oued Aārkoub	Ârkoub	Chefchaouen	140	35°15.943'N, 04°50.631'W	River of the middle course
Oued Abou Brar	Douar Abou Brar, Parc National de Talassentane	Chefchaouen	1254	35°10.977'N, 05°08.005'W	River of the higher course
Oued Amsa	Amsa	Tétonan	13	35°31.593'N, 05°13.821'W	Downstream of the river
Oued El Kanar	Stetħare	Chefchaouen	50	35°17.233'N, 04°59.639'W	Downstream of the river
Oued Amsmil (Site 1)	Jbel Bouhachem	Larache	1059	35°15.614'N, 05°25.943'W	River of the higher course
Oued Amsmil (Site 3)	Jbel Bouhachem	Larache	1134	35°15.635'N, 05°26.134'W	River of the higher course
Oued at 15 Km from Fifi	Bouzhatte	Chefchaouen	1256	35°00.805'N, 05°12.365'W	River of the higher course
Oued Azila	Azila, Jbel Tidghine	Al Hoċċima	1601	34°52.028'N, 04°32.609'W	River of the higher course
Oued El Kebir	Route Vers Beni Yedder	Tétonan	89	35°27.351'N, 05°25.796'W	River of the middle course
Oued El Koub	Souk El Had	Chefchaouen	124	35°01.298'N, 05°12.365'W	Waterfall
Oued Farda	Akchour	Chefchaouen	394	35°14.147'N, 05°10.576'W	River of the higher course
Oued]nane Niche	Jnane Niche	Chefchaouen	36	35°17.019'N, 04°51.233'W	River of the middle course
Oued Loukous	Souk El Had	Chefchaouen	148	35°01.317'N, 05°25.246'W	River of the middle course
Oued Kelħaa	Akchour	Chefchaouen	303	35°14.268'N, 05°10.452'W	River of the higher course

Site	Locality	Province	Altitude (m)	Geographical coordinates	Type of habitat
Oued Madissouka.	Madissouka, Parc National de Tassastane	Chefchaouen	1367	35°10.622'N, 05°08.400'W	River of the higher course
Oued Majjou (Hafa meqlouba)	Majjou	Chefchaouen	825	35°06.175'N, 05°10.836'W	River of the higher course
Oued Majjou (Pont)	Majjou	Chefchaouen	786	35°06.730'N, 05°11.309'W	River of the higher course
Oued Majjou (Zaouiet El Habiyyne)	Zaouiet El Habiyyne	Chefchaouen	800	35°06.784'N, 05°11.784'W	River of the higher course
Oued Mila	Route de Ouazzane	Ouazzane	172	35°02.086'N, 05°22.133'W	River of the middle course
Oued Ouara	Ikedjouen	Chefchaouen	680	35°3.987'N, 05°14.045'W	River of the middle course
Oued Ouringa	Jebha	Chefchaouen	17	35°11.438'N, 04°41.150'W	Downstream of the river
Oued Sidi Yahya Aârab	Sidi Yahya Aârab	Chefchaouen	62	35°17.545'N, 04°53.503'W	River of the middle course
Oued Taïda	Taïda	Larache	505	35°22.250'N, 05°32.355'W	River of the middle course
Oued Tamerte	Belota	Chefchaouen	114	34°57.607'N, 05°32.061'W	River of the middle course
Oued Tizelkhte	Larbaa Beni Hassan	Tétouan	728	35°18.600'N, 05°23.097'W	Stream of the middle course
Oued Tkaraâ	Jbel Bouhachem	Larache	959	35°16.063'N, 05°25.829'W	River of the higher course
Oued Zarka	Kirané	Tétonian	52	35°32.412'N, 05°20.393'W	River of the middle course
Oued Zendoula	15 km North Ouazzane	Ouazzane	108	34°55.707'N, 05°31.942'W	River of the middle course
Seguia Mtahen	Mtahen, Jbel Bouhachem	Larache	983	35°16.226'N, 05°26.212'W	Small stream
Beni Snassen					
Grotte du Chaneau	Zegzel, Beni Snassen	Berkane	427	34°50.447'N, 02°21.532'W	Irrigation canal
Oued Tafoughalt	Tafoughalt, Beni Snassen,	Berkane	751	34°48.941'N, 02°24.471'W	River of the Middle course
Oued Beni Ouachekradi	Beni Snassen	Berkane	534	34°50.974'N, 02°16.217'W	River of the middle course
Middle Atlas Mts					
Barrage Allal El Fassi	Gîte Aït Ayoub	Sefrou	537	33°55.446'N, 4°40.558'W	Dam
High Atlas Mts					
Oued Sidi Fares	Sidi Fares, Parc National de Toubkal	Marrakech	1807	31°14.029'N, 07°52.813'W	River of the higher course
Assif Haouz	Imilil, Parc National De Toubkal	Marrakech	1896	31°07.424'N, 7°55.083'W	Stream
Sahara					
Oued Zag	Zag	Assa-Zag	400	28°01.344'N, 09°17.926'W	River of the middle course

Systematic part

Limoniidae: Chioneinae

Cheilotrichia (Empeda) cinerascens (Meigen, 1804)

Material examined. Oued Ouara, 1♂, 2♀♀, 23.XI.2012; Oued Amsa, 2♂♂, 13.XII.2013; Oued Majjou (Hafa meqlouba), 1♂, 10.V.2014; Grotte du Chameau, 1♂, 3♀♀, 24.XI.2014; Aïn Quanquben, 1♂, 28.IV.2015 (sweep net).

Distribution. Europe, Morocco, Georgia, Turkey, Cyprus, Israel, Iran. First record from Beni Snassen.

Cheilotrichia (Empeda) fuscohalterata (Strobl, 1906)

Material examined. Daya Fifi, 1♂, 23.XI.2012; Tributary of Oued El Fondak, 2♂♂, 10.II.2013, 1♂, 10.IV.2013; Barrage Ajras, 2♂♂, 10.II.2013; Daya Mghara, 6♂♂, 14.II.2013; Oued Zarka, 1♂, 23.II.2015; Oued Tizekhte, 1♂, 2♀♀, 28.II.2015 (sweep net).

Distribution. Italy, Portugal, Spain, First records from Morocco and North Africa.

Cheilotrichia (Empeda) minima (Strobl, 1898)

Material examined. Oued Zendoula, 2♂♂, 31.V.2013; Oued Jnane Niche, 1♂, 1♀, 25.IV.2015 (sweep net).

Distribution. Known from Europe (except for the north) and the south of the Palaearctic as far east as Kyrgyzstan. First records from Morocco and North Africa.

Ellipteroides (Ellipteroides) lateralis (Macquart, 1835)

Material examined. Daya Afrate, 3♂♂, 18.IV.2015 (sweep net).

Distribution. Southern part of Europe, Morocco, Lebanon, Israel and Turkey. First record from the Rif.

Erioconopa diuturna (Walker, 1848)

Material examined. Daya Jbel Zemzem, 1♂, 23.IV.2014; Oued Mezine, 4♂♂, 18.IV.2015; Aïn Bab Tariouente, 2♂♂, 2♀♀, 28.IV.2015 (sweep net).

Distribution. Europe, Morocco, Turkey. First record from the Rif.

***Erioptera (Erioptera) fuscipennis* Meigen, 1818**

Material examined. Aïn El Ma Bared, 2♂♂, 30.III.2012; Oued Amsemil (Site 1), 1♂, 1♀, 27.IV.2012; Oued Majjou (Pont), 2♂♂, 15.V.2012; Oued Tkaraâ, 1♂, 30.V.2013; Daya near Aïn Afersiw, 3♂♂, 1♀, 11.VI.2013; Aïn Afersiw, 3♂♂, 1♀, 11.VI.2013; Oued Abou Bnar, 2♂♂, 18.V. 2014; Oued Majjou (Zaouiet El habtiyne), 9♂♂, 6♀♀, 12.VII.2014; Daya Amsemil, 2♂♂, 2.XI.2014; Daya Aïn Jdioui, 12♂♂, 5♀♀, 28.III.2015; Daya Afrate, 2♂♂, 2♀♀, 18.IV.2015; Oued Mezine, 1♂, 18.IV.2015 (sweep net).

Distribution. Europe, Morocco, Algeria, Azerbaijan, Iran.

***Erioptera (Erioptera) lutea lutea* Meigen, 1804**

Material examined. Aïn Boughaba, 1♂, 24.V.2013 (sweep net).

Distribution. Europe, Transcaucasia, Turkey, Israel, ?Iran, Russia, Kazakhstan, Uzbekistan, Tajikistan, Kyrgyzstan. First records from Morocco and North Africa.

***Gonomyia (Gonomyia) abscondita* Lackschewitz, 1935**

Material examined. Maison forestière, 3♂♂, 21.IV.2015 (sweep net).

Distribution. Austria, Bulgaria, Czech Rep., Finland, Germany, Great Britain, Italy, Netherlands, Romania, Slovakia, Sweden, Ukraine. First records from Morocco and North Africa.

***Gonomyia (Gonomyia) subtenella* (Meigen, 1818),**

Material examined. Oued Beni Ouachekradi, 2♂♂, 30.XII.2014, 1♀, 8.I.2015 (reared); Oued Jnane Niche, 1♂, 25.IV.2015; Oued Sidi Yahya Aârab, 1♂, 3♀♀, 25.IV.2015 (sweep net).

Distribution. Czech Rep., Lithuania, Macedonia, Slovakia, Sweden, Morocco (High Atlas), Georgia, Azerbaijan, Iran. First records from Rif and Beni Snassen.

***Gonomyia (Gonomyia) sicula* Lackschewitz, 1940**

Material examined. Oued El Kebir, 1♂, 17.IV.2013; Daya Jbel Zemzem, 7♂♂, 23.IV.2014; Aïn Sidi Brahim Ben Arrif (Site 2), 1♂, 25.IV.2014; Daya Tazia, 1♂, 25.IV.2014; Aïn El Maounzil, 1♂, 21.IV.2015 (sweep net).

Distribution. Italy, Spain, Tunisia, Turkey. First records from Morocco.

***Hoplolabis (Parilisia) sororcula* (Lackschewitz, 1940)**

Material examined. Barrage Ajras, 3♂♂, 1♀, 10.II.2013; Oued Ouringa, 1♂, 19.IV.2013; Oued El Kanar, 1♂, 14.VI.2013; Oued Majjou (Hafa meqlouba), 1♂, 10.V.2014 (sweep net).

Distribution. France, Germany, Italy (incl. Sicily), Portugal, Romania, Slovakia, Spain, Switzerland, Ukraine, Morocco. First records from the Rif.

***Idiocera (Idiocera) ampullifera* (Stary, 1979)**

Material examined. Oued Zag, 1♂, 1♀, 22.V.2015 (sweep net).

Distribution. Egypt, Israel. First record from Morocco.

***Idiocera (Idiocera) pulchripennis* (Loew, 1856)**

Material examined. Oued Jnane Niche, 1♂, 2♀♀, 19.IV.2013, 3♂♂, 1♀, 14.VI.2013; Oued Aârkoub, 1♂, 19.IV.2013, 1♂, 16.III.2014; Aïn Jdioui, 4♂♂, 2♀♀, 28.III.2015 (sweep net).

Distribution. Southern part of Europe, Canary Islands, Morocco, Algeria, Egypt, Transcaucasia, Turkey, Cyprus, Israel, Iran, Central Asia. First record from the Rif.

***Idiocera (Idiocera) sziladyi* (Lackschewitz, 1940)**

Material examined. Oued Zarka, 1♂, 22.V.2014 (light trap).

Distribution. Europe (except for northern countries), Algeria, Egypt, Yemen. First record from Morocco.

***Ilisia maculata* (Meigen, 1804)**

Material examined. Oued Tafoughalt, 2♂♂, 25.XI.2014; Oued Majjou (Hafa meqlouba), 1♂, 27.IV.2015 (sweep net).

Distribution. Europe, Transcaucasia, Turkey, Iran. First records from Morocco and North Africa.

***Molophilus (Molophilus) obscurus* (Meigen, 1818)**

Material examined. Tributary of Oued Ouara, 1♂, 27.IV.2013, 5♂♂, 6.V.2014; Tributary of Oued Majjou, 2♂♂, 03.V.2013; Daya El Ânassar, 1♂, 24.V.2013; Daya Jbel Zemzem, 2♂♂, 23.IV.2014; Aïn El Ma Bared, 1♂, 1♀, 6.V.2014; Daya Rmali,

1♂, 17.V.2014; Oued Amsemlil (Site 1), 1♂, 2.XI.2014; Aïn Mâaze, 1♂, 1.XI.2014; Daya Afrate, 9♂♂, 18.IV.2015; Daya Mtahen, 1♂, 25.V.2015 (sweep net).

Distribution. Europe, Morocco (High Atlas), Transcaucasia, Turkey, Cyprus, Lebanon, Israel. First records from the Rif.

Molophilus (Molophilus) propinquus propinquus (Egger, 1863)

Material examined. Oued Ouara, 1♂, 7.I.2013 (reared); Oued Farda, 3♂♂, 24.IV.2013; Tributary of Oued Taïda, 1♂, 25.IV.2014; Oued Majjou (Hafa meqlouba), 1♂, 27.IV.2015 (sweep net).

Distribution. Europe, Morocco (High Atlas), Georgia, Turkey. First records from the Rif.

Molophilus (Molophilus) testaceus Lackschewitz, 1940

Material examined. Daya Amsemlil, 1♂, 23.V.2012, 1♂, 1♀, 30.V.2013; Daya Mtahen, 2♀♀, 23.V.2012; Marje El kheyl, 3♂♂, 5♀♀, 30.V.2013; Oued Tkaraâ, 1♂, 30.V.2013; Daya near Aïn Afersiw, 1♂, 11.VI.2013; Daya Mezine, 1♂, 11.VI.2013; Daya Tazia, 1♂, 25.IV.2014; Daya Rmali, 1♂, 17.V.2014 (sweep net). Daya Amsemlil, 1♂, 4♀♀, 9.VII.2013; Daya near Aïn Afersiw, 4♂♂, 1♀, 2.VII.2013 (reared).

Distribution. Only known from Portugal and Spain. First records from Morocco and North Africa.

Symplecta (Symplecta) hybrida (Meigen, 1804)

Material examined. Barrage Ajras, 1♂, 10.II.2013; Barrage Allal El Fassi, 1♂, 15.IV.2014; Oued El Kanar, 1♀, 14.VI.2013, 1♂, 25.IV.2015; Oued Aârkoub, 1♂, 16.III.2014; Oued Jnane Niche, 1♀, 25.IV.2015 (sweep net).

Distribution. Nearctic (Canada, USA, Greenland); widespread in Palaearctic, including North Africa, Central Asia, Mongolia, as far east as North Korea and Japan; Oriental (India, Nepal, Pakistan). First records from the Rif and the Middle Atlas.

Symplecta (Trimicra) pilipes (Fabricius, 1787)

Material examined. Tributary of Oued Hachef, 2♂♂, 2♀♀, 14.II.2013; Oued Aârkoub, 1♂, 19.IV.2013; Oued El Kanar, 2♂♂, 2♀♀, 14.VI.2013; Oued Mezine, 1♂, 18.IV.2015 (sweep net); Aïn Chouk, 1♂, 3.II.2015 (reared).

Distribution. The only cosmopolitan cranefly, widespread in all zoogeographical regions. First records from the Rif.

***Tasiocera (Dasymolophilus) murina* (Meigen, 1818)**

Material examined. Oued Farda, 1♂, 28.III.2013; Oued Amsemilil (Site 3), 4♂♂, 30.V.2013 (sweep net).

Distribution. Europe, Transcaucasia, Turkey. First records from Morocco and North Africa.

Limoniidae: Limnophilinae***Austrolimnophila (Austrolimnophila) latistyla* Stary, 1977**

Material examined. Oued Majjou (Hafa meqlouba), 1♂, 27.VI.2015 (sweep net).

Distribution. Croatia, France, Greece, Italy, Portugal, Spain. First record from Morocco and North Africa.

***Dicranophragma (Brachylinnophila) nemorale* (Meigen, 1818)**

Material examined. Aïn Sidi Brahim Ben Arrif (Site 1), 1♂, 25.IV.2014; Tributary Oued Ouara, 2♀♀, 27.IV.2013; Oued Majjou (Hafa meqlouba), 1♂, 10.V.2014 (sweep net).

Distribution. Widespread in the Palaearctic. First records from the Rif.

***Euphylidorea crocotula* (Seguy, 1941)**

Material examined. Aïn Bab Tariouente, 3♂♂, 28.IV.2015 (sweep net).

Distribution. Spain (Granada), Morocco (High Atlas). First record from the Rif.

***Euphylidorea (Euphylidorea) dispar* (Meigen, 1818)**

Material examined. Oued at 15 Km from Fifi, 1♀, 26.III.2014 (sweep net).

Distribution. Widespread in Europe, recorded from Russia. First records from Morocco and North Africa.

***Hexatoma (Hexatoma) bicolor* (Meigen, 1818)**

Material examined. Oued at 15 Km from Fifi, 1♂, 27.IV.2013, 1♀, 26.III.2014, 3♂♂, 6.V.2014; Tributary of Oued Ouara, 1♀, 27.IV.2013; Oued Madissouka,

1♂, 18.V.2014; Oued Majou (Hafa meqlouba), 1♀, 10.V.2014; Aïn Quanquben, 1♂, 28.IV.2015; Oued Tkaraâ, 2♂♂, 7.V.2015; Oued Tamerte, 1♂, 1♀, 7.V.2015 (sweep net).

Distribution. Europe, Georgia, Turkey. First records from Morocco and North Africa.

Pseudolimnophila (Pseudolimnophila) sepium (Verrall, 1886)

Material examined. Daya Tazia, 11♂♂, 8♀♀, 23.IV.2012; Aïn Sidi Brahim Ben Arrif (2), 2♂♂, 2♀♀, 17.IV.2013; Aïn Afersiw, 1♀, 11.VI.2013; Oued Majou (Hafa meqlouba), 1♂, 10.V.2014 (sweep net). Daya Tazia, 1♂, 2♀♀, 14.V.2013; Oued Taïda, 1♂, 12.V.2014; Aïn Sidi Brahim Ben Arrif (Site 2), 2♂♂, 1♀, 26.V.2014 (reared).

Distribution. Europe, Morocco (Rif), Transcaucasia, Turkey, Turkmenistan, Kyrgyzstan, Afghanistan.

Limoniidae: Limoniinae

Dicranomyia (Dicranomyia) affinis (Schummel, 1829)

Material examined. Oued Amsemil (Site 1), 1♂, 27.IV.2012; Daya Mtahen, 10♂♂, 1♀, 27.IV.2012; Oued Tkaraâ, 1♂, 27.IV.2012; Marj El Kheyl, 2♀♀, 30.V.2013; Aïn Sidi Brahim Ben Arrif (Site 2), 5♂♂, 1♀, 17.IV.2013, 3♂♂, 1♀, 25.IV.2014; Aïn Sidi Brahim Ben Arrif (Site 1), 10♂♂, 5♀♀, 25.IV.2014; Daya Tazia, 14♂♂, 4♀♀, 25.IV.2014; Aïn El Maounzil, 6♂♂, 4♀♀, 21.IV.2015; Aïn El Malâab, 1♂, 1♀, 21.IV.2015; Daya near Aïn Afersiw, 5♂♂, 3♀♀, 18.IV.2015; Daya Mtahen, 4♀♀, 7.V.2015; Aïn Bab Tariouente, 4♂♂, 28.IV.2015 (sweep net).

Distribution. Great Britain, Ireland, Poland, Morocco (High Atlas).

Dicranomyia (Dicranomyia) chorea (Meigen, 1818)

Material examined. Oued Sidi Fares, 1♂, 7.V.2011; Oued Farda, 2♂♂, 2♀♀, 13.II.2013, 7♂♂, 4♀♀, 23.IV.2013; Oued Kelâa, 3♂♂, 13.II.2013; Aïn Ras El Ma, 1♂, 3.V.2013; Oued Jnane Niche, 2♂♂, 2♀♀, 25.IV.2015; Cascade Chrafate, 1♂, 2♀♀, 28.IV.2015; Maison forestière, 2♀♀, 21.IV.2015; Oued El Kanar, 1♂, 25.IV.2015 (sweep net). Tributary of Oued Zarka, 4♂♂, 1♀, 9.XII.2013 (reared).

Distribution. Nearctic (Canada), Europe, Morocco, Transcaucasia, Turkey, Cyprus, Israel, Iran. First records from the Rif.

***Dicranomyia (Dicranomyia) goritiensis* (Mik, 1864)**

Material examined. Oued Majjou (Hafa meqlouba), 1♂, 3.V.2013, 2♂♂, 10.V.2014; Cascade Chrafate, 1♀, 24.V.2013; Oued El Koub, 1♂, 31.V.2013; Cascade Zarka, 1♂, 14.XI.2013; Âounsal Aheramen, 1♂, 2♀♀, 10.V.2014, 2♂, 27.IV.2015; Maison forestière, 1♂, 21.IV.2015 (sweep net).

Distribution. Europe, Morocco, Algeria, Georgia, Turkey, Israel. First records from the Rif.

***Dicranomyia (Dicranomyia) longicollis* (Macquart, 1846)**

Material examined. Oued Aârate, 4♂♂, 3♀♀, 26.III.2014; Barrage Moulay Bouchta, 2♂♂, 5.IV.2014; Barrage Allal El Fassi, 1♂, 15.IV.2014; Oued El Kebir, 1♀, 23.IV.2014 (sweep net).

Distribution. Portugal, Spain, Morocco, Algeria. First records from the Rif and the Middle Atlas.

***Dicranomyia (Dicranomyia) modesta* (Meigen, 1818)**

Material examined. Oued Farda, 1♂, 24.IV.2013; Oued El Kanar, 4♂♂, 3♀♀, 14.VI.2013, 5♂♂, 10♀♀, 25.IV.2015; Oued Jnane Niche, 2♂♂, 16.III.2014; Oued Sidi Yahya Aârab, 4♂♂, 1♀, 25.IV.2015 (sweep net).

Distribution. Nearctic (Canada, USA, Greenland), widespread over Palaearctic, including Central Asia and Mongolia, as far east as Far East of Russia and Japan. First records from Morocco and North Africa.

***Dicranomyia (Dicranomyia) novemmaculata* (Strobl, 1906)**

Material examined. Tributary of Oued el Fondak, 1♂, 10.II.2013; Oued Aârate, 9♂♂, 4♀♀, 26.III.2014 (sweep net).

Distribution. Gibraltar, Portugal, Spain, Algeria. First records from Morocco.

***Dicranomyia (Melanolimonia) hamata* Becker, 1908**

Material examined. Tributary of Oued El Kebir, 2♂♂, 1♀, 18.III.2011; Oued Aârate, 2♂♂, 26.III.2014; Aïn Sidi Brahim Ben Arrif (Site 1), 2♂♂, 1♀ 25.IV.2014; Daya Tazia, 1♂, 25.IV.2014 (sweep net).

Distribution. France, Portugal, Spain, Turkey. First records from Morocco and North Africa.

***Dicranomyia (Dicranomyia) ventralis* (Schummel, 1829)**

Material examined. Lake Badriouen, 2♂♂, 26.IV.2011 (sweep net).

Distribution. Europe, Egypt, Azerbaijan, Turkey, Israel, Iran, Russia, Kazakhstan, Kyrgyzstan, Afghanistan, North Korea, India. First records from Morocco and North Africa.

***Dicranomyia (Melanolimonia) morio* (Fabricius, 1787)**

Material examined. Seguia Mtahen, 1♂, 30.V.2013; Daya near Aïn Afersiw, 2♀♀, 11.VI.2013; Daya Jbel Zemzem, 1♂, 23.IV.2014 (sweep net).

Distribution. Europe, Morocco, Transcaucasia, Turkey, Iran, Mongolia. First records from the Rif.

***Dicranoptycha fuscescens* (Schummel, 1829)**

Material examined. Oued Mlilah, 3♀♀, 31.V.2013; Oued Loukous, 4♂♂, 1♀, 6.V.2015; Oued Zendoula, 1♀, 6.V.2015 (sweep net).

Distribution. Europe, Morocco, Algeria, Transcaucasia, Turkey, Cyprus, Lebanon, Israel, ?Kazakhstan, Mongolia.

***Helius (Helius) hispanicus* Lackschewitz, 1928**

Material examined. Oued Amsemil (Site 3), 1♀, 30.V.2013 (sweep net).

Distribution. Great Britain, Spain, Portugal, Morocco (High Atlas), Georgia, Azerbaijan, Turkey, Cyprus, Iran. First record from the Rif.

***Helius (Helius) pallirostris* Edwards, 1921**

Material examined. Aïn Chouk, 1♂, 1♀, 25.II.2015 (reared).

Distribution. Europe, Tunisia, Azerbaijan, Israel, Iran, Central Asia. First record from Morocco.

***Limonia nubeculosa* (Meigen, 1804)**

Material examined. Tributary of Oued El Fondak, 1♀, 10.II.2013; Aïn Ras El Ma, 1♀, 3.V.2013; Aïn Boughaba, 3♂♂, 4♀♀, 24.V.2013; Oued Azila, 1♀, 27.VI.2013; Aounsar Aheramen, 2♂♂, 1♀, 10.V.2014; Aïn Takhnjoute, 1♂, 17.V.2014; Mai-

son forestière, 1♂, 4♀♀, 17.V.2014; Oued Madissouka, 1♂, 18.V.2014; Aïn Quanquaben, 1♀, 28.IV.2015 (sweep net).

Distribution. Nearctic (Canada, USA), widespread over Palaearctic, including North Africa, Central Asia, and Mongolia, as far east as Far East of Russia, North Korea, and Japan. Recorded from Morocco in the High Atlas. First records from the Rif.

Limonia phragmitidis (Schrink, 1781)

Material examined. Aïn Quanquaben, 1♂, 27.VI.2013 (sweep net).

Distribution. Widespread in the West Palaearctic, also known from Russia, Kazakhstan and Kyrgyzstan. Recorded from Morocco (Middle Atlas). First record from the Rif.

Pediciidae

Dicranota (Paradicranota) landrocki Czizek, 1931

Material examined. Oued Ouara, 2♂♂, 22.XI.2013; Assif Haouz, 1♂, 17.IV.2014; Oued Taïda, 1♀, 25.IV.2014; Aïn Sidi Brahim Ben Arrif (Site 2), 2♂♂, 25.IV.2014; Âounsar Aheramen, 1♂, 10.V.2014, 1♂, 27.IV.2015; Oued Tizekhte, 1♂, 28.II.2015; Oued Mezine, 1♂, 18.IV.2015; Maison forestière, 32♂♂, 2♀♀, 21.IV.2015; Aïn Bab Tariouente, 1♂, 28.IV.2015 (sweep net).

Distribution. Europe, Russia (North Caucasus), Morocco (Rif), Transcaucasia Lebanon, Tajikistan. First record from the High Atlas.

Dicranota (Ludicia) claripennis (Verrall, 1888)

Material examined. Oued Amsemilil (Site 1), 1♂, 16.XII.2013; Maison forestière, 1♂, 21.IV.2015 (sweep net).

Distribution. Austria, Belgium, France, Germany, Great Britain, Ireland, Italy, Netherlands, Spain. First records from Morocco and North Africa.

Tricyphona (Tricyphona) immaculata (Meigen, 1804)

Material examined. Maison forestière, 1♂, 21.IV.2015; Daya Mtahen, 1♂, 7.V.2015 (sweep net).

Distribution. Europe, Transcaucasia, Turkey, Lebanon, West Siberia, Central Asia. First records from Morocco and North Africa.

Discussion

Altogether a number of 570 specimens, most of which (521) belongs to the Limoniidae distributed in 20 genera, whereas only 49 specimens belong to the Pediciidae distributed in 2 genera.

This study provides an important contribution to the short-palped crane flies fauna of North Africa and Morocco, particularly to the Rif region where 35 species were recorded from the first time. The most abundant species was *Dicranomyia (Dicranomyia) affinis*, with 86 specimens, followed successively by, *Erioptera (Erioptera) fuscipennis* (56), *Dicranota (Paradicranota) landrocki* (45), *Dicranomyia (Dicranomyia) chorea* (35), *Pseudolimnophila (Pseudolimnophila) sepium* (32), *Dicranomyia (Dicranomyia) modesta* (30) and *Molophilus (Molophilus) testaceus* (29). The most frequently collected species were *Erioptera (Erioptera) fuscipennis* (12 sites), *Dicranomyia (Dicranomyia) affinis* (11 sites), *Molophilus (Molophilus) obscurus* (10 sites), followed by *Dicranomyia (Dicranomyia) chorea*, *Limonia nubeculosa* and *Dicranota (Paradicranota) landrocki*, all three collected in 9 sites.

18 species of Limoniidae and 2 species of Pediciidae are newly recorded from Morocco, of which 15 species are provided to be new to North Africa. Those new records extend the distribution of the species to most southern localities of the western Palearctics in Africa. *Idiocera (Idiocera) ampullifera* represents the most southern record (Sahara) of the Limoniidae in Morocco.

Including our results, the total number of species of Limoniidae and Pediciidae known from Morocco now increases from 53 species to 73 (see checklist).

These new data of crane flies fauna reflect the variety of suitable habitats prospected, such as are springs, mountain rivers and streams, and stagnant water mainly belonging to forest areas. This suggests that many more species and subspecies can yet be found in Morocco, that enjoys a wide variety of geographical and climatic properties.

Check-list of the Moroccan short-palped crane flies

Limoniidae

Chioneinae

Baeoura ebenina Starý, 1981

Baeoura staryi Driauach & Belqat, 2015

Cheilotrichia (Empeda) cinerascens (Meigen, 1804)

Cheilotrichia (Empeda) fuscohalterata* (Strobl, 1906)

Cheilotrichia (Empeda) minima* (Strobl, 1898)

Ellipterooides (Ellipterooides) lateralis (Macquart, 1835)

Ellipterooides (Protogonomyia) alboscutellatus (Von Roser, 1840)

Ellipterooides (Protogonomyia) hutsoni (Starý, 1971)

Eriocnopa diuturna (Walker, 1848)

Eriocnopa symplectoides (Kuntze, 1914)

- Erioptera (Erioptera) fuscipennis* Meigen, 1818
- Erioptera (Erioptera) lutea lutea* Meigen, 1804***
- Erioptera (Erioptera) transmarina* Bergroth, 1889
- Gonomyia (Gonomyia) abscondita* Lackschewitz, 1935***
- Gonomyia (Gonomyia) subtenella* Savchenko, 1972
- Gonomyia (Gonomyia) sicula* Lackschewitz, 1940***
- Gonomyia (Gonomyia) tenella* (Meigen, 1818)
- Hoplolabis (Parilisia) obtusiapex* (Savchenko, 1982)
- Hoplolabis (Parilisia) punctigera* (Lackschewitz, 1940)
- Hoplolabis (Parilisia) sororcula* (Lackschewitz, 1940)
- Idiocera (Euptilosstena) jucunda* (Loew, 1873)
- Idiocera (Idiocera) ampullifera* (Stary, 1979)***
- Idiocera (Idiocera) pulchripennis* (Loew, 1856)
- Idiocera (Idiocera) sziladyi* (Lackschewitz, 1940)***
- Ilisia maculata* (Meigen, 1804)***
- Molophilus (Molophilus) ibericus* Stary, 2011
- Molophilus (Molophilus) obscurus* (Meigen, 1818)
- Molophilus (Molophilus) propinquus propinquus* (Egger, 1863)
- Molophilus (Molophilus) testaceus* Lackschewitz, 1940***
- Symplecta (Symplecta) hybrida* (Meigen, 1804)
- Symplecta (Trimicra) pilipes* (Fabricius, 1787)
- Tasiocera (Dasymolophilus) murina* (Meigen, 1818)***

Dactylolabinae

- Dactylolabis (Dactylolabis) symplectoidea* Egger, 1863

Limnophilinae

- Austrolimnophila (Austrolimnophila) latistyla* Stary, 1977***
- Dicranophragma (Brachylimnophila) adjunctum* (Walker, 1848)
- Dicranophragma (Brachylimnophila) nemorale* (Meigen, 1818)
- Eloeophila maroccana* Stary, 2009
- Euphylidorea (Euphylidorea) crocotula* (Seguy, 1941)
- Euphylidorea (Euphylidorea) dispar* (Meigen, 1818)***
- Euphylidorea (Euphylidorea) lineola* (Meigen, 1804)
- Hexatoma (Hexatoma) bicolor* (Meigen, 1818)***
- Hexatoma (Hexatoma) gaedii* (Meigen, 1830)
- Pseudolimnophila (Pseudolimnophila) sepium* (Verrall, 1886)

Limoniinae

- Dicranomyia (Dicranomyia) affinis* (Schummel, 1829)
- Dicranomyia (Dicranomyia) chorea* (Meigen, 1818)
- Dicranomyia (Dicranomyia) didyma* (Meigen, 1804)
- Dicranomyia (Dicranomyia) goritiensis* (Mik, 1864)
- Dicranomyia (Dicranomyia) longicollis* (Macquart, 1846)

- Dicranomyia (Dicranomyia) lutea* Meigen
Dicranomyia (Dicranomyia) mitis (Meigen, 1830)
Dicranomyia (Dicranomyia) modesta (Meigen, 1818)*
Dicranomyia (Dicranomyia) novemmaculata (Strobl, 1906)*
Dicranomyia (Dicranomyia) ventralis (Schummel, 1829)*
Dicranomyia (Glochina) sericata (Meigen, 1830)
Dicranomyia (Melanolimonia) hamata* Becker, 1908
Dicranomyia (Melanolimonia) morio (Fabricius, 1787)
Dicranomyia majuscula Pierre, 1924
Dicranoptyla fuscescens (Schummel, 1829)
Geranomyia caloptera (Mik, 1867)
Geranomyia obscura Strobl, 1900
Helius (Helius) hispanicus Lackschewitz, 1928
Helius (Helius) pallirostris* Edwards, 1921
Limonia flavipes (Fabricius, 1787)
Limonia hercegovinae (Strobl, 1898)
Limonia macrostigma (Schummel, 1829)
Limonia nubeculosa Meigen, 1804
Limonia phragmitidis (Schrank, 1781)

Pediciidae

Pediciinae

- Dicranota (Dicranota) bimaculata* (Schummel, 1829)
Dicranota (Dicranota) irregularis Pierre, 1922
Dicranota (Paradicranota) candelisequa Stary, 1981
Dicranota (Paradicranota) landrocki Czizek, 1931
Dicranota (Ludicia) claripennis* (Verrall, 1888)
Tricyphona (Tricyphona) immaculata (Meigen, 1804)

Acknowledgements

The authors sincerely thank Dr. Jaroslav Starý (Olomouc, Czech Republic) for his valuable help concerning the identification of some species.

References

- Driauach O, Belqat B (2015) A new species of the genus *Baeoura* from Morocco, with a key to the West Palaearctic species (Diptera: Tipuloidea: Limoniidae). ZooKeys 532: 99–105. doi: 10.3897/zookeys.532.5994
- Driauach O, Belqat B, De Jong H (2013) A first checklist of the short-palped crane flies (Diptera: Limoniidae, Pediciidae) of Morocco. Boletín de la Sociedad Entomológica Aragonesa 53: 187–190.

- Eiroa E (2000) Primera cita de *Dicranoptycha fusescens* (Schummel, 1829) para Marruecos (Diptera: Limoniidae). Boletín de la Asociación Española de Entomología 24: 204.
- De Jong H, Oosterbroek P, Gelhaus J, Reusch H, Young C (2008) Global biodiversity of craneflies (Insecta, Diptera: Tipulidae or Tipulidae sensu lato) in freshwater. Hydrobiologia 595: 457–467. doi: 10.1007/s10750-007-9131-0
- Oosterbroek P (2015) Catalogue of the Craneflies of the World (Insecta, Diptera, Nematocera, Tipuloidea). <http://ccw.naturalis.nl/index.php> [version 26 November 2015]
- Pârvu C, Răzvan PM, Răzvan Z (2006) Faunistic data on some dipteran families (Insecta: Diptera) from Morocco. Travaux du Muséum d'Histoire Naturelle "Grigore Antipa" 49: 271–281.
- Pierre C (1922a) Nematocera Polyneura recueillis au Maroc par M. Charles Alluaud (1919–1920) (Insectes Diptères). Bulletin de la Société des Sciences naturelles du Maroc 1: 21–24.
- Pierre C (1922b) Nematocera Polyneura recueillis au Maroc par M. Charles Alluaud (2e liste, 1920–1921) (Insectes Diptères). Bulletin de la Société des Sciences naturelles du Maroc 1: 148–151.
- Pierre C (1924) Nematocera Polyneura receuillis au Maroc par M. Charles Alluaud (3e liste, 1922–1923) (Insectes Diptères). Bulletin de la Société des Sciences naturelles du Maroc 4: 198–201.
- Reusch H, Oosterbroek P (1997) Diptera Limoniidae and Pediciidae, Short-palped Crane Flies. In: Nilsson A (Ed.) Aquatic Insects of North Europe 2: 105–132.
- Savchenko EN, Oosterbroek P, Starý J (1992) Family Limoniidae. Catalogue of Palaearctic Diptera 1: 183–369.
- Séguy E (1941a) Récoltes de R. Paulian et A. Villiers dans le haut Atlas marocain, 1938, (XVIIe note) Diptères. Revue Française d'Entomologie 8: 25–33.
- Séguy E (1941b) Diptères receuillis par M. Berland dans le sud marocain. Annales de la Société Entomologique de France 110: 1–23.
- Starý J (1971) A new palaearctic representative of the subgenus *Protogonomyia* Alexander (Diptera, Tipulidae). Acta Entomologica Bohemoslovaca 68: 319–321.
- Starý J (1974) The identity of *Gonomyia (Idiocera) sexguttata* (Diptera, Tipulidae). Acta Entomologica Bohemoslovaca 71: 136–140.
- Starý J (2006) *Hoplolabis (Parilisia)* species related to *H. (P.) punctigera* (Lackschewitz, 1940) and *H. (P.) spinosa* (Nielsen, 1953) with the description of a new species (Diptera, Limoniidae). Studia dipterologica 13: 115–125.
- Starý J (2009) West Palaearctic species of the genus *Eloeophila* (Diptera: Limoniidae). European Journal of Entomology 106: 425–440. doi: 10.14411/eje.2009.054
- Starý J (2011) Descriptions and records of the Palaearctic *Molophilus* Curtis (Diptera, Limoniidae). Zootaxa 2999: 45–62.
- Starý J, Kubík S, Barták M (2005) Limoniidae. In: Barták M, Kubík S (Eds) Diptera of Podyjí National Park and its environs. Ceska Zemedelska Univerzita v Praze, 24–32.
- Starý J, Oosterbroek P (2008) New records of West Palaearctic Limoniidae, Pediciidae and Cylindrotomidae (Diptera) from the collections of the Zoological Museum, Amsterdam. Zootaxa 1922: 1–20.
- Vaillant F (1956) Recherches sur la faune madicole (Hygropetriques.l.) de France, de Corse et d'Afrique de Nord. Mémoires du Museum National d'Histoire Naturelle (N.S.), Série A, Zoologie 11: 1–257.