

Contributions to the knowledge of the genus *Horaeomorphus* Schaufuss (Coleoptera, Staphylinidae, Scydmaeninae) in mainland China

De-Yao Zhou¹, Su-Jiong Zhang², Li-Zhen Li¹

1 Department of Biology, College of Life and Environmental Sciences, Shanghai Normal University, 100 Guilin Road, Shanghai, 200234, P. R. China **2** Forestry Bureau of Pan'an County, Pan'an 322300, Zhejiang, China

Corresponding author: Li-Zhen Li (scydmaeninae@163.cn)

Academic editor: P. Stoev | Received 10 December 2015 | Accepted 12 February 2016 | Published 15 March 2016

<http://zoobank.org/2427CCB8-B274-4D96-83DE-391125C5F8BC>

Citation: Zhou D-Y, Zhang S-J, Li L-Z (2016) Contributions to the knowledge of the genus *Horaeomorphus* Schaufuss (Coleoptera, Staphylinidae, Scydmaeninae) in mainland China. *ZooKeys* 572: 51–70. doi: 10.3897/zookeys.572.7474

Abstract

Five new species of the ant-like stone beetle genus *Horaeomorphus* Schaufuss (Scydmaeninae: Glandulariini) from China are described: *H. hainanicus* sp. n., *H. biwenxuani* sp. n., *H. pengzhong* sp. n., *H. huijiayaoi* sp. n. and *H. punctatus* sp. n. The previously unknown male of *H. chinensis* Franz, 1985 is now discovered, and its aedeagus and metatrochanter are illustrated. The latter species is newly recorded from Zhejiang. Three females from Guangxi are also recorded, but their identity remains unconfirmed until associated males become available. A key to *Horaeomorphus* of mainland China is included.

Keywords

Scydmaeninae, Glandulariini, *Horaeomorphus*, new species, new records, Oriental, China

Introduction

The Australo-Oriental genus *Horaeomorphus* Schaufuss currently comprises 59 species distributed in Malaysia, Singapore, Thailand, Nepal, Vietnam, Japan, continental China, Taiwan, Laos, the Philippines, Indonesia, Fiji, Australia, and Madagascar (Franz 1985; Jałoszyński 2002, 2003, 2004, 2006, 2009, 2012, 2014a, 2014b; Jałoszyński and Nomura 2004, 2008; Jałoszyński et al. 2007; Vít 2004); among them 21 Madagascan species were transferred from previous *Euconnus* (*Anthicimorphus*) together with the Australian type species of *Anthicimorphus*, but they should be treated as species *incertae sedis* within Glandulariini (Jałoszyński 2014b). Until now, only one species from mainland China has been described.

Among asian glandulariine (= former Cyrtoscydmini, see Newton 2015) genera, *Horaeomorphus* is characterized by the mesoventral intercoxal process being shorter and less elevated than the mesocoxae, the metaventral intercoxal process with two long spines projecting posteriorly, the presence of a small pit at the posteromesal margin of each supra-antennal tubercle, pronotal base with distinct median pit in addition to lateral pits, and each elytron with two foveae connected by a U-shaped groove extending anteriorly; many species have the male trochanters modified (Jałoszyński 2015).

Recent examination of unsorted material in our collection revealed five new species of *Horaeomorphus* from Hainan, Guangxi, Yunnan and Xizang. Moreover, during our recent expedition, two males of *H. chinensis* Franz, 1985 were discovered in the type locality, Fujian: Guadun (=Kuatun), which makes it possible to add a description of male characters. This species is also newly recorded from Zhejiang: Baishanzu. Three females from Guangxi are recorded; they belong to a group of species characterized by a strongly convex and broad body, but their identities remain unknown until associated males become available. A key to *Horaeomorphus* of mainland China is included.

Material and methods

All material treated in this study is housed in the Insect Collection of Shanghai Normal University, Shanghai, China (SNUC).

The collecting data are quoted verbatim. Each type specimen bears the following label: 'HOLOTYPE [red] (or PARATYPE [yellow]), ♂ (or ♀), *Horaeomorphus* + specific name sp. n., det. Zhou & Zhang, 2016, SNUC'.

The following abbreviations are applied: AeL—length of the median lobe of aedeagus in ventral view; AnL—length of the antennae; BL—length of the body (= HL + PL + EL); EI—elytral index (= EL / EW); EL—length of the elytra along the suture, from the base of scutellum to the apex; EW—maximum width of the elytra; HW—width of the head across eyes; HL—length of the head from the anterior clypeal margin to the occipital constriction; PL—length of the pronotum along the midline; PWb—width of the pronotum at base; PWm—maximum width of the pronotum; SpL—length of the spermatheca.

Taxonomy

Horaeomorphus chinensis Franz, 1985

Figs 1A, 6B–E, 6G–K

Horaeomorphus chinensis Franz, 1985: 116.

Material examined. 2 ♂♂, 4 ♀♀, labeled ‘China: Fujian Province, Wuyishan City [武夷山市], Tongmu Village [桐木村], Guadun [挂墩], 27°44'03"N, 117°38'40"E, decaying log, 1178m, 2.x.2015, Yan, Tu, Shen, Jiang & Zhou leg.’; 2 ♂♂, 2 ♀♀, labeled ‘China: Zhejiang, Lishui City [丽水市], Qingyuan Hsien [庆元县], Baishanzu [百山祖], Huangpi Swamp [黄皮湿地], 27°49'38"N, 119°11'29"E, decaying log, 1429m, 25.iv.2015, Song & Yan leg.’; 1 ♀, labeled ‘China: S.Zhejiang, Qingyuan, Mt nr. Liyang Village [栗洋村], 27°49'38"N, 119°11'22"E, leaf litter, sifted, 29.iv.2014, 990–1160m, Peng, Song, Yan & Yu leg.’

Description of male. Males (Fig. 1A) similar to females in external morphology, but metatrochanter (Fig. 6J) modified, with distal part protruded; slightly longer than half of metafemur, apex spiculate. BL 2.52–2.76 mm; HL 0.40–0.43 mm; HW 0.53–0.58 mm; AnL 1.05–1.17 mm, relative lengths of antennomeres: 1.0 : 0.9 : 1.4 : 1.1 : 1.1 : 1.0 : 1.0 : 1.0 : 1.1 : 1.1 : 2.0; PL 0.75–0.81 mm; PWb 0.46–0.52 mm; PWm 0.65–0.71 mm; EL 1.37–1.53 mm; EW 0.89–0.94 mm; EI 1.54–1.63.

Aedeagus (Fig. 6C–E) elongate, AeL 0.48 mm; endophallus (Fig. 6H–I) very complicated, with large bell-shaped central complex surrounded at each side by weakly sclerotized, elongate lateral structures; parameres (Fig. 6G) slender, longer than median lobe, each with three apical setae and two subapical setae.

Distribution. Eastern China: Fujian, Zhejiang (new provincial record).

Comments. *Horaeomorphus chinensis* shares many characters with several congeners: *H. punctifrons* Jałoszyński, 2006 (Malaysia: Pahang), *H. jeraianus* Jałoszyński, 2006 (Malaysia: Kedah), *H. jaechi* Jałoszyński, 2006 (Malaysia: Sarawak), *H. tiomanensis* Jałoszyński, 2006 (Malaysia: Tioman Is.), *H. caverniventris* Jałoszyński, 2006 (Malaysia: Pahang), *H. endauensis* Jałoszyński & Nomura, 2007 (Malaysia: Pahang), *H. imitator* Jałoszyński, 2009 (the Philippines: Mindanao), *H. solodownikovi* Jałoszyński, 2014 (Laos: Champasak), *H. sakishimanus* Jałoszyński, 2002 (Japan: Iriomote-jima and Ishigaki-jima islands), *H. hainanicus* sp. n. (China: Hainan) and *H. pengzhongii* sp. n. (China: Yunnan), all of them have elongate and convex habitus, the pronotum with three pits connected by a groove, the base of elytra barely wider than the basal margin of pronotum and protruded, recurved metatrochanters in males. Among these species, *H. hainanicus* sp. n. and *H. pengzhongii* sp. n. have a similar shape of the aedeagus, structures of endophallus and slender parameres each with two subapical setae, as those in *H. chinensis*. The character combination of the pronotum and elytra with a dense and distinct punctation, the recurved metatrochanters without expansions on their ventral margins in male, the parameres longer than median lobe and the structures of the endophallus can be used to distinguish *H. chinensis* from the two species mentioned above.



Figure 1. Habitus of *Horaeomorphus* species. **A** *H. chinensis*, male **B** *Horaeomorphus* sp., female. Scale bars: 1.0 mm.

Bionomics. Specimens from Guadun were collected by sifting material from an ant nest (Fig. 10B) in a relatively dry rotten trunk in a bamboo forest; four specimens from Baishanzu were collected from under bark of a rotten tree.

***Horaeomorphus* sp.**

Figs 1B, 6A, 6F

Material examined. 3 ♀♀, labeled ‘China: Guangxi Prov., Shangsi County [上思县], Shiwandashan [十万大山], alt. 300–400m, 23.iv.2011, Peng, Zhai & Zhu leg.’.

Description. Body (Fig. 1B) large, strongly convex, BL 2.48–2.51 mm; HL 0.37–0.38 mm; HW 0.50 mm; antennae (Fig. 6A) relatively short, AnL 0.99 mm; relative lengths of antennomeres: 1.0 : 0.8 : 1.2 : 1.0 : 1.0 : 0.9 : 0.9 : 0.9 : 1.0 : 1.0 : 2.2; PL 0.74–0.76 mm; PWm 0.59–0.61 mm; PWb 0.46–0.47 mm; EL 1.35–1.39 mm; EW 1.00–1.03 mm; EI 1.30–1.38.

Spermatheca (Fig. 6F) elongate, SpL 0.13mm; with longitudinal groove at middle.

Comments. These females have strongly convex and broadened elytra (EI 1.30–1.38), distance between humeral calli wider than the width of the pronotum at base, large and deep punctures sharply delimited from background in the center of pronotal disc and anterior third of elytra. The shape of the spermatheca is similar to that of *H. caverniventris* Jąłoszyński, 2006 (Malaysia: Pahang), but females of the Malaysian species are larger (2.79–3.12 mm). Therefore the three specimens almost certainly belong to a new species, but a male must be found for formal description.

Distribution. Southern China: Guangxi.

***Horaeomorphus hainanicus* D.-Y. Zhou & S.-J. Zhang, sp. n.**

<http://zoobank.org/754810B2-86FD-4684-8B7C-A4981F223105>

Figs 2, 4

Type material (20 ♂♂, 25 ♀♀). **Holotype:** CHINA: ♂, labeled 'China: Hainan, Ledong Hsien [乐东县], Jianfengling [尖峰岭] N.R., Mingfenggu [鸣凤谷], 18°44'30"N, 108°50'29"E, rainforest, decaying log from a colony termite nest, 995 m, 23.i.2015, Peng, Yin, Tu, Song, Shen, Zhou, Yan, Wang leg.'. **Paratypes:** 7 ♂♂, 14 ♀♀, same locality as holotype; 1 ♀, same locality as holotype, except 'alt. 950m, 30-IV-2012, PAN Y.H. & LI W. R. leg.'; 6 ♂♂, 7 ♀♀, labeled 'China: Hainan, Qiongzong Hsien [琼州县], Limu Mt [黎母山], nr. residence, path to peak, 19°10'04"N, 109°44'45"E, decaying log, 1000 m, 31.i.2015, Peng, Yin, Tu, Song, Shen, Zhou, Yan, Wang leg.'; 2 ♂♂, 1 ♀, same locality as previously except '01.ii.2015'; 1 ♂, 1 ♀, same locality as previously except 'nr. residence, 19°10'04"N, 109°44'45"E, decaying log, 625m, 29.i.2015'; 1 ♂, same locality as previously except 'path to Limu Temple, 19°08'09"N, 109°45'46"E, 580–760m, 29.i.2015, Peng, Yin, Tu, Song, Shen leg.'; 2 ♂♂, 1 ♀, labeled 'China: Hainan, Wuzhi Shan [五指山] N. R., nr.reservoir, 18°53'10"N, 109°36'11"E, 500m, 22.iv.2012, leaf litter, sifted, Peng& Dai leg.'.

Diagnosis. *Horaeomorphus hainanicus* can be readily separated from all other congeners by its moderately large (2.53–3.08 mm) and elongate body, fine punctuation on pronotal disc and elytra, rounded apices of metatrochanters in male, parameres each with 3–5 apical setae and two subapical setae, unique structure of endophallus and shape of spermatheca.

Description. Male. BL 2.77–3.08 mm; body (Fig. 2A) large, strongly convex, reddish brown, legs and palpi slightly lighter. Head broadest at large, finely faceted and moderately convex eyes, HL 0.40–0.48 mm, HW 0.58–0.68 mm; tempora rounded and slightly shorter than eye in dorsal view; vertex strongly transverse and weakly con-



Figure 2. Habitus of *Horaemorphus hainanicus* sp. n. **A** male **B** female. Scale bars: 1.0 mm.

vex, with pair of small but distinct pits located near posterior margins of supra-antennal tubercles; frons weakly convex; supra-antennal tubercles strongly raised. Punctuation on vertex and frons fine and inconspicuous; setae moderately long, sparse. Antennae (Fig. 4A) short, AnL 1.21–1.24 mm, relative lengths of antennomeres: 1.0 : 1.1 : 1.7

: 1.5 : 1.4 : 1.5 : 1.2 : 1.1 : 1.2 : 1.1 : 2.2. Pronotum inversely subtrapezoidal, widest near anterior third, PL 0.87–0.94 mm, PWm 0.72–0.81 mm, PWb 0.52–0.58 mm; anterior margin rounded, sides narrowing toward base; hind angles obtuse and blunt; posterior collar delimited from disc by deep and narrow transverse groove connecting three small pits. Punctuation on disc as fine as that on frons and vertex; dorsal surface glossy; setation moderately long. Elytra oval and elongate, moderately convex; widest near anterior third, narrowing toward apices. EL 1.50–1.66 mm, EW 0.92–1.04 mm, EI 1.59–1.63. Humeral calli distinct. Punctures more distinct than those on pronotum, sharply marked and separated by spaces 3–4× as wide as puncture diameters; setation moderately dense. Hindwings fully developed. Metatrochanter (Fig. 4I) modified, with distal part protruded and recurved; as long as half of metafemur, apex rounded. Aedeagus (Fig. 4B–D) elongate, AeL 0.53 mm; endophallus (Fig. 4F–G) very complicated, with large bell-shaped central complex surrounded at each side by weakly sclerotized, elongate lateral structures; parameres (Fig. 4E) slender, longer than median lobe, each with 3–5 apical setae and two subapical setae.

Female. Similar to male, with slightly smaller body and unmodified metatrochanter. BL 2.53–2.85 mm, HL 0.39–0.42 mm, HW 0.57–0.62 mm, AnL 1.11–1.20 mm, relative lengths of antennomeres: 1.0 : 1.1 : 1.5 : 1.4 : 1.4 : 1.3 : 1.1 : 1.1 : 1.2 : 1.2 : 2.2. PL 0.76–0.88 mm, PWm 0.63–0.74 mm, PWb 0.49–0.54 mm; EL 1.39–1.54 mm, EW 0.84–0.97 mm, EI 1.58–1.64, spermatheca (Fig. 4H) ovoid, slightly elongate, SpL 0.12 mm.

Comments. *Horaeomorphus hainanicus* is similar to many congeners (see comments at *H. chinensis*), and has a similar endophallic structures and the shape of spermatheca as *H. chinensis*. The relatively larger body (2.77–3.08 mm in males), the pronotal and elytral surface glossy, protruded metatrochanters with rounded apices and subtle differences in the endophallus can be used to readily separate this species from *H. chinensis*.

Bionomics. Specimens of the type series were collected from ant and termite nest material in rotten wood and under bark of standing rotten logs in rainforests of Hainan.

Distribution. Southern China: Hainan.

Etymology. The specific epithet refers to the province where the type locality of the new species lies.

***Horaeomorphus punctatus* D.-Y. Zhou & S.-J. Zhang, sp. n.**

<http://zoobank.org/B832FF57-E3B1-44C3-9D4F-87C3471AA98B>

Figs 3, 5

Type material (1 ♂, 1 ♀). **Holotype:** CHINA: ♂, labeled ‘China: Yunnan Prov., Yingjiang Hsien [盈江县], Xima [昔马], 1500–1650m, 20–22.v.2013, Wen-Xuan Bi leg.’. **Paratype:** 1 ♀, same locality as holotype.

Diagnosis. *Horaeomorphus punctatus* can be readily separated from all other congeners by its moderately large (2.07–2.36 mm) and stout body, broad pronotum with



Figure 3. Habitus of *Horaeomorphus punctatus* sp. n. **A** male **B** female. Scale bars: 0.5 mm.

a row of three pits connected by a shallow, barely notable groove; dense and coarse punctation on elytra, unmodified metatrochanters in male, aedeagal parameres lacking setae at apices and shape of spermatheca.

Description. Male. BL 2.36 mm; body (Fig. 3A) moderately large, flattened, dark reddish-brown, legs and palpi slightly lighter. Head broadest at large, finely faceted and moderately convex eyes, HL 0.34 mm, HW 0.46 mm; tempora shorter than eye in dorsal view, in anterior third nearly parallel, then strongly bent and in posterior third nearly transverse to long axis of head; vertex strongly transverse and weakly convex,




Figure 4. Diagnostic characters of *Horaeomorphus hainanicus* sp. n. **A** Left antenna of male, in dorsal view **B** Aedeagus, in ventral view **C** Same, in lateral view **D** Same, in dorsal view **E** Apical portion of paramere, enlarged **F** Endophallus, enlarged, in ventral view **G** Same, schematic **H** Spermatheca, in lateral view **I** Left metatrochanter of male, in ventral view. Scale bars: 0.5 mm (**A**); 0.2 mm (**B**, **C**, **D**); 0.04 mm (**E**); 0.1 mm (**F**, **G**, **H**); 0.3 mm (**I**).




Figure 5. Diagnostic characters of *Horaemorphus punctatus* sp. n. **A** Left antenna of male, in dorsal view **B** Aedeagus, in ventral view **C** Same, in lateral view **D** Same, in dorsal view **E** Apical portion of paramere, enlarged **F** Endophallus, enlarged, in ventral view **G** Same, schematic **H** Spermatheca, in lateral view **I** Left metatrochanter of male, in ventral view. Scale bars: 0.3 mm (**A**); 0.2 mm (**B**, **C**, **D**); 0.03 mm (**E**); 0.08 mm (**F**, **G**); 0.04 mm (**H**); 0.3 mm (**I**).




Figure 6. Diagnostic characters of *Horaeomorphus* species. (**A, F** *Horaeomorphus* sp. **B–E, G–K** *H. chinensis*) **A** Left antenna of male, in dorsal view **B** Left antenna of female, in dorsal view **C** Aedeagus, in ventral view **D** Same, in lateral view **E** Same, in dorsal view **F** Spermatheca, in lateral view **G** Apical portion of paramere, enlarged **H** Endophallus, enlarged, in ventral view **I** Same, schematic **J** Left metatrochanter of male, in ventral view **K** Spermatheca with bursa copulatrix, in lateral view. Scale bars: 0.5 mm (**A, B**); 0.2 mm (**C, D, E**); 0.1 mm (**F, H, I**); 0.04 mm (**G**); 0.3 mm (**J, K**).

with pair of small but distinct pits located near posterior margins of supra-antennal tubercles; frons weakly convex; supra-antennal tubercles strongly raised. Punctures on vertex and frons fine and inconspicuous; setation moderately long, sparse. Antennae (Fig. 5A) short, AnL 0.87 mm, relative lengths of antennomeres: 1.0 : 0.9 : 1.0 : 0.9 : 0.9 : 0.8 : 0.8 : 0.8 : 0.9 : 1.0 : 1.5. Pronotum broad, nearly circular, slightly longer than wide, widest at middle, PL 0.76 mm, PWm 0.72 mm, PWb 0.47 mm; posterior collar short, well delimited from disc by constriction and row of three shallow pits connected by shallow groove; punctation dense and coarse; setation moderately long. Elytra broad, flattened and distinctly impressed in middle at about anterior third; EL 1.26 mm, EW 0.84 mm, EI 1.50. Punctures coarse, more distinct than those on pronotum, sharply marked and separated by spaces as wide as diameters of punctures; setation long, moderately dense, erect to suberect. Hindwings fully developed. Metatrochanter (Fig. 5I) unmodified, with distinct ventral edge. Aedeagus (Fig. 5B–D) relatively slender, with median lobe strongly narrowing toward apex, AeL 0.57 mm; endophallus (Fig. 5F–G) relatively small and complicated, with sclerotized central portion and two moderately darkly sclerotized and curved structures; parameres (Fig. 5E) slender, minimally longer than median lobe, without apical setae.

Female. Similar to male, but with smaller body and less contractive pronotal base. BL 2.07 mm, HL 0.33 mm, HW 0.44 mm, AnL 0.75 mm, relative lengths of antennomeres: 0.8 : 0.8 : 0.9 : 0.8 : 0.7 : 0.7 : 0.6 : 0.7 : 0.6 : 1.0 : 1.2. PL 0.63 mm, PWm 0.58 mm, PWb 0.44 mm; EL 1.11 mm, EW 0.74 mm, EI 1.50, spermatheca (Fig. 5H) spherical, SpL 0.08 mm, with complicated internal structures.

Comments. *Horaeomorphus punctatus* is similar to *H. mesaios* Jałoszyński & Nomura, 2004 (Vietnam: Ninh Binh), *H. valdepunctatus* Franz, 1984 (Malaysia: Pahang), *H. sarawakensis* Franz, 1992 (W Malaysia) and *H. samosirensis* Jałoszyński, 2009 (Indonesia: Sumatra) in habitus, and is also very similar to *H. mesaios* in the shape of aedeagus and structures of endophallus; but it can be identified by its relatively large body (2.07–2.36 mm), the pronotum with three basal pits; unmodified metafemora, non-protruded metatrochanters in males and achaetous parameral apex.

Distribution. Southern China: Yunnan.

Etymology. The specific epithet refers to the coarse punctation of the elytra.

***Horaeomorphus pengzhongi* D.-Y. Zhou & S.-J. Zhang, sp. n.**

<http://zoobank.org/F7F96BFA-C563-45EF-AFCF-5F6539869D30>

Fig. 7

Type material (1 ♂). **Holotype:** CHINA: ♂, labeled ‘China: Yunnan, Baoshan City [保山市], Tengchong County [腾冲县], Mingguang Town [明光镇], Zizhi Village [自治乡], 25°18'24"N, 98°48'22"E, 1230 m, 24.vi.2013, Dai, Song& Peng leg.’.

Diagnosis. *Horaeomorphus pengzhongi* can be readily separated from all other congeners by its moderately large (3.00 mm) and elongate body, sparse and fine punctation on pronotal disc and elytra, protruded metatrochanter with expansion on its




Figure 7. *Horaeomorphus pengzhongi* sp. n. **A** male, dorsal habitus **B** Right antenna of male, in dorsal view **C** Aedeagus, in ventral view **D** Same, in lateral view **E** Same, in dorsal view **F** Apical portion of paramere, enlarged **G** Endophallus, enlarged, in ventral view **H** Same, schematic **I** Left metatrochanter of male, in ventral view. Scale bars: 1 mm (**A**); 0.5 mm (**B**); 0.2 mm (**C, D, E**); 0.04 mm (**F**); 0.1 mm (**G, H**); 0.3 mm (**I**).

ventral edge in male, parameres each with four apical setae and two subapical setae and unique structures of endophallus.

Description. Male. BL 3.00 mm; body (Fig. 7A) large, strongly convex, reddish brown, legs and palpi slightly lighter. Head broadest at finely faceted and slightly convex eyes, HL 0.47 mm, HW 0.63 mm; tempora rounded, about as long as length of eye in dorsal view; vertex strongly transverse and weakly convex, with pair of small but distinct pits located near posterior margins of supra-antennal tubercles; frons weakly convex; supra-antennal tubercles strongly raised. Punctures on vertex and frons dense and coarse; setae moderately long, sparse. Antennae (Fig. 7B) short, AnL 1.31 mm, relative lengths of antennomeres: 1.0 : 0.8 : 1.8 : 1.4 : 1.1 : 1.1 : 1.0 : 1.1 : 1.2 : 1.1 : 2.2. Pronotum oval, convex, distinctly longer than wide, widest near anterior 2/5, PL 0.97 mm, PWm 0.79 mm, PWb 0.53 mm; anterior margin rounded, sides narrowing

toward base; hind angles obtuse and blunt; base with 3 large and deep pits connected by narrow groove. Punctuation on disc sparse and fine; dorsal surface glossy; setation moderately long. Elytra oval and elongate, moderately convex; widest near anterior 2/5, narrowing toward apices. EL 1.57 mm, EW 1.0 mm, EI 1.56. Humeral calli distinct. Punctures fine, more distinct than those on pronotum, separated by spaces 3–4× as wide as puncture diameters; setation moderately dense. Hindwings fully developed. Metatrochanter (Fig. 7I) modified, with distal portion straight and apical 1/7 recurved, as long as half of metafemur, with expansion on ventral margin, apex rounded. Aedeagus (Fig. 7C–E) elongate, AeL 0.53 mm; endophallus (Fig. 7G–H) very complicated, with large bell-shaped central complex surrounded at each side by weakly sclerotized, elongate lateral structures; parameres (Fig. 7F) slender, subequal in length to median lobe, each with four apical setae and two subapical setae.

Female. Unknown.

Comments. *Horaeomorphus pengzhongi* is similar to many congeners (see comments of *H. chinensis*); the relatively larger body (3.00 mm), elongated metatrochanters each with the ventral expansion in male, each paramere with two subapical setae and structures of endophallus are clearly different and can be used to identify this new species.

Distribution. Southern China: Yunnan.

Etymology. This species is dedicated to Zhong Peng, one of the collectors of the type specimen.

***Horaeomorphus biwenxuani* D.-Y. Zhou & S.-J. Zhang, sp. n.**

<http://zoobank.org/E5D8A881-C40E-4321-BAAE-415074D29AEF>

Fig. 8

Type material (1 ♂). **Holotype: CHINA:** ♂, labeled ‘China: Xizang Prov., Cuona County [措那县], Lexiang [勒乡], alt. 2500m, 15.vii.2012, Wen-Xuan Bi leg.’.

Diagnosis. *Horaeomorphus biwenxuani* can be readily separated from all other congeners by its moderately large (2.78 mm) and elongate body, small pronotum lacking basal groove, with a row of three dorsal pits, subtriangular metatrochanter with distal edge produced into a short acute spine, and slender aedeagus with a complicated and strongly asymmetrical endophallus.

Description. Male. BL 2.78 mm; body (Fig. 8A) large, slightly convex, dark reddish-brown, legs and palpi slightly lighter. Head broadest at large, finely faceted and moderately convex eyes, HL 0.41 mm, HW 0.58 mm; tempora rounded, about as long as length of eye in dorsal view; vertex strongly transverse and weakly convex, with pair of small but distinct pits located near posterior margins of supra-antennal tubercles; frons weakly convex; supra-antennal tubercles strongly raised. Punctuation on vertex and frons sparse, small but distinct; setae moderately long, sparse. Antennae (Fig. 8B) short, AnL 1.18 mm, relative lengths of antennomeres: 0.9 : 1.0 : 1.6 : 1.3 : 1.1 : 1.0 : 0.8 : 0.9 : 1.0 : 1.1 : 1.7. Pronotum short, longer than wide, widest slightly behind anterior fourth, PL 0.77 mm, PWm 0.65 mm, PWb 0.53 mm; pronotal margin rounded near anterior 2/3, then nearly




Figure 8. *Horaeomorphus biwenxuani* sp. n. **A** male, dorsal habitus **B** Left antenna of male, in dorsal view **C** Aedeagus, in ventral view **D** Same, in lateral view **E** Same, in dorsal view **F** Apical portion of paramere, enlarged **G** Endophallus, enlarged, in ventral view **H** Same, schematic **I** Left metatrochanter of male, in ventral view. Scale bars: 1 mm (**A**); 0.5 mm (**B**); 0.2 mm (**C, D, E**); 0.04 mm (**F**); 0.1 mm (**G, H**); 0.3 mm (**I**).

straight up to sub-basal constriction; base with row of three dorsal pits and pair of lateral impressions located in constriction. Punctuation on disc sparse and fine; dorsal surface glossy; setation moderately long. Elytra elongate, more convex than pronotum, distinctly impressed in middle at about anterior third; widest near anterior $2/5$, narrowing toward apices. EL 1.59 mm, EW 0.99 mm, EI 0.59. Humeral calli distinct. Punctures fine, more distinct than those on pronotum, sharply marked and separated by spaces 3–4× as wide as puncture diameters; setation moderately dense. Hindwings fully developed. Metatrochanter (Fig. 8I) short, subtriangular, distal edge produced into short acute spine. Aedeagus (Fig. 8C–E) slender, AeL 0.55 mm; endophallus (Fig. 8G–H) strongly asymmetrical, with curved axial component protruding from posterior complicated structure surrounded by two lateral Λ -shaped structures; parameres (Fig. 8F) very slender with broadened apical parts, slightly shorter than median lobe, each with nine apical and subapical setae.

Female. Unknown.

Comments. This new species with remarkably long legs has subtriangular metatrochanters in males, each with a sharp distal edge. This unique character can be found also in all four known Nepalese congeners: *H. obrus* Vít, 2004, *H. deharvengi* Vít, 2004, *H. himalayensis* Franz, 1974 and *H. nepalensis* Franz, 1973 (Franz 1974; Vít 2004), but so far has not been recorded in *Horaeomorphus* outside the Himalayas. However, a relatively small body (2.78 mm; among Himalayan species only *H. deharvengi* can be smaller than 3mm) and strongly asymmetrical endophallus are clearly different from characters of the Nepalese species. An asymmetrical endophallus also occurs in *H. deformatus* Jałoszyński, 2006 (W Malaysia: Kuala Terengganu), *H. pseudosabahensis* Jałoszyński, 2006 (E Malaysia: Sabah, Sarawak) and *H. minor* Jałoszyński, 2009 (the Philippines: Bukidnon, Mindanao), but its structure is distinctly different from that in *H. biwenxuani*.

Distribution. Western China: Xizang.

Etymology. This species is dedicated to Wen-Xuan Bi, who collected the type specimen.

***Horaeomorphus bujiayaoi* D.-Y. Zhou & S.-J. Zhang, sp. n.**

<http://zoobank.org/85FD7AC1-06A4-41EA-8F3D-B8A2E2FB221C>

Fig. 9

Type material (1 ♂). **Holotype:** CHINA: ♂, labeled 'China: Guangxi Prov., Jinxiu County [金秀县], Mt.Lianhuashan [莲花山], alt. 1000–1150m, 30.vii.2011, Jia-Yao Hu leg.'

Diagnosis. *Horaeomorphus bujiayaoi* can be readily separated from all other congeners by its moderately large (2.53 mm), short pronotum with five basal pits connected by a shallow groove, unmodified metatrochanters, median lobe of aedeagus with a blade-shaped, asymmetrical apex bent at an obtuse angle in relation to the long axis of aedeagus and asymmetrical parameres each with ten apical and subapical setae.

Description. Male. BL 2.53 mm; body (Fig. 9A) large, flattened, dark reddish-brown, legs and palpi slightly lighter. Head broadest at large, finely faceted, and moderately convex eyes, HL 0.36 mm, HW 0.50 mm; tempora rounded but not bent, about as long as length of eye in dorsal view; vertex strongly transverse and weakly convex, with pair of small but distinct pits located near posterior margins of supra-antennal tubercles; frons weakly convex; supra-antennal tubercles strongly raised. Punctuation on vertex and frons sparse, small but distinct; setae moderately long, sparse. Antennae (Fig. 9B) short, AnL 1.01 mm, relative lengths of antennomeres: 1.2 : 1.0 : 1.3 : 1.0 : 1.0 : 1.0 : 0.9 : 0.9 : 1.0 : 1.0 : 1.9. Pronotum inversely subtrapezoidal, flattened, longer than wide, widest near middle, PL 0.76 mm, PWm 0.61 mm, PWb 0.53 mm; anterior margin rounded; lateral margins rounded near anterior 2/3, then nearly straight up to sub-basal constriction; base with five pits connected by shallow groove. Punctuation on disc sparse and fine; dorsal surface glossy; setation moderately




Figure 9. *Horaeomorphus hujiayaoi* sp. n. **A** male, dorsal habitus **B** Left antenna of male, in dorsal view **C** Aedeagus, in ventral view **D** Same, in lateral view **E** Same, in dorsal view **F** Apical portion of paramere, enlarged **G** Endophallus, enlarged, in ventral view **H** Same, schematic **I** Left metatrochanter of male, in ventral view. Scale bars: 1 mm (**A**); 0.5 mm (**B**); 0.2 mm (**C, D, E**); 0.04 mm (**F**); 0.1 mm (**G, H**); 0.3 mm (**I**).

long. Elytra elongate, more convex than pronotum, distinctly impressed in middle at about anterior third; widest slightly before middle, narrowing toward apices. EL 1.41 mm, EW 0.93 mm, EI 1.51. Humeral calli distinct. Punctures fine, more distinct than those on pronotum, especially on impressed area, sharply marked and separated by spaces 2–4× as wide as puncture diameters; setation moderately dense. Hindwings fully developed. Metatrochanter (Fig. 9I) unmodified. Mesotibiae slightly curved and with inner margin expanded near middle to form broad subtriangular tooth. Aedeagus (Fig. 9C–E) moderately elongate, median lobe with a strongly asymmetrical blade-shaped apex bent at obtuse angle in relation to long axis of aedeagus, AeL 0.59 mm; endophallus (Fig. 9G–H) with complicated system of variously sclerotized




Figure 10. A Distribution of *Horaeomorphus* in mainland China **B** Habitat of *H. chinensis* at Guadun.

structures; parameres (Fig. 9F) asymmetrical, of unequal lengths, each with ten apical and subapical setae.

Female. Unknown.

Comments. This new species has the aedeagal median lobe with an apical blade-shaped projection bent to the left in ventral view and parameres of unequal lengths, characters shared with *H. obrus* Vít, 2004 (Nepal: Janakpur, Bagmati). *Horaeomorphus hujiayaoi* can be unambiguously separated from its Nepalese congener by the much smaller body (2.53 mm vs 4.3 mm in *H. obrus*) and different structures of the endophallus.

Distribution. Southern China: Guangxi.

Etymology. This species is dedicated to Jia-Yao Hu, who collected the type specimen.

Key to species of *Horaeomorphus* in mainland China

- 1 Basal pits on pronotum connected by a shallow and barely notable groove ...2
- Basal pits on pronotum connected by a deep and distinct groove..... 3

- 2 Pronotum broad, nearly circular; elytral punctures coarse, sharply marked and separated by spaces as wide as puncture diameters..... *H. punctatus* sp. n. (Yunnan: Yingjiang)
- Pronotum small; elytral punctation fine and sparse, elytra glossy *H. biwenxuani* sp. n. (Xizang: Cuona)
- 3 Pronotum with five basal pits..... 4
- Pronotum with three basal pits 5
- 4 Body slender and flattened, EI > 1.5 *H. bujiayaoi* sp. n. (Guangxi: Jinxiu)
- Body stout and convex, EI < 1.4 *H. sp.* (Guangxi: Shangsi)
- 5 BL < 2.5 mm; punctation of pronotal disc and elytra dense and coarse..... *H. chinensis* Franz, 1985 (N.Fujian, S.Fujian)
- BL > 2.5 mm; punctation of pronotal disc and elytra sparse and fine, surface glossy 6
- 6 Pronotum inversely subtrapezoidal, its lateral margins sharply bent at anterior third and strongly constricted near posterior fourth *H. hainanicus* sp.n. (Hainan)
- Pronotum oval, its anterior margin and lateral margins evenly rounded together *H. pengzhong* sp.n. (Yunnan: Tengchong)

Acknowledgments

We thank Paweł Jałoszyński (Wrocław, Poland) and two anonymous reviewers for critically reading the manuscript. We would like to thank Wen-Xuan Bi (Shanghai, China) for his generous gift of specimens. The present study was supported by the National Science Foundation of China (No. 31501874).

References

- Franz H (1985) Neue und ungenügend bekannte Scydmaeniden (Coleoptera) aus Taiwan, Fukien und Thailand. Mitteilungen der Münchner Entomologische Gesellschaft 74: 91–128.
- Jałoszyński P (2002) First record of the Genus *Horaeomorphus* Schaufuss (Coleoptera, Scydmaenidae) from Japan, with Description of a New Species. Bulletin of National Science Museum Tokyo (Series A) 28: 223–232.
- Jałoszyński P (2003) Taxonomical Notes on Southeast Asiatic Species of *Horaeomorphus* Schaufuss (Coleoptera, Scydmaenidae), with Description of a New Species from Taiwan. Bulletin of National Science Museum Tokyo (Series A) 29(2): 107–117.
- Jałoszyński P (2004) The first record of *Horaeomorphus* Schaufuss (Coleoptera, Scydmaenidae) from the Philippines, with description of *H. blattnyi* sp. nov. Revue suisse de Zoologie 111(4): 785–789. doi: 10.5962/bhl.part.80269
- Jałoszyński P (2006) Revision of *Horaeomorphus* Schaufuss of East Malaysia, Singapore, and Sunda Islands (Coleoptera, Scydmaenidae). Genus 17(1): 19–66.

- Jałoszyński P (2009) Four new species of *Horaeomorphus* Schaufuss from Oriental Region (Coleoptera: Scydmaenidae). Raffles Bulletin of Zoology 57(2): 297–303.
- Jałoszyński P (2012) Taxonomic status of *Allohoraeomorphus* Franz (Coleoptera, Staphylinidae, Scydmaeninae). Zootaxa 3313: 62–68.
- Jałoszyński P (2014a) Oriental *Horaeomorphus* Schaufuss: a new species, synonymic notes and distributional synopsis (Coleoptera: Staphylinidae: Scydmaeninae). Raffles Bulletin of Zoology 62: 83–88.
- Jałoszyński P (2015) *Schuelkelia* gen. n., a new eastern Palaearctic ant-like stone beetle, with synopsis of Eurasian genera of Cyrtoscydmini (Coleoptera: Staphylinidae: Scydmaeninae). Zootaxa 4007(3): 343–369. doi: 10.11646/zootaxa.4007.3.3
- Jałoszyński P (2014c) ‘The curse of *Horaeomorphus*’: taxonomy of misplaced Australian Cyrtoscydmini (Coleoptera: Staphylinidae: Scydmaeninae). Zootaxa 3828(1): 1–76. doi: 10.11646/zootaxa.3828.1.1
- Jałoszyński P, Nomura S (2004) A new species of *Horaeomorphus* Schaufuss (Coleoptera, Scydmaenidae) from Vietnam. Elytra 32(1): 65–70.
- Jałoszyński P, Nomura S (2008) Two new species of *Horaeomorphus* Schaufuss (Coleoptera, Scydmaenidae) from Sabah, Borneo. Elytra 36(1): 159–166.
- Jałoszyński P, Nomura S, Idris AG (2007) A new *Horaeomorphus* from the Malay Peninsula. Elytra 35(1): 307–311.
- Newton AF (2015) Cyrtoscydmini Schaufuss, 1889 Replaced by Glandulariini Schaufuss, 1889 (Coleoptera: Staphylinidae: Scydmaeninae). The Coleopterists Bulletin, 69(4): 758–759.
- Vít S (2004) Notes on *Horaeomorphus* occurring in Nepal (Coleoptera, Scydmaenidae). Acta Societas Zoologicae Bohemicae 68: 287–299.

Erratum

Zhou DY, Li LZ (2015) Discovery of the genus *Loeblites* Franz (Coleoptera: Staphylinidae: Scydmaeninae) in China, with description of a new species. Zootaxa 3986(3): 393–396. doi: 10.11646/zootaxa.3986.3.8

The generic name *Loeblites* occurs in the text in a miswriting as *Syndicus* (Page 394, figure caption, line 1).