

Two new species of cricket genus *Anaxiphomorpha* Gorochov, 1987 (Orthoptera, Trigonidiidae, Trigonidiinae) in China

Zhi Xin He¹, Li Bin Ma¹

¹ Shaanxi Normal University, 710119 Xi'an, China

Corresponding author: Li Bin Ma (libinma@snnu.edu.cn)

Academic editor: T. Robillard | Received 22 September 2021 | Accepted 3 November 2021 | Published 29 November 2021

<http://zoobank.org/6B382A8F-0F1B-439B-99D5-200D43E94075>

Citation: He ZX, Ma LB (2021) Two new species of cricket genus *Anaxiphomorpha* Gorochov, 1987 (Orthoptera, Trigonidiidae, Trigonidiinae) in China. ZooKeys 1073: 13–20. <https://doi.org/10.3897/zookeys.1073.75598>

Abstract

Two new species, *Anaxiphomorpha nonggangensis* **sp. nov.** and *Anaxiphomorpha manereserratus* **sp. nov.**, are reported from Guangxi Province, China. Descriptions and illustrations for the new species and a key to all known species of *Anaxiphomorpha* are provided.

Keywords

Southern China, swordtail crickets, taxonomy, Trigonidiini, tropics

Introduction

The genus *Anaxiphomorpha* was established with *Anaxiphomorpha brachyapodemalis* Gorochov, 1987 as the type species. *Anaxiphomorpha* is recognized by a smaller body size, yellow coloration and special genitalia structure (epiphallus and ectoparamere possess multiple lobes or branches).

To date, seven species have been reported worldwide (Cigliano et al. 2021). Apart from two species, *Anaxiphomorpha brachyapodemalis* Gorochov, 1987 and *Anaxiphomorpha longiapodemalis* Gorochov, 1987 reported from Vietnam, the other five, *Anaxiphomorpha biserratus* Liu & Shi, 2015, *Anaxiphomorpha brevisparamerus* Liu & Shi, 2015, *Anaxiphomorpha longiserratus* Liu & Shi, 2015, *Anaxiphomorpha serratiprotuberus* Liu & Shi, 2015 and *Anaxiphomorpha hexagona* Ma, 2018 have been

recorded from China. After comparing our new materials with the species of this genus, we concluded that two Chinese species are new for science. The distribution of *Anaxiphomorpha* species worldwide including the new species is also presented (Fig. 1).

Materials and methods

All specimens were collected at night with a sweep net. Specimens were preserved in ethanol during field work, and pinned and dry-preserved in the laboratory. Male genitalia were dissected from softened specimens. Photomicrographs of genitalia were collected using Toup-Cam Digital camera and bundled software (ToupTek, Hangzhou, China). Photographs of specimens were obtained using a VHX–6000 digital microscope (Keyence, Osaka, Japan).

Measurements

All specimens were measured using a ToupCam Digital camera and bundled software (ToupTek, Hangzhou, China). All the measurements are in millimeters (mm).

Abbreviations

- BL

body length (from head to apical hindwing);
- PL

pronotal length;
- TL

tegmen length;
- HFL

hind femur length;
- OL

ovipositor length.

The specimens are deposited at the Museum of Flora and Fauna of Shaanxi Normal University, Xi'an, China (SNNU).

Taxonomy

Genus *Anaxiphomorpha* Gorochov, 1987

Type species. *Anaxiphomorpha brachyapodemalis* Gorochov, 1987

Key to known species of *Anaxiphomorpha*

- 1

Dorsally viewed, epiphalllic lateral lobes apically acute.....

2
- Dorsally viewed, epiphalllic lateral lobes somewhat blunt of apex

3
- 2

Epiphalllic transverse bridge and ectoparamere short ... *A. brachyapodemalis*
- Epiphalllic transverse bridge and ectoparamere long *A. longiapodemalis*
- 3

Epiphalllic lateral lobes bifurcated as six significant branches *A. hexagona*
- Not as above and laterally viewed as following

4
- 4

Apex of epiphalllic lateral lobes almost straight *A. brevisparamerus*
- Apex of epiphalllic lateral lobes curved.....

5

- 5 Apex of epiphallic lateral lobes upward curved 6
– Apex of epiphallic lateral lobes downward curved 8
6 Epiphallic lateral lobes boot-like *A. nonggangensis* sp. nov.
– Epiphallic lateral lobes rod-like 7
7 Epiphallic lateral lobes proximally bearing a protuberance
..... *A. manereserratus* sp. nov.
– Not as above *A. longiserratus*
8 Epiphallic lateral lobes long, medially raised and apically acute
..... *A. serratiprotuberus*
– Epiphallic lateral lobes short and apically blunt..... *A. biserratus*

***Anaxiphomorpha nonggangensis* He & Ma, sp. nov.**

<http://zoobank.org/960B75C1-468B-4D24-BADE-06F8D7F93D2E>

Figs 1, 2, 3A, B, 4A–C

Material examined. Holotype. CHINA: Male, Guangxi, Longzhou, Nonggang National Natural Reserve, 5.V.2019, 22.46°N, 106.96°E, Libin Ma & Tao Zhang leg.
Paratypes. 7 males and 4 females, same information as holotype (SNNU).




Figure 1. Known distribution of *Anaxiphomorpha* species, including the new species *A. nonggangensis* sp. nov. and *A. manereserratus* sp. nov..



Figure 2. Habitus (alive) of *A. nonggangensis* sp. nov. on leaf **A** male **B** female (photography: Zhang, Tao).

Diagnosis. Male (Figs 2A, 3A). Body size small. Head small and pubescent, slightly wider than anterior margin of pronotum; frontal rostrum as wide as antennal scape; eyes large and protruding forwards; apical three joints of maxillary palpi distinctly elongate, 5th joint truncated apically. Pronotum transverse, strongly widened posteriorly and distinctly longer than width of anterior one. Tegmina extending slightly over apex of abdomen, armed with one oblique vein; mirror slightly elongate; hindwings absent. Fore tibia armed with a large long-oval external tympanum, and without internal tympanum. Hind tibia bearing three dorsal spurs on two sides respectively, and bearing two apical spurs inside and three outside.

Genitalia (Fig. 4A–C). Lateral lobes of epiphallus slightly longer than median lobes, and possessing several teeth at outer margin, not narrowed apically in dorsal view, gradually narrowed apically in lateral view; median lobes shaped as boot and abruptly narrowed apically in lateral view. Etcoparameres transversely and truncated apically.




Figure 3. Habitus photographs. **A, B** *A. nonggangensis* sp. nov. **C, D** *A. manereserratus* sp. nov. **A, C** Male **B, D** Female. Scale bar: 1 mm.

Female (Figs 2B, 3B). Body slightly smaller than male. Tegmen rather convex, armed with five regular veins on dorsal field. Ovipositor blade-shaped.

Coloration. Body colored yellow. Dorsal area of head ornamented with four brown longitudinal stripes in ventral view, anterior half of abdomen colored dark brown in male or small part of middle abdomen colored dark in female. Apex of each hind femur bearing a small dark spot on two sides respectively. Ovipositor ventrally colored brown to dark brown and remainder yellowish.

Measurements. Male. BL 5.17–6.42, PL 0.98–1.19, TL 3.73–4.54, HFL 3.27–3.96. **Female.** BL 4.50–5.63, PL 0.99–1.17, TL 3.87–4.42, OL 2.22–2.67.

Etymology. The name refers its type locality, Nonggang National Natural Reserve.

Distribution. China (Guangxi) (Fig. 1).

Remarks. This species is very similar to *A. brevisparamerus* in the ectoparameres of the male genitalia, but different in the epiphallic lateral lobes of this new species, which are shaped as boot with an acute apex.

***Anaxiphomorpha manereserratus* He & Ma, sp. nov.**

<http://zoobank.org/9B5380BF-9334-4480-B796-4FE6DF673DB1>

Figs 1, 3C–D, 4D–F, 5

Material examined. Holotype. CHINA: Male, Guangxi, Jingxi, Longbang, 22.87°N, 106.32°E, 1.V.2019, Libin Ma & Tao Zhang leg. **Paratypes.** 10 males and 2 females, same information as holotype (SNNU).

Description. Male (Figs 3C, 5A). Body size small. Head small and pubescent, slightly wider than anterior margin of pronotum; frontal rostrum 1.2 times wider than




Figure 4. Male genitalia **A–C** *A. nonggangensis* sp. nov. **D–F** *A. manereserratus* sp. nov. **A, D** dorsal views **B, E** lateral views **C, F** ventral views.

antennal scape; eyes large and protruding forwards; apical three joints of maxillary palpi distinctly elongate, 5th joint truncated apically. Pronotum transverse, strongly widened posteriorly, and slightly wider than the anterior. Tegmina extending slightly over apex of abdomen, armed with one oblique vein; mirror slightly elongate; hindwings absent. Fore tibia armed with a large long-oval external tympanum; internal tympanum absent. Hind tibia bearing three dorsal spurs on two sides respectively, and bearing two apical spurs inside and three outside.

Genitalia (Fig. 4D–F). Lateral lobes of epiphallus slightly longer than median lobes in lateral view and possessing horned protuberances at inner margin proximally and outer margin medially, bearing several teeth at inner and outer margin and not narrowed apically in lateral view. Ectoparameres short and serrated apically.



Figure 5. Habitus (alive) of *A. manereserratus* sp. nov. on leaf **A** male **B** female (photography: Zhang, Tao).

Female (Figs 3D, 5B). Body slightly smaller than male. Tegmen rather convex, armed with five regular veins on dorsal field. Ovipositor blade-shaped.

Coloration. Body colored yellow. Dorsal area of head ornamented with four brown longitudinal stripes. Apex of each hind femur bearing a small dark spot in two sides respectively. Ovipositor ventrally colored brown to dark brown and remainder yellowish.

Measurements. Male. BL 4.92–6.05, PL 0.94–1.15, TL 3.42–4.09, HFL 3.56–4.17. **Female.** BL 4.65–4.94, PL 0.87–0.90, TL 3.21–3.33, OL 1.892.03.

Etymology. The name refers to the epiphallic median lobes almost as long as the lateral lobes.

Distribution. China (Guangxi) (Fig. 1).

Remarks. This species is very similar to *A. serratiprotuberus* and *A. longiserratus* in dorsal and ventral views of the male genitalia, but different in lateral view (epiphallallic lateral lobes of the new species possessing horned protuberances at the inner margin proximally and the outer margin medially, and the ectoparameres of this new species are acute apically).

Acknowledgements

We thank staff of Nonggang National Nature Reserve, for their help in our field work. This work is supported by National Natural Science Foundation of China (No. 31750002, 32070474) and the Fundamental Research Funds for the Central Universities (GK202003049).

References

- Cigliano MM, Otte D, Eades DC, Braun H (2021) Orthoptera species file online. Version 5.0/5.0. <http://Orthoptera.SpeciesFile.org> [accessed 6 September 2021]
- Gorochov AV (1987) On the fauna of crickets of the subfamily Gryllinae (Orthoptera, Gryllidae) of eastern Indochina. In: Medvedev LN (Ed.) Insect Fauna of Vietnam. Nauka, Moscow, 5–22.
- Liu HY, Shi FM (2015) First record of the genus *Anaxiphomorpha* Gorochov (Orthoptera: Gryllidae) from China, with description of four new species. Zootaxa 3918: 433–438 <https://doi.org/10.11646/zootaxa.3918.3.7>
- Ma LB (2018) *Anaxiphomorpha hexagona* sp. n., a new, unusual swordtail cricket collected from Motuo County, Xizang, China (Orthoptera: Grylloidea; Trigonidiidae; Trigonidiinae). Zootaxa 4418: 197–200. <https://doi.org/10.11646/zootaxa.4418.2.10>