

Description of one new species of the genus *Orthozona* Hampson, 1895 (Lepidoptera, Erebidae, Herminiinae) from China

Ting-Ting Zhao¹, Xin-Yu Zhang², Hui-Lin Han^{1,3,4}

1 School of Forestry, Northeast Forestry University, Harbin, 150040, China **2** Simianshan Forest Resource Service Center, Jiangjin District, Chongqing, 402296, China **3** Key Laboratory of Sustainable Forest Ecosystem Management-Ministry of Education, Northeast Forestry University, Harbin, 150040, China **4** Northeast Asia Biodiversity Research Center, Northeast Forestry University, Harbin, 150040, China

Corresponding author: Hui-Lin Han (hanhuilin@aliyun.com)

Academic editor: Pavel Stoev | Received 28 December 2021 | Accepted 22 February 2023 | Published 13 March 2023

<https://zoobank.org/AA538D0F-56CF-4A53-9626-B685BDFF0E64>

Citation: Zhao T-T, Zhang X-Y, Han H-L (2023) Description of one new species of the genus *Orthozona* Hampson, 1895 (Lepidoptera, Erebidae, Herminiinae) from China. ZooKeys 1153: 65–72. <https://doi.org/10.3897/zookeys.1153.79856>

Abstract

A new species of the genus *Orthozona* Hampson, 1895, *O. parallelilineata* sp. nov., is described from China. The new species is illustrated with images of adults and genitalia, and it is compared to similar species, *O. quadrilineata* and *Paracolax curvilineata*. A distribution map of this new species is also presented.

Keywords

Herminine moths, new species, southwest China, taxonomy

Introduction

The genus *Orthozona* was erected by Hampson (1895) with *Madopa quadrilineata* Moore, 1882 as the type species from Darjiling, India. The genus has included as many as five species, namely *Orthozona bilineata* Wileman, 1915; *O. curvilineata* Wileman, 1915; *O. quadrilineata* (Moore, 1882); *O. karapina* Strand, 1920, and *O. rufilineata* (Hampson, 1895) (Poole 1989; Beccaloni et al. 2003). However, *O. bilineata* and *O. curvilineata* were transferred to *Paracolax* Hübner, [1825] by Owada (1992a) and

Owada et al. (2021). To date, two species are known to occur in China: *O. karapina* and *O. quadrilineata* (Owada 1992b; Chen 1999; Wu et al. 2013). Smetacek and Kitching (2012) recorded a male specimen of *O. quadrilineata* from northwestern India and illustrated a female syntype of *O. quadrilineata* from Darjeeling, but they did not show the male and female genitalia. In the present study, a new species is described from Xizang Autonomous Region, China. Further study is needed to understand the relatedness of *Orthozona* with *Paracolax*.

Materials and methods

Specimens were collected in Xizang Autonomous Region, China, using a 220 V/450 W mercury lamp and a DC black light. Standard methods for dissection of the genitalia and preparation of the slide mounts were used (Kononenko and Han 2007). Photographs of the adults were taken with a Nikon D700 camera; genitalia slides were photographed with an Olympus photo-microscope, composited with Helicon Focus, and further processed with Adobe Photoshop CS6. The types of the two new species are deposited in the collection of Northeast Forestry University, Harbin, China.

Abbreviations for institutional collections are as follows:

- NEFU Northeast Forestry University (Harbin, China);
NSMT National Museum of Nature and Science (Tsukuba, Japan).

Taxonomic account

Genus *Orthozona* Hampson, 1895

Orthozona Hampson, 1895, 94. Type species: *Madopa quadrilineata* Moore, 1882.

Diagnosis. In morphology, both *Orthozona* and *Paracolax* Hübner, [1825] share some characters as follows: the color of the forewings is ochre to dark ochre, the antemedial and postmedial line are distinct; the distal part of valva are tongue-like, the saccular process are developed; and the ductus bursae have a pair of sclerotized stripes. However, these genera can be distinguished by the following features: the antemedial and postmedial lines in *Orthozona* are inwardly oblique or slightly curved but always parallel with each other, whereas the two lines in *Paracola* are oblique or wavy and seldom parallel; the reniform spots in *Orthozona* are not obvious, while these spots in *Paracola* are obvious; the saccular process of *Orthozona* is finger-shaped, slightly separated from the valva, and slightly sclerotized, whereas the saccular process in *Paracola* is variously shaped, e.g. digitiform, conical, truncate, etc.; and the corpus bursae of *Orthozona* is sac-like, with a single signum, while the corpus bursae of *Paracola* is very long, more than twice as long as the ductus bursae, and with or without signum (Owada 1992a; Chen 1999; Smetacek and Kitching 2012; Wu 2014).

Description. The proboscis is developed; the labial palpus in males is sickle-shaped, mostly covered with scale tufts ventrally; labial palpus in females with the 2nd segments straight and the 3rd segments upturned; the antenna is filiform. Thorax: quite stout; the forewing is broad, with the outer margin broad and slightly excurred; in many species of this genus, the medial line of the hindwing is indistinct and the subterminal line is slightly arched. Abdomen: slender, slightly lighter than the thorax; the uncus is slender, with a hooked apex; the tegumen is narrow and triangular; the saccus is U-shaped; the valva is simple and weakly sclerotized, with saccular process; the vesica is covered minutely granular and bears a basal cornutus; in females, the analis papili is short, the apophyses posteriores and anteriores are moderate in length, the ductus bursae is short, the corpus bursae have more extensive microspines, and a signum is present.

***Orthozona parallelilineata* Zhao, Zhang & Han, sp. nov.**

<https://zoobank.org/5DE18C80-120E-445A-A501-54900F5F7D91>


Figs 1, 2, 7, 9, 12–14

Material examined. **Holotype:** ♀, CHINA; Xizang Autonomous Region, Linzhi City, Lulang Town; 19.VIII.2014; H.L. Han leg.; genitalia No. zxy-0132-2; coll. NEFU.

Paratypes: 1 ♀, CHINA; Xizang Autonomous Region, Linzhi City, Pailong Countryside; 22–23.IX.2011; H.L. Han leg.; genitalia No. ztt-5280-2; 1 ♀ • Xizang Autonomous Region, Linzhi City, Nadengzuo Village; 14–15.VIII.2014; H.L. Han leg.; genitalia No. ztt-5286-2; 3 ♂♂, 2 ♀♀ • Xizang Autonomous Region, Linzhi City, Nadengzuo Village; 17.VIII.2014; H.L. Han leg.; genitalia No. zxy-0099-2, zxy-0103-1, ztt-5278-2, ztt-5281-1, ztt-5284-1; 3 ♀♀, 1 ♂ • Xizang Autonomous Region, Linzhi City, Mount Sejila; 20.VIII.2014; H.L. Han leg.; genitalia No. hhl-5279-1, hhl-5282-2, hhl-5283-2, hhl-5291-2; 1 ♀ • Xizang Autonomous Region, Linzhi City, Mount Sejila; 22.IX.2016; Z.H. Pan leg.; genitalia No. hhl-5289-2; 3 ♂♂ • Xizang Autonomous Region, Linzhi City, 13.VIII.2017; H.L. Han leg.; genitalia No. hhl-5285-1, hhl-5288-1, hhl-5293-1; coll. NEFU.


Diagnosis. *O. parallelilineata* sp. nov. (Figs 1, 2) is superficially similar to *O. quadrilineata* (Figs 3, 4) and *P. curvilineata* (Figs 5, 6) but can be separated from these species by the following characters. In the male genitalia (Fig. 7), the valva is narrower (in *P. curvilineata*, the cucullus is roundish and broader); the sacculus process is approximately 3/4 the length of valva, narrow, and the distal part is thin, finger-shaped (in *P. curvilineata*, the length of sacculus processes is obviously less than 1/2 of the valva, and the sclerotized is apically pointed); the saccus is narrower (in *P. curvilineata*, the saccus is broader); the phalli are slightly longer (in *P. curvilineata*, the phalli are shorter and stouter).

The characters of female genitalia (Fig. 9) are very similar to those of *O. quadrilineata* (Fig. 10) and *P. curvilineata* (Fig. 11); the corpus bursae is covered with sparse spinules (in *O. quadrilineata* and *P. curvilineata*, the spinules are more dense); the ductus bursae has two long, sclerotized longitudinal bands, which is similar to that of *P. curvilineata* (in *O. quadrilineata*, the ductus bursae has a small sclerotized band).




Figures 1–6. *Orthozona* and *Paracolax* spp., adults **1** *O. parallelilineata* sp. nov., female, holotype, Xizang **2** *O. parallelilineata* sp. nov., male, paratype, Xizang **3** *O. quadrilineata* (Moore, 1882), female, Sikkim (from Dr Mamoru Owada) **4** *O. quadrilineata* (after Smetacek and Kitching 2012) **5** *P. curvilineata* (Wileman, 1915), male, Taiwan (from Dr Shipher Wu) **6** *P. curvilineata*, female, Taiwan (from Dr Shipher Wu).

Description. Adult (Figs 1, 2). Wingspan 37–39 mm in both sexes. Head: dark brown; labial palpus sickle-shaped in male, extended forward in female; antenna filiform. Thorax: patagium and tegula dark brown; mesothorax and metathorax light ochre-grey. Forewing light ochre-brown, diffused with brown scales; antemedial line dark brown, excurved, with a deep, excurved dent in Cu_2 –1A+2A area; medial line inwardly oblique, dark blackish brown, broadly band-shaped, slightly curved inwards; postmedial line narrow, brown, waved on the vein, with a large, excurved dent at the costal margin region, and a large, incurved dent in Cu_2 –1A+2A area; subterminal line inwardly oblique, dark blackish brown, broadly band-shaped, slightly curved inwards, parallel to the medial line, and running from the apex to inner margin close to the tornus; terminal line grey, narrow; orbicular spot small, brown, distinct, encircled with



Figures 7–11. Genitalia of *Orthozona* and *Paracolax* spp. **7** *O. parallelilineata* sp. nov., paratype, genit. prep. zxy-0103-1 **8** *P. curvilineata* (Wileman, 1915), Taiwan (from Dr Mamoru Owada, genit. prep. NSMT4187, in NSMT) **9** *O. parallelilineata* sp. nov., holotype, genit. prep. zxy-0132-2 **10** *O. quadrilineata* (Moore, 1882), Sikkim (from Dr Mamoru Owada, genit. prep. NSMT4373, in NSMT) **11** *P. curvilineata* (from Dr Mamoru Owada).

a halo; reniform spot oval; a flap with a fringe of scales underlapping the forewing costa below over ca 2/3 of its length in male. Hindwing slightly lighter than forewing; media line brown, more prominent in inner margin region; postmedial line very narrow,



Figures 12–14. Distribution and collecting sites of *Orthozona* spp. **12** distribution map: a = Mount Sejila; b = Nadengzuo Village; c = Lulang Town; d = Layue Village; e = Pailong Countryside **13** collection site at Nadengzuo Village, Linzhi City, Xizang Autonomous Region **14** collection site in Luluang countryside, Linzhi City, Xizang Autonomous Region.

almost not visible; subterminal line broad, blackish brown to brown, gradually noticeable from costal to inner margin; discal spot obscure. Both forewing and hindwing with a discal spot on the ventral side, respectively. Abdomen: light ochre-grey, rather slender, terminus with yellow tufts.

Male genitalia (Fig. 7). Tegumen broad, slightly longer than vinculum. Vinculum narrow, U-shaped. Saccus with a rounded and rough central plate. Valva narrow,

insole-shaped, gradually widening from base to cucullus; sacculus moderately sclerotized, narrow, band-shaped, ca 3/4 length of valva; sacculus process sclerotized, narrow, finger-shaped, weakly split from the valva; costa flattened, tapering from basal to terminal part, ca 2/3 length of valva; cucullus smoothly arched. Uncus long and slender, tapering from basal part to top, curved at 1/3 from base, apically sharp. Juxta inverted V-shaped, slightly concavea on both sides, papillate and sclerotized apically. Phallus straight; coecum narrow, ca 1/3 length of aedeagus; carina with a short and narrow spine, and with a sclerotized and short band ventrally. Vesica has four membranous diverticula, densely covered with small grains; basal part of vesica with a reversed, curved, saw-toothed cornutus; subbasal diverticulum is the largest.

Female genitalia (Fig. 9). Ostium bursae broad and straight. Dorsal side of 8th segment with a pair of subcircular, slightly sclerotized areas, connected to base of apophysis anterior. Ductus bursae flattened, shorter than 1/2 length of corpus bursae, with two long, sclerotized longitudinal bands ca 2/3 length of ductus bursae. Corpus bursae membranous; anterior part densely covered with small granulations, with an oval signum covered with small, stout spines; posterior half smooth, covered with longitudinal wrinkles. Apophysis anterior thicker and longer than apophysis posterior. Papilla analis thick and short.

Distribution. China (Xizang: Linzhi) (Fig. 12). This species occurs in grasslands around coniferous and broad-leaved mixed forests in the Xizang Autonomous Region (Figs 13, 14). It flies in the dry season. Specimens were captured with UV light in August.

Etymology. The species is named for the parallel medial and subterminal lines.

Remarks. *Orthozona quadrilineata* and *Paracolax curvilineata* are very similar in external appearance and genitalia in both sexes. However, they are not discussed further here due to the poor condition of their materials; the relationship between these three species deserves in-depth study when the materials become available. In this paper, one female of *O. quadrilineata* was collected by Dr M. Owada on 25 September 1983 in Sikkim, India (genitalia slide no. NSMT4373).

Acknowledgements

This research was supported by grants from the National Natural Science Foundation of China (No. 31872261), the project of the Northeast Asia Biodiversity Research Center (NABRI202203) and the Fundamental Research Funds for the Central Universities (No. 2572019CP11, 2572021DJ08, 2572021AW28). We sincerely thank Dr Shipher Wu (Biodiversity Research Center, Academia Sinica, Taipei) for providing the images of *P. curvilineata*, and Dr. Mamoru Owada (NSMT, Tsukuba) for the images of *O. quadrilineata* and *P. curvilineata*. We also would like to thank Dr Alberto Zilli (Natural History Museum, London, UK) for his valuable comments and precise linguistic adjustments.

References

- Beccaloni GW, Scoble MJ, Robinson GS, Pitkin B (2003) The Global Lepidoptera Names Index (Lep Index). <http://www.nhm.ac.uk/entomology/lepinde> [Accessed 31 March 2011]
- Chen YX (1999) Lepidoptera: Noctuidae: Fauna Sinica (Insecta) 16. Science Press, Beijing, 1596 pp.
- Hampson GF (1895) The Fauna of British India Including Ceylon and Burma. Moths (Vol. 2). Taylor and Francis, London, 546 pp. <https://doi.org/10.5962/bhl.title.100745>
- Kononenko VS, Han HL (2007) Atlas Genitalia of Noctuidae in Korea (Lepidoptera). In: Park K-T (Ed.) Insects of Korea (Series 11). Junhaeng-Sa, Seoul, 464 pp.
- Moore F (1882) Descriptions of Indian Lepidoptera Heterocera from the collection of the late Mr. W.S. Atkinson [part 2]. In: Hewitson WC, Moore F (Eds) (1879–1888) Descriptions of New Indian Lepidopterous Insects from the Collection of the Late Mr. W.S. Atkinson. M.A., F.L.S., &c. Asiatic Society of Bengal, Calcutta, 89–198.
- Owada M (1992a) Synonymic notes on the herminiine moths (Noctuidae) of Japan with descriptions of three new species. *Tinea* 13: 183–203.
- Owada M (1992b) Herminiinae. In: Heppner JB, Inoue H (Eds) Lepidoptera of Taiwan (Vol. 1). Part 2: Checklist. Association for Tropical Lepidoptera, Gainesville, 172–173.
- Owada M, Wu S, Fu CM (2021) Herminiinae. In: Fu CM, Owada M, Shih LC, Lin HH (Eds) Moths of Nanheng Part 2. Endemic Species Research Institute, Nantou, 261–331.
- Poole RW (1989) Noctuidae. In: Heppner JB (Ed.) Lepidopterorum Catalogus (New Series), 118. E.J. Brill/Flora & Fauna Publications, Leiden, 745 pp.
- Smetacek P, Kitching IL (2012) The hitherto undescribed male of *Orthozona quadrilineata* (Moore, 1882) (Lepidoptera: Erebidae). *Journal of Threatened Taxa* 4(14): 3366–3368. <https://doi.org/10.11609/JoTT.o2749.3366-8>
- Wileman AE (1915) New species of Heterocera from Formosa. *Entomologist* 48: 12–40.
- Wu S (2014) Systematics of Herminiinae of the Palaearctic and Oriental regions, with a taxonomic revision of the Taiwanese fauna (Lepidoptera, Erebidae). PhD thesis, National Taiwan University, Department of Entomology, College of Bioresources and Agriculture, [ix +] 357 pp.
- Wu S, Fu CM, Owada M (2013) Erebidae: Herminiinae. In: Fu CM, Ronkay L, Lin HH (Eds) Moths of Hehuanshan. Endemic Species Research Institute, Nantou, 333–363.