

A new species of Metopiinae (Hymenoptera, Ichneumonidae) parasitizing lepidopteran larvae in China

Ying Zhang¹, Mao-Ling Sheng¹, Zi-Cheng Xiong²

1 General Station of Forest Pest Management, State Forestry Administration, Shenyang 110034, P.R. China
2 Forest Pest Control and Quarantine Station of Etuoke, Inner Mongolia Autonomous Region 016100, P.R. China

Corresponding author: *Mao-Ling Sheng* (shengmaoling@163.com)

Academic editor: *B. Santos* | Received 4 February 2016 | Accepted 19 February 2016 | Published 15 March 2016

<http://zoobank.org/52AB4358-8972-4DB8-922E-33BA01DAF783>

Citation: Zhang Y, Sheng M-L, Xiong Z-C (2016) A new species of Metopiinae (Hymenoptera, Ichneumonidae) parasitizing lepidopteran larvae in China. ZooKeys 572: 71–79. doi: 10.3897/zookeys.572.8031

Abstract

A new species of Metopiinae, *Trieces etuokensis* Sheng, **sp. n.**, is described and illustrated. Specimens were reared from two species of Lepidoptera: *Bazaria turensis* (Ragonot, 1887) (Pyrilidae) from Balong, Dulan, Qinghai Province, and an unidentified psychid (Psychidae) from Mukainor, Etuoke, Inner Mongolia Autonomous Region, China. The new species is characterized by a yellow face and clypeus, fore and middle femora and hind tibia mainly black, antennae slightly longer than head and mesosoma combined, with 17 flagellomeres, occipital carina entirely absent, and the hind femur being compressed, 2.5 times as long as maximum width.

Keywords

Trieces, new species, *Bazaria turensis*, Pyrilidae, Psychidae, China

Introduction

Trieces Townes, 1946 (Hymenoptera, Ichneumonidae, Metopiinae) comprises 68 species (Tolkanitz 2010, Yu et al. 2012), of which 13 are from the Eastern Palaearctic Region (Tolkanitz 2010). No species of *Trieces* have been reported from China to date. The genus is characterized mainly by areolet absent, mesopleural suture indistinct or

absent, lateral carina of first tergite weak or obsolescent, second tergite with complete median and sublateral longitudinal carinae, basal portion of sublateral longitudinal and median carinae of third tergite present, and epipleura of third to fifth tergites almost absent (Townes 1971, Gauld et al. 2002, Tolkanitz 1987, 2010).

The known hosts of *Triece* Townes mainly belong to the lepidopteran families Elachistidae (Bradley 1974), Geometridae (Petrice, et al. 2004), Tortricidae (Walley 1969) and Yponomeutidae (Gencer 2003, Yu et al. 2012).

Herein one new species of *Triece* is reported, reared from the pupae of *Bazaria turensis* (Pylalidae) and an unidentified psychid (Psychidae).

Materials and methods

Mature larvae of the host, *Bazaria turensis* (Ragonot, 1887) were collected on 28 August 2013 in a forest from where there had been an outbreak lasting at least three years, and brought to the laboratory. The forest is a shrubbery (Fig. 1) composed of *Nitraria tangerorum* Bobrov, *Lycium chinense* Miller var. *potaninii* (Pojarkova) A.M. Lu and *Kalidium foliatum* (Pallas) Moquin-Tandon, located in Dulan County, 36°09.65'N; 97°27.42'E, Qinghai Province. Mature larvae were maintained in a nylon cage at room temperature, and after pupating were stored individually in glass tubes (60 × 6 mm) with a piece of filter paper dipped in distilled water to maintain moisture and plugged tightly with absorbent cotton. After the emergence of moths and parasitoids was complete, remaining pupae were dissected to record their condition (i.e. status of moths, and parasitism).

Mature larvae of the psychid moth were collected in 16 September 2014 from a scrub where there had been an outbreak lasting at least fourth years, and brought to the laboratory. The scrub (Fig. 2) is composed of *Caragana intermedia* Kuang & H.C. Fu and located in Mukainor, 39°33.71'N; 108°40.24'E, Etuoke, Inner Mongolia Autonomous Region.

Images were taken using a Leica M205A Stereomicroscope with LAS Montage MultiFocus. Morphological terminology is mostly based on Gauld (1991).

Type specimens are deposited in the Insect Museum, General Station of Forest Pest Management (GSFPM), State Forestry Administration, People's Republic of China.

Results

Triece Townes, 1946

Triece Townes, 1946. Boletín de Entomología Venezolana, 5:60. Type-species: *Exochus texanus* Cresson.

Diagnosis. Interantennal process forming a triangular projection in front of antennal sockets. Anterior spur of mid tibia 0.25× to 0.65× length of posterior spur. Meso-



Figure 1. Habitat of *Trieces etuokensis* Sheng, sp. n. The shrubbery in Balong, Dulan County, Qinghai Province.



Figure 2. Habitat of *Trieces etuokensis* Sheng, sp. n. The scrub in Mukainor, Etuoke, Inner Mongolia Autonomous Region.

pleural suture indistinct or absent. Areolet absent. Lateral carina of first tergite weak or obsolescent. Second tergite with complete median and sublateral longitudinal carinae. Basal portion of sublateral longitudinal and median carinae of third tergite present. Epipleura of third to fifth tergites almost absent.

In Tolkanitz's (2010) key to the species of Palaearctic fauna, the new species can be inserted as follows:

- 14 (15) Head in dorsal view widened posteriorly behind eyes. Occipital carina obsolete. Metapleura as in Figs 3, 15. Face and clypeus of female reddish brown (Figs 5, 4). 2.7. (Male unknown). Russia (Amurskaya Province)....
..... *T. femoralis* Tolkanitz

- 15 (14) Head in dorsal view not widened posteriorly behind eyes, parallel-sided. Metapleura as in figs 3, 16. Occipital carina absent. Face, clypeus, and genae of female yellow **15(a, b)**
- 15a (15b) Antenna not longer than head and mesosoma combined; fore wing with vein 1cu-a slightly distal of 1-M, nearly interstitial; length of hind femur 2.2× its width; hind tibia reddish yellow; antennae reddish yellow. (Male unknown). Mongolia..... ***T. pumicatus* Tolkanitz**
- 15b (15a) Antenna longer than head and mesosoma combined; fore wing with vein 1cu-a distinctly distal of 1-M, distance between them 0.4× length of 1cu-a; hind femur distinctly compressed, 2.5× as long as maximum width; hind tibia mainly black; basal ventral profile of antennae red brown, basal dorsal profile blackish brown; apical portion brownish black. China (Inner Mongolia Autonomous Region) ***T. etuokensis* Sheng, sp. n.**

***Trieces etuokensis* Sheng, sp. n.**

<http://zoobank.org/23C8E597-4794-4069-B60C-665573C8159C>

Figs 3–12

Etymology. The specific name is derived from the holotype locality.

Material examined. Holotype female reared from pupa of unidentified psychid moth on 27 October 2014, CHINA: Mukainor, 1476m, Etuoke, Inner Mongolia Autonomous Region, Mao-Ling Sheng. Paratypes: 1 male, same data as holotype except 24 October 2014. 1 female reared from pupa of *Bazararia turensis* (Ragonot, 1887) on 2 October 2013, China: Balong, 2857m, Dulan, Qinghai Province, Mao-Ling Sheng.

Diagnosis. Malar space approximately as long as basal width of mandible. Ocular-ocellar line as long as diameter of posterior ocellus. Antenna slightly longer than head and mesosoma combined, with 17 flagellomeres. Occipital carina entirely absent. Metapleuron shiny, impunctate, lower posterior portion with distinct wrinkles. Hind femur compressed, 2.5× as long as maximum width. Face and clypeus yellow. Fore and middle femora and hind tibia mainly black.


Description. Female. Body length approximately 2.8 mm. Fore wing length approximately 2.2 mm.

Head. Inner margins of eyes distinctly convergent downwards. Face (Fig. 4) with dense fine punctures, distance between punctures 0.2–0.5× diameter of puncture. Clypeus with denser punctures than that of face, distance between punctures approximately 0.2× diameter of puncture, apical portion slightly concave; apical margin almost truncate. Mandible small, weakly narrowed toward apex; upper tooth distinctly longer than lower tooth. Malar area flat, with fine punctures. Malar space approximately as long as basal width of mandible. Gena in dorsal view approximately as long as width of eye, almost smooth, with sparse, indistinct fine punctures, scarcely convergent posteriorly. Vertex (Fig. 5) and frons almost shiny, with indistinct fine punctures. Postocellar line 1.2× as long as ocular-ocellar line. Ocular-ocellar line approximately as long as



Figures 3–5 *Triecces etuokensis* Sheng, sp. n. Holotype. Female **3** Habitus, lateral view **4** Head, anterior view **5** Head, dorsal view. Scale bars: 0.5 mm (**3**); 0.1 mm (**4, 5**).

diameter of posterior ocellus. Upper portion of frons slightly convex, lower portion slightly concave. Antenna (Fig. 6) slightly longer than head and mesosoma combined, with 17 flagellomeres; each flagellomere longer than wide. First flagellomere 2.2× as long as maximum width, 1.1× as long as third flagellomere. Occipital carina absent.



Figures 6–12. *Trieces etuokensis* Sheng, sp. n. Holotype. Female **6** Antenna **7** Mesoscutum and scutellum **8** Mesosoma, lateral view **9** Fore wing **10** Propodeum **11** Tergites 1–4, dorsal view **12** Apical portion of metasoma, lateral view. Scale bars: 0.1 mm (**6, 7, 8, 10**); 0.2 mm (**9, 11, 12**)

Mesosoma. Lateral concavity of pronotum smooth, shiny, upper-posterior portion slightly rough, with dense shallow fine punctures. Epomia indistinct. Mesoscutum (Fig. 7) smooth, shiny, anterior portion slightly convex, posterior flat; with dense distinct punctures, distance between punctures 0.2–2.0× their diameter. Notaulus absent. Scutellum almost flat, shiny, with sparse, indistinct, fine punctures; lateral carina reaching apex. Postscutellum so short it resembles a transverse carina. Mesopleuron (Fig. 8) with sparse fine punctures. Speculum with fine indistinct punctures. Mesopleural fovea vestigial. Upper end of epicnemial carina reaching anterior margin of mesopleuron, at dorsal 0.75 of posterior margin of pronotum. Metapleuron almost flat, shiny, postero-dorsal portion with few hairs, lower posterior portion with distinct wrinkles. Juxtacoxal carina strong. Wings (Fig. 9) slightly infusate. Fore wing with vein 1cu-a strongly inclivous, distal to 1-M by about 0.4× length of 1cu-a. 2m-cu straight, distinctly inclivous, distal to 2rs-m by about 1.4× length of 2rs-m. Hind wing vein 1-cu 2.0× as long as cu-a. Hind femur compressed, 2.5× as long as maximum width. Hind tibia gradually widened apically. Posterior spur of hind tibia approximately 2.75× as long as anterior spur. Ratio of length of hind tarsomeres 1:2:3:4:5 is 2.8:1.2:1.0:0.8:1.6. Propodeum (Fig. 10) with strong apical transverse and complete longitudinal carinae. Pleural areas with dense, distinct, fine punctures. Area petiolaris with longitudinal wrinkles. Remainder of propodeum smooth, shiny, with indistinct fine punctures. Propodeal spiracle small, circular.

Metasoma (Figs 11, 12). First tergite approximately 0.75× as long as apical width, with dense punctures; median dorsal, dorsolateral, ventrolateral and sublateral longitudinal carinae complete; apical half with distinct median longitudinal carina. Second tergite (Fig. 11) 1.27× as long as apical width, densely punctate, median and sublateral carinae complete. Third tergite 0.78× as long as apical width; basal 0.7 with dense punctures, apical 0.3 with sparser punctures and smoother than basal 0.7; basal 0.4 with median longitudinal carina; basal 0.3 with lateral longitudinal carinae. Basal 0.6 of fourth tergite densely punctate, apical 0.4 shiny, gradually impunctate. Basal 0.4 of fifth tergite with dense punctures, apical 0.6 gradually impunctate. Ovipositor (Fig. 12) tapered from base to apex, with a large, shallow notch.

Color (Fig. 3). Black, except as follows. Face (Fig. 4) except upper median light brown spot, malar area and clypeus yellow. Maxillary and labial palpi yellow brown. Anterior profile of pedicel dark brown; basal ventral profile of antenna red-brown, basal dorsal profile darkish brown; apical portion brownish black. Anterior and posterior profiles of fore femur, tibia and tarsus, basal and apical portions of mid tibia, mid tarsus, all trochanters more or less, tegula brown to dark brown. Pterostigma (Fig. 9) blackish. Wing veins dark brown.

Male. Body length approximately 3.1 mm. Fore wing length approximately 2.5 mm. Antenna with 22 flagellomeres. Very similar to the female, except with hind first tarsomere yellow, apical portion pale brown yellow.

Hosts. One female was reared from pupa of *Bazaria turensis* (Ragonot, 1887) (Lepidoptera: Pyralidae). One female and one male were reared from unidentified species of Psychidae (Lepidoptera) collected as mature larvae but details of development and emergence unknown.

Host plants. *Caragana intermedia* Kuang & H.C. Fu (Leguminosae), *Nitraria tangutorum* Bobrov (Zygophyllaceae), *Kalidium foliatum* (Pallas) Moquin-Tandon (Amaranthaceae).

Remarks. This new species is similar to *T. pumicatus* Tolkanitz, 2010 and can be distinguished from the latter by the following combination of characters: antenna slightly longer than head and mesosoma combined, with 17 flagellomeres (female); fore wing with vein 1cu-a distinctly distal of 1-M, distance between them 0.4× length of 1cu-a; hind femur compressed, 2.5× as long as maximum width; lateral longitudinal carinae of tergite 3 distinct on anterior 0.3; fore and middle femora, hind tibia mainly black; basal ventral profile of antennae red brown, basal dorsal profile darkish brown; apical portion brownish black. The same characters for *Trieces pumicatus* Tolkanitz are as follows: antenna not longer than head and mesosoma combined, with 20 flagellomeres (female); fore wing with vein 1cu-a slightly distal of 1-M, nearly interstitial; length of hind femur 2.2× its width; lateral longitudinal carinae of tergite 3 vanishing behind its middle; fore and mid legs, hind tibia reddish yellow, fore and mid femora slightly darkened on outer side; antenna reddish yellow.

Acknowledgements

The authors are deeply grateful to Drs. Gavin Broad (Department of Life Sciences, the Natural History Museum, London, UK) and Bernardo Santos (American Museum of Natural History, New York, USA) for reviewing this manuscript, and Yan-Ling Zhang (Director of Forestry Pest Control and Quarantine Station of Dulan, Qinghai, China) for her help in the course of exploration in Qinghai Province. This research was supported by the “Twelfth Five-year” National Science and Technology Support Program of China (Grant No. 2012BAD19B0701) and the National Natural Science Foundation of China (NSFC, No. 31372246, No. 31070585).

References

- Bradley GA (1974) Parasites of forest Lepidoptera in Canada. Subfamily Metopiinae and Pimplinae (Hymenoptera: Ichneumonidae). Part 1. Environment Canada, Canadian Forestry Service, Publication No.1336, 99 pp.
- Gauld ID (1991) The Ichneumonidae of Costa Rica, 1. Introduction, keys to subfamilies, and keys to the species of the lower Pimpliform subfamilies Rhyssinae, Poemeniinae, Acaenitinae and Cylloceriinae. *Memoirs of the American Entomological Institute* 47: 1–589.
- Gauld ID, Sithole R, Gómez JU, Godoy C (2002) The Ichneumonidae of Costa Rica. 4. *Memoirs of the American Entomological Institute* 66: 1–768.
- Gencer L (2003) The parasitoids of *Yponomeuta malinellus* Zeller (Lepidoptera: Yponomeutidae) in Sivas. *Turkish Journal of Zoology* 27(1): 43–46.

- Petrice TR, Strazanac JS, Butler L (2004) A survey of hymenopteran parasitoids of forest Macrolepidoptera in the central Appalachians. *Journal of Economic Entomology* 97(2): 451–459. doi: 10.1093/jee/97.2.451
- Tolkanitz VI (1987) Parasitic Hymenoptera. Ichneumonidae - Metopiinae. *Fauna Ukraina* 11(2): 1–212.
- Tolkanitz VI (2010) Ichneumon flies of the genus *Triece* (Hymenoptera, Ichneumonidae, Metopiinae) in the Palearctic Fauna. *Entomological Review* 90(4): 465–472. doi: 10.1134/S001387381004007X
- Townes HK (1971) The genera of Ichneumonidae, Part 4. *Memoirs of the American Entomological Institute* 17: 1–372.
- Walley GS (1969) A synopsis of the Nearctic species of the onitis group of the genus *Triece* (Hymenoptera: Ichneumonidae). *Canadian Entomologist* 101: 1092–1106. doi: 10.4039/Ent1011092-10
- Yu DS, van Achterberg C, Horstmann K (2012) Taxapad 2012 – World Ichneumonoidea 2011. Taxonomy, Biology, Morphology and Distribution. On USB Flash drive. Ottawa, Ontario, Canada. www.taxapad.com