

Replacement name for a Panamic bivalve (Mollusca, Bivalvia, Cyrenidae)

Eugene V. Coan¹, Paul Valentich-Scott¹

¹ Santa Barbara Museum of Natural History, 2559 Puesta del Sol, Santa Barbara, CA 93105, USA

Corresponding author: Eugene V. Coan (genecoan@gmail.com)

Academic editor: Graham Oliver | Received 3 March 2022 | Accepted 28 April 2022 | Published 23 June 2022

<http://zoobank.org/7778CC28-4E54-4CE4-A030-8C5C2A8A0F28>

Citation: Coan EV, Valentich-Scott P (2022) Replacement name for a Panamic bivalve (Mollusca, Bivalvia, Cyrenidae). ZooKeys 1108: 11–13. <https://doi.org/10.3897/zookeys.1108.83037>

Thomas A. Neubauer of the Systematics & Biodiversity Lab, Justus Liebig University, Giessen, Germany, has called to our attention that *Cyrena acuta* Prime, 1861, published in October, is a junior primary homonym of *Cyrena acuta* Ludwig, 1861, published in January that same year.

Prime's species, currently known as *Polymesoda acuta* (Prime, 1861: 355), ranges from Costa Rica to Ecuador, where it occurs intertidally in mangrove areas (Coan and Valentich-Scott 2012: 464–465). The holotype of this species is deposited in the Museum of Comparative Zoology, USA, Harvard University (MCZ 176951) (Johnson 1959: 441). In Prime's original description the type locality was given only as Central America.

Cyrena acuta Ludwig (1861: 197–199, pl. 72, figs 15, 16) was described from the early Miocene of Münzenberg, Hesse, Germany, where it occurs with other fresh and brackish-water species (Kadolsky 2008). The species is presently considered a junior synonym of *Falsocorbicula faujasii* (Deshayes, 1830: 51) [originally *Cyrena*] (Ott et al. 2009). It also remains in the Cyrenidae.

The International Code of Zoological Nomenclature's (1999) Article 23.9 [Reversal of Precedence] does not apply in that Prime's name has been used less than ten times in the last 150 years since its publication, although Ludwig's name has seen little mention because it has been long regarded as a junior synonym.

We hereby rename *Polymesoda acuta* as *Polymesoda neubaueri* Coan & Valentich-Scott, 2022 (Fig. 1). We restrict the type locality to Costa Rica, Guanacaste Province, Lower Río Tempisque; 10.2583°N, 85.2644°W; intertidal zone, because Prime's locality could have been on either the Atlantic or Pacific coast of Central America and there is no additional information accompanying the holotype (International Code of Zoological Nomenclature (1999: Article 76A). Material from this locality in the Santa Barbara Museum of Natural History collection matches the holotype.




Figure 1. Holotype of *Cyrene acuta* Prime, 1861, renamed herein as *Polymesoda neubaueri* nom. nov. (MCZ 176951), length 41 mm, height 35 mm. **A** exterior of right valve **B** exterior of left valve **C** dorsal view of both valves **D** interior of left valve **E** interior of right valve **F** close up of hinge of left valve **G** close up of hinge of right valve.

References

- Coan EV, Valentich-Scott P (2012) Bivalve seashells of tropical west America. Marine bivalve mollusks from Baja California to northern Perú. Santa Barbara Museum of Natural History, Monographs 6: [xv +] 1258 pp.
- Deshayes GP (1830) Encyclopédie méthodique. Histoire naturelle des vers. Panckoucke, Paris, vol. 2(1): [vii +] 256 pp.
- International Code of Zoological Nomenclature (1999) Fourth edition. International Trust for Zoological Nomenclature, London, UK. <http://www.iczn.org/iczn/index.jsp>
- Johnson RI (1959) The types of Corbiculidae and Sphaeriidae (Mollusca: Pelecypoda) in the Museum of Comparative Zoology, and a bio-bibliographic sketch of Temple Prime, an early specialist of the group. Bulletin of the Museum of Comparative Zoology 120(4): 429–479.
- Kadolsky D (2008) Mollusks from the Late Oligocene of Oberleichtersbach (Rhön Mountains, Germany). Part 1: Overview and preliminary biostratigraphical, palaeoecological and palaeogeographical conclusions. Courier Forschungsinstitut Senckenberg 260: 89–101.
- Ludwig R (1861) [January]. Süßwasser-Bivalven aus der Wetterauer Tertiär-Formation. Palaeontographica 8(6): 195–199 [pl. 72]. www.biodiversitylibrary.org/item/43687#page/9/mode/1up
- Ott W, Kadolsky D, Wiesner E (2009) Von einer Lagune zum “trockenen Kalkhügel”: Geologischer Untergrund und Fossilien des Lohwaldes bei Offenbach am Main. Abhandlungen des Offenbacher Vereins für Naturkunde 10: 113–213.
- Prime T (1861) [18 October]. Diagnoses d'espèces nouvelles. Journal de Conchyliologie 9[(3)4](4): 354–357. [https://ams.wildapricot.org/resources/Documents/2400%20Years%20of%20Malacology%202022/AMS2400collations\(Journals\).pdf](https://ams.wildapricot.org/resources/Documents/2400%20Years%20of%20Malacology%202022/AMS2400collations(Journals).pdf)