

# An updated, illustrated inventory of the marine fishes of the US Virgin Islands

D. Ross Robertson<sup>1</sup>, Carlos J. Estapé<sup>2</sup>, Allison M. Estapé<sup>2</sup>,  
Lee Richter<sup>3</sup>, Ernesto Peña<sup>1</sup>, Benjamin Victor<sup>4,5</sup>

**1** *Smithsonian Tropical Research Institute, Balboa, Panama* **2** *197 Gulfview Drive, Islamorada, Florida, 33036, USA* **3** *National Park Service, 1300 Cruz Bay Creek, St. John, VI 00830, Virgin Islands, USA* **4** *Ocean Science Foundation, 4051 Glenwood, Irvine, CA 92604, USA* **5** *Guy Harvey Research Institute, Nova Southeastern University, 8000 North Ocean Drive, Dania Beach, FL 33004, USA*

Corresponding author: D. Ross Robertson ([robertsondr@si.edu](mailto:robertsondr@si.edu))

Academic editor: Caleb McMahan | Received 14 March 2022 | Accepted 9 May 2022 | Published 1 June 2022

<http://zoobank.org/F2333822-5B4F-45F4-8172-8E794AFBC316>

**Citation:** Robertson DR, Estapé CJ, Estapé AM, Richter L, Peña E, Victor B (2022) An updated, illustrated inventory of the marine fishes of the US Virgin Islands. *ZooKeys* 1103: 79–122. <https://doi.org/10.3897/zookeys.1103.83795>

## Abstract

The US Virgin Islands (USVI) include St. John and St. Thomas on the Puerto Rican Platform (PRP) and St. Croix, isolated by 2000 m deep water 45 km south of that platform. Previous inventories of the marine fishes of these islands include a comprehensive 2014 checklist of the fishes of St. Croix and a list of the fishes of the PRP produced in 2000. The latter list noted the locations of many records of the plateau's fishes, allowing the construction of a combined inventory for St. John and St. Thomas. Those two islands are treated here as a single faunal unit because they are only 3.5 km apart on a shared shallow shelf with various islets and reefs in between. Here we provide updated information on those two USVI (St. Croix and St. John-Thomas) marine fish faunas. The additions to the St. Croix and St. John-Thomas inventories presented here are based on a combination of information from the two sources indicated above, more recent publications dealing with those faunas, a review of location records on various online sources of biogeographic data, and voucher photographs taken of fishes in the field by authors of this paper and other citizen scientists. This assessment increased the known fauna of St. Croix by 7.5% to 585 species. The inventory for St. John-Thomas increased by 39.9% from 401 species on the 2000 PRP list to 561 with the inclusion of records from other sources. On-site mtDNA (COI) barcodes are available for approximately one-third of the species of the St. John-Thomas fauna, but for only one species collected at St. Croix. A set of underwater photographs of 372 species (34 of them representing the sole record of a species) from St. John-Thomas and of 11 shallow-water species added to the St. Croix fauna is included. These represent occurrence vouchers and also are intended to facilitate future work that builds on the present compendium.

**Keywords**

Biodiversity, checklist, citizen science, DNA-barcode, photographic voucher, SCUBA survey

**Introduction**

The United States Virgin Islands (USVI) comprise a US territory adjacent to Puerto Rico, in the northeast Caribbean, that includes three large, inhabited islands, St. John, St. Thomas and St. Croix, and approximately 50 smaller islands and cays around them. The former two are situated only 3.5 km apart, in the center of the Puerto Rico Plateau (PRP), which has an area approximately twice the 9,100 km<sup>2</sup> of Puerto Rico Island and extends ~ 150 km eastwards from Puerto Rico. St. Croix is located south of St. John and St. Thomas, on its own insular platform, which is separated by 45 km of deep water from the southern edge of the PRP.

The fish fauna of St. Croix was comprehensively reviewed by Smith-Vaniz and Jelks (2014), who built upon an older list by Clavijo et al. (1980), using their own extensive collections of shallow fishes of the Buck Island Reef National Monument on the northern side of St. Croix (Smith-Vaniz et al. 2006), and a review of literature and examination of specimens of fishes collected at St. Croix that are lodged in various museums. In 2000, George Dennis produced an extensive (244 page; 500+ sources cited) U.S. Geological Survey report based on collections and observational records for marine and brackish-water fish from Puerto Rico, St. John and St. Thomas, and other islands on the PRP. Although never formally published in a scientific journal, and no longer available through the USGS source cited by Dennis et al. (2004), that compendium is available online (Dennis 2000).

Here we add new information to update the 2014 list for St. Croix and assemble an inventory for St. John and St. Thomas that includes and expands on data for those two islands contained in Dennis (2000). We extracted the additional information from museum records in online sources of biogeographic data, publications produced since Dennis (2000), digital images of live fishes obtained at the USVI, plus our recent collections and mtDNA barcode records obtained from the database BOLD. The great majority of the species in this compendium are marine, plus we include a small number of species found in fresh to brackish waters.

**Materials and methods****Study sites**

St. Croix is a 215 km<sup>2</sup> island in the northeast corner of the Caribbean. It is isolated by ~ 45 km of deep water from the Puerto Rican Platform (**PRP**). Other islands of the Lesser Antilles chain lie within ~ 150 km to the east and southeast of St. Croix. The surrounding

shallow (above ~ 150 m depth) shelf of St. Croix, extending almost 20 km eastward, has approximately the same area as the island. In addition to exposed and sheltered coral reefs and soft bottoms, the island has extensive areas of seagrasses and mangroves.

St. John (area 50 km<sup>2</sup>) and St. Thomas (area 83 km<sup>2</sup>) are situated in the center of the shallow (to ~ 150 m deep) tongue of the PRP that extends 150 km eastwards from Puerto Rico. St. Thomas is closest to and 64 km from the main island of Puerto Rico. St. John and St. Thomas are separated from each other by only 3.5 km of water shallower than 20 m deep, with scattered islets and shallow reefs in between them. They have a similar range of habitats as St. Croix, with large areas of both sheltered and deeper shelf-edge coral reefs, rocky shores, seagrass beds and mangroves. Due to their proximity and similarity of habitats we treat them here as a single unit (hereafter St. John-Thomas). The shallow PRP associated with St. John-Thomas extends ~ 25 km north and ~ 15 km south of those islands and covers an area of ~ 2,100 km<sup>2</sup> (Rohmann et al. 2005).

Suppl. material 2: File S1 shows the bathymetry of bottom habitats on the above-150 m shelves of the USVI. The shelf area of the St. John-Thomas EEZ is not only much larger than that of St. Croix but also differs from the latter in containing a much greater diversity of areas of different depths. There are large expanses, in both absolute and relative terms, of habitat between 40–60 m deep to the north of St. Thomas and to the south of both islands. In contrast, most of the smaller shelf of St. Croix is shallower than 20 m deep.

## Data sources

We reviewed and cited only publications from which we extracted information relating to the USVI fishes that were published after those cited by Dennis (2000) for St. John-Thomas, and after that by Smith-Vaniz and Jelks (2014) for St. Croix, plus a few earlier publications that contained additional relevant information.

Smith-Vaniz and Jelks (2014) published a comprehensive, annotated checklist of 544 fishes known from St. Croix. That checklist was based, in large part, on the yield of fishes from 106 rotenone stations obtained by Smith-Vaniz et al. (2006) and by later workers to document the shallow cryptobenthic fauna. That 2014 list identified questionable records, a few of which, as we show, have turned out to be valid. Smith-Vaniz and Jelks (2014). That checklist also excluded deep-water fishes not found above 200 m as well as Exocoetids and Myctophids. For completeness we have included any such species recorded by other sources among the additions noted here. We used the 2014 list of valid species and reviewed fishes listed by other surveys: a SCUBA study of the shallower parts (30–50 m depth) of a mesophotic coral ecosystem at the eastern end of the shelf (García-Sais et al. 2014); two JSL submersible dives off St. Croix to 30–600 m (Nelson and Appeldoorn (1985); and two ROV dives off St. Croix at depths greater than 800 m (Quattrini et al. 2017). In addition, we reviewed the records of fish species from St. Croix available from various online sources: the aggregators GBIF (<https://www.gbif.org/>), FishNet2 (<http://www.fishnet2.net/>), iDigBio (<https://www.idigbio.org/portal>), OBIS (<https://obis.org/>) and Vertnet (<http://vertnet.org/>), and the American

Museum of Natural History (AMNH; <https://www.amnh.org/research/vertebrate-zoology/ichthyology>). Those searches were made within a quadrat with latitudinal limits of 17.62°N to 17.85°N, and longitudinal limits of -64.4°W to -65.0°W, encompassing St. Croix and all of its platform. The sources of St. Croix records produced by those online searches were evaluated and museum records within the known geographic range of various species were accepted. Evaluation of individual records is necessary because aggregator information includes significant numbers of erroneous records.


Finally, the list includes shallow-reef fishes photographed by authors AME and CJE during a month spent at the island from 19 December 2020 to 13 January 2021. Suppl. material 3: File S2A presents a list, with georeferenced locations, of the 11 dive sites at which they together made 25 dives (total 47 hours duration per person) during that period (see also Fig. 1B and Suppl. material 4: File S3, a Google Earth © KMZ file that shows, for each of those sites, its location and georeferenced coordinates, and the number of dives and total dive time spent at that site). These photographs document a few species not previously recorded at the island, plus several not accepted by Smith-Vaniz and Jelks (2014) due to a lack of reliable information.

For St. John-Thomas we extracted a list of 401 species listed at those islands by Dennis (2000) and reviewed various publications dealing with fish records at and near those islands that were subsequently produced. Finally, we also used the same online data sources as for St. Croix (see above) to obtain records of fishes from the part of the Exclusive Economic Zone of the USVI that includes St. John-Thomas and extends between the northern and southern edges of the PRP. That irregularly shaped EEZ was obtained from [Marineregions.org](http://Marineregions.org), which provides a standard set of global maps of EEZs (<https://www.marineregions.org/eezsearch.php>).

CJE and AME spent six months between 3 November 2020 and 29 May 2021 diving at both islands and photographing fishes to obtain voucher images of as many members of those islands' marine fish fauna as possible. File S2A presents a list, with georeferenced locations, of their dive sites at St. John (37) and St. Thomas (12), at which they made 113 joint dives (involving multiple dives at some sites) totaling 221 hours per person and 37 dives totaling 37 hours per person, respectively. Fig. 1A is a map with those 49 dive sites at St. John-Thomas indicated and File S2 provides additional information. Fig. 1A (and see File S2B) also indicates the location of sites from other sources at which additional species not recorded by CJE and AME were documented photographically by other divers.

## Reef-associated bony fishes of the USVI

Greater Caribbean (GC) reef systems have reef-fish faunas that are dominated by members of typical, shallow-reef families of bony fishes extending down to depths of ~ 250–300 m (Baldwin et al. 2018). Here we focus on species belonging to those families, which have traditionally been viewed as reef fishes. We classed species living entirely or largely below 40 m depth as belonging to the deep-reef subset. Species classed here as shallow include both species restricted to depths shallower than 40 m



**Figure 1.** **A** dive sites generating fish-occurrence data at St. John and St. Thomas islands. Dive sites of CJE and AME are indicated by red symbols, and of other sources of voucher photographs by blue symbols. Note that some close-proximity sites are indicated by a single symbol. Symbols at the northern and southern edges of Fig. 1A are representative only, as their latitudes are outside the area of the map **B** dive sites of CJE and AME generating data at St. Croix. See Suppl. material 3: File S2A, B and Suppl. material 4: File S3 for further information. Base map in both cases: NOAA Chart 25641.

and those with depth ranges that extend above and below that level. These reef-associated fishes include not only benthic and demersal species found on hard-reef substrata, but also pelagic fishes that facultatively associate with reefs and benthic and demersal species that live on soft bottoms within and immediately around the fringes of reefs. Benthic species (e.g., eels, flatfishes) are restricted to life on and in different types of substrata, while demersal species (e.g., snappers and grunts) use both substratum habitats and the water column. Cryptobenthic species are visually cryptic and typically small. We followed Brandl et al. (2018) in classifying families dominated by small cryptobenthic coral-reef species as Core Coral Reef Fish families (CCRFs).

We also evaluate the ecological and zoogeographic composition of the two USVI fish faunas (St. Croix and St. John-Thomas) compared to the complete checklist of the regional fauna of reef-associated bony fishes, which includes 992 species in 342 genera and

84 families (Robertson and Tornabene 2021). These aspects of the fauna of the USVI are also compared with results from another recent comprehensive survey of the fish fauna of nearby Sint Eustatius, which is 170 km from St. Croix (Robertson et al. 2020).

### mtDNA-barcode coverage of fishes collected in the USVI and Puerto Rico

Relatively few small marine locations have been comprehensively sampled for fish DNA barcoding, i.e., tissues sequenced for the mtDNA COI marker as a standard for identifying fishes, as compiled in the Barcode of Life Database, BOLD (Ward et al. 2009). Notably, BOLD not only includes a wide variety of projects, most of which are publicly available, but also regularly harvests all available COI sequences from GenBank. In contrast, GenBank does not harvest from BOLD, and BOLD sequences are generally submitted to GenBank only by request. As a result, only a fraction (~ 15% for GC fishes) of COI sequences on BOLD also are present on GenBank, despite its widespread use as the sole source for barcoding studies. BOLD further differs from GenBank by applying quality control to sequences and taxon identifications as data is entered, including sequences harvested from GenBank. It also has post-hoc quality control via a tagging and comment option on individual records. BOLD also includes a large number of private sequences, which can be assessed to a limited degree (with some metadata removed) via the BIN portal, which compiles all records, public and private, within a lineage, assigns a code, and presents some statistics, especially variance and nearest neighbor distances, as well as countries of origin.

The BOLD BIN code is a key advance enabling the compilation and comparison of mtDNA barcoding lists, since it supplies an independent identifier for a monophyletic genetic lineage, which is not the same as a species name. BOLD creates **BINs** (Barcode Index Numbers) by clustering barcode sequences algorithmically. The BIN often represents a particular species, but there are many exceptions to the “one-species, one BIN” concept: either multiple BINs per species, indicating genetically divergent populations within species (usually allopatric, but not always), a subset of which are putative new cryptic species awaiting morphological confirmation; or shared BINs by two or more species that retain shared or closely related haplotypes due to a short time since speciation, to incomplete lineage sorting, or to a small degree of hybridization.

Our broad assessment suggests that BOLD has a BIN that can be assigned (with widely varying degrees of confidence) to ~ 900 species of shallow-dwelling, reef-associated bony fishes from the GC. A list of sequences obtained in a particular area is obtained from BOLD by using a vector map in its search engine. The resulting list is from public projects (including all GenBank COI sequences), as well as whichever private projects the user has permission to access (often granted by an email request to the source of the sequence). In our case, we have been given access to all of the larger private projects in the region and barcodes for the vast majority (~ 90%) of sequence records in BOLD that could be evaluated in their respective BINs. The list of records from the geographic-area search on BOLD are individual sequences with metadata (including GenBank number if a sequence has one) and photographs of specimens (when available), together with a link to the BIN code to which it belongs. The species name originally submitted for

each is preserved, and the accuracy of the assignment can be assessed by examining the BIN to which it belongs, which has details on the various names applied to sequences in the BIN and by whom and where they were obtained. Accuracy assessments are critical, especially for more obscure species, since a “majority rules” decision is often inaccurate due to multiple identifications by inexperienced contributors, the tendency to repeat the species-level identification made by others as a shortcut, and the practice of assigning species-level names to submitted records that are from eggs, larvae, isolated tissue, or fish-market specimens. GenBank records are harvested by BOLD with whatever name is assigned in GenBank, often a preliminary one from submission, rather than the one later corrected or published in the subsequent literature.

## Results

### The island faunas

St. Croix: The checklist of Smith-Vaniz and Jelks (2014) included 544 species from 280 genera in 94 families. We obtained records of 41 species (belonging to 39 genera and 35 families; see Table 1) that were not included on that checklist, an increase of 7.5% in the number of species. Those new records included 19 deep-living species, six of them (11.1% of all deep species at St. Croix) resulting solely from observations by the JSL submersible (Nelson and Appeldoorn 1985; García-Sais et al. 2014) and an ROV (Remotely Operated Vehicle; Quattrini et al. 2017). It should be noted that almost all of that group belong to very deep taxa specifically excluded by Smith-Vaniz and Jelks (2014) from their list, which was focused primarily on shallower fishes. The remaining 22 species are shallow-water, reef-associated fishes. Ten of the latter group were photographed by AME and CJE (Table 1; Suppl. material 1: Plate S1). These additions include three species (*Eucinostomus melanopterus*, *Coryphopterus glaucofrenum* and *Opistognathus macrogathus*) that Smith-Vaniz and Jelks (2014) referred to but did not include in their checklist due to lack of confirmed records. Records of two mobulid rays consisted of identified photographs/videos provided by Mantatruster.org (<https://www.mantatruster.org/>) that were inspected by DRR. The list (Table 1, which includes source information) also includes records from museum collections that provide online data directly or indirectly through aggregators, which were included if consistent with the known geographic range of each of those species.

St. John-Thomas: Table 2 presents a list of species recorded from those islands together with the source(s) of each record (images, publications, DNA barcodes, or online museum records) and which species have a voucher image in the supplementary plates (Suppl. material 1: Plates S2–S18). In addition, for uncommon species (those encountered by AME, CJE, LR, or third-party photographers at three or fewer dive sites) the names of the sites at which those uncommon species were found are included, to aid future investigations. Dennis (2000) also included information on species that were collected using the ichthyocide Rotenone (see Table 2). Smith-Vaniz and Jelks (2014) list for St. Croix also included some species recorded at these St. John-Thomas as a result of collections using

**Table 1.** Species of fishes added to the St. Croix checklist of fishes of Smith-Vaniz and Jelks (2014).

Scientific name	Common name	Deep	Image plate	Literature source	Online source
<b>Antennariidae</b>					
<i>Fowlerichthys ocellatus</i> (Bloch & Schneider, 1801)	Ocellated Frogfish				TNHCI
<b>Bathygadidae</b>					
<i>Gadomus arcuatus</i> (Goode & Bean, 1886)	Doublethread Grenadier	yes		6	
<b>Blenniidae</b>					
<i>Hypleurochilus pseudoaequipinnis</i> Bath, 1994	Oyster Blenny		S1		
<b>Bramidae</b>					
<i>Eumegistus brevorti</i> (Poey, 1860)	Tropical Pomfret	yes			FIMNH
<b>Chaenopsidae</b>					
<i>Emblemariopsis leptocirris</i> Stephens, 1970	Fine-cirrus Blenny		S1		
<b>Chimaeridae</b>					
<i>Chimaera cubana</i> Howell Rivero, 1936	Cuban Chimaera	yes		1	
<b>Etmopteridae</b>					
<i>Etmopterus hillianus</i> (Poey, 1861)	Caribbean Lantern Shark	yes			FIMNH
<b>Exocoetidae</b>					
<i>Cheilopogon melanurus</i> (Valenciennes, 1847)	Atlantic Flyingfish				CF
<i>Cypselurus comatus</i> (Mitchill, 1815)	Clearwing Flyingfish				CF
<b>Gempylidae</b>					
<i>Lepidocybium flavobrunneum</i> (Smith, 1843)	Escolar	yes			NOAA
<i>Nesiarchus nasutus</i> Johnson, 1862	Black Gemfish	yes			NMNH
<b>Gerreidae</b>					
<i>Eucinostomus melanopterus</i> (Bleeker, 1863)	Flagfin Mojarra		S1	5,7*	
<b>Gobiesocidae</b>					
<i>Acyrtus lanthanum</i> Conway, Baldwin & White, 2014	Orange-spotted Clingfish				FIMNH
<b>Gobiidae</b>					
<i>Coryphopterus glaucofraenum</i> Gill, 1863	Bridled Goby		S1	2,5,7*	
<i>Coryphopterus kuna</i> Victor, 2007	Kuna Goby		S1		
<i>Oxyurichthys stigmaphius</i> (Mead & Böhlke, 1958)	Spotfin Goby		S1		NOAA
<b>Kyphosidae</b>					
<i>Kyphosus cinerascens</i> (Forsskål, 1775)	Topsail Seachub		S1		
<b>Macrouridae</b>					
<i>Nezumia aequalis</i> (Günther, 1878)	Atlantic Blacktip Grenadier	yes		6	
<b>Malakichthyidae</b>					
<i>Verilus pseudomicrolepis</i> (Schultz, 1940)	False-smallscale Bass	yes			CAS
<b>Mobulidae</b>					
<i>Mobula cf birostris</i>	Giant Manta			4	
<i>Mobula tarapacana</i> (Philippi, 1892)	Sicklefin Devil Ray			4	
<b>Muraenidae</b>					
<i>Gymnothorax nigromarginatus</i> (Girard, 1858)	Blackedge Moray				CAS
<b>Nemichthyidae</b>					
<i>Nemichthys curvirostris</i> (Strömman, 1896)	Spottedbelly Snipe Eel	yes		6	
<b>Neoscopelidae</b>					
<i>Neoscopelus microchir</i> Matsubara, 1943	Shortfin Blackchin	yes		6	
<b>Ophichthidae</b>					
<i>Myrophis punctatus</i> Lütken, 1852	Speckled Worm Eel				MCZ
<b>Ophidiidae</b>					
<i>Monomitopus agassizii</i> (Goode & Bean, 1896)	Threespine Cusk-eel	yes			MCZ

Scientific name	Common name	Deep	Image plate	Literature source	Online source
<b>Opistognathidae</b>					
<i>Opistognathus macrognathus</i> Poey, 1860	Banded Jawfish		S1	5,7*	
<b>Paralichthyidae</b>					
<i>Syacium micrurum</i> Ranzani, 1842	Channel Flounder		S1		
<b>Peristediidae</b>					
<i>Peristedion longispatha</i> Goode & Bean, 1886	Widehead Armored Searobin	yes			MCZ
<b>Pleuronectidae</b>					
<i>Poecilopsetta inermis</i> (Breder, 1927)	Unarmed Deepwater Dab	yes			CAS, NMNH
<b>Polymixiidae</b>					
<i>Polymixia nobilis</i> Lowe, 1836	Noble Beardfish	yes		3	
<b>Scombroptidae</b>					
<i>Scombroptus oculatus</i> (Poey, 1860)	Atlantic Scombroptus	yes			FIMNH
<b>Sparidae</b>					
<i>Calamus calamus</i> (Valenciennes, 1830)	Saucereye Porgy			5	
<b>Squalidae</b>					
<i>Cirrhitigaleus asper</i> (Merrett, 1973)	Roughskin Spiny Dogfish	yes			FIMNH
<b>Stomiidae</b>					
<i>Borostomia mononema</i> (Regan & Trewavas, 1929)	Sickle Snaggletooth	yes		8	
<b>Synagropidae</b>					
<i>Synagrops bellus</i> (Goode & Bean, 1896)	Blackmouth Bass	yes		6	
<b>Syngnathidae</b>					
<i>Hippocampus erectus</i> Perry, 1810	Lined Seahorse				NCSM
<b>Synodontidae</b>					
<i>Synodus foetens</i> (Linnaeus, 1766)	Inshore Lizardfish				ANSP
<i>Trachinocephalus myops</i> (Forster, 1801)	Snakefish		S1		
<b>Trachipteridae</b>					
<i>Zu cristatus</i> (Bonelli, 1820)	Scalloped Ribbonfish	yes		8	
<b>Tripterygiidae</b>					
<i>Enneanectes quadra</i> Victor, 2017	Squaretail Triplefin				FIMNH

**Notes:** Deep – restricted to depths below 40 m. Image Plate – see Suppl. material 1: Plate S1 for voucher images. Literature source – 1 Bunckley-Williams and Williams (2004); 2 Garcia-Sais et al. (2014); 3 Nelson and Appeldoorn (1985); 4 Mantatrust.org; 5 Pittman et al. (2008); 6 Quattrinni et al. (2017); 7 Smith-Vaniz and Jelks (2014) (asterisk indicates a species that was discussed by not included by those authors); 8 Clavijo et al. (1980). Online source - TNHCi (University of Texas at Austin, Biodiversity Center, Ichthyology collection; FIMNH (Florida Museum of Natural History); CF (Biological observations from the Dana Expedition Reports); NOAA (National Oceanographic and Atmospheric Administration); CAS (California Academy of Sciences); MCZ (Museum of Comparative Zoology); NMNH (National Museum of Natural History); NCSM (North Carolina State Museum of Natural Sciences); ANSP (Academy of Natural Sciences of Philadelphia). *Coryphopterus*: Smith-Vaniz et al. (2014) concluded that *C. tortugae*, but not *C. glaucofrenum*, was present at St. Croix. However, CJE and AME photographed both species at St. Croix, illustrated in Suppl. material 1: Plate S1.

that ichthyocide. Two ROV dives of Quattrinni et al. (2017) and four dives (including one to only 50 m depth on the PRP a little to the north of St. Thomas) by the JSL submersible at St. John-Thomas (Nelson and Appeldoorn 1985; Garcia-Sais 2005) yielded 75 species records. Of those 19 were of deep-living species, with 14 (28%) representing sole-source records of the 50 deep-living fishes currently known to occur at St. John-Thomas.

**Table 2.** Checklist of the fishes of St. John-Thomas islands.

Scientific name	Common name	Image Plate	Literature source	Online source	Uncommon (site code)	Ichthyocide	DNA
<b>Acanthuridae</b>							
<i>Acanthurus chirurgus</i> (Bloch, 1787)	Doctorfish	S2	2,4,8	1		1	
<i>Acanthurus coeruleus</i> Bloch & Schneider, 1801	Blue Tang	S2	2,4,5,8	1		1	YES
<i>Acanthurus tractus</i> Poey, 1860	Northern Ocean Surgeonfish	S2	2,4,5,8	1		1	YES
<b>Achiridae</b>							
<i>Gymnachirus nudus</i> Kaup, 1858	Flabby Sole	S2	2,11	1	SJ5, SJ18, SJ25		YES
<b>Aetobatidae</b>							
<i>Aetobatus narinari</i> (Euphrasen, 1790)	Spotted Eagle Ray	S2	2	1			
<b>Albulidae</b>							
<i>Albula goreensis</i> Valenciennes, 1847	Senegalese Bonefish				NOAA-BOLD		YES
<i>Albula vulpes</i> (Linnaeus, 1758)	Bonefish		2,4	1			YES
<b>Anguillidae</b>							
<i>Anguilla rostrata</i> (Lesueur, 1817)	American Eel		6	1			
<b>Antennariidae</b>							
<i>Antennarius multiocellatus</i> (Valenciennes, 1837)	Longlure Frogfish	S2	2	1		1	
<i>Antennarius paucinadiatus</i> Schultz, 1957	Dwarf Frogfish		2	1			
<i>Histrio histrio</i> (Linnaeus, 1758)	Sargassumfish	S2	12		O22		
<b>Apogonidae</b>							
<i>Apogon aurolineatus</i> (Mowbray, 1927)	Bridle Cardinalfish	S2	2,4	1			YES
<i>Apogon binotatus</i> (Poey, 1867)	Barred Cardinalfish	S2	2,4	1		1	
<i>Apogon lachneri</i> Böhlke, 1959	Whitestar Cardinalfish	S2	2,4	1	SJ2	1	
<i>Apogon maculatus</i> (Poey, 1860)	Flamefish	S2	2,4	1		1	YES
<i>Apogon phenax</i> Böhlke & Randall, 1968	Mimic Cardinalfish	S2	2,11			1	
<i>Apogon planifrons</i> Longley & Hildebrand, 1940	Pale Cardinalfish	S2	2	1		1	
<i>Apogon pseudomaculatus</i> Longley, 1932	Twospot Cardinalfish		2,4	1			
<i>Apogon quadrisquamatus</i> Longley, 1934	Sawcheek Cardinalfish	S2	2,4	1	SJ22, SJ25	1	YES
<i>Apogon robinsi</i> Böhlke & Randall, 1968	Roughlip Cardinalfish		2			-1	
<i>Apogon townsendi</i> (Breder, 1927)	Belted Cardinalfish	S2	2,4	1		1	YES
<i>Astrapogon puncticulatus</i> (Poey, 1867)	Blackfin Cardinalfish	S2	2	1			YES
<i>Astrapogon stellatus</i> (Cope, 1867)	Conchfish	S2	2,4	1	SJ5, SJ13		YES
<i>Paroncheilus affinis</i> (Poey, 1875)	Bigtooth Cardinalfish		2	1			

Scientific name	Common name	Image Plate	Literature source	Online source	Uncommon (site code)	Ichthyocide	DNA
<i>Phaeoptyx conklini</i> (Silvester, 1915)	Freckled Cardinalfish	S2	2	1		1	YES
<i>Phaeoptyx pigmentaria</i> (Poey, 1860)	Dusky Cardinalfish	S2	2	1		1	YES
<i>Phaeoptyx xenus</i> (Böhlke & Randall, 1968)	Sponge Cardinalfish	S2	2	1		1	YES
<i>Zapogon evermanni</i> (Jordan & Snyder, 1904)	Oddscales Cardinalfish	S2			SJ22		YES
<b>Atherinidae</b>							
<i>Atherina harringtonensis</i> Goode, 1877	Reef Silverside		2	1		1	YES
<i>Atherinomorus stipes</i> (Müller & Troschel, 1848)	Hardhead Silverside	S2	2,6	1		1	
<b>Aulostomidae</b>							
<i>Aulostomus maculatus</i> Valenciennes, 1841	Atlantic Trumpetfish	S2	2,4	1		1	
<b>Balistidae</b>							
<i>Balistes capricornis</i> Gmelin, 1789	Gray Triggerfish	S3	2				
<i>Balistes vetula</i> Linnaeus, 1758	Queen Triggerfish	S3	2,4,5,8	1		1	YES
<i>Canthidermis sufflamen</i> (Mitchill, 1815)	Ocean Triggerfish	S3	2	1	SJ33		
<i>Melichthys niger</i> (Bloch, 1786)	Black Durgon	S3	2,4	1	SJ33		
<i>Xanthichthys ringens</i> (Linnaeus, 1758)	Sargassum Triggerfish	S3	2,5	1	SJ33		
<b>Belontiidae</b>							
<i>Ablennes hians</i> (Valenciennes, 1846)	Barred Needlefish	S3					
<i>Platybelone argalus argalus</i> (Lesueur, 1821)	Keeltail Needlefish	S3	2	1		1	
<i>Strongylura timucu</i> (Walbaum, 1792)	Timucú		2,6	1			
<i>Tylosurus acus</i> (Lacepède, 1803)	Atlantic Agujón				FIMNH, MCZ		
<i>Tylosurus crocodilus</i> (Péron & Lesueur, 1821)	Houndfish	S3	2	1			
<b>Blenniidae</b>							
<i>Entomacrodus nigricans</i> Gill, 1859	Pearl Blenny	S3	2	1		1	YES
<i>Hyppleurochilus pseudoaequipinnis</i> Bath, 1994	Oyster Blenny	S3	2,11	1			YES
<i>Hyppleurochilus springeri</i> Randall, 1966	Orangespotted Blenny	S3	2	1			
<i>Hypsoblennius invemar</i> Smith-Vaniz & Acero P., 1980	Tessellated Blenny	S3	11	1	ST11		YES
<i>Ophioblennius macclurei</i> (Silvester, 1915)	Redlip Blenny	S3	2,4	1		1	YES
<i>Parablennius marmoratus</i> (Poey, 1876)	Seaweed Blenny	S3	2,4	1		1	YES
<i>Scartella cristata</i> (Linnaeus, 1758)	Molly Miller	S3	2,4	1		1	YES
<b>Bothidae</b>							
<i>Bothus lunatus</i> (Linnaeus, 1758)	Peacock Flounder	S3	2,4	1		1	
<i>Bothus maculiferus</i> (Poey, 1860)	Mottled Flounder	S3			SJ3, SJ5, SJ28		

Scientific name	Common name	Image Plate	Literature source	Online source	Uncommon (site code)	Ichthyocide	DNA
<i>Bothus ocellatus</i> (Agassiz, 1831)	Eyed Flounder	S3	2,4	1			
<i>Bothus robinsi</i> Topp & Hoff, 1972	Twospot Flounder		2,3				
<b>Bythitidae</b>							
<i>Calamopteryx goslinei</i> Böhlke & Cohen, 1966	Longarm Brotula		2				-1
<i>Grammonus claudei</i> (de la Torre y Huerta, 1930)	Reef-cave Brotula		2	1			-1
<i>Petrotyx sanguineus</i> (Meek & Hildebrand, 1928)	Redfin Brotula		2	1			-1
<b>Callionymidae</b>							
<i>Callionymus bairdi</i> Jordan, 1888	Lancer Dragonet	S3	2,4	1			YES
<i>Chalinops pauciradiatus</i> (Gill, 1865)	Spotted Dragonet	S3	2	1	SJ28, SJ3, SJ5		YES
<b>Carangidae</b>							
<i>Alectis ciliaris</i> (Bloch, 1787)	African Pompano	S4	2	1	ST1, SJ13		
<i>Caranx bartholomaei</i> Cuvier, 1833	Yellow Jack	S4	2,4	1			
<i>Caranx crysos</i> (Mitchill, 1815)	Blue Runner	S4	2,4	1			
<i>Caranx hippos</i> (Linnaeus, 1766)	Crevalle Jack	S4			SJ29		
<i>Caranx latus</i> Agassiz, 1831	Horse-eye Jack	S4	2,6	1			
<i>Caranx lugubris</i> Poey, 1860	Black Jack	S4	2,4,5,8	1	SJ33		
<i>Caranx ruber</i> (Bloch, 1793)	Bar Jack	S4	2,4,8	1			1
<i>Chloroscombrus chrysurus</i> (Linnaeus, 1766)	Atlantic Bumper		2				
<i>Decapterus macarellus</i> (Cuviers, 1833)	Mackerel Scad	S4	2	1			
<i>Decapterus punctatus</i> (Cuvier, 1829)	Round Scad	S4	2	1			
<i>Decapterus tabl</i> Berry, 1968	Redtail Scad	S4			SJ11		
<i>Elagatis bipinnulata</i> (Quoy & Gaimard, 1825)	Rainbow Runner	S4	2	1	SJ33		
<i>Oligoplites saurus saurus</i> (Bloch & Schneider, 1801)	Leatherjack		2	1			
<i>Selar crumenophthalmus</i> (Bloch, 1793)	Bigeye Scad	S4	2	1	SJ13		
<i>Selene brownii</i> (Cuvier, 1816)	Caribbean Moonfish		2	1			
<i>Selene vomer</i> (Linnaeus, 1758)	Lookdown				FIMNH		
<i>Seriola dumerili</i> (Risso, 1810)	Greater Amberjack		2,5				
<i>Seriola rivoliana</i> Valenciennes, 1833	Almaco Jack	S4	2	1	SJ16, SJ23		
<i>Trachinotus falcatus</i> (Linnaeus, 1758)	Permit	S4	2	1	SJ22, SJ23		
<i>Trachinotus goodei</i> Jordan & Evermann, 1896	Palometta	S4	2,4	1	SJ23, SJ15		
<b>Carcharhinidae</b>							
<i>Carcharhinus acronotus</i> (Poey, 1860)	Blacknose Shark	S4	1,2,10	1	SJ35, SJ27, ST7		
<i>Carcharhinus falciformis</i> (Müller & Henle, 1839)	Silky Shark	S4			1, O1		
<i>Carcharhinus galapagensis</i> (Snodgrass & Heller, 1905)	Galapagos Shark		2				
<i>Carcharhinus limbatus</i> (Müller & Henle, 1839)	Blacktip Shark		1,2	1			

Scientific name	Common name	Image Plate	Literature source	Online source	Uncommon (site code)	Ichthyocide	DNA
<i>Carcharhinus longimanus</i> (Poey, 1861)	Oceanic Whitetip Shark			NMNH			
<i>Carcharhinus perezii</i> (Poey, 1876)	Reef Shark	S4	2,10	1	SJ13		
<i>Carcharhinus plumbeus</i> (Nardo, 1827)	Sandbar Shark			ANSP			
<i>Negaprion brevirostris</i> (Poey, 1868)	Lemon Shark	S4	1,2,6,10	1	SJ12, O2		
<i>Rhizoprionodon porosus</i> (Poey, 1861)	Caribbean Sharpnose Shark		1,2,10	1			
<b>Centrophoridae</b>							
<i>Centrophorus uyato</i> (Rafinesque, 1810)	Little Gulper Shark			CAS			
<b>Centropomidae</b>							
<i>Centropomus ensiferus</i> Poey, 1860	Swordspine Snook		6	1			
<i>Centropomus undecimalis</i> (Bloch, 1792)	Common Snook	S4	2,6	1			
<b>Chaenopsidae</b>							
<i>Acanthemblemaria aspera</i> (Longley, 1927)	Roughhead Blenny	S5	2	1	ST3		YES
<i>Acanthemblemaria maria</i> Böhlke, 1961	Secretary Blenny	S5	4	1			YES
<i>Acanthemblemaria spinosa</i> Metzelaar, 1919	Spinyhead Blenny	S5	2,4	1		1	YES
<i>Chaenopsis limbaughii</i> Robins & Randall, 1965	Yellowface Pikeblenny	S5	2,4	1			YES
<i>Chaenopsis ocellata</i> Poey, 1865	Bluethroat Pikeblenny		2,4	1			
<i>Coralliozetus cardonae</i> Evermann & Marsh, 1899	Twinhorn Blenny	S5	11	1			YES
<i>Emblemaria pandionis</i> Evermann & Marsh, 1900	Sailfin Blenny	S5	2,4	1			YES
<i>Emblemaria vitta</i> Williams, 2002	Ribbon Blenny	S5	2,3	1	ST6	-1	YES
<i>Emblemariaopsis bahamensis</i> Stephens, 1961	Blackhead Blenny	S5		1			YES
<i>Emblemariaopsis carib</i> Victor, 2010	Carib Blenny		2	1		-1	YES
<i>Emblemariaopsis leptocirris</i> Stephens, 1970	Fine-cirrus Blenny	S5	2,11			-1	YES
<i>Emblemariaopsis ruetzleri</i> Tyler & Tyler, 1997	Ruetzler's Blenny			BOLD, NMNH			YES
<i>Lucayablennius zingaro</i> (Böhlke, 1957)	Arrow Blenny	S5			SJ18, SJ19		
<b>Chaetodontidae</b>							
<i>Chaetodon capistratus</i> Linnaeus, 1758	Foureye Butterflyfish	S5	2,4,5,8	1		1	YES
<i>Chaetodon ocellatus</i> Bloch, 1787	Spotfin Butterflyfish	S5	2,4	1			
<i>Chaetodon sedentarius</i> Poey, 1860	Reef Butterflyfish	S5	2,4,5,8	1			
<i>Chaetodon striatus</i> Linnaeus, 1758	Banded Butterflyfish	S5	2,4	1		1	
<i>Prognathodes aculeatus</i> (Poey, 1860)	Longsnout Butterflyfish	S5	2,5,8	1			
<i>Prognathodes guyanensis</i> (Durand, 1960)	Guyana Butterflyfish		2,5,8,11				

Scientific name	Common name	Image Plate	Literature source	Online source	Uncommon (site code)	Ichthyocide	DNA
<b>Chaunacidae</b>							
<i>Chaunax pxtus</i> Fowler, 1946	Uniform Gaper		5				
<i>Chaunax suttkusi</i> Caruso, 1989	Pale-cavity Gaper			CAS			
<b>Chlopsidae</b>							
<i>Chilorhinus suensonii</i> Lütken, 1852	Seagrass Eel		2	1			
<i>Kaupichthys hyporoides</i> (Strömman, 1896)	False Moray		2	1		-1	
<i>Kaupichthys nuchalis</i> Böhlke, 1967	Collared Eel		2,11	1			
<b>Chlorophthalmidae</b>							
<i>Parasudis truculenta</i> (Goode & Bean, 1896)	Longnose Greeneye		5				
<b>Cichlidae</b>							
<i>Oreochromis mossambicus</i> (Peters, 1852)	Mozambique Tilapia		6	1			
<b>Cirrhitidae</b>							
<i>Amblycirrhitus pinos</i> (Mowbray, 1927)	Redspotted Hawkfish	S5	2,4	1		1	
<b>Clupeidae</b>							
<i>Harengula clupeola</i> (Cuvier, 1829)	False Pilchard		2	1			YES
<i>Harengula humeralis</i> (Cuvier, 1829)	Redear Sardine	S5	2	1	SJ28, SJ13		YES
<i>Harengula jaguana</i> Poey, 1865	Scaled Sardine			FIMNH			
<i>Opisthonema oglinum</i> (Lesueur, 1818)	Atlantic Thread Herring			FIMNH			YES
<i>Sardinella aurita</i> Valenciennes, 1847	Spanish Sardine			FIMNH			
<b>Congridae</b>							
<i>Ariosoma balearicum</i> (Delaroche, 1809)	Bandtooth Conger		2				
<i>Conger triporceps</i> Kanazawa, 1958	Manytooth Conger		4	1			
<i>Heteroconger longissimus</i> Günther, 1870	Brown Garden Eel	S5	2,4	1			
<i>Xenomystax bidentatus</i> (Reid, 1940)	Rabbit Conger			NMNH			
<b>Coryphaenidae</b>							
<i>Coryphaena equiselis</i> Linnaeus, 1758	Pompano Dolphinfish			ROM			
<i>Coryphaena hippurus</i> Linnaeus, 1758	Dolphinfish	S5	2	1			
<b>Cynoglossidae</b>							
<i>Symphurus anawak</i> Robins & Randall, 1965	Caribbean Tonguefish		2	1		1	
<b>Dactylopteridae</b>							
<i>Dactylopterus volitans</i> (Linnaeus, 1758)	Flying Gurnard	S5	4	1			YES
<b>Dactyloscopidae</b>							
<i>Dactyloscopus comptus</i> Dawson, 1982	Ornamented Stargazer		2,11	1			
<i>Dactyloscopus crossotus</i> Starks, 1913	Bigeye Stargazer			AMNH			
<i>Dactyloscopus poeyi</i> Gill, 1861	Shortchin Stargazer			FIMNH			
<i>Dactyloscopus tridigitatus</i> Gill, 1859	Sand Stargazer	S5	2	1		1	
<i>Gillellus greylae</i> Kanazawa, 1952	Arrow Stargazer		2			-1	
<i>Gillellus uranidea</i> Böhlke, 1968	Warteye Stargazer		2			-1	YES

Scientific name	Common name	Image Plate	Literature source	Online source	Uncommon (site code)	Ichthyocide	DNA
<i>Platygillellus rubrocinctus</i> (Longley, 1934)	Saddle Stargazer						
<b>Dasyatidae</b>							
<i>Hypanus americanus</i> (Hildebrand & Schroeder, 1928)	Southern Stingray	S5	1,2,4,10	1			
<b>Diodontidae</b>							
<i>Chilomycterus antennatus</i> (Cuvier, 1816)	Bridled Burrfish	S5	2,4	1	SJ18		
<i>Chilomycterus antillarum</i> Jordan & Rutter, 1897	Web Burrfish		2	1			
<i>Diodon holocanthus</i> Linnaeus, 1758	Balloonfish	S5	2,4	1	SJ11, SJ13	-1	
<i>Diodon hystrix</i> Linnaeus, 1758	Porcupinefish	S5	2,4	1		1	
<b>Echeneidae</b>							
<i>Echeneis naucrates</i> Linnaeus, 1758	Sharksucker	S6	2,4	1	SJ19, SJ23		YES
<i>Echeneis neucratoides</i> Zuiew, 1789	Whitefin Sharksucker	S6		1			
<i>Remora remora</i> (Linnaeus, 1758)	The Remora	S6		1	O3		YES
<b>Eleotridae</b>							
<i>Dormitorator maculatus</i> (Bloch, 1792)	Fat Sleeper	S6	6	1	SJ10		
<i>Eleotris perniger</i> (Cope, 1871)	Smallscaled Spinycheek Sleeper	S6	6	1	SJ10		
<i>Erotelis smaragdus</i> (Valenciennes, 1837)	Emerald Sleeper		6	1			
<i>Gobiomorus dormitor</i> Lacepède, 1800	Bigmouth Sleeper	S6		1			
<b>Elopidae</b>							
<i>Elops smithi</i> McBride, Rocha, Ruiz-Carus & Bowen, 2010	Malacho		2,6				YES
<b>Engraulidae</b>							
<i>Anchoa lyolepis</i> (Evermann & Marsh, 1900)	Dusky Anchovy		2	1			YES
<b>Ephippidae</b>							
<i>Chaetodipterus faber</i> (Broussonet, 1782)	Atlantic Spadefish	S6	2,4	1	SJ18, ST2		
<b>Epigonidae</b>							
<i>Epigonus pandionis</i> (Goode & Bean, 1881)	Caudal-ring Deepwater Cardinalfish			CAS			
<b>Exocoetidae</b>							
<i>Cheilopogon exsiliens</i> (Linnaeus, 1771)	Bandwing Flyingfish		2	1			
<i>Exocoetus obtusirostris</i> Günther, 1866	Oceanic Two-wing Flyingfish			MCZ			
<i>Hirundichthys affinis</i> (Günther, 1866)	Fourwing Flyingfish		2				
<i>Hirundichthys speculiger</i> (Valenciennes, 1847)	Mirrorwing Flyingfish		2	1			
<i>Prognichthys occidentalis</i> Parin, 1999	Bluntnose Flyingfish	S6					YES
<b>Fistulariidae</b>							
<i>Fistularia tabacaria</i> Linnaeus, 1758	Bluespotted Cornetfish	S6	2		O4		

Scientific name	Common name	Image Plate	Literature source	Online source	Uncommon (site code)	Ichthyocide	DNA
<b>Galeocerdonidae</b>							
<i>Galeocerdo cuvier</i> (Peron & Lesueur, 1822)	Tiger Shark		10				
<b>Gempylidae</b>							
<i>Epinnula magistralis</i> Poey, 1854	Domine		5	1			
<b>Gerreidae</b>							
<i>Eucinostomus argenteus</i> Baird & Girard, 1855	Spotfin Mojarra		2	1			YES
<i>Eucinostomus gula</i> (Quoy & Gaimard, 1824)	Silver Jenny	S6	4	1	SJ18, SJ13, SJ3		
<i>Eucinostomus harengulus</i> Goode & Bean, 1879	Tidewater Mojarra	S6		1	SJ28		
<i>Eucinostomus havana</i> (Nichols, 1912)	Bigeye Mojarra			FIMNH			
<i>Eucinostomus jonesii</i> (Günther, 1879)	Slender Mojarra	S6	4,6		SJ28		
<i>Eucinostomus lefroyi</i> (Goode, 1874)	Mottled Mojarra	S6			SJ28, SJ21		
<i>Eucinostomus melanopterus</i> (Bleeker, 1863)	Flagfin Mojarra	S6	4	1	SJ28		
<i>Eugerres brasiliensis</i> (Cuvier, 1830)	Brazilian Mojarra		6,11	1			
<i>Gerres cinereus</i> (Walbaum, 1792)	Yellowfin Mojarra	S6	2,4,6	1			
<b>Ginglymostomatidae</b>							
<i>Ginglymostoma cirratum</i> (Bonnaterre, 1788)	Nurse Shark	S6	1,2,4,10	1			
<b>Gobiesocidae</b>							
<i>Acyrtops amplicirrus</i> Briggs, 1955	Flarenostril Clingfish		2				
<i>Acyrtops beryllinus</i> (Hildebrand & Ginsburg, 1927)	Emerald Clingfish		2	1			
<i>Acyrtus artius</i> Briggs, 1955	Papillate Clingfish		2				
<i>Acyrtus rubiginosus</i> (Poey, 1868)	Red Clingfish	S6		1	SJ23, SJ13, SJ5		YES
<i>Arcos nudus</i> (Linnaeus, 1758)	Padded Clingfish	S6	2	1	SJ23	1	
<i>Gobiesox nigripinnis</i> (Peters, 1859)	Dark-finned Clingfish	S6	2	1	SJ29		
<i>Gobiesox punctulatus</i> (Poey, 1876)	Stippled Clingfish	S6	2	1		1	YES
<i>Tomicodon cryptus</i> Williams & Tyler, 2003	Cryptic Clingfish	S6					YES
<i>Tomicodon fasciatus</i> (Peters, 1859)	Barred Clingfish		2	1		1	
<i>Tomicodon leurodiscus</i> Williams & Tyler, 2003	Smooth-suckered Clingfish		11	1			
<i>Tomicodon reitzae</i> Briggs, 2001	Accidental Clingfish	S6					YES
<i>Tomicodon rhabdodus</i> Smith-Vaniz, 1969	Antillean Clingfish	S6			O24		
<i>Tomicodon rupestris</i> (Poey, 1860)	Barred Clingfish		11	1			
<b>Gobiidae</b>							
<i>Awaous banana</i> (Valenciennes, 1837)	River Goby	S7		1	SJ10		
<i>Barbulifer ceuthoecus</i> (Jordan & Gilbert, 1884)	Bearded Goby		2	1			YES
<i>Bathygobius antillensis</i> Tornabene, Baldwin & Pezold, 2010	Antilles Frillfin	S7			SJ36		YES

Scientific name	Common name	Image Plate	Literature source	Online source	Uncommon (site code)	Ichthyocide	DNA
<i>Bathygobius curacao</i> (Metzelaar, 1919)	Notchtongue Goby		11	1			YES
<i>Bathygobius lacertus</i> (Poey, 1860)	Checkerboard Frillfin			FIMNH			YES
<i>Bathygobius mystacium</i> Ginsburg, 1947	Island Frillfin	S7			SJ21, SJ19		YES
<i>Bathygobius soporator</i> (Valenciennes, 1837)	Frillfin Goby		2,6,11	1			YES
<i>Bollmannia boqueronensis</i> Evermann & Marsh, 1899	White-eye Goby	S7	4		SJ19		YES
<i>Cerdale floridana</i> Longley, 1934	Pugjaw Wormfish	S7	2	1	SJ23	1	
<i>Coryphopterus alloides</i> Böhlke & Robins, 1960	Barfin Goby		2	1		-1	
<i>Coryphopterus dicrus</i> Böhlke & Robins, 1960	Colon Goby	S7	2,4	1		1	YES
<i>Coryphopterus eidolon</i> Böhlke & Robins, 1960	Pallid Goby	S7	2,4	1		1	YES
<i>Coryphopterus glaucofraenum</i> Gill, 1863	Bridled Goby	S7	2,4	1		1	YES
<i>Coryphopterus hyalinus</i> Böhlke & Robins, 1962	Glass Goby	S7	2	1		-1	YES
<i>Coryphopterus kuna</i> Victor, 2007	Kuna Goby	S7			SJ5, SJ12		
<i>Coryphopterus lipernes</i> Böhlke & Robins, 1962	Peppermint Goby	S7	2,4	1	ST6		YES
<i>Coryphopterus personatus</i> (Jordan & Thompson, 1905)	Masked Goby	S7	2	1		1	YES
<i>Coryphopterus thrix</i> Böhlke & Robins, 1960	Bartail Goby		2	1		1	YES
<i>Coryphopterus tortugae</i> (Jordan, 1904)	Sand Goby	S7		1			YES
<i>Coryphopterus venezuelae</i> Cervigón, 1966	Venezuela Goby	S7		1			YES
<i>Ctenogobius boleosoma</i> (Jordan & Gilbert, 1882)	Darter Goby	S7	6	1	SJ28		YES
<i>Ctenogobius saepepallens</i> (Gilbert & Randall, 1968)	Dash Goby	S7	2,4	1			YES
<i>Ctenogobius smaragdus</i> (Valenciennes, 1837)	Emerald Goby		11				
<i>Ctenogobius stigmaturus</i> (Goode & Bean, 1882)	Spottail Goby	S7			SJ28		YES
<i>Elacatinus chancei</i> (Beebe & Hollister, 1933)	Shortstripe Goby	S7	2,4	1			YES
<i>Elacatinus evelynae</i> (Böhlke & Robins, 1968)	Sharknose Goby	S7	2,4	1		1	YES
<i>Elacatinus prochilos</i> (Böhlke & Robins, 1968)	Broadstripe Goby	S7		1			YES
<i>Evorthodus lyricus</i> (Girard, 1858)	Lyre Goby		6	1			
<i>Ginsburgellus novemlineatus</i> (Fowler, 1950)	Nineline Goby	S7		1	SJ23, SJ5		YES
<i>Gnatholepis thompsoni</i> Jordan, 1904	Goldspot Goby	S7	2,4	1		1	YES
<i>Gobionellus oceanicus</i> (Pallas, 1770)	Highfin Goby	S7		1	SJ28		
<i>Gobiosoma grosvenori</i> (Robins, 1964)	Rockcut Goby		4	1			

Scientific name	Common name	Image Plate	Literature source	Online source	Uncommon (site code)	Ichthyocide	DNA
<i>Lophogobius cyprinoides</i> (Pallas, 1770)	Crested Goby	S8	6	1	SJ28		
<i>Lythrypnus elason</i> Böhlke & Robins, 1960	Dwarf Goby	S8	2	AMNH	ST5	1	YES
<i>Lythrypnus minimus</i> Garzón-Ferreira & Acero P, 1988	Pygmy Goby	S8					YES
<i>Lythrypnus nesiotus</i> Böhlke & Robins, 1960	Island Goby	S8	2	1	SJ34	1	YES
<i>Lythrypnus spilus</i> Böhlke & Robins, 1960	Bluegold Goby	S8			ST3		
<i>Microgobius carri</i> Fowler, 1945	Seminole Goby	S8	2,4	1	SJ19, SJ25	1	YES
<i>Microgobius signatus</i> Poey, 1876	Signal Goby	S8		1	SJ28, SJ22, SJ3		YES
<i>Nes longus</i> (Nichols, 1914)	Orangespotted Goby	S8	4	1			YES
<i>Oxyurichthys stigmaliops</i> (Mead & Böhlke, 1958)	Spotfin Goby	S8	4	1	SJ5, SJ19, SJ28		
<i>Palatogobius paradoxus</i> Gilbert, 1971	Mauve Goby		2,11	1			
<i>Priolepis hipoliti</i> (Metzelaar, 1922)	Rusty Goby	S8	2,4	1		1	
<i>Pilotris celsa</i> Böhlke, 1963	Highspine Goby		2	1			
<i>Ptereleotris helenae</i> (Randall, 1968)	Hovering Dartfish	S8	2,4	1			
<i>Risor ruber</i> (Rosén, 1911)	Tusked Goby	S8	2	1		1	YES
<i>Sicydium plumieri</i> (Bloch, 1786)	Sirajo Goby	S8	6	1	SJ10		YES
<i>Sicydium punctatum</i> Perugia, 1896	Spotted Algae-eating Goby	S8		1	SJ10		YES
<i>Tigriogobius dilepis</i> (Robins & Böhlke, 1964)	Orangesided Goby		4	1			
<i>Tigriogobius multifasciatus</i> (Steindachner, 1876)	Greenbanded Goby	S8	2	1			YES
<i>Tigriogobius pallens</i> (Ginsburg, 1939)	Semiscaled Goby	S8			SJ23		
<i>Tigriogobius saucrus</i> (Robins, 1960)	Leopard Goby	S8	2	1		1	YES
<b>Grammatidae</b>							
<i>Gramma linki</i> Starck & Colin, 1978	Yellowcheek Basslet		2,5,8			1	
<i>Gramma loreto</i> Poey, 1868	Fairy Basslet	S8	2,4	1			YES
<b>Haemulidae</b>							
<i>Anisotremus surinamensis</i> (Bloch, 1791)	Black Margate	S9	2,4,5,8	1		1	YES
<i>Anisotremus virginicus</i> (Linnaeus, 1758)	Porkfish	S9	2,5,6,8	1			YES
<i>Brachygenys chrysargyrea</i> (Günther, 1859)	Smallmouth Grunt	S9	2,4	1		1	YES
<i>Emmelichthys atlanticus</i> Schultz, 1945	Bonnetmouth	S9	2		ST8		
<i>Haemulon album</i> Cuvier, 1830	Margate	S9	2,4	1	SJ7		
<i>Haemulon aurolineatum</i> Cuvier, 1830	Tomtate	S9	2,4,5,8	1		1	YES
<i>Haemulon carbonarium</i> Poey, 1860	Caesar Grunt	S9	2,4	1		1	
<i>Haemulon flavolineatum</i> (Desmarest, 1823)	French Grunt	S9	2,4,5,8	1		1	YES
<i>Haemulon macrostoma</i> Günther, 1859	Spanish Grunt	S9	2,4	1		1	
<i>Haemulon melanurum</i> (Linnaeus, 1758)	Cottonwick	S9	2	1	O5		YES

Scientific name	Common name	Image Plate	Literature source	Online source	Uncommon (site code)	Ichthyocide	DNA
<i>Haemulon parra</i> (Desmarest, 1823)	Sailors Choice	S9	2,4	1	SJ1, SJ21		YES
<i>Haemulon plumieri</i> (Lacepède, 1801)	White Grunt	S9	2,4	1		1	YES
<i>Haemulon sciurus</i> (Shaw, 1803)	Bluestriped Grunt	S9	2,4,5	1		1	YES
<i>Haemulon striatum</i> (Linnaeus, 1758)	Striped Grunt		2,4	1			YES
<i>Haemulon vittatum</i> (Poey, 1860)	Boga	S9	2,4	1	ST6, ST8, ST2	1	
<b>Hemiramphidae</b>							
<i>Euleptorhamphus velox</i> Poey, 1868	Flying Halfbeak			MCZ			
<i>Hemiramphus balao</i> Lesueur, 1821	Balao			MCZ			
<i>Hemiramphus brasiliensis</i> (Linnaeus, 1758)	Ballyhoo	S9	2	1			
<i>Hyporhamphus unifasciatus</i> (Ranzani, 1841)	Atlantic Silverstripe Halfbeak		2	1			
<b>Hexanchidae</b>							
<i>Heptranchias perlo</i> (Bonnaterre, 1788)	Sharpnose Sevengill Shark			FIMNH			
<i>Hexanchus vitulus</i> Springer & Waller, 1969	Atlantic Sixgill Shark			FIMNH			
<b>Holocentridae</b>							
<i>Holocentrus adscensionis</i> (Osbeck, 1765)	Squirrelfish	S9	2,4,5,8	1		1	YES
<i>Holocentrus rufus</i> (Walbaum, 1792)	Longspine Squirrelfish	S9	2,4,5,8	1		1	
<i>Myripristis jacobus</i> Cuvier, 1829	Blackbar Soldierfish	S9	2,4,5,8	1		1	
<i>Neoniphon coruscum</i> (Poey, 1860)	Reef Squirrelfish	S9	2,4,5,8	1		1	YES
<i>Neoniphon marianus</i> (Cuvier, 1829)	Longjaw Squirrelfish	S9	2,4,5,8	1		1	
<i>Neoniphon vexillarium</i> (Poey, 1860)	Dusky Squirrelfish	S9	2,4	1		1	
<i>Ostichthys trachypoma</i> (Günther, 1859)	Bigeye Soldierfish		2,5,8	1			
<i>Plectrypops retrospinis</i> (Guichenot, 1853)	Cardinal Soldierfish	S9	2,5,8	1	SJ9, SJ22, ST3	1	
<i>Sargocentron bullisi</i> (Woods, 1955)	Deepwater Squirrelfish		2,11	1			
<b>Ipnopidae</b>							
<i>Bathypterois bigelowi</i> Mead, 1958	Spottail Tripodfish			CAS			
<i>Bathypterois phenax</i> Parr, 1928	Blackfin Spiderfish		9				
<i>Bathypterois viridensis</i> (Roule, 1916)	Twobanded Tripodfish		9				
<i>Ipnops murrayi</i> Günther, 1878	Grیدهye Fish		9				
<b>Istiophoridae</b>							
<i>Istiophorus platypterus</i> (Shaw, 1792)	Sailfish	S9	2				
<i>Kajikia albigata</i> (Poey, 1860)	White Marlin	S9	2				
<i>Makaira nigricans</i> Lacepède, 1802	Blue Marlin		2				YES
<i>Tetrapturus pfluegeri</i> Robins & de Sylva, 1963	Longbill Spearfish		2				

Scientific name	Common name	Image Plate	Literature source	Online source	Uncommon (site code)	Ichthyocide	DNA
<b>Kyphosidae</b>							
<i>Kyphosus cinerascens</i> (Forsskål, 1775)	Topsail Seachub	S10					
<i>Kyphosus sectatrix</i> (Linnaeus, 1758)	Bermuda Chub	S10	2,4	1			
<i>Kyphosus vaigiensis</i> (Quoy & Gaimard, 1825)	Yellow Chub	S10		1			
<b>Labridae</b>							
<b>Labrinae</b>							
<i>Bodianus rufus</i> (Linnaeus, 1758)	Spanish Hogfish	S10	2,4,5,8	1			YES
<i>Clepticus parvae</i> (Bloch & Schneider, 1801)	Creole Wrasse	S10	2,4,5,8	1			YES
<i>Decodon puellaris</i> (Poey, 1860)	Red Hogfish		2	1			
<i>Doratonotus megalepis</i> Günther, 1862	Dwarf Wrasse		2	1			
<i>Halichoeres bivittatus</i> (Bloch, 1791)	Slippery Dick	S10	2,4	1		1	YES
<i>Halichoeres caudalis</i> (Poey, 1860)	Painted Wrasse			NOAA			
<i>Halichoeres cyanocephalus</i> (Bloch, 1791)	Yellowcheek Wrasse		2	1			
<i>Halichoeres garnoti</i> (Valenciennes, 1839)	Yellowhead Wrasse	S10	2,4	1		1	YES
<i>Halichoeres maculipinna</i> (Müller & Troschel, 1848)	Clown Wrasse	S10	2,4	1		1	
<i>Halichoeres pictus</i> (Poey, 1860)	Rainbow Wrasse	S10	2,4	1		1	
<i>Halichoeres poeyi</i> (Steindachner, 1867)	Blackear Wrasse	S10	2,4	1			
<i>Halichoeres radiatus</i> (Linnaeus, 1758)	Puddingwife	S10	2,4	1		1	YES
<i>Lachnolaimus maximus</i> (Walbaum, 1792)	Hogfish	S10	2,4,5,8	1			
<i>Thalassoma bifasciatum</i> (Bloch, 1791)	Bluehead	S10	2,4	1		1	
<i>Xyrichtys martinicensis</i> Valenciennes, 1840	Rosy Razorfish	S10	2,4	1			
<i>Xyrichtys novacula</i> (Linnaeus, 1758)	Pearly Razorfish	S10	2,4	1			YES
<i>Xyrichtys splendens</i> Castelnau, 1855	Green Razorfish	S10	2,4	1			
<b>Scarinae</b>							
<i>Cryptotomus roseus</i> Cope, 1871	Bluelip Parrotfish	S10	2,4	1			YES
<i>Scarus coelestinus</i> Valenciennes, 1840	Midnight Parrotfish	S10	2	1	O6	1	
<i>Scarus coeruleus</i> (Edwards, 1771)	Blue Parrotfish		2,4	1		1	
<i>Scarus guacamaia</i> Cuvier, 1829	Rainbow Parrotfish	S10	2,4	1	SJ28, SJ33, O2		
<i>Scarus iseri</i> (Bloch, 1789)	Striped Parrotfish	S10	2,4	1		1	YES
<i>Scarus taeniopterus</i> Lesson, 1829	Princess Parrotfish	S10	2,4,5,8	1		1	YES
<i>Scarus vetula</i> Bloch & Schneider, 1801	Queen Parrotfish	S10	2,4	1		1	YES
<i>Sparisoma atomarium</i> (Poey, 1861)	Greenblotch Parrotfish	S11	2,4	1			
<i>Sparisoma aurofrenatum</i> (Valenciennes, 1840)	Redband Parrotfish	S11	2,4,5,8	1		1	YES
<i>Sparisoma chrysopterygum</i> (Bloch & Schneider, 1801)	Redtail Parrotfish	S11	2,4	1		1	YES
<i>Sparisoma radians</i> (Valenciennes, 1840)	Bucktooth Parrotfish	S11	2,4	1		1	YES
<i>Sparisoma rubripinne</i> (Valenciennes, 1840)	Yellowtail Parrotfish	S11	2,4	1		1	YES

Scientific name	Common name	Image Plate	Literature source	Online source	Uncommon (site code)	Ichthyocide	DNA
<i>Sparisoma viride</i> (Bonnaterre, 1788)	Stoplight Parrotfish	S11	2,4,5,8	1		1	YES
<b>Labrisomidae</b>							
<i>Brockius albigenys</i> Beebe & Tee-Van, 1928	Whitecheek Blenny		DNA		Berry Bay, St. John		YES
<i>Brockius nigricinctus</i> (Howell Rivero, 1936)	Spotcheek Blenny	S11		1	SJ21		YES
<i>Gobioclinus bucciferus</i> (Poey, 1868)	Puffcheek Blenny	S11	2	1			YES
<i>Gobioclinus filamentosus</i> (Springer, 1960)	Quillfin Blenny	S11	3,11	1	O7		YES
<i>Gobioclinus gobio</i> (Valenciennes, 1836)	Palehead Blenny	S11	2	1		1	YES
<i>Gobioclinus guppyi</i> (Norman, 1922)	Mimic Blenny	S11	2	1		-1	YES
<i>Gobioclinus bairdianus</i> (Beebe & Tee-Van, 1928)	Longfin Blenny	S11	2	1	SJ12	1	YES
<i>Labrisomus cricota</i> Sazima, Gasparini & Moura, 2002	Mock Blenny	S11			SJ10		
<i>Labrisomus nuchipinnis</i> (Quoy & Gaimard, 1824)	Hairy Blenny	S11	2,4	1		1	YES
<i>Malacoctenus aurolineatus</i> Smith, 1957	Goldline Blenny	S11	2,4	1		1	YES
<i>Malacoctenus boehlkei</i> Springer, 1959	Diamond Blenny	S11	2,4	1		1	YES
<i>Malacoctenus erdmani</i> Smith, 1957	Imitator Blenny	S11			SJ23		YES
<i>Malacoctenus gilli</i> (Steindachner, 1867)	Dusky Blenny	S11	2,4	1			YES
<i>Malacoctenus macropus</i> (Poey, 1868)	Rosy Blenny	S11	2,4	1			YES
<i>Malacoctenus triangulatus</i> Springer, 1959	Saddled Blenny	S11	2,4	1		1	YES
<i>Malacoctenus versicolor</i> (Poey, 1876)	Barfin Blenny	S11	2,4	1	SJ23, SJ12		YES
<i>Nemaclinus atelestos</i> Böhlke & Springer, 1975	Threadfin Blenny		2,11	1			
<i>Paraclinus barbatus</i> Springer, 1955	Goatee Blenny		2,11				
<i>Paraclinus cingulatus</i> (Evermann & Marsh, 1899)	Coral Blenny		2				
<i>Paraclinus fasciatus</i> (Steindachner, 1876)	Banded Blenny	S11	2		SJ12		
<i>Paraclinus nigripinnis</i> (Steindachner, 1867)	Blackfin Blenny	S11	2	1	SJ12		YES
<i>Starksia culebrae</i> (Evermann & Marsh, 1899)	Culebra Blenny	S11	2	1	ST2, SJ13		YES
<i>Starksia hassi</i> Klausewitz, 1958	Ringed Blenny	S11	2,11	1	SJ24	1	
<i>Starksia lepicoelia</i> Böhlke & Springer, 1961	Blackcheek Blenny		2	1		1	
<i>Starksia nanodes</i> Böhlke & Springer, 1961	Dwarf Blenny		2	1			
<i>Starksia williamsi</i> Baldwin & Castillo, 2011	Williams's Blenny	S11			SJ2, SJ13		YES
<i>Stathmonotus gymnodermis</i> Springer, 1955	Naked Blenny		2	1			
<i>Stathmonotus stabli</i> (Evermann & Marsh, 1899)	Southern Eelgrass Blenny		2	1			

Scientific name	Common name	Image Plate	Literature source	Online source	Uncommon (site code)	Ichthyocide	DNA
<b>Latilidae</b>							
<i>Caulolatilus cyanops</i> Poey, 1866	Blackline Tilefish		2				
<b>Lobotidae</b>							
<i>Lobotes surinamensis</i> (Bloch, 1790)	Atlantic Tripletail	S11	2	1	O18		
<b>Lutjanidae</b>							
<i>Apsilus dentatus</i> Guichenot, 1853	Black Snapper		2				
<i>Etelis oculatus</i> (Valenciennes, 1828)	Queen Snapper	S12	2,5,8				YES
<i>Lutjanus analis</i> (Cuvier, 1828)	Mutton Snapper	S12	2,4,5,8	1			YES
<i>Lutjanus apodus</i> (Walbaum, 1792)	Schoolmaster	S12	2,4,5,6,8	1		1	YES
<i>Lutjanus buccanella</i> (Cuvier, 1828)	Blackfin Snapper	S12	2,5,8	1			YES
<i>Lutjanus cyanopterus</i> (Cuvier, 1828)	Cubera Snapper	S12	2,4	1			YES
<i>Lutjanus griseus</i> (Linnaeus, 1758)	Gray Snapper	S12	2,4,6	1		1	
<i>Lutjanus jocu</i> (Bloch & Schneider, 1801)	Dog Snapper	S12	2,4,5,8	1			YES
<i>Lutjanus mahogoni</i> (Cuvier, 1828)	Mahogany Snapper	S12	2,4	1		1	YES
<i>Lutjanus purpureus</i> (Poey, 1866)	Caribbean Red Snapper		2	1			
<i>Lutjanus synagris</i> (Linnaeus, 1758)	Lane Snapper	S12	2,4	1			YES
<i>Lutjanus vivanus</i> (Cuvier, 1828)	Silk Snapper	S12	2,5,8	1			YES
<i>Ocyurus chrysurus</i> (Bloch, 1791)	Yellowtail Snapper	S12	2,4	1		1	YES
<i>Pristipomoides macrophthalmus</i> (Müller & Troschel, 1848)	Cardinal Snapper		2				
<i>Rhomboplites aurorubens</i> (Cuvier, 1829)	Vermilion Snapper	S12	2		SJ20		
<b>Malacanthidae</b>							
<i>Malacanthus plumieri</i> (Bloch, 1786)	Sand Tilefish	S12	2,4,5,8	1		1	
<b>Megalopidae</b>							
<i>Megalops atlanticus</i> Valenciennes, 1847	Tarpon	S12	2,6	1			
<b>Mobulidae</b>							
<i>Mobula birostris</i> (Walbaum, 1792)	Giant Manta	S12	2				
<i>Mobula cf. birostris</i>	Caribbean Manta	S12				SJ12	
<b>Monacanthidae</b>							
<i>Aluterus monoceros</i> (Linnaeus, 1758)	Unicorn Filefish	S12				O23	
<i>Aluterus schoepfii</i> (Walbaum, 1792)	Orange Filefish	S12		1		SJ34	
<i>Aluterus scriptus</i> (Osbeck, 1765)	Scrawled Filefish	S12	4	1			
<i>Cantherbines macrocerus</i> (Hollard, 1853)	Whitespotted Filefish	S12	2	1			YES
<i>Cantherbines pullus</i> (Ranzani, 1842)	Orangespotted Filefish	S12	2,4	1		1	
<i>Monacanthus ciliatus</i> (Mitchill, 1818)	Fringed Filefish	S12	2,4	1			YES
<i>Monacanthus tuckeri</i> Bean, 1906	Slender Filefish	S12	2,4	1			
<i>Stephanolepis hispidus</i> (Linnaeus, 1766)	Planehead Filefish				FIMNH		
<i>Stephanolepis setifer</i> (Bennett, 1831)	Pygmy Filefish		2				
<b>Moringuidae</b>							
<i>Moringua edwardsi</i> (Jordan & Bollman, 1889)	Spaghetti Eel		2	1			-1

Scientific name	Common name	Image Plate	Literature source	Online source	Uncommon (site code)	Ichthyocide	DNA
<b>Mugilidae</b>							
<i>Dajaus monticola</i> (Bancroft, 1834)	Mountain Mullet	S13	6	1	SJ10		
<i>Mugil curema</i> Valenciennes, 1836	White Mullet	S13	2,6		SJ21		YES
<i>Mugil rubrioculus</i> Harrison, Nirchio, Oliveira, Ron & Gaviria, 2007	Redeye Mullet	S13	DNA				YES
<i>Mugil trichodon</i> Poey, 1875	Fantail Mullet			ROM			
<b>Mullidae</b>							
<i>Mulloidichthys martinicus</i> (Cuvier, 1829)	Yellow Goatfish	S13	2,4,6	1		1	YES
<i>Pseudupeneus maculatus</i> (Bloch, 1793)	Spotted Goatfish	S13	2,4,5,8	1		1	
<b>Muraenidae</b>							
<i>Echidna catenata</i> (Bloch, 1795)	Chain Moray	S13	2,4	1	SJ21, SJ10	1	
<i>Enchelycore carychroa</i> Böhlke & Böhlke, 1976	Chestnut Moray	S13	2	1	SJ5	1	
<i>Enchelycore nigricans</i> (Bonnaterre, 1788)	Viper Moray	S13	2	1	SJ9	1	
<i>Gymnothorax conspersus</i> Poey, 1867	Saddled Moray			ANSP			
<i>Gymnothorax funebris</i> Ranzani, 1839	Green Moray	S13	2,4	1		1	YES
<i>Gymnothorax maderensis</i> (Johnson, 1862)	Sharktooth Moray		2	1			
<i>Gymnothorax miliaris</i> (Kaup, 1856)	Goldentail Moray	S13	2,4	1		1	
<i>Gymnothorax moringa</i> (Cuvier, 1829)	Spotted Moray	S13	2,4			1	YES
<i>Gymnothorax vicinus</i> (Castelnau, 1855)	Purplemouth Moray	S13	2,4	1		1	
<i>Uropterygius macularius</i> (Lesueur, 1825)	Marbled Moray	S13	2	1		1	
<b>Myctophidae</b>							
<i>Centrobranchus nigroocellatus</i> (Günther, 1873)	Roundnose Lanternfish			ROM			
<b>Neoscopelidae</b>							
<i>Neoscopelus macrolepidotus</i> Johnson, 1863	Largescale Blackchin			CAS			
<b>Nomeidae</b>							
<i>Penes cyanophrys</i> Valenciennes, 1833	Freckled Driftfish		2	1			
<b>Ogcocephalidae</b>							
<i>Ogcocephalus nasutus</i> (Cuvier, 1829)	Shortnose Batfish		2	1			
<i>Ogcocephalus pumilus</i> Bradbury, 1980	Dwarf Batfish			CAS			
<b>Ophichthidae</b>							
<i>Abliia egmontis</i> (Jordan, 1884)	Key Worm Eel		2	1			
<i>Aprognathodon platyventris</i> Böhlke, 1967	Stripe Eel		2	1			
<i>Callecbelys guineensis</i> (Osório, 1893)	Shorttail Snake Eel		11	1			
<i>Echiophis intertinctus</i> (Richardson, 1848)	Spotted Spoon-nose Eel		2				
<i>Ichthyapus ophioneus</i> (Evermann & Marsh, 1900)	Surf Eel			FIMNH			
<i>Myrichthys breviceps</i> (Richardson, 1848)	Sharptail Eel	S13	2		SJ13		

Scientific name	Common name	Image Plate	Literature source	Online source	Uncommon (site code)	Ichthyocide	DNA
<i>Myrichthys ocellatus</i> (Lesueur, 1825)	Goldspotted Eel		2	1			
<i>Myrophis anterodorsalis</i> McCosker, Böhlke & Böhlke, 1989	Longfin Worm Eel	S13			SJ28		
<i>Myrophis platyrhynchus</i> Breder, 1927	Broadnose Worm Eel		2	1			YES
<i>Myrophis punctatus</i> Lütken, 1852	Speckled Worm Eel		2,11	1			
<b>Ophidiidae</b>							
<i>Brotula barbata</i> (Bloch & Schneider, 1801)	Atlantic Bearded Brotula		2	1		-1	
<i>Lepophidium pheromystax</i> Robins, 1960	Upsilon Cusk-eel		2	1			
<i>Luciobrotula corethromycter</i> * Cohen, 1964	Broomnose Cusk-eel		9				
<i>Ophidion bolbrookii</i> Putnam, 1874	Bank Cusk-eel		2,3,11	1		-1	
<i>Parophidion schmidti</i> (Woods & Kanazawa, 1951)	Dusky Cusk-eel			1			
<i>Xyelacyba myersi</i> * Cohen, 1961	Gargoyle Cusk-eel		9				
<b>Opistognathidae</b>							
<i>Lonchopisthus micrognathus</i> (Poey, 1860)	Swordtail Jawfish	S13	4	1	SJ28, SJ19		YES
<i>Opistognathus aurifrons</i> (Jordan & Thompson, 1905)	Yellowhead Jawfish	S13	2,4	1			YES
<i>Opistognathus macrognathus</i> Poey, 1860	Banded Jawfish	S13	2,4,11	1	SJ5, SJ13, SJ19		
<i>Opistognathus maxillosus</i> Poey, 1860	Mottled Jawfish	S13	2	1	SJ5, SJ13, SJ19	1	
<i>Opistognathus whitehursti</i> (Longley, 1927)	Dusky Jawfish	S13		1	SJ12		
<b>Ostraciidae</b>							
<i>Acanthostracion polygonium</i> Poey, 1876	Honeycomb Cowfish	S13	2	1			
<i>Acanthostracion quadricornis</i> (Linnaeus, 1758)	Scrawled Cowfish	S13	2	1			
<i>Lactophrys bicaudalis</i> (Linnaeus, 1758)	Spotted Trunkfish	S13	2,4	1			
<i>Lactophrys trigonus</i> (Linnaeus, 1758)	Trunkfish	S13	2,4	1			
<i>Lactophrys triquetus</i> (Linnaeus, 1758)	Smooth Trunkfish	S13	2,4	1		1	
<b>Paralichthyidae</b>							
<i>Citharichthys cornutus</i> (Günther, 1880)	Horned Whiff				FMNH		
<i>Citharichthys uhleri</i> Jordan, 1889	Voodoo Whiff				FMNH		
<i>Cyclosetta fimbriata</i> (Goode & Bean, 1885)	Spotfin Flounder	S14	2	1	SJ12, O14		
<i>Syacium micrurum</i> Ranzani, 1842	Channel Flounder		2	1			YES
<b>Parazenidae</b>							
<i>Cyttopsis rosea</i> (Lowe, 1843)	Red Dory		5				
<b>Pempheridae</b>							
<i>Pempheris schomburgkii</i> Müller & Troschel, 1848	Glassy Sweeper	S14	2,4	1	SJ13, ST3, SJ15		YES

Scientific name	Common name	Image Plate	Literature source	Online source	Uncommon (site code)	Ichthyocide	DNA
<b>Poeciliidae</b>							
<i>Poecilia reticulata</i> Peters, 1859	Guppy	S14		1	SJ10		
<b>Polymixiidae</b>							
<i>Polymixia lowei</i> Günther, 1859	Beardfish			FIMNH, CAS			
<i>Polymixia nobilis</i> Lowe, 1836	Noble Beardfish		5,8				
<b>Polynemidae</b>							
<i>Polydactylus virginicus</i> (Linnaeus, 1758)	Barbu			FIMNH			
<b>Pomacanthidae</b>							
<i>Centropyge argi</i> Woods & Kanazawa, 1951	Cherubfish	S14	2,4,5,8	1	O21		
<i>Holocanthus ciliaris</i> (Linnaeus, 1758)	Queen Angelfish	S14	2,4	1		1	YES
<i>Holocanthus tricolor</i> (Bloch, 1795)	Rock Beauty	S14	2,4,5,8	1		1	
<i>Pomacanthus arcuatus</i> (Linnaeus, 1758)	Gray Angelfish	S14	2,4,5,8	1		1	
<i>Pomacanthus paru</i> (Bloch, 1787)	French Angelfish	S14	2,4,5	1		1	
<b>Pomacentridae</b>							
<i>Abudefduf saxatilis</i> (Linnaeus, 1758)	Sergeant Major	S14	2,4,6	1		1	YES
<i>Abudefduf taurus</i> (Müller & Troschel, 1848)	Night Sergeant	S14	2,4	1		1	
<i>Azurina cyanea</i> (Poey, 1860)	Blue Chromis	S14	2,4,8	1		1	YES
<i>Azurina multilineata</i> (Guichenot, 1853)	Brown Chromis	S14	2,4,5,8	1		1	YES
<i>Chromis insolata</i> (Cuvier, 1830)	Sunshinefish	S14	2,5,8	1	O20		
<i>Microspathodon chrysurus</i> (Cuvier, 1830)	Yellowtail Damsel	S14	2,4,5	1		1	
<i>Stegastes adustus</i> (Troschel, 1865)	Dusky Damsel	S14	2,4,6	1		1	
<i>Stegastes diencaeus</i> (Jordan & Rutter, 1897)	Longfin Damsel	S14	2,4	1			YES
<i>Stegastes leucostictus</i> (Müller & Troschel, 1848)	Beaugregory	S14	2,4	1		1	YES
<i>Stegastes partitus</i> (Poey, 1868)	Bicolor Damsel	S14	2,4,5,8	1		1	YES
<i>Stegastes planifrons</i> (Cuvier, 1830)	Threespot Damsel	S14	2,4	1		1	YES
<i>Stegastes xanthurus</i> (Poey, 1860)	Cocoa Damsel	S14	2,4	1		1	YES
<b>Priacanthidae</b>							
<i>Heteropriacanthus cruentatus</i> (Lacepède, 1801)	Glasseye Snapper	S15	2,4	1		1	YES
<i>Priacanthus arenatus</i> Cuvier, 1829	Bigeye	S15	2	1	SJ24		
<i>Pristigenys alta</i> (Gill, 1862)	Short Bigeye		2	1			
<b>Rachycentridae</b>							
<i>Rachycentron canadum</i> (Linnaeus, 1766)	Cobia	S15			ST3		
<b>Rhincodontidae</b>							
<i>Rhincodon typus</i> Smith, 1828	Whale Shark	S15					
<b>Rivulidae</b>							
<i>Kryptolebias marmoratus</i> (Poey, 1880)	Mangrove Rivulus		6	1			

Scientific name	Common name	Image Plate	Literature source	Online source	Uncommon (site code)	Ichthyocide	DNA
<b>Sciaenidae</b>							
<i>Cornula batabana</i> (Poey, 1860)	Blue Croaker		2,11	1			
<i>Eques lanceolatus</i> (Linnaeus, 1758)	Jackknife-fish	S15	2,4	1	SJ30		
<i>Eques punctatus</i> Bloch & Schneider, 1801	Spotted Drum	S15	2,4	1		1	
<i>Odontoscion dentex</i> (Cuvier, 1830)	Reef Croaker	S15	2,4	1		1	
<i>Pareques acuminatus</i> (Bloch & Schneider, 1801)	High-hat	S15	2,4	1		1	YES
<i>Umbrina coroides</i> Cuvier, 1830	Sand Drum		2	1			
<b>Scomberesocidae</b>							
<i>Scomberesox saurus</i> (Walbaum, 1792)	Atlantic Saury				KU		
<b>Scombridae</b>							
<i>Acanthocybium solandri</i> (Cuvier, 1832)	Wahoo	S15	2				
<i>Euthynnus alletteratus</i> (Rafinesque, 1810)	Little Tunny	S15	2	1			YES
<i>Katsuwonus pelamis</i> (Linnaeus, 1758)	Skipjack Tuna	S15	2				
<i>Scomberomorus brasiliensis</i> Collette, Russo & Zavala-Camin, 1978	Serra		2	1			
<i>Scomberomorus cavalla</i> (Cuvier, 1829)	King Mackerel	S15	2		SJ4, ST6		
<i>Scomberomorus regalis</i> (Bloch, 1793)	Cero	S15	2,4	1			
<i>Thunnus albacares</i> (Bonnaterre, 1788)	Yellowfin Tuna	S15	2				
<i>Thunnus atlanticus</i> (Lesson, 1831)	Blackfin Tuna	S15	2	1			
<b>Scorpaenidae</b>							
<i>Pontinus castor</i> Poey, 1860	Longsnout Scorpionfish		2	1			
<i>Pterois volitans</i> (Linnaeus, 1758)	Red Lionfish	S15		1			YES
<i>Scorpaena albifimbria</i> Evermann & Marsh, 1900	Coral Scorpionfish	S15	2,11	1	O8		
<i>Scorpaena bergii</i> Evermann & Marsh, 1900	Goosehead Scorpionfish				FIMNH		
<i>Scorpaena brasiliensis</i> Cuvier, 1829	Barbfish		2,11	1			
<i>Scorpaena calcarata</i> Goode & Bean, 1882	Smoothhead Scorpionfish		2,11	1			
<i>Scorpaena grandicornis</i> Cuvier, 1829	Plumed Scorpionfish		2,6	1			
<i>Scorpaena inermis</i> Cuvier, 1829	Mushroom Scorpionfish	S15	2	1	SJ5		
<i>Scorpaena plumieri</i> Bloch, 1789	Spotted Scorpionfish	S15	2,4	1		1	
<i>Scorpaenodes caribbaeus</i> Meek & Hildebrand, 1928	Reef Scorpionfish	S15	2	1	SJ34, SJ23, SJ13	1	
<b>Serranidae</b>							
<i>Alphesthes afer</i> (Bloch, 1793)	Mutton Hamlet	S16	2	1	SJ23	1	
<i>Bullisichthys caribbaeus</i> Rivas, 1971	Pugnose Bass		5,8				
<i>Cephalopholis cruentata</i> (Lacepède, 1802)	Graysby	S16	2,4,5,8	1		1	YES
<i>Cephalopholis fulva</i> (Linnaeus, 1758)	Coney	S16	2,4,5,8	1		1	
<i>Diplectrum bivittatum</i> (Valenciennes, 1828)	Dwarf Sand Perch	S16	2	1		1	YES
<i>Diplectrum formosum</i> (Linnaeus, 1766)	Sand Perch		4	1			

Scientific name	Common name	Image Plate	Literature source	Online source	Uncommon (site code)	Ichthyocide	DNA
<i>Epinephelus adscensionis</i> (Osbeck, 1765)	Rock Hind	S16	2,4	1	SJ22, SJ15	1	
<i>Epinephelus guttatus</i> (Linnaeus, 1758)	Red Hind	S16	2,4,5,8	1		1	
<i>Epinephelus itajara</i> (Lichtenstein, 1822)	Atlantic Goliath Grouper	S16	2				
<i>Epinephelus morio</i> (Valenciennes, 1828)	Red Grouper	S16	2	1			
<i>Epinephelus striatus</i> (Bloch, 1792)	Nassau Grouper	S16	2,4,5,8	1		1	YES
<i>Hypoplectrus aberrans</i> Poey, 1868	Yellowbelly Hamlet	S16	2,4	1		1	
<i>Hypoplectrus chlorurus</i> (Cuvier, 1828)	Yellowtail Hamlet	S16	2,4,5,8	1			
<i>Hypoplectrus guttavarius</i> (Poey, 1852)	Shy Hamlet	S16	2,4	1	SJ19, ST6		
<i>Hypoplectrus indigo</i> (Poey, 1851)	Indigo Hamlet	S16	2,4	1			
<i>Hypoplectrus nigricans</i> (Poey, 1852)	Black Hamlet	S16	2,4	1		1	
<i>Hypoplectrus puella</i> (Cuvier, 1828)	Barred Hamlet	S16	2,4	1		1	
<i>Hypoplectrus unicolor</i> (Walbaum, 1792)	Butter Hamlet	S16	2,4	1		1	
<i>Hyporhamphus mystacinus</i> (Poey, 1852)	Misty Grouper		2,8				
<i>Liopropoma mowbrayi</i> Woods & Kanazawa, 1951	Cave Basslet		2,5				
<i>Liopropoma rubre</i> Poey, 1861	Peppermint Basslet	S16	2,4	1	ST1, SJ9, SJ13	1	
<i>Mycteroperca acutirostris</i> (Valenciennes, 1828)	Western Comb Grouper		2	1			
<i>Mycteroperca bonaci</i> (Poey, 1860)	Black Grouper	S17	2	1	SJ33, O9, O10		
<i>Mycteroperca interstitialis</i> (Poey, 1860)	Yellowmouth Grouper	S17	2,4,5	1	SJ7		YES
<i>Mycteroperca tigris</i> (Valenciennes, 1833)	Tiger Grouper	S17	2,5,8	1	O11, O12, O13	1	
<i>Mycteroperca venenosa</i> (Linnaeus, 1758)	Yellowfin Grouper	S17	2,4,5,8	1		1	YES
<i>Paranthias furcifer</i> (Valenciennes, 1828)	Atlantic Creolefish	S17	2,5,8	1	SJ33		
<i>Pronotogrammus martinicensis</i> (Guichenot, 1868)	Roughtongue Bass		5				
<i>Rypticus bistrispinus</i> (Mitchill, 1818)	Freckled Soapfish	S17			O14		
<i>Rypticus carpenteri</i> Baldwin & Weigt, 2012	Slope Soapfish	S17					
<i>Rypticus saponaceus</i> (Bloch & Schneider, 1801)	Greater Soapfish	S17	2,4	1		1	
<i>Rypticus subbifrenatus</i> Gill, 1861	Spotted Soapfish		2	1		1	
<i>Schultzea beta</i> (Hildebrand, 1940)	School Bass	S17	2	1	O19		YES
<i>Serraniculus pumilio</i> Ginsburg, 1952	Pygmy Sea Bass	S17	11	1	SJ19		YES
<i>Serranus annularis</i> (Günther, 1880)	Orangeback Bass	S17	2,11	1	O17		
<i>Serranus baldwini</i> (Evermann & Marsh, 1899)	Lantern Bass	S17	2,4	1	SJ32, SJ12, SJ22		YES
<i>Serranus luciopercanus</i> Poey, 1852	Crosshatch Bass		2,5,8				
<i>Serranus phoebe</i> Poey, 1851	Tattler		2	1			
<i>Serranus tabacarius</i> (Cuvier, 1829)	Tobaccofish	S17	2,4	1			YES
<i>Serranus tigrinus</i> (Bloch, 1790)	Harlequin Bass	S17	2,4	1		1	
<i>Serranus tortugarum</i> Longley, 1935	Chalk Bass	S17	2,4,5	1			YES

Scientific name	Common name	Image Plate	Literature source	Online source	Uncommon (site code)	Ichthyocide	DNA
<b>Sparidae</b>							
<i>Archosargus rhomboidalis</i> (Linnaeus, 1758)	Sea Bream	S17	2,8	1	SJ13, SJ3		
<i>Calamus bajonado</i> (Bloch & Schneider, 1801)	Jolthead Porgy	S17	2	1			
<i>Calamus calamus</i> (Valenciennes, 1830)	Saucereye Porgy	S17	2,4	1			
<i>Calamus penna</i> (Valenciennes, 1830)	Sheepshead Porgy	S17	2,4	1			
<i>Calamus pennatula</i> Guichenot, 1868	Pluma Porgy	S17	2,4	1			YES
<i>Calamus proridens</i> Jordan & Gilbert, 1884	Littlehead Porgy			CMN			
<i>Diplodus caudimaculatus</i> (Poey, 1860)	Silver Porgy	S17	2,4,11	1	ST6		
<b>Sphyraenidae</b>							
<i>Sphyraena barracuda</i> (Edwards, 1771)	Great Barracuda	S17	2,4,5,6,8	1			YES
<i>Sphyraena borealis</i> DeKay, 1842	Sennet	S17	2	1	SJ13, SJ12, SJ21		
<b>Sphyrnidae</b>							
<i>Sphyrna lewini</i> (Griffith & Smith, 1834)	Scalloped Hammerhead		10	1			
<i>Sphyrna mokarran</i> (Rüppell, 1837)	Great Hammerhead		10				
<b>Spratelloididae</b>							
<i>Jenkinsia lamprotaenia</i> (Gosse, 1851)	Dwarf Herring		2,6	1		1	YES
<i>Jenkinsia parvula</i> Cervigón & Velazquez, 1978	Shortstriped Round Herring		2				
<i>Jenkinsia stolifera</i> (Jordan & Gilbert, 1884)	Shortband Herring		2				
<b>Squalidae</b>							
<i>Squalus cubensis</i> Howell Rivero, 1936	Cuban Dogfish			FMNH			
<b>Sternoptychidae</b>							
<i>Sonoda paucilampa</i> Grey, 1960	Deepsea Hatchetfish			NMNH			
<b>Stomiidae</b>							
<i>Astronesthes similis</i> Parr, 1927	Similar Snaggletooth			NMNH			
<b>Syngnathidae</b>							
<i>Amphelikurtus dendriticus</i> (Barbour, 1905)	Pipehorse	S18			SJ12		
<i>Bryx dunckeri</i> (Metzelaar, 1919)	Pugnose Pipefish	S18	2	1	SJ13	1	YES
<i>Cosmocampus brachycephalus</i> (Poey, 1868)	Crested Pipefish		2			1	
<i>Cosmocampus elucens</i> (Poey, 1868)	Shortfin Pipefish	S18	2,4	1	SJ19		
<i>Cosmocampus profundus</i> (Herald, 1965)	Deepwater Pipefish		2				
<i>Halicampus crinitus</i> (Jenyns, 1842)	Banded Pipefish	S18			SJ34, SJ13, SJ22		
<i>Hippocampus erectus</i> Perry, 1810	Lined Seahorse		11	1			YES
<i>Hippocampus reidi</i> Ginsburg, 1933	Longsnout Seahorse	S18	4	1	SJ19		YES
<i>Microphis lineatus</i> (Kaup, 1856)	Opposum Pipefish	S18			O23		
<i>Pseudophallus mindii</i> (Meek & Hildebrand, 1923)	Freshwater Pipefish		11				

Scientific name	Common name	Image Plate	Literature source	Online source	Uncommon (site code)	Ichthyocide	DNA
<i>Syngnathus caribbaeus</i> Dawson, 1979	Caribbean Pipefish	S18	2		SJ21		
<i>Syngnathus dawsoni</i> (Herald, 1969)	Antillean Pipefish		2,4,11	1			
<i>Syngnathus pelagicus</i> Linnaeus, 1758	Sargassum Pipefish			ROM			
<b>Synodontidae</b>							
<i>Saurida brasiliensis</i> Norman, 1935	Largescale Lizardfish		2				
<i>Saurida suspicio</i> Breder, 1927	Doubtful Lizardfish	S18	2	1	SJ5, SJ13		YES
<i>Synodus foetens</i> (Linnaeus, 1766)	Inshore Lizardfish	S18	2	1	SJ5, SJ13	1	YES
<i>Synodus intermedius</i> (Spix & Agassiz, 1829)	Sand Diver	S18	2,4	1		1	YES
<i>Synodus poeyi</i> Jordan, 1887	Offshore Lizardfish		2				
<i>Synodus synodus</i> (Linnaeus, 1758)	Red Lizardfish	S18	2	1	SJ11, SJ21	1	
<i>Trachinocephalus myops</i> (Forster, 1801)	Snakefish			CAS			
<b>Tetraodontidae</b>							
<i>Canthigaster rostrata</i> (Bloch, 1786)	Sharppose Puffer	S18	2,4,5,8	1		1	
<i>Sphoeroides spengleri</i> (Bloch, 1785)	Bandtail Puffer	S18	2,4	1		1	YES
<i>Sphoeroides testudineus</i> (Linnaeus, 1758)	Checkered Puffer	S18	2,4,6	1	O15	1	
<b>Triakidae</b>							
<i>Mustelus canis</i> (Mitchill, 1815)	Smooth Dogfish			FIMNH			
<b>Triglidae</b>							
<i>Peristedion longispatha</i> Goode & Bean, 1886	Widehead Armored Searobin			CAS			
<b>Tripterygiidae</b>							
<i>Enneanectes altivelis</i> Rosenblatt, 1960	Lofty Triplefin	S18	2	1		1	
<i>Enneanectes atrorus</i> Rosenblatt, 1960	Blackedge Triplefin		2,11	1			
<i>Enneanectes boehlkei</i> Rosenblatt, 1960	Roughhead Triplefin	S18	2	1		-1	YES
<i>Enneanectes jordani</i> (Evermann & Marsh, 1899)	Mimic Triplefin	S18	2	1	SJ21		
<i>Enneanectes matador</i> Victor, 2013	Matador Triplefin	S18		1			YES
<b>Xiphiidae</b>							
<i>Xiphias gladius</i> Linnaeus, 1758	Swordfish	S18					

**Notes:** Image voucher – supplementary plate number is given; photographer name is imbedded in each image. Literature source – 1 DeAngelis et al. (2008); 2 Dennis (2000); 3 Dennis et al. (2004); 4 Friedlander et al. (2013); 5 Garcia-Sais (2005); 6 Loftus (2003); 7 Mantatrust.org pers. comm. to DRR; 8 Nelson and Appledorn (1985); 9 Quatrinni et al. (2017); 10 Recksiek et al. (2006), 11 Smith-Vaniz and Jelks (2014); 12 Rogers et al. (2010). Online source – 1 indicates that an aggregator source exists, with the source named whenever it represents the sole voucher: AMNH (American Museum of Natural History); NOAA (National Oceanographic and Atmospheric Administration); BOLD (Barcode of Life); FIMNH (Florida Museum of Natural History); MCZ (Museum of Comparative Zoology); NMNH (National Museum of Natural History); ANSP (Academy of Natural Sciences of Philadelphia); CAS (California Academy of Sciences); ROM (Royal Ontario Museum); KUBI (University of Kansas Biodiversity Institute); CMN (Canadian Museum of Nature). Uncommon – species seen at 3 or less named sites by CJE and AME (see Suppl material 3: File S2a, b (for site codes) and Suppl. material 4: File S3). Ichthyocide – species collected by this method as noted in Dennis (2000); parentheses indicate ichthyocide was the only collection method noted by Dennis (2000). Gobiidae – we follow Thacker (2009) in including *Cerdale* and *Ptereleotris* among the Gobiidae. *Hypoplectrus* – we follow Puebla et al. (2022) in treating *H. maculiferus* as a synonym of *H. aberrans*.

**Table 3.** Fishes from St. John-Thomas recorded by different sources.

Types of fish taxa recorded	Species	Genera	Families
<b>Total from all sources</b>	561	296	108
<b>From Literature sources All</b>	451	251	89
Dennis 2000 All	401	216	79
Sole source is Dennis 2000	164	126	55
Sources other than Dennis 2000	50	44	25
<b>From Online sources All</b>	453	253	97
Online sources only	50	46	42
<b>From Images All</b>	371	20	73
Images only	34	29	20
<b>Deep species All sources</b>	49	44	33
Recorded by Dennis 2000	19	18	13
<b>Uncommon shallow species</b>	138	104	45
<b>Ichthyocide Collection All</b>	173	99	45
Ichthyocide only	18	15	11
<b>mtDNA BARCODES</b>	<b>Species</b>	<b>Genera</b>	<b>Families</b>
St. John-Thomas	156	93	41
Sole record is from barcode data	1	1	1
Puerto Rico	90	50	25
St. John-Thomas but not Puerto Rico	113	61	24
Puerto Rico but not St. John-Thomas	47	18	8
St. Croix	1	1	1
British Virgin Islands	3	2	1
All sites combined	207	112	49

**Notes:** Data sources (literature, online sources, images) are listed in Table 2. Deep species are those exclusively or typically found below 40 m depth. Uncommon shallow species are those found at 1–3 sites by CJE, AME, LR, and third-party photographers as indicated in Table 2. Ichthyocide collection: recorded as being collected with rotenone by a source cited by Dennis (2000). Ichthyocide only: the only collection method listed for a species from St. John-Thomas by Dennis (2000). DNA barcodes: (see Suppl. material 7: File S6). The single DNA Barcoded species collected at St. Croix (see Suppl. material 7: File S6) is not known from St. John-Thomas. The St. John-Thomas species count includes four identified only to genus. DNA barcode data for *Pterois volitans* are not included in this table.

Dennis (2000) listed 401 species from 216 genera and 79 families from those two islands (Table 2). We found records of an additional 159 species, producing an increase of 39.7% in the number of species, 37.0% in the number of genera and 36.7% in the number of families known from there (Table 3). The additions include 34 species for which the only source is a voucher image, 50 species recorded in post-2000 publications, and 49 species recorded only by online sources of museum (and other) data (Table 3). Of the 561 in Table 2, 24.6% were uncommon. Although 30.1% were collected using rotenone, species accounts by Dennis (2000) mentioned no other collecting method for only 10.4% of that subgroup of species. The 561 include three non-natives to the area (*Oreochromis niloticus*, *Poecilia reticulata* and *Pterois volitans*), 11 freshwater/estuarine species (*Anguilla rostratus*, *Dormitator maculatus*, *Eleotris perniger*, *Gobiomorus dormitor*, *Awaous banana*, *Sicydium plumieri*, *Sicydium punctatum*, *Dajaus monticola*, *Microphis lineatus* and *Pseudophallus mindii*) and 547 marine species native to the GC.

**Table 4.** Taxonomic comparisons of St. John-Thomas and St. Croix marine fish faunas.

Site	Species	Genera	Families
<b>Both US Virgin Islands</b>			
Entire fauna (n)	679	345	122
Shallow fishes (n)	590	279	90
Deep fishes (n)	89	77	54
<b>St. John-Thomas</b>			
Entire Fauna (n)	547	283	105
Percent of USVI fauna	80.6	82.0	86.0
Percent only at St. John-Thomas	19.3	15.5	10.5
Shallow fishes (n)	497	245	86
Percent of USVI shallow fauna	84.2	86.6	94.5
Percent only at St. John-Thomas	13.0	7.4	1.9
Deep fishes (n)	50	44	34
Percent of USVI deep fauna	56.2	57.1	63.0
Percent only at St. John-Thomas	70.0	50.0	26.5
<b>St. Croix</b>			
Entire fauna (n)	573	301	112
Percent of USVI fauna	84.5	87.2	91.8
Percent only at St. Croix	23.4	20.4	15.5
Shallow fishes (n)	519	256	88
Percent of USVI fauna	88.0	91.8	97.8
Percent only at St. Croix	18.3	13.1	2.7
Deep fishes (n)	54	50	39
Percent of USVI deep fauna	61.4	64.9	62.2
Percent only St. Croix	72.2	60.0	41.0

**Notes:** USVI fauna = combined fauna of St. John-Thomas and St. Croix, with exotic and primarily freshwater species excluded. Some genera and families have a deep member in one site but not the other, which affects USVI totals for deep and shallow genera and families. Shallow fishes: species exclusively or commonly found shallower than 40 m. Deep fishes: species exclusively or largely found deeper than 40 m (see methods for further details).

### Comparative taxonomic composition of the USVI fish faunas (Table 4, Suppl. material 5: File S4)

The species richness of the USVI marine fauna (i.e., the combined St. John-Thomas plus St. Croix faunas) was 15–20% greater than that of either of the two insular faunas (Table 4). Those two faunas had slightly higher relative rates of richness of genera and families than of species. The larger size of the USVI fauna of species derives from ~ 1/5 of species in each insular fauna not being present in the other, with lower proportions of genera and families also being recorded only at one of the two islands. Relative faunal richness at all three taxonomic levels and the relative abundance of taxa present at only one island were ~ 5% higher at St. Croix than St. John-Thomas. In both island faunas the relative representation of species, genera, and families in the entire USVI fauna was substantially greater among shallow species than deep species. The deep fauna was much smaller than the shallow fauna at each island and there was much less overlap in occurrence of species, genera, and families between the two insular deep faunas than between their shallow faunas (Table 4).

**Table 5.** Abundance of ecotypes of reef-associated bony fishes in the Greater Caribbean and the USVI.

	Region	St. John-Thomas	St. Croix
<b>All species (n)</b>	992	470	493
<b>Pelagic species % of fauna</b>	8.0	<b>10.4</b>	<b>10.3</b>
<b>Non-pelagic species % of fauna</b>	92.0	89.6	89.7
Demersal species %	34.6	<b>46.3</b>	<b>45.0</b>
Benthic species %	<b>65.4</b>	53.7	55.0
Cryptobenthic species %	<b>64.6</b>	53.0	54.3
Small cryptobenthic species %	<b>42.6</b>	31.6	32.5
CCRF species %	<b>45.9</b>	36.3	35.7
<b>SHALLOW NON-PELAGIC SPECIES (n)</b>	772	400	424
<b>Percent of fauna</b>	84.6	<b>95.0</b>	<b>95.9</b>
Demersal species %	34.9	<b>45.3</b>	<b>44.0</b>
Benthic species %	<b>65.1</b>	54.7	56.0
Cryptobenthic species %	<b>64.0</b>	54.3	55.2
Small cryptobenthic species %	<b>42.5</b>	33.3	34.0
CCRF species %	<b>46.0</b>	37.5	37.3
<b>DEEP NON-PELAGIC SPECIES (n)</b>	141	21	18
<b>Percent of fauna</b>	<b>15.4</b>	5.0	4.2
Demersal species %	33.3	<b>66.7</b>	<b>66.7</b>
Benthic species %	<b>66.7</b>	33.3	33.3
Cryptobenthic species %	<b>66.7</b>	33.3	33.3
Small cryptobenthic species %	<b>43.3</b>	4.8	0
CCRF species %	<b>45.4</b>	19.0	0

**Notes:** Data for the region pattern are from Robertson and Tornabene (2021), for St. Croix are from Robertson et al. (2022), and for St. John-Thomas are in File S5. Bold percentages indicate whether the value(s) for either the region or the USVI islands were > 5% higher than the value(s) for the other group in each case.

**Table 6.** Zoogeographic composition of the USVI and Sint Eustatius faunas. Percentage of species in each category. Non-native species are not included.

Site (n)	Northwest Atlantic	Western Atlantic	Trans-Atlantic	Atlantic & Indo-Pacific
St. Croix (534)	41.6	33.9	13.9	10.6
St. John-Thomas (558)	39.5	36.5	14.0	10.0
Sint Eustatius (406)	41.1	33.3	15.3	10.3

**Notes:** St. Croix data are from Smith-Vaniz and Jelks 2014. Sint Eustatius values are from Robertson et al. (2020). St. John-Thomas values are from the present study. Northwest Atlantic = Greater Caribbean, with or without range extensions to the north of that region. Western Atlantic = Northwestern Atlantic + Brazil. Trans-Atlantic = anywhere in the western Atlantic + any of the islands of the central Atlantic and/or the Eastern Atlantic. Atlantic & Indo-Pacific = Anywhere in the Western Atlantic + anywhere in the Indo-Pacific.

## Ecotypic structure of the USVI reef-fish faunas vs. the region (Table 5, Suppl. material 6: File S5)

We compared the ecotypic structure of the St. John-Thomas and St. Croix faunas of reef-associated fishes with that of the GC fauna (see Robertson and Tornabene 2021). Both St. Croix and St. John-Thomas have faunas that are almost half the size of the total regional

fauna, with the listed St. John-Thomas fauna being ~ 5% smaller than that of St. Croix (Table 5). Compared to the GC fauna both islands have slightly higher percentages of pelagic species, distinctly higher percentages of demersal species, and correspondingly lower percentages of benthic, cryptobenthic, small cryptobenthic, and CCRF species. These differences for non-pelagic types apply to each entire USVI fauna, and to both shallow- and deep-reef subgroups of those faunas. Both USVI sites also have markedly lower relative abundances (~ 1/3) of deep-reef species than the regional fauna. The relative abundances of different ecotypes are remarkably similar at both islands, except for the presence of a few deep cryptobenthic and CCRF species detected only at St. John-Thomas.

### Zoogeographic structure of the USVI faunas (Table 6)

The zoogeographic structures of the faunas of the two USVI sites and nearby Sint Eustatius are quite similar (Table 6). Species that are endemic to the Greater Caribbean and, in a few cases, surrounding areas are the largest group in all three faunas, with West Atlantic species also found in Brazil being the second largest by a small margin in each case. The two smallest groups in each case are Trans-Atlantic and Atlantic & Indo-Pacific. The ranks of those four groups are the same in all three faunas, a measure of their strong similarities.

### mtDNA-Barcode Coverage (Tables 2, 3; Suppl. material 7: File S6)

Table 2 indicates which members of the St. John-Thomas fauna have mtDNA-barcode sequences on the BOLD database derived from specimens collected at that site. Table 3 presents a summary of taxa that have sequences obtained from St. John-Thomas, Puerto Rico, the British Virgin Islands and St. Croix, singly and in combination. File S6 provides technical information about those barcode data for the various species. We obtained local DNA-barcodes for 156 fish species in 156 BINs from St. John-Thomas, with one additional from St. Croix, and three additional species from around the British Virgin Islands (total 160 species). Of these, two are only from GenBank records harvested by BOLD, and 10 are added from specimens collected in offshore larval plankton tows described in Lamkin et al. (2009). We obtained 91 species records (including one non-native, *Pterois volitans*) for Puerto Rico, 44 of them shared with the Virgin Islands. Of the 91, 27 are added from Harms-Tuohy et al. (2016), 14 from GenBank records harvested by BOLD, and seven from other sources, including the University of Kansas (UKFBJ), Smithsonian (Birmingham/Lessios; BSMUA & BSOPA), the Guy Harvey Research Institute (Hanner et al. 2011; EBFSF), and the Museum and Art Gallery of the Northern Territory (GOBY) in Australia.

The available DNA-barcode sequence records from specimens collected at St. John-Thomas represent coverage of 27.8% of the species, 31.4% of the genera and 38% of the families of fishes known from that site. Barcode records represent the sole source of information on the presence of one species known from those islands and are also available for another four species currently identifiable only to genus. Distinctly fewer species have been barcoded from fish taken at Puerto Rico, and there are almost no

such data available from either St. Croix or the British Virgin Islands. Barcode records from Puerto Rico and the British Virgin Islands exist for 52 species occurring in St. John-Thomas but not sequenced from there, bringing the total PRP DNA-barcoded species to 36.5% of St. John-Thomas fauna. All but seven of the 200 barcoded species are reef-associated bony fishes. The vast majority (98.5%) of barcoded species are shallow forms. Deep-living species are especially under-represented among the barcoded forms: only three of 51 such species have barcode data (File S6).

## Discussion

### St. Croix

The species records we have added increased the size of that island's fauna by 7.5%. Almost a third of the additions arise from voucher photographs of shallow-reef species photographed by CJE and AME (and provided by Mantatrust.org). Those include several not accepted by Smith-Vaniz and Jelks (2014) due to inadequate information available at that time. Cryptobenthic fishes, which, by definition, are generally difficult to observe, are a major component of Greater Caribbean reef-fish faunas, including that of St. Croix. Such species comprised all but one of those added by CJE and AME. The exception, *Kyphosus cinerascens*, may have been misidentified previously, since the taxonomic status and global distributions of members of the genus were only comprehensively reviewed by Knudsen and Clements (2016), after Smith-Vaniz and Jelks (2014) published their checklist. Almost half the additions were deep-living species, one third of which were recorded only by submersible or ROV, with the remainder coming from online and literature records.

The process of obtaining location records is an ongoing one for online aggregators, which have vastly increased the amounts of data they host during the last half decade. Although the aggregators offer such information, and are involved in collaborative data sharing, such sharing is sufficiently incomplete that it is necessary to examine records from multiple aggregators to obtain a comprehensive picture of all the data available for any particular site. Even “old” data becomes newly available on the aggregators from time to time and needs to be included in faunal inventories of well-studied sites. The increase in faunal size, although not large in percentage terms, demonstrates the utility of citizen-science efforts to provide photographic vouchers, of reviews of submersible and ROV studies of deep-living fishes, and of periodic evaluations of information available online from aggregators.

### St. John-Thomas

Although the 401 species list for this site extracted from Dennis (2000) was substantial (74% the size of Smith-Vaniz and Jelks (2014) count for St. Croix), our use of the same methods as those that produced an increase in the St. Croix fauna produced a much larger increase in the St. John-Thomas fauna: 40% vs. 7.5% for St. Croix. Dennis (2000) was

the sole source for 29% of species recorded in our expanded list of the St. John-Thomas fauna. Records from additional sources brought the size of the St. John-Thomas fauna to within 5% of the size of the St. Croix fauna. Citizen-scientists' photographic records accounted for 22% of the new additions and data only available from online databases for 33%, while other literature sources provided the sole records for 32% of the additional species. Multiple types of sources accounted for the remaining 13% of new records.

### The size, and taxonomic- and ecotypic structure of the two USVI marine faunas

Both insular marine faunas are over 80% the size of the combined USVI fauna in terms of species richness. Species found at only one of the two islands represent ~ 20% of each fauna. For shallow species the size of each insular fauna is 85–90% that of the combined fauna, with correspondingly lower rates of occurrence at only one island. Two factors may contribute to these differences between the island faunas: variation in ecological conditions between the islands and inadequate sampling. The possibility of differing ecological conditions seems small as both islands have the same range of large-scale habitat types, although those vary in abundance between the islands. The shelf area of St. John-Thomas is close to 10 times the size of the St. Croix shelf, yet the former has the smaller known fauna. At both islands the great majority of sampling has occurred in quite shallow water, often very close to shore in the case of St. John-Thomas. Shelf habitats likely are under-sampled at both islands, strongly so at St. John-Thomas, where there are large areas of habitat between 40–60 m depth some distance from the islands on both the northern and southern parts of the PRP. At St. Croix most shallow sampling has occurred in and near the Buck Island Reef National Monument, rather than spread around different parts of the platform and different sides of the island. Hence both insular faunas likely are larger than indicated here, particularly in the case of St. John-Thomas.

Review of the two USVI marine species lists show that species not shared between the two islands are distributed through many genera and families (Suppl. material 5: File S4; Table 4). None are endemic to either USVI island and single-island endemics are rare amongst the Greater Caribbean fauna and limited to highly isolated islands such as Cayman. Most species in that region have geographic ranges much larger than the USVIs. The larger size of the St. Croix fauna, particularly of cryptobenthic species can be attributed to a greater effort to find such species. This was done using rotenone during two intensive sampling campaigns that occurred ~ 40 y after rotenone sampling at St. John-Thomas, plus some subsequent minor efforts in the shallow part of a Buck Island Reef National Monument that, in its entirety constitutes ~ 1/3 of the St. Croix insular platform: 46% (262) of the native marine species known from St. Croix are shallow species collected using rotenone (Smith-Vaniz and Jelks 2014), vs. 31.7% (173) of such species from St. John-Thomas. Later sampling by Pittman et al. (2008) at the same small, shallow St. Croix site as studied by Smith-Vaniz et al. (2006) added 10.9% more species to the tally of the first two series of collections. Smith-Vaniz and Jelks (2014) produced a list of 41 species from 22 families that, at that time, were known from St. John-Thomas but not St. Croix. Since then, five of the 35 shallow species on that table have been added to the St. Croix fauna (Table 1 here), together with two others that were listed as unconfirmed

by those authors. Photographic sampling of shallow reef fishes at St. John-Thomas by CJE, AME and other citizen scientists, by itself increased the size of the fauna registered by Dennis (2000) by 8.5%. Finally, the species composition of local reef-fish faunas can change substantially through time at intensively sampled sites, for varying reasons (e.g., see changes registered by Starck et al. 2017 over a 50y period), highlighting the utility of temporally dispersed sampling. With further sampling many shallow species currently known from only one of the USVI should be expected to be found at the other, in which case the shallow faunas of each island would be 10–15% larger than the current figures.

The deep-species fauna represents only 13.1% of the entire (shallow plus deep) USVI fauna and deep species exhibit much lower rates of faunal overlap between the two islands than occurs among shallow species. The two islands have experienced low rates of exploration of deep habitats, particularly deep reefs, by submersibles and ROVs, which were limited to observational studies. The few ROV (Quattrini et al. 2017) and submersible dives (Nelson and Appeldoorn 1985; Garcia-Sais 2005) were the sole source of only 11.1% and 28% of records of deep fishes at St. Croix and St. John-Thomas, respectively. The edges of the insular platforms of the two USVIs are < 50 km apart and the suite of deep species involved have ranges much larger than the area occupied by the USVI. Low levels of sampling can account for the small size of both USVI deep faunas, particularly the deep-reef component, and to the low level of overlap between the deep faunas of the two islands.

At both USVI sites the deep-reef species represent only 4.2–5% of the entire local reef-fish fauna, i.e., ~ 1/3 of the percentage for the GC regional fauna (Robertson et al. 2022). In contrast, when intensive submersible collecting and observations have been aimed specifically at assessing the diversity of deep-reef fish faunas, such as has occurred at other Caribbean islands (Curacao, Roatan and Sint Eustatius), the inventory of deep-reef species at individual islands has increased ~ 9 fold, with such species representing 16% of the entire (shallow plus deep) reef-fish fauna at the most intensively sampled island (Robertson et al. 2022), i.e., more than three times the level at each USVI. Similar sampling at both USVI undoubtedly will increase the absolute and relative sizes of their deep-reef faunas. Smith-Vaniz and Jelks (2014) concluded that there was no indication at the time of their study that the St. Croix fauna had reached asymptotic size. The additions reported here and patterns of variation in faunal composition between the two islands support that view for St. John-Thomas as well as St. Croix.

Reef-associated bony fishes comprised 84% and 91%, respectively, of the faunas of St. John-Thomas and St. Croix, and the St. John-Thomas reef-fish fauna was 94.3% the size of the equivalent fauna of St. Croix. The ecotypic structure of those two USVI reef-fish faunas was very similar, with both differing from the broad structure of the GC regional fauna by having larger proportions of pelagic and demersal species that are readily visible to observers and correspondingly smaller proportions of cryptic species. Similarities in the zoogeographic structures and sizes of the two USVI faunas support the view that both can be considered to be sampled with a similar level of efficiency, at least in terms of their shallow faunas.

## mtDNA-barcode coverage

In terms of the availability of DNA-barcodes for marine fishes, the Greater Caribbean currently is the most well-sampled large marine biogeographic region in the tropics, with ~ 90% of the shore-fishes barcoded and up to 95% of the shallow reef-associated species (Victor et al. 2015). However, several specific locations account for the vast majority of sequences. Those include Florida, Yucatan (Mexico), Belize, Panama, and Curacao; with species lists published for Yucatan by Valdez-Moreno et al. (2010) and lists for additional locations in Weigt et al. (2012). The Puerto Rican Plateau has been only lightly sampled, with information derived mostly from older collections by author BV at St. John-Thomas and Puerto Rico, and from a set of lionfish stomach contents from La Parguera in Puerto Rico sequenced by Harms-Tuohy et al. (2016). The latter identified 39 species from 16 families. A few additional sequences come from open-ocean sampling for larvae around the USVI, by Lamkin et al. (2009). The older collections from St. Thomas and Puerto Rico were collected by BV for recruitment and otolith studies as well as some taxonomic reviews (e.g., the genera *Coryphopterus* and *Emblemariopsis*). The recent additions of 19 species from St. John were collected by CJE and AME mainly for DNA confirmation of the species identification of diagnostic underwater photographs that serve as vouchers here, mostly of cryptobenthic fishes. No collections at St. John-Thomas or elsewhere on the PRP that provided DNA barcodes were expressly made for assembling an inventory of fish species- hence the absence of some of the most abundant and widespread shallow reef fishes in the barcode list presented here (e.g., the Bluehead Wrasse, *Thalassoma bifasciatum*).

We cannot directly compare barcode coverage of fishes at St. John-Thomas with that at other intensively barcoded locations noted above because neither the number of barcoded species nor the local species inventory have been comparably evaluated at any of those sites. The results of the present assessment of DNA-barcode coverage for the USVI and the remainder of the PRP highlight the usefulness of the DNA-barcode database for ancillary projects. Accumulating sequences for unrelated purposes, such as taxonomic reviews, stomach-content studies, larval or e-DNA surveys (environmental DNA, where water is sampled for dissolved DNA sequences), augments the general DNA-barcode coverage for specific biogeographic regions and helps confirm species identifications for faunal surveys.

## Permits

Collections from St. John in 2021 were made under National Park Service Collecting Permit VIIS-2021-SCI-0006: We especially thank Thomas Kelley (Virgin Islands NP/ Virgin Islands Coral Reef NM) for facilitating that work.

## Acknowledgements

CJE and AME greatly appreciate the assistance, knowledge, photographic contributions, and camaraderie of various people and organizations in the USVI: Alasdair Dunlap-Smith for logistical support, help with collections and photographic contributions; Miguel Andres Goolishian Hernandez for assistance with diving and photo contributions; the owners and staff of Low Key Watersports' on St. John and to St. Croix Ultimate Bluewater Adventures on St. Croix; Bogie's Villa for logistical support at St. John. Thanks to Caroline Rogers, Chris Rapchick, Serenity Mitchell, Spencer Riley, Max Koestenblatt, Cristina Kessler, Jenny Keith, Nicole Krampitz, Sara Richter and Jamie Irving (see File S2C) for contributing voucher images, to Rafe Boulon for information about USVI fishes and to Guy Stevens and Nicole Pelletier of Mantatrust.org for providing photographic documentation of mobulid rays at the USVI. We also appreciate the cooperation of J. Lamkin, as well as T. Girard, E. Malka, A. Shiroza, L. Vasquez-Yeomans, and G. Zapfe for providing information on larvae from the NOAA SEFSC NMFS Miami Larval Fish collection (funded by NOAA's Coral Reef Conservation Program, NOAA NMFS SEFSC, and University of Miami's Cooperative Institute for Marine and Atmospheric Studies). Thanks also to Andrew Bentley (Biodiversity Institute, University of Kansas), Amy Driskell (LAB, National Museum of Natural History), Chelsea Harms-Tuohy (Isla Mar Research Expeditions, Rincon, PR) and Eric Post (FFWCC, Fish and Wildlife Research Institute, FL) for access to barcodes and information and discussion for assigning them to species.

## References

- Baldwin CC, Tornabene L, Robertson DR (2018) Below the mesophotic. *Scientific Reports* 8: e4920. <https://doi.org/10.1038/s41598-018-23067-1>
- Brandl SJ, Goatley CHR, Bellwood DR, Tornabene L (2018) The hidden half: Ecology and evolution of cryptobenthic fishes on coral reefs. *Biological Reviews of the Cambridge Philosophical Society* 93(4): 1846–1873. <https://doi.org/10.1111/brv.12423>
- Bunckley-Williams L, Williams EH (2004) New Locality, Depth, and Size Records and Species Character Modifications of Some Caribbean Deep-Reef/Shallow Slope Fishes and a New Host and Locality Record for the Chimaera Cestodarian. *Caribbean Journal of Science* 40: 88–119.
- Clavijo IE, Yntema JA, Ogden JC (1980) An annotated list of the fishes of St. Croix, U.S. Virgin Islands. West Indies Lab, Special Publication, 2<sup>nd</sup> edn. West Indies Laboratory, Christiansted, 49 pp. <https://doi.org/10.5281/zenodo.5510695>
- DeAngelis BM, McCandless CT, Kohler NE, Recksiek CW, Skomal GB (2008) First characterization of shark nursery habitat in the United States Virgin Islands: Evidence of habitat partitioning by two shark species. *Marine Ecology Progress Series* 358: 257–271. <https://doi.org/10.3354/meps07308>
- Dennis GD (2000) Annotated checklist of shallow-water marine fishes from the Puerto Rico Plateau including Puerto Rico, Culebra, Vieques, St. Thomas, St. John, Tortola, Virgin Gorda and Anegada. Florida Caribbean Science Center, Gainesville, [244 + 26] 286 pp. <https://doi.org/10.5281/zenodo.5770763>

- Dennis GD, Hensley D, Colin PL, Kimmel JJ (2004) New Records of Marine Fishes from the Puerto Rican Plateau. *Caribbean Journal of Science* 40: 70–87. <https://doi.org/10.5281/zenodo.5512853>
- Friedlander AM, Jeffrey CFG, Hile SD, Pittman SJ, Monaco ME, Caldow C [Eds] (2013) Coral reef ecosystems of St. John, U.S. Virgin Islands: Spatial and temporal patterns in fish and benthic communities (2001–2009). NOAA Technical Memorandum 152. Silver Spring, MD, 150 pp. <https://repository.library.noaa.gov/view/noaa/789>
- García-Sais JR (2005) Inventory and Atlas of Corals and Coral Reefs, with Emphasis on Deep-Water Coral Reefs from the U. S. Caribbean EEZ Final Report; Caribbean Fishery Management Council San Juan, Puerto Rico, 214 pp. [http://sedarweb.org/docs/wsuff/S26\\_RD\\_02\\_deep\\_reefs\\_report\\_2005.pdf](http://sedarweb.org/docs/wsuff/S26_RD_02_deep_reefs_report_2005.pdf)
- García-Sais JR, Williams SM, Sabater-Clavell J, Esteves R, Carlo M (2014) Mesophotic benthic habitats and associated reef communities at Lang Bank, St. Croix, USVI. Final Report; Caribbean Fishery Management Council San Juan, Puerto Rico, 124 pp.
- Hanner R, Floyd R, Bernard A, Collette BB, Shivji M (2011) DNA barcoding of billfishes. *Mitochondrial DNA* 22(sup1, S1): 27–36. <https://doi.org/10.3109/19401736.2011.596833>
- Harms-Tuohy CA, Schizas NV, Appeldoorn RS (2016) Use of DNA metabarcoding for stomach content analysis in the invasive lionfish *Pterois volitans* in Puerto Rico. *Marine Ecology Progress Series* 558: 181–191. <https://doi.org/10.3354/meps11738>
- Knudsen SW, Clements KC (2016) World-wide species distributions in the family Kyphosidae (Teleostei: Perciformes). *Molecular Phylogenetics and Evolution* 101: 252–266. <https://doi.org/10.1016/j.ympev.2016.04.037>
- Lamkin JT, Gerard TL, Malca E, Shiroza A, Muhling BA, Davis N, Fuenmayor F, Whitecraft S, Johns L, Smith R, Melo N, Rawson G, Idrisi N, Smith T, Brown K (2009) USVI larval reef fish supply study: 2007-08 report. Coral Reef Conservation Program (U.S.). U.S. Dept. of Commerce, National Oceanic and Atmospheric Administration, Southeast Fisheries Science Center, Miami, 43 pp. <https://repository.library.noaa.gov/view/noaa/548>
- Loftus WF (2003) Inventory of fishes in inland fresh and brackish-water habitats of Virgin Island National Park. Final Report, U.S. Inventory and Monitoring Program 3207–24F29 VIIS-1102, 52 pp. <https://irma.nps.gov/DataStore/Reference/Profile/2175778>
- Nelson WR, Appeldoorn RS (1985) Cruise Report R/V Seward Johnson. A Submersible Survey of the Continental Slope of Puerto Rico and the U.S. Virgin Islands, 1–23 October 1985, 76 pp. [https://link.springer.com/chapter/10.1007/978-3-319-92735-0\\_7](https://link.springer.com/chapter/10.1007/978-3-319-92735-0_7)
- Pittman SJ, Hile SD, Jeffrey CFG, Caldow C, Kendall MS, Monaco ME, Hillis-Starr Z (2008) Fish assemblages and benthic habitats of Buck Island Reef National Monument (St. Croix, U.S. Virgin Islands) and the surrounding seascape: A characterization of spatial and temporal patterns. NOAA Technical Memorandum NOS NCCOS 71. Silver Spring, MD, 96 pp. [https://coastalscience.noaa.gov/data\\_reports/fish-assemblages-and-benthic-habitats-of-buck-island-reef-national-monument-st-croix-u-s-virgin-islands-and-the-surrounding-seascape-a-characterization-of-spatial-and-temporal-patterns-2/](https://coastalscience.noaa.gov/data_reports/fish-assemblages-and-benthic-habitats-of-buck-island-reef-national-monument-st-croix-u-s-virgin-islands-and-the-surrounding-seascape-a-characterization-of-spatial-and-temporal-patterns-2/)
- Puebla O, Coulmance F, Estape CJ, Morgan Estape A, Robertson DR (2022) A review of 263 years of taxonomic research on *Hypoplectrus* (Perciformes: Serranidae), with a redescription of *Hypoplectrus affinis* (Poey, 1861). *Zootaxa* 5093(2): 101–141. <https://doi.org/10.11646/zootaxa.5093.2.1>

- Quattrini AM, Demopoulos AWJ, Singer R, Roa-Varon A, Chaytor JD (2017) Demersal fish assemblages on seamounts and other rugged features in the northeastern Caribbean. Deep-sea Research. Part I, Oceanographic Research Papers 123: 90–104. <https://doi.org/10.1016/j.dsr.2017.03.009>
- Recksiek C, Wetherbee BM, DeAngelis B (2006) Assessment of the Status of Shark Populations in the USVI. Final Report. University of Rhode Island, Kingston, 22 pp. [https://www.ncei.noaa.gov/data/oceans/coris/library/NOAA/CRCP/project/1413/NA04NMF4630343\\_FinalReport\\_shark\\_usvi.pdf](https://www.ncei.noaa.gov/data/oceans/coris/library/NOAA/CRCP/project/1413/NA04NMF4630343_FinalReport_shark_usvi.pdf)
- Robertson DR, Tornabene L (2021) Reef-associated Bony Fishes of the Greater Caribbean: a Checklist (Version 4) [Data set]. Zenodo. <https://doi.org/10.5281/zenodo.5592149>
- Robertson DR, Estapé CJ, Estapé AM, Peña E, Tornabene L, Baldwin CC (2020) The marine fishes of St Eustatius Island, northeastern Caribbean: an annotated, photographic catalog. Zookeys 1007: 145–180. <https://doi.org/10.3897/zookeys.1007.58515>
- Robertson DR, Tornabene L, Lardizabal CC, Baldwin CC (2022) Submersibles greatly enhance research on the diversity of deep-reef fishes in the Greater Caribbean. Frontiers in Marine Science 8: e800250. <https://doi.org/10.3389/fmars.2021.800250>
- Rogers CS, Pietsch TW, Randall JE, Arnold RJ (2010) The Sargassum Frog-fish (*Histrio histrio* Linnaeus) Observed in Mangroves in St. John, U.S. Virgin Islands. Coral Reefs 29(3): 577. <https://doi.org/10.1007/s00338-010-0636-z>
- Rohmann SO, Hayes JJ, Newhall RC, Monaco ME, Grigg RW (2005) The area of potential shallow-water tropical and subtropical coral ecosystems in the United States. Coral Reefs 24(3): 370–383. <https://doi.org/10.1007/s00338-005-0014-4>
- Smith-Vaniz W, Jelks HL (2014) Marine and inland fishes of St. Croix, U. S. Virgin Islands: an annotated checklist. Zootaxa 3803: 1–120. <https://doi.org/10.11646/zootaxa.3803.1.1>
- Smith-Vaniz W, Jelks HL, Rocha LA (2006) Relevance of cryptic fishes in biodiversity assessments: A case study at Buck Island Reef National Monument, St. Croix. Bulletin of Marine Science 79: 17–48. <https://www.ingentaconnect.com/content/umrsmas/bull-mar/2006/00000079/00000001/art00002>
- Starck WA, Estapé CJ, Morgan Estapé A (2017) The fishes of Alligator Reef and environs in the Florida Keys: A half-century update. Journal of the Ocean Science Foundation 27: 74–117. <https://doi.org/10.5281/ZENODO.851651>
- Thacker CE (2009) Phylogeny of Gobioidae and placement within Acanthomorpha, with a new classification and investigation of diversification and character evolution. Copeia 2009(1): 93–104. <https://doi.org/10.1643/CI-08-004>
- Valdez-Moreno M, Vásquez-Yeomans L, Elías-Gutiérrez M, Ivanova NV, Hebert PDN (2010) Using DNA barcodes to connect adults and early life stages of marine fishes from the Yucatan Peninsula, Mexico: Potential in fisheries management. Marine and Freshwater Research 61(6): 655–671. <https://doi.org/10.1071/MF09222>
- Victor BC, Valdez-Moreno M, Vasquez-Yeomans L (2015) Status of DNA Barcoding Coverage for the Tropical Western Atlantic Shore-fishes and Reef Fishes. DNA Barcodes 3(1): 89–93. <https://doi.org/10.1515/dna-2015-0011>

Ward RD, Hanner R, Hebert PDN (2009) The campaign to DNA barcode all fishes, FISH-BOL. *Journal of Fish Biology* 74(2): 329–356. <https://doi.org/10.1111/j.1095-8649.2008.02080.x>

Weigt LA, Baldwin CC, Driskell A, Smith DG, Ormos A, Reyley EA (2012) Using DNA Barcoding to Assess Caribbean Reef Fish Biodiversity: Expanding Taxonomic and Geographic Coverage. *PLoS ONE* 7(7): e41059. <https://doi.org/10.1371/journal.pone.0041059>

## Supplementary material 1

### Plates S1–S18

Authors: D. Ross Robertson, Carlos J. Estapé, Allison M. Estapé, Lee Richter, Ernesto Peña, Benjamin Victor

Data type: images (jpg, images in ZIP arhiv)

Explanation note: Fishes of St. Croix (Plate S1), fishes of St. John-Thomas (Plates S2–S18).

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/zookeys.1103.83795.suppl1>

## Supplementary material 2

### File S1

Authors: D. Ross Robertson, Carlos J. Estapé, Allison M. Estapé, Lee Richter, Ernesto Peña, Benjamin Victor

Data type: image (jpg file)

Explanation note: Bathymetry of the US Virgin Islands.

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/zookeys.1103.83795.suppl2>

## Supplementary material 3

### File S2

Authors: D. Ross Robertson, Carlos J. Estapé, Allison M. Estapé, Lee Richter, Ernesto Peña, Benjamin Victor

Data type: GPS data (excel file)

Explanation note: File S2A: Georeferencing coordinates and site codes for dive sites of authors Carlos and Allison Estapé at St John, St Thomas and St. Croix during 2021. File S2B: Georeferencing coordinates and site codes for dive sites used by non-author photographers at St John-Thomas. File S2C: Names and emails of third party Citizen Scientists who provided voucher images of various St John-Thomas fishes.

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/zookeys.1103.83795.suppl3>

## Supplementary material 4

### File S3

Authors: D. Ross Robertson, Carlos J. Estapé, Allison M. Estapé, Lee Richter, Ernesto Peña, Benjamin Victor

Data type: GPS data (kmz. file)

Explanation note: KMZ file of USVI dive sites.

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/zookeys.1103.83795.suppl4>

## Supplementary material 5

### File S4

Authors: D. Ross Robertson, Carlos J. Estapé, Allison M. Estapé, Lee Richter, Ernesto Peña, Benjamin Victor

Data type: occurrences (excel file)

Explanation note: File S4. Native marine fish faunas of St. John-Thomas and St. Croix.

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/zookeys.1103.83795.suppl5>

## Supplementary material 6

### File S5

Authors: D. Ross Robertson, Carlos J. Estapé, Allison M. Estapé, Lee Richter, Ernesto Peña, Benjamin Victor

Data type: occurrences (excel file)

Explanation note: Ecological Characteristics of Reef-Associated Bony Fishes from St John-Thomas. See Methods of paper for details.

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/zookeys.1103.83795.suppl6>

## Supplementary material 7

### File S6

Authors: D. Ross Robertson, Carlos J. Estapé, Allison M. Estapé, Lee Richter, Ernesto Peña, Benjamin Victor

Data type: genomic (excel file)

Explanation note: File S6: mtDNA-Barcode information for fishes from islands on the Puerto Rico Platform (St John-Thomas, Puerto Rico and the British Virgin Islands) and St. Croix. For coding of differently colored highlighting see bottom of table. For explanations of "Reef Associated" and "Deep" see main text.

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/zookeys.1103.83795.suppl7>