

# A new species and three newly recorded species of Tetrastichinae (Hymenoptera, Eulophidae) from China

Wen-Jian Li<sup>1,2</sup>, Cheng-De Li<sup>3</sup>

**1** Jiangsu Provincial Key Laboratory of Coastal Wetland Bioresources and Environmental Protection, School of Wetland, Yancheng Teachers University, Yancheng, 224007, China **2** Jiangsu Key Laboratory for Bioresources of Saline Soils, School of Wetland, Yancheng Teachers University, Yancheng, 224007, China **3** School of Forestry, Northeast Forestry University, Harbin, 150040, China

Corresponding author: Cheng-De Li ([lichengde0608@sina.com](mailto:lichengde0608@sina.com))

Academic editor: Zachary Lahey | Received 22 July 2022 | Accepted 28 October 2022 | Published 24 November 2022

<https://zoobank.org/17D6DA06-6B46-426E-8363-5CB93EF4E207>

**Citation:** Li W-J, Li C-D (2022) A new species and three newly recorded species of Tetrastichinae (Hymenoptera, Eulophidae) from China. ZooKeys 1131: 197–215. <https://doi.org/10.3897/zookeys.1131.90688>

## Abstract

Five species of five genera in Tetrastichinae (Hymenoptera, Eulophidae) from China are reviewed, including one new species, *Mestocharella qingdaoensis* **sp. nov.**, and three new country record species: *Nesolynx thymus* (Girault, 1916), *Holcotetrastichus rhusaces* (Walker, 1839), and *Peckelachertus diprioni* Yoshimoto, 1970. New distributional data for *Ceratoneura indi* Girault, 1917 are provided.

## Keywords

Chalcidoidea, genera, parasitoids, taxonomy

## Introduction

The subfamily Tetrastichinae (Hymenoptera, Eulophidae) is one of the largest groups of Chalcidoidea (Graham 1987; LaSalle 1994). Species are distributed in almost all geographic realms and play a vital role in terrestrial ecosystems (Graham 1987; LaSalle 1994). Most species of Tetrastichinae are parasitic; they attack species from approximately 1000 families in 10 different orders of Insecta (Graham 1987; LaSalle 1994). Also, some species, such as *Leptocybe invasa* Fisher and LaSalle, 2004, are phytophagous and live in galls produced by their hosts.

Unfortunately, Chinese species of Tetrastichinae are poorly investigated compared to other countries and regions (Kostjukov 1978, 2000; Graham 1987, 1991; Bouček 1988; LaSalle 1994; Narendran 2007). In the early stage, foreign entomologists reported several Tetrastichinae from Guangdong, Macao, and Taiwan in China (Perkins 1912; Timberlake 1921; Miwa and Sonan 1935). With more research on parasitic wasps, Chinese entomologists realized the importance of this faunal group: Liao et al. (1984) reported 201 economically important insect species of China including eight species of Tetrastichinae; Yang (1996) systematically investigated parasitic wasps on bark beetles from China and reported 141 species including 16 species of Tetrastichinae; Zhu and Huang (2001) investigated Eulophidae from Zhejiang province and reported 12 species of Tetrastichinae; Zhu and Huang (2002) investigated Eulophidae from Guangxi province and reported 23 species of Tetrastichinae; Zhang et al. (2007) investigated Eulophidae from south Gansu and Qinlin Mountains and reported 14 species of Tetrastichinae; Yang et al. (2015) systematically investigated parasitic wasps on forest defoliators, reporting 115 species including 20 species belonging to four genera of Tetrastichinae. Subsequently, there are many more reports of new species and records of Tetrastichinae (Yang 1989; Sheng and Wang 1992, 1993; Sheng 1995; Sheng and Zhao 1995; Sheng and Zhu 1998; Yang and Wei 2003; Jiao et al. 2006; Zhang et al. 2009; Wu et al. 2009; Li et al. 2014; Yang et al. 2014; Feng et al. 2016; Li et al. 2016; Song et al. 2017; Li and Li 2020, 2021; Song et al. 2020; Guo et al. 2022; Ning et al. 2022). In terms of Tetrastichinae species richness, there is an obvious imbalance among provinces of China. Most southern provinces have more species than northern provinces, such as 27 species in Guangxi Province compared with just two species in Ningxia Province. Therefore, there is still much to study, and knowledge to be gained, about this group in China.

## Materials and methods

Specimens were collected by sweep netting and yellow-pan trapping. They were preserved and were dissected and mounted in Canada balsam following the method of Noyes (1982), or fixed on triangular cards. Photographs were taken with a digital CCD camera attached to an Olympus BX51 compound microscope and a AOSVI HK-830 microscope. Most measurements were made from slide-mounted specimens using an eye-piece reticule with an Olympus CX21 microscope. Terminology follows Gibson et al. (1997) and the following abbreviations are used:

- F1–4** (flagellomeres 1–4);
- POL** (minimum distance between lateral ocelli);
- OOL** (minimum distance between lateral ocellus and eye margin);
- OD** (longest diameter of a lateral ocellus);
- MV** (marginal vein);
- STV** (stigmatal vein);
- SMV** (submarginal vein);
- PMV** (postmarginal vein).

All the specimens listed below are deposited in Northeast Forestry University (NEFU), Harbin, China.

## Species accounts

The genus *Mestocharella* (Eulophidae, Tetrastichinae) was erected by Girault (1913) with *Mestocharella feralis* Girault, 1913 as the type species. It is a small genus with 12 valid species worldwide (Noyes 2019) and only one species occurring in China, *M. javensis* (Kamijo 1994). Because the propodeum is different from the propodeum of all the other species included in *Mestocharella*, *M. deltoidea* Khan, Agnihotri & Sushil, 2005 and *M. indica* Jaikishan Singh & Khan, 1995 probably do not belong to the genus (Narendran 2007).

*Mestocharella* is a unique genus and can be distinguished from Tetrastichinae by the following characteristics: malar sulcus present; antenna slender, one anellus, funicle with four segments and clava bi-segmented in female; funicle 4-segmented and clava 3-segmented in male; pronotum long, collar with or without transverse carina; axillae not so advanced; dorsellum with a median carina; propodeum long, with a large sub-pentagonal area; spiracles small; gastral petiole conspicuous, strongly carinate; gaster usually shorter than mesosoma.

The species of *Mestocharella* can be divided into three species groups: the *kumatai*, *feralis*, and *javensis* groups (Kamijo 1994). The species of *Mestocharella* are parasitic on Lepidoptera (Bouček 1988; Kamijo 1994).

### *Mestocharella qingdaoensis* sp. nov.


<https://zoobank.org/BA195019-145C-4A7E-838E-BAEB094AF9FB>

Figs 1–4

**Type material.** *Holotype*, female [on card], CHINA, Shandong Province, Qingdao City, Mount Xiao Zhu, 18–20.V.2014, Guo-Hao Zu, Si-Zhu Liu, by yellow pan trapping (deposited in NEFU). *Paratypes*, 1 female [on slide], same data as holotype (deposited in NEFU).

**Diagnosis.** Female. Body mainly brownish, head and posterior half of mesoscutum and axillae yellow; propodeum median carina not forked anteriorly; plicae distinct but not connected with median carina; forewing SMV with three dorsal setae, MV 6.9–7.3× as long as STV. *Mestocharella qingdaoensis* belongs to the *kumatai* group (Kamijo 1994) in that the pronotal collar is without transverse carina, and it is similar to *M. kumatai* Kamijo, 1994. However, it can be separated from *M. kumatai* by the following characteristics: head yellow (vs blackish); mid-lobe of mesoscutum without median line (vs vague); median carina of propodeum not forked anteriorly (vs always forked); plicae distinct but not connected with median carina (connected by anterior oblique carinae); forewing SMV with three dorsal setae (vs five).

**Description. Female.** Body length 1.8–1.9 mm, mainly yellow (Figs 1, 2). Head yellow, eyes deep reddish brown, ocellus yellowish white; antenna scape yellowish, pedi-



**Figures 1–4.** *Mestocharella qingdaoensis* sp. nov., holotype, female **1** habitus, dorsal view **2** habitus, lateral view. Scale bars: 500  $\mu$ m **3** antenna, lateral view **4** fore and hind wings, dorsal view. Scale bars: 100  $\mu$ m.

cel and flagellum yellowish brown. Metasoma mainly brownish with posterior half of mesoscutum and axillae yellow; wings hyaline, venation yellowish brown; legs yellow, tarsomere IV of all legs dark brown. Mesosoma brownish with basal 1/3 yellowish brown.

Head in dorsal view, nearly as broad as mesosoma,  $2.5\text{--}2.6\times$  ( $2.5\times$ ) as broad as long; vertex with setae shorter than OD, POL  $1.3\times$  OOL, OOL  $2. \times$  OD. Face depressed slightly, without median line; torulus with lower edge above the ventral edge of eyes; eyes separated by  $1.45\times$  their height. Malar sulcus present; malar space  $0.6\times$  as long as eye height. Mouth cavity  $1.4\times$  as wide as malar space; clypeus with anterior margin bidentate; mandible tridentate. Antenna (Fig. 3) scape  $5\times$  as long as broad with shorter setae on dorsal and ventral side; two anelli, first anellus slightly transverse, second anellus lamellar; pedicel  $2.2\text{--}2.3\times$  as long as broad, shorter than F1; F1–F4:  $3.0\times$ ,

3.4×, 3.2×, 2.3× as long as broad respectively; clava 3.2× as long as broad, ca as broad as F3, bi-segmented; flagellum with long whorled setae.

Metasoma relatively long, 1.7–1.8× (1.8×) as long as broad. Pronotum subconical, 3.15× as broad as long, ~0.6× as long as mid-lobe of mesoscutum; collar rounded anteriorly and without transverse carina. Mid-lobe of mesoscutum with extremely fine reticulation; without median line; 3 adnotaular setae in one row on each side. Scutellum ca. as long as broad; submedian grooves shallow but distinct enclosing a space ~2.9× as long as broad, sublateral grooves distinct without weak costulae; anterior setae situated before middle distinctly. Dorsellum ~3× as broad as long, with a weak median carina. Propodeum subpentagonal area broad, smooth, without reticulation, median carina distinct and thin, not forked anteriorly; plicae distinct but not connecting with median carina; spiracle small, circular; callus with 2 setae. Forewing (Fig. 4) 2.2× as long as broad, SMV with 3 dorsal setae; costal cell shorter than MV, MV 6.9–7.3× (7.3×) as long as STV with front edge 12–15 setae; STV short with a long uncus; speculum small, nearly closed posteriorly, subcubital line of setae not reaching to distal edge of speculum. Legs slender, spur of metatibia 0.5× as long as length of metabasitarsus.

Gastral petiole long with several transverse weak carinae anteriorly and 3 or 4 longitudinal strong carinae. Gaster 1.2–1.4× as long as broad, shorter than mesosoma; ovipositor 0.5× as long as gaster and slightly exerted at apex of gaster, tip of hypopygium situated at basal 4/5 of gaster.

**Male.** Unknown.

**Host.** Unknown.

**Distribution.** China (Shandong).

**Etymology.** The epithetic *qingdao* refers to the place where the species collected.

### *Nesolynx* Ashmead, 1905

**Note.** The genus *Nesolynx* was erected by Ashmead (1905) with *Nesolynx flavipes* Ashmead, 1905 as the type species. Bouček (1988) proposed *Aceratoneurella* Girault, 1917, *Ceratotrastichus* Girault & Dodd, 1913, and *Omphalomomyia* Girault, 1913 as synonyms of *Nesolynx*. It is a characteristic genus with 17 species recorded worldwide (Noyes 2019), but only one species, *Nesolynx thymus* (Girault, 1916), is found in China. It is distributed in tropical and subtropical countries, in the warmer parts of the temperate zones of Europe, Africa, Asia, Australia, and the Pacific islands (Bouček 1988). It can be distinguished from Tetrastichinae particularly by the mid-lobe of mesoscutum bearing dense setae and without a median line (Bouček 1988). The species are parasitoids of various groups of Diptera and Lepidoptera (Bouček 1988).

### *Nesolynx thymus* (Girault, 1916), new record from China

Figs 5–10

*Omphalomomyia thymus* Girault, 1916: 485.

*Omphalomomyia thymus javae* Girault, 1917: 7 (subspecies). [Synonymized by Bouček 1977: 404].

*Buonaparteia aeniceps* Girault, 1924: 5. Syntypes. [Synonymized by Bouček 1988: 697].

*Syntomosphyrum obscuriceps* Ferrière, 1940: 138. [Synonymized by Bouček 1977: 404].

*Omphalomomyia* [sic] *thymus*: Thompson 1955: 292.

*Nesolynx thymus*: Bouček 1977: 404.

**Material examined.** 7 females: [1 female on slide], Henan Province, Xinyang City, Mount Yan, Temple Xianyin, 6–7.VIII.2015, Hui Geng, Zhi-Guang Wu, Yan Gao, by yellow pan trapping; [1 female on slide], Hainan Province, Changjiang County, Mount Bawanglin, 15–17.V.2019, Wen-Jian Li, Jun Wu, by yellow pan trapping; [1 female on slide], Hainan Province, Haikou City, Hainan University, 27–29.VI.2019, Yu-Ting Jiang, by yellow pan trapping; [2 females on cards], Yunnan Province, Yuanjiang County, 26–28.XI.2020, Jun Wu, Jun-Jie Fan; Ming-Rui Li, Gang Fu, by yellow pan trapping; [2 females on cards], Yunnan Province, Shuangjiang County, 21.IV.2013, Xiang-Xiang Jin, Guo-Hao Zu, Chao Zhang, by sweeping. (All deposited in NEFU).


**Diagnosis. Female.** Body mainly yellow (Figs 5, 6); upper face, vertex, gena, and occiput dark green with metallic reflections, lower face yellow (Fig. 7); gaster yellow with black sides. Mesosoma with dense setae on mid-lobe of mesoscutum, especially a pair of long black setae posteriorly similar to setae on scutellum; propodeum with median carina distinct, cup-shaped. Gaster 1.6–1.8× as long as broad.

**Male.** Unknown.

**Hosts.** Not known from China. Non-Chinese records include *Musca domestica* Linnaeus, 1758, *Exorista bombycis* (Louis, 1880), *Bombyx mori* Linnaeus, 1758 (Bansude et al. 2010), *Argyrophylax leefmansii* Baranov, 1933, *Bessa remota* (Aldrich, 1925), *Chaetogena bezziana* Baranov, 1934, *Nephantis serinopa* Meyrick, 1905, *Artona catoxantha* Hampson, 1892 (Herting 1978), *Exorista sorbillans* (Wiedemann, 1830) (Kumar et al. 1991), *Ptychomyia remota* Aldrich, 1925, *Cadurcia leefmansii* Baranov, 1933 (Lever 1964), *Zaratha* sp. (Bouček 1988), *Sturmiopsis inferens* Townsend, 1916,



**Figures 5–7.** *Nesolynx thymus* (Girault), female **5** habitus, dorsal view **6** habitus, lateral view. Scale bars: 500  $\mu$ m **7** head, frontal view. Scale bar: 200  $\mu$ m.



**Figures 8–10.** *Nesolynx thymus* (Girault), female **8** legs, lateral view, from left to right: fore, mid, and hind legs **9** antenna, lateral view **10** fore and hind wings, dorsal view. Scale bars: 100  $\mu\text{m}$ .

*Chilo auricilius* Dudgeon, 1905 (Varma 1989), *Cnaphalocrocis medinalis* (Guenée, 1854) (Talgeri and Dalaya 1971), *Maruca testulalis* (Geyer, 1832), Tachinidae unspecified sp. (Narendran 2007), *Apanteles artonae* (Rehwer, 1926) (Herting 1977).

**Distribution.** China (Henan, Yunnan, Hainan); Bangladesh (Rahman 1989), Myanmar (Husain and Khan 1986), Indonesia (Bouček 1988), Malaysia (Lever 1964), India, and Sri Lanka (Narendran 2007).

**Comments.** The species can be easily identified by the unique color of head.

### *Holcotetrastichus* Graham, 1987

**Note.** This is a small genus erected by Graham (1987), with *Cirrospilus rhosaces* Walker, 1839 as the type species. Only two species have been described: *Holcotetrastichus manaliensis* Graham, 1991 and *Holcotetrastichus rhosaces* (Walker, 1839). It can be distinguished from other Tetrastichinae especially by the strong transverse costulae in deep broad sublateral grooves and the hypopygium reaching nearly the tip of the gaster (Graham 1987). The species are parasitoids of some species of *Cassida* (Coleoptera, Chrysomelidae) (Graham 1991).

### *Holcotetrastichus rhosaces* (Walker, 1839), new record from China

Figs 11–20

*Cirrospilus rhosaces* Walker, 1839: 293.

*Cirrospilus racilla* Walker, 1839: 312. [Synonymised by Graham 1961: 37].

*Tetrastichus racilla*: Walker 1848: 149.


*Tetrastichus rhosaces*: Walker 1848: 147.

*Aprostocetus rhosaces*: Graham 1961: 37.


*Holcotetrastichus rhosaces*: Graham 1991: 272; Narendran 2007: 120.

*Holcotetrastichus rhosaceus* [sic]: Boyadzhiev 2000: 27.


**Material examined.** 9 females and 2 males: [2 females on slides], Liaoning Province, Anshan City, Mount Qianshan, 23.VI.2013, Hui Geng, Zhi-Guang Wu, Yan Gao, Si-Zhu Liu, by sweeping; [1 female on slide], Jiangxi Province, Yichun City, Mount Guanshan, 22–24.VIII.2018, Xiang-Xiang Jin, Wang-Ming Li, by yellow-pan trapping; [1 female and 1 male on slides, 1 female and 1 male on cards], Qinghai Province, Prefecture Huangnan, Forestry Station Maixiu, 26–29.VIII.2019, Ming-Rui Li, by yellow pan trapping; [2 females on cards], Jinlin Province, County Wangqing, Forestry



**Figure 11.** *Holcotetrastichus rhosaces* (Walker), female, habitus, lateral view. Scale bar: 500  $\mu\text{m}$ .



**Figures 12–17.** *Holcotetrastichus rhosaces* (Walker), female **12** head, frontal view **13** antenna, lateral view **14** mesosoma, dorsal view **15** fore and hind wings, dorsal view **16** metasoma, ventral view **17** legs, lateral view, from left to right: fore, mid, and hind legs. Scale bars: 100  $\mu\text{m}$ .



**Figures 18–20.** *Holcotetrastichus rhosaces* (Walker), male **18** antenna, lateral view **19** fore and hind wings, dorsal view **20** legs, lateral view, from left to right: fore, mid, and hind legs. Scale bars: 100  $\mu$ m.

Station Qinhe, 8.VII.2013, Ye Chen, Zhi-Guang Wu, by sweeping; [2 females on cards], Heilongjiang Province, City Heihe, Park Beishang, 22.VII.2020, Ming-Rui Li, by sweeping. (All deposited in NEFU).

**Diagnosis. Female.** Body black, with weak metallic reflections (Fig. 11). Mesosoma (Fig. 14) with mid-lobe of mesoscutum weakly reticulate, 2 adnotaular setae in single row on each side, median line indicated only posteriorly; sublateral grooves of scutellum deep and broad with strong transverse costulae, submedian grooves rather weak. Forewing (Fig. 15) broad, 2.0 $\times$  as long as broad, SMV with 2 dorsal setae, MV

2.8–3.2 times length of STV, PMV distinctly short. Gaster (Fig. 16) with hypopygium almost reaching tip of gaster.

**Male.** Antenna (Fig. 18) with scape broad, ventral plaque 0.7 length of scape; F1 shorter than F2; each segment of funicle with whorl setae reaching well beyond the tip of the segment.

**Hosts.** Unknown from China. Non-Chinese records include *Cassida deflorate* Sufrian, 1844, *Cassida murraea* Linnaeus, 1767, *Cassida nebulosa* Linnaeus, 1758, *Cassida nobilis* Linnaeus, 1758, *Cassida rubiginosa* Mueller, 1776, *Cassida viridis* Linnaeus, 1758, *Cassida vittate* Villers, 1789 (Graham 1991), *Cassida piperata* Hope, 1842 (Nagasawa et al. 2003).

**Distribution.** China (Heilongjiang, Liaoning, Jilin, Qinghai, Jiangxi); Austria, Czech Republic, Czechoslovakia, France, Germany, Hungary, Ireland, Italy, Moldova, Romania, Switzerland, United Kingdom (Graham 1991), Bulgaria (Boyadzhiev 2000), Netherlands (Gijswijt 2003), Poland (Domenichini 1966), Russia (Yegorenkova et al. 2007), Sweden (Hansson 1991), Japan (Ikeda 1997), and United States of America (Boyadzhiev 2000).

**Comments.** Most species we collected had weak metallic reflections compared to the species reported by Graham (1991).

### *Peckelachertus* Yoshimoto, 1970

**Note.** This is a small genus with only two known species worldwide (Noyes 2019): *P. diprioni* Yoshimoto, 1970 and *P. anglicus* Graham, 1977. Both of these were transferred from the subfamily Elachertinae to Tetrastichinae by Graham (1977). The genus can be distinguished from other Tetrastichinae especially by having the PMV equally or nearly as long as STV and scutellum without submedian grooves (Graham 1977).

### *Peckelachertus diprioni* Yoshimoto, 1970, new record from China


Figs 21–26

*Peckelachertus diprioni* Yoshimoto, 1970: 909.

*Peckelachertus diprioni*: Graham 1977: 47.

**Material examined.** 2 females. [2 females on slides], China, Heilongjiang Province, Shangzhi City, Mount Laoyeling, 9.VII.2015, Ye Chen, Chao Zhang, by sweeping.

**Diagnosis. Female.** Body dark brown, without metallic reflections. Head with anterior margin of clypeus truncate, without any teeth, malar sulcus present and distinct. Antenna with pedicel 1.8–1.9× as long as broad, F11.6× as long as broad. Mesosoma (Fig. 14) 1.5× as long as broad, mid-lobe of mesoscutum with 2 adnotaular setae in single row on each side, median line absent; scutellum submedian grooves absent or indicated at posterior half, anterior pair of setae situated near anterior margin of scutel-



**Figures 21–26.** *Peckelachertus diprioni* Yoshimoto, female **21** head, frontal view **22** antenna, lateral view **23** mesosoma, dorsal view **24** fore and hind wings, dorsal view **25** metasoma, ventral view **26** legs, lateral view, from bottom to top: fore, mid, and hind legs. Scale bars: 100  $\mu$ m.

lum. Forewing (Fig. 15), 2.2 $\times$  as long as broad, SMV with 4 dorsal setae, the length of PMV as long as STV.

**Male.** Unknown from Chinese material.

**Hosts.** Unknown from China. Non-Chinese records include *Gilpinia frutetorum* (Fabricius, 1793) (LaSalle 1994), *Gilpinia pallida* (Klug, 1812) (Graham 1977).

**Distribution.** China (Heilongjiang); Finland, Canada (Graham 1977).

**Comments.** Graham (1977) pointed out that Yoshimoto's description of genus *Peckelachertus* and of its type species *P. diprioni* are not correct in some respects and proposed some remarks after examining material. Our specimens agree well with the remarks by Graham (1977).

### *Ceratoneura* Ashmead, 1849

**Note.** The genus *Ceratoneura* was erected with *Ceratoneura petiolata* Ashmead, 1894 as the type species by subsequent designation of Ashmead (1904). Ikeda (2001) revised of the world species of *Ceratoneura* in detail, describing five new species and redescribing six known species. It is a small genus with 12 species recorded worldwide (Noyes 2019), but only one species *Ceratoneura indi* Girault, 1917 has been reported from China (Ikeda 2001). This genus can be distinguished from other Tetrastichinae especially by the strongly sclerotized body and the face with conspicuous striae radiating from the mouth. The species are parasitoids of various groups of Diptera and Lepidoptera (Bouček 1988).


### *Ceratoneura indi* Girault, 1917

Figs 27–31


*Ceratoneura indi* Girault, 1917: 10.

*Ceratoneura indica* Rohwer, 1921: 127. [Synonymized by Bouček 1988: 670].

**Material examined.** 7 females: [1 female on slide and 2 females on cards], China, Zhejiang Province, County Panan, Mount Dapan, 30.VI.–2.VII.2019, Jun Wu, Jun-Jie Fan, by yellow pan trapping; [4 females on cards], China, City Chongqing, Mount



**Figures 27, 28.** *Ceratoneura indi* Girault, female **27** habitus, lateral view. Scale bar: 500  $\mu$ m **28** head, frontal view. Scale bar: 100  $\mu$ m.



**Figures 29–34.** *Ceratoneura indi* Girault, female **29** head, frontal view **30** antenna, lateral view **31** mesosoma, dorsal view **32** fore and hind wings, dorsal view **33** metasoma, ventral view **34** legs, lateral view, from left to right: fore, mid, and hind legs. Scale bars: 100  $\mu\text{m}$ .

Simianshan, Village Hongdong, 26.VII.2019, Ting-Ting Zhao, Shu-Cheng Deng, by sweeping. (All deposited in NEFU).

**Diagnosis. Female.** Body black, strongly sclerotized. Face with conspicuous striae radiating from mouth, torulus with lower margin distinctly above the level of ventral margin of eyes. Mesosoma with mid-lobe of mesoscutum weakly reticulate, 4 or 5 adnotaular setae in single row on each side, median line absent. Forewing  $2.2\text{--}2.3\times$  as long as broad, SMV with 3 dorsal setae, speculum large. Petiole distinct,  $0.4\text{--}0.5\times$  as long as propodeum. Gaster  $1.7\text{--}2.0\times$  as long as broad.

**Male.** Unknown for Chinese material.

**Hosts.** Unknown from China. Non-Chinese records include *Asphondylia sphaera* Monzen, 1937 (Ikeda 2001).

**Distribution.** China (Zhejiang, Chongqing, Hong Kong), Japan, India, Malaysia, New Caledonia, Papua New Guinea, Sri Lanka.

**Comments.** Ikeda (2001) reported only one specimen from Hong Kong, and we add seven additional specimens from Zhejiang and Chongqing, which are new locality records for China.

## Acknowledgements

We are grateful to Dr Xiang-Xiang Jin, Dr Si-Zhu Liu, Dr Hui Geng, Dr Guo-Hao Zu, Dr Ye Chen, Dr Ming-Rui Li, Miss Ting-Ting Zhao, Miss Shu-Cheng Deng, Miss Yan Gao, Mr Jun Wu, Mr Jun-Jie Fan, Mr Gang Fu, and Mr Chao Zhang for specimen collections.

## References

- Ashmead WH (1904) Classification of the chalcid flies of the superfamily Chalcidoidea, with descriptions of new species in the Carnegie Museum, collected in South America by Herbert H. Smith. *Memoirs of the Carnegie Museum* 1(4): i–xi, 225–551. [39 pls] <https://doi.org/10.5962/p.234821>
- Ashmead WH (1905) Additions to the recorded hymenopterous fauna of the Philippine Islands, with descriptions of new species. *Proceedings of the United States National Museum* 28(1413): 957–971. <https://doi.org/10.5479/si.00963801.28-1413.957>
- Bansude V, Berde V, Chaphalkar S (2010) Emergence potential of *Nesolynx thymus* Girault and environmental impact. *Karnataka Journal of Agricultural Sciences* 23(1): 115–117.
- Bouček Z (1977) Taxonomic studies on some Eulophidae (Hym.) of economic interest mainly from Africa. *Entomophaga* 21(4): 401–414. <https://doi.org/10.1007/BF02371639>
- Bouček Z (1988) Australasian Chalcidoidea (Hymenoptera). A Biosystematic Revision of Genera of Fourteen Families, with a Reclassification of Species. CAB International, Wallingford /Cambrian News, Aberystwyth, 832 pp.

- Boyadzhiev P (2000) Unknown species of family Eulophidae to the fauna of Bulgaria from the Rhodopes (Hymenoptera: Chalcidoidea). *Acta Zoologica Bulgarica* 52(2): 25–29.
- Domenichini G (1966) Hym. Eulophidae. Palaearctic Tetrastichinae. Index of Entomophagous Insects, Vol. 1 (Eds: Delucchi V, Remaudière G) Le François, Paris, 101 pp.
- Feng MX, Cao HX, Hao HH, Wang W, Cheng LS (2016) A new record species of *Quadrastichus* (Hymenoptera: Eulophidae) from China. *Redai Zuowu Xuebao* 37(3): 582–585.
- Ferrière C (1940) On some parasites and hyperparasites of *Artona catoxantha*, Hmps. *Bulletin of Entomological Research* 31(2): 131–139. <https://doi.org/10.1017/S0007485300004910>
- Gibson GAP, Huber JT, Woolley JB (1997) Annotated keys to the genera of Nearctic Chalcidoidea (Hymenoptera). National Research Council Research Press, Ottawa, 794 pp.
- Gijswijt MJ (2003) Naamlijst van de Nederlandse bronswespen (Hymenoptera: Chalcidoidea). *Nederlandse Faunistische Mededelingen* 18: 17–79.
- Girault AA (1913) Australian Hymenoptera Chalcidoidea – IV. *Memoirs of the Queensland Museum* 2: 140–296. <https://doi.org/10.5962/bhl.title.9562>
- Girault AA (1916) New Javanese chalcidoid Hymenoptera. *Proceedings of the United States National Museum* 51(2161): 479–485. <https://doi.org/10.5479/si.00963801.2161.479>
- Girault AA (1917) New Javanese Hymenoptera. Private publication, Washington, 12 pp.
- Graham MWR de V (1961) The genus *Aprostocetus* Westwood *sensu lato* (Hym., Eulophidae) notes on the synonymy of European species. *Entomologist's Monthly Magazine* 97: 34–64.
- Graham MWR de V (1977) Systematic position of *Peckelachertus* Yoshimoto (Hym., Eulophidae) and description of a new species from Britain. *Systematic Entomology* 2(1): 45–47. <https://doi.org/10.1111/j.1365-3113.1977.tb00356.x>
- Graham MWR de V (1987) A reclassification of the European Tetrastichinae (Hymenoptera: Eulophidae), with a revision of certain genera. *Bulletin of the British Museum (Natural History) [Entomology]* 55(1): 1–392.
- Graham MWR de V (1991) A reclassification of the European Tetrastichinae (Hymenoptera: Eulophidae): revision of the remaining genera. *Memoirs of the American Entomological Institute* 49: 1–322.
- Guo SH, Kang N, Li Q, Hu HY (2022) Species and identification of *Dzhanokmenia* in China (Hymenoptera: Eulophidae). *Sichuan Journal of Zoology* 41(1): 63–73.
- Hansson C (1991) A catalogue of Chalcidoidea described by C.G. Thomson, with a checklist of Swedish species. *Entomologica Scandinavica (Supplement No 38)*: 1–70.
- Herting B (1977) A Catalogue of Parasites and Predators of Terrestrial Arthropods. Section A. Host or Prey/Enemy. Volume IV. Hymenoptera. Commonwealth Agricultural Bureaux, Institute of Biological Control, Farnham Royal, iii + 206 pp.
- Herting B (1978) A Catalogue of Parasites and Predators of Terrestrial Arthropods. Section A. Host or Prey/Enemy. Volume V. Neuroptera, Diptera, Siphonaptera. Commonwealth Agricultural Bureaux, Commonwealth Institute of Biological Control, Farnham Royal, 1–156.
- Husain T, Khan MY (1986) Family Eulophidae. In: Subba Rao BR, Hayat M (Eds) *The Chalcidoidea (Insecta: Hymenoptera) of India and the adjacent countries*. *Oriental Insects* 20: 211–245. <https://doi.org/10.1080/00305316.1986.10433730>
- Ikeda E (1997) Three species of Tetrastichinae (Hymenoptera, Eulophidae), newly recorded from Japan. *Japanese Journal of Entomology* 65(1): 186–192.

- Ikeda E (2001) A revision of the world species of *Ceratoneura* Ashmead (Hymenoptera, Eulophidae). *Insecta Matsumurana* 58: 27–50.
- Jiao Y, Chen ZL, Yu DJ, Kang L, Yang WD (2006) A new record genus and new record species of Eulophidae (Hymenoptera) in continental China. *Entomotaxonomia* 28(1): 69–74.
- Kamijo K (1994) A revision of *Mestocharella* (Hymenoptera, Eulophidae), with descriptions of six new species. *Japanese Journal of Entomology* 62(4): 747–762.
- Kostjukov VV (1978) Hymenoptera II. Chalcidoidea 13. Eulophidae (Tetrastichinae). *Opredeliteli Nasekomykh Evropeyskoy Chasti SSR* 3: 430–467.
- Kostjukov VV (2000) Nadsem. Chalcidoidea 46. Sem. Eulophidae. *Opredeliteli Nasekomikh Dalinego Vostoka Rossii* 4(4): 582–601.
- Kumar P, Kishore R, Jayaprakas CA, Sengupta K (1991) Parasitoids of uzi fly, *Exorista sorbillans* Wiedermann (Diptera: Tachinidae), XII: studies on the efficiency of *Nesolynx thymus* (Girault) at the field level. *Indian Journal of Sericulture* 30(2): 161–163.
- LaSalle J (1994) North American genera of Tetrastichinae (Hymenoptera: Eulophidae). *Journal of Natural History* 28(1): 109–236. <https://doi.org/10.1080/00222939400770091>
- Lever RJAW (1964) Notes on some parasites, hyperparasites and predators of coconut pests in Malaya. *FAO Plant Protection Bulletin* 12(2): 42–43.
- Li WJ, Li CD (2020) A new species of *Oomyzus* Rondani (Hymenoptera, Eulophidae) and first record of *O. gallerucae* (Fonscolombe) from China, with a key to Chinese species. *ZooKeys* 950: 41–49. <https://doi.org/10.3897/zookeys.950.48795>
- Li WJ, Li CD (2021) Two new species of *Neotrichoporoides* Girault (Hymenoptera, Eulophidae) from China and a key to Chinese species. *ZooKeys* 1023: 61–79. <https://doi.org/10.3897/zookeys.1023.61580>
- Li XX, Xu ZJ, Zhu CD, Zhao JN, He YY (2014) A new phytophagous eulophid wasp (Hymenoptera: Chalcidoidea: Eulophidae) that feeds within leaf buds and cones of *Pinus massoniana*. *Zootaxa* 3753(4): 391–397. <https://doi.org/10.11646/zootaxa.3753.4.8>
- Li Q, Wang C, Hu HY, Kostjukov VV, LaSalle J, Zhu CD (2016) Descriptions of three new species of *Dzhanokmenia* (Hymenoptera: Eulophidae) from China. *Zootaxa* 4121(4): 447–457. <https://doi.org/10.11646/zootaxa.4121.4.5>
- Liao DX, Li XL, Pang XF, Cheng TL (1984) Economic Insect Fauna of China. Fasc. 34. Hymenoptera: Chalcidoide (1). Science Press, Beijing, 241 pp.
- Miwa Y, Sonan J (1935) Description of a new egg parasite of *Melanauster chinensis* Först. from Formosa. *Transactions of the Natural History Society of Formosa* 25(146): 406–407.
- Nagasawa A, Konno Y, Matsuda K (2003) New record of *Holcotetrastichus rhesaces* (Walker) (Hymenoptera: Eulophidae), a hymenopterous parasitoid of *Cassida nebulosa* L. and *C. piperata* Hope (Coleoptera: Chrysomelidae), from Japan. *Japanese Journal of Entomology* 6(3): 117–118. [New Series]
- Narendran TC (2007) Indian chalcidoid parasitoids of the Tetrastichinae (Hymenoptera: Eulophidae). *Records of the Zoological Survey of India, Occasional Paper* 272: 1–390.
- Ning HF, Cao HX, Zhu CD (2022) Notes on the genus *Aceratoneuromyia* Girault (Hymenoptera: Eulophidae). *Insects* 13(5): 450. <https://doi.org/10.3390/insects13050450>
- Noyes JS (1982) Collecting and preserving chalcid wasps (Hymenoptera: Chalcidoidea). *Journal of Natural History* 16(3): 315–334. <https://doi.org/10.1080/00222938200770261>

- Noyes JS (2019) Universal Chalcidoidea Database. <http://www.nhm.ac.uk/chalcidoidea> [Accessed on 2022-7-1]
- Perkins RCL (1912) Parasites of insects attacking sugar cane. Bulletin of the Hawaiian Sugar Planters' Association Experiment Station 10: 1–27. [Entomology Series]
- Rahman SM (1989) Some observations on *Nesolynx thymus* Girault (Hymenoptera: Eulophidae), and endoparasite of *Tricholyga bomycis* Beck (Diptera: Tachinidae). Annals of Entomology, Dehra Dun 7(1): 27–30.
- Sheng JK (1995) Two new species of Tetrastichinae from China. Acta Agriculturae Universitatis Jiangxiensis 17(1): 21–24.
- Sheng JK, Wang GH (1992) Four species of Tetrastichinae including a new record to China. Jiangxi Plant Protection 2(15): 35.
- Sheng JK, Wang GH (1993) One new species and two new records of parasitoids of *Phyllocnistis citrella* Stainton (Hymenoptera: Eulophidae) from Jiangxi. Acta Agriculturae Universitatis Jiangxiensis 15(1): 36–39.
- Sheng JK, Zhao FX (1995) A new species of *Aprostocetus* from China (Hymenoptera: Eulophidae: Tetrastichinae). Entomologia Sinica 2(4): 308–310. <https://doi.org/10.1111/j.1744-7917.1995.tb00052.x>
- Sheng JK, Zhu XF (1998) Two new species of the genus *Oomyzus* Rondani (Hymenoptera: Eulophidae: Tetrastichinae). Acta Zootaxonomica Sinica 23(3): 313–315.
- Song LW, Cao LM, Li XP, Yang ZQ, Chen YQ (2017) A new species of *Baryscapus* (Hymenoptera: Eulophidae) parasitizing pupae and larvae of two *Dioryctria* species (Lepidoptera: Pyralidae). Annals of the Entomological Society of America 110(3): 288–292. <https://doi.org/10.1093/aesa/saw099>
- Song HT, Fei MH, Li BP, Zhu CD, Cao HX (2020) A new species of *Oomyzus* Rondani (Hymenoptera, Eulophidae) reared from the pupae of *Coccinella septempunctata* (Coleoptera, Coccinellidae) in China. ZooKeys 953: 49–60. <https://doi.org/10.3897/zookeys.953.53175>
- Talgeri GM, Dalaya VP (1971) New record of parasites of paddy leaf roller. Research Journal of Mahatma Phule Agricultural University 2(2): 156–158.
- Timberlake PH (1921) Description of a new species of *Ootetrastichus* from Formosa. Proceedings of the Hawaiian Entomological Society 4: 557–564.
- Varma A (1989) Record of a hyperparasitoid *Nesolynx thymus* (Girault) on *Sturmiopsis inferens* Townsend in Jagadhari area of Haryana State. Indian Journal of Plant Protection 17(1): 103.
- Walker F (1839) Monographia Chalciditum. (Continued.). Entomological Magazine 3(5): 465–496. <https://doi.org/10.5962/bhl.title.67725>
- Walker F (1848) List of the Specimens of Hymenopterous Insects in the Collection of the British Museum, Part 2. E. Newman, London, [i–iv +] 99–237.
- Wu YJ, Jiang XJ, Li DW, Luo JT, Zhou GF, Chang MS, Yang ZQ (2009) *Leptocybe invasa*, a new invasive forest pest making galls on twigs and leaves of eucalyptus trees in China (Hymenoptera: Eulophidae). Linze Kexue 45(7): 161–163.
- Yang ZQ (1989) A new genus and species of Eulophidae (Hymenoptera: Chalcidoidea) parasitizing *Hyphantria cunea* (Drury) (Lepidoptera: Arctiidae) in China. Entomotaxonomia 11(1–2): 117–130.

- Yang ZQ (1996) Parasitic wasps on bark beetles in China (Hymenoptera). Science Press, Beijing, 363 pp.
- Yang ZQ, Wei JR (2003) Two new species of *howardi* species group in the genus *Tetrastichus* (Hymenoptera: Eulophidae) parasitizing fall webworm from China. *Linye Kexue* 39(5): 67–73.
- Yang MM, Lin YC, Wu YJ, Fisher N, Saimanee T, Sangtongpraow B, Zhu CD, Chiu WCH, LaSalle J (2014) Two new *Aprostocetus* species (Hymenoptera: Eulophidae: Tetrastichinae), fortuitous parasitoids of invasive eulophid gall inducers (Tetrastichinae) on *Eucalyptus* and *Erythrina*. *Zootaxa* 3846(2): 261–272. <https://doi.org/10.11646/zootaxa.3846.2.6>
- Yang ZQ, Yao YX, Cao LM (2015) Chalcidoidea parasitizing forest defoliators (Hymenoptera). Science Press, Beijing, 296 pp.
- Yegorenkova EN, Yefremova ZA, Kostjukov VV (2007) Contributions to the knowledge of tetrastichine wasps (Hymenoptera, Eulophidae, Tetrastichinae) of the middle Volga region. *Entomologicheskoe Obozrenie* 86(4): 781–796. <https://doi.org/10.1134/S0013873807090084>
- Yoshimoto CM (1970) A new eulophid parasite (Hym., Chalcidoidea) from eggs of the nursery pine sawfly *Diprion frutetorum* (Hym., Diprionidae). *Canadian Entomologist* 102(7): 908–910. <https://doi.org/10.4039/Ent102908-7>
- Zhang YZ, Ding L, Huang HR, Zhu CD (2007) Eulophidae fauna (Hymenoptera, Chalcidoidea) from south Gansu and Qinling mountain areas, China. *Acta Zootaxonomica Sinica* 32(1): 6–16.
- Zhang YP, Li DS, Zhao YC, Huang SH, Zhang BX (2009) *Aceratoneuromyia indica* (Silvestri), a new recorded species of parasitoid for *Bactrocera dorsalis* (Hendel) in China and its parasitic efficiency. *Chinese Journal of Biological Control* 25(2): 106–111.
- Zhu CD, Huang DW (2001) A taxonomic study on Eulophidae from Zhejiang, China (Hymenoptera: Chalcidoidea). *Acta Zootaxonomica Sinica* 26(4): 533–547.
- Zhu CD, Huang DW (2002) A taxonomic study on Eulophidae from Guangxi, China (Hymenoptera: Chalcidoidea). *Acta Zootaxonomica Sinica* 27(3): 583–607.