

A new species of the genus *Podocerus* from the Seto Inland Sea, Japan (Crustacea, Amphipoda, Podoceridae)

Takanobu Arai¹, Yujiro Ohno¹, Ko Tomikawa²

1 Saijyo Agricultural High School, 3-16-1 Kagamiyama, Higashihiroshima 739-0046, Japan **2** Graduate School of Humanities and Social Sciences, Hiroshima University, 1-1-1 Kagamiyama, Higashihiroshima 739-8524, Japan

Corresponding author: Ko Tomikawa (tomikawa@hiroshima-u.ac.jp)

Academic editor: Rachael Peart | Received 2 August 2022 | Accepted 19 September 2022 | Published 8 November 2022

<https://zoobank.org/2377C791-A776-4794-A26E-CA354919A921>

Citation: Arai T, Ohno Y, Tomikawa K (2022) A new species of the genus *Podocerus* from the Seto Inland Sea, Japan (Crustacea, Amphipoda, Podoceridae). ZooKeys 1128: 99–109. <https://doi.org/10.3897/zookeys.1128.91155>

Abstract

A new podoceric amphipod, *Podocerus setouchiensis* **sp. nov.**, is described from the Etajima Island, the Seto Inland Sea, Japan. This new species differs from its congeners by the dorsal carination of pereonites and pleonites, and form of the antenna 1, gnathopods 1 and 2, uropods 1 and 2, and telson. Nucleotide sequence data of the mitochondrial cytochrome *c* subunit I (COI) from a paratype of *Podocerus setouchiensis* **sp. nov.** is provided for future molecular systematic studies.

Keywords

COI, intertidal, podoceric, *Podocerus setouchiensis*, systematics

Introduction

Podocerus Leach, 1814, is an amphipod crustacean genus belonging to the family Podoceridae Leach, 1814 and is cosmopolitan in world seas (Barnard and Karaman 1991). So far, 63 species of *Podocerus* have been described worldwide (Horton et al. 2022). In Japan, Nagata (1965b) recorded *P. inconspicuus* (Stebbing, 1888) from the sandy mud bottom of the Seto Inland Sea. Hughes (2012) noted morphological differences between Nagata's (1965b) description of *P. inconspicuus* recorded from Japan

and the original description of this species, and considered them to be distinct species. Yamato (1992) described *P. umigame* Yamato, 1992 from green algae growing on the shell of the loggerhead sea turtle *Caretta caretta*. However, this podocerid species is now considered a synonym of *P. chelonophilus* (Chevreux & Guerne, 1888) (Baldinger 2001; Kilgallen 2009; Hughes 2016). Recently, Tomikawa et al. (2019) described *P. jinbe* Tomikawa, Yanagisawa & Vader, 2019, a unique species that lives in the mouth of a whale shark (Tomikawa et al. 2019).

The Seto Inland Sea is the largest inland sea in Japan, surrounded by the three of four largest islands in Japan (excluding Okinawa), Honshu, Shikoku, and Kyushu, with more than 700 islands and a rich marine ecosystem. More than 90 species of amphipods have been reported from the Seto Inland Sea (Nagata 1965a; Hirayama 1987; Tomikawa et al. 2016). During our field survey of shallow-water amphipods in the Seto Inland Sea, an undescribed species of *Podocerus* was collected from coasts of Hiroshima and Okayama Prefectures. In this study, this undescribed species is described and illustrated. In addition, DNA sequence data will be provided for future taxonomic studies based on molecular data.

Materials and methods

Sampling and morphological observation

Specimens were collected using a hand net and fixed in 99% ethanol on-site (Fig. 1). Some specimens were frozen, then fixed and preserved with polyvinyl alcohol.

All appendages were dissected using insect pins in 80% ethanol and mounted in gum-chloral medium on glass slides using a stereomicroscope (Olympus SZX7). Slides were examined using a light microscope (Nikon Eclipse Ni), with appendages illustrated using a camera lucida. Bodies were dehydrated through a graded ethanol series, and dried using hexamethyldisilazane (HMDS) (Nation 1983). They were then sputter coated with gold and observed using scanning electron microscopy (SEM, JSM-6510LV). Body length was measured from the rostrum tip to the telson base, along the dorsal curvature to the nearest 0.1 mm. The specimens have been deposited in the National Museum of Nature and Science, Tsukuba (**NSMT**).

PCR and DNA sequencing

Genomic DNA extraction from body or appendage muscle followed Tomikawa et al. (2014). The cytochrome c oxidase subunit I (COI) gene [LCO1490 and HCO2198 (Folmer et al. 1994)] primer set was used for PCR and cycle sequencing reactions. PCR reactions and DNA sequencing were performed following Tomikawa et al. (2017). Sequences obtained from both strands of the gene segments were edited using MEGA11 (Tamura et al. 2021). DNA sequences have been deposited with the International Nucleotide Sequence Database Collaboration (INSDC) through the DNA Data Bank of Japan (DDBJ).




Figure 1. Map showing sampling locality and habitat of *Podocerus setouchiensis* sp. nov. **A** Japan **B** Seto Inland Sea **C** the type locality, Tsuruduki, Etajima, Hiroshima Prefecture, Japan. Circles indicate sampling localities.

Systematics

Suborder Senticaudata Lowry & Myers, 2013

Infraorder Corophiida Leach, 1814

Superfamily Caprelloidea Leach, 1814

Family Podoceridae Leach, 1814

Genus *Podocerus* Leach, 1814

***Podocerus setouchiensis* sp. nov.**

<https://zoobank.org/0A9432E7-8420-431C-8FA8-243EC74DFDF3>

Figs 2–5

New Japanese name: Setouchi-doronomi

Material examined. Holotype: NSMT-Cr 30866, male (5.3 mm), intertidal zone of Tsuruduki, Etajima, Hiroshima Prefecture, Japan (34.1462°N, 132.4399°E), collected by K. Tomikawa on 18 April 2019. **Paratypes:** NSMT-Cr 30867 (female 4.6 mm), NSMT-Cr 30868 (male 6.0 mm, G1527), NSMT-Cr 30869 (male 4.7 mm), NSMT-Cr 30870 (male 4.2 mm), data same as for holotype; NSMT-Cr 30871, 2 males (5.9 mm, 5.1 mm), intertidal zone of Gokan, Tamano, Okayama Prefecture, Japan (34.5263°N, 133.9893°E), collected by H. Ogawa on 9 April 2019.

Diagnosis. Body weakly rugose; pereonites 6–7 and pleonites 1–2 with dorsal carina. Head dorsally smooth. Antenna 1 accessory flagellum 1-articulate. Antenna 2 flagellar article 1 elongate. Uropod 3 rami with setae. Telson shorter than wide, with 2 long robust setae apically, lower margin with short lateral setae.

Description (male, holotype, NSMT-Cr 30866). Body (Fig. 2A–C) weakly rugose; pereonites 6–7 and pleonites 1–2 with dorsal carina. Head (Fig. 2A, C) dorsally

smooth; rostrum absent; lateral cephalic lobe squarish. Gnathopod 2 palm of propodus slightly convex with long plumose setae. Uropods 1 and 2 with distoventral projection.

Antenna 1 (Fig. 3A, B) length $0.8\times$ body length; length ratio of peduncular articles 1–3 1.0: 2.1: 2.0; peduncular article 1 subquadrate, with long setae on posterior margin; peduncular articles 2 and 3 with 9 and 8 clusters of long setae on posterior margins, respectively; primary flagellum 6-articulate, $0.6\times$ peduncular articles 1–3 combined, article 1 long, $3.2\times$ article 2; accessory flagellum slender, 1-articulate. Antenna 2 (Fig. 3C, D) $1.5\times$ antenna 1; peduncular article 4 with long setae on posterior margin; peduncular article 5, $1.4\times$ article 4, with short setae on anterior and posterior margins; flagellum 4-articulate, $0.2\times$ peduncular articles 1–5 combined, article 1 slightly longer than articles 2–4 combined.

Upper lip (Fig. 3E) oval, ventral margin weakly concave, with minute setae. Lower lip (Fig. 3F) outer lobe broad, setulose; inner lobes distinct. Left and right mandibles (Fig. 3G, H) with 5-dentate incisor; molar process small, non-tritulative, with a short plumose seta apically; accessory setal row with 3 setae; palp 3-articulate, length ratio of articles 1–3 1.0: 2.7: 1.9, article 1 bare, article 2 with 20 setae on ventral margin and submargin, ventral margin of article 3 lined with plumose setae, inner surface of article 3 with cluster of setae. Maxilla 1 (Fig. 3I) inner plate indistinct; outer plate rectangular with 9 serrate robust setae; palp article 2 bearing 5 robust and 6 slender setae distally. Maxilla 2 (Fig. 3J) with broad outer plate longer than inner plate, inner and outer plates bearing long plumose setae on apical margin. Maxilliped (Fig. 3K–M) inner plate subrectangular, 3 small robust setae on apical margin and 1 on subapical margin; outer plate slightly exceeding half of palp article 2, medial margin with robust setae and long plumose setae; palp 4-articulate.

Gnathopod 1 (Fig. 3N) coxa slender, subtriangular, longer than broad; basis length $3.7\times$ width, lacking setae on anterior and posterior margins; carpus $2.0\times$ broad, ventral margin weakly lobate with long setae; propodus subtriangular, length $1.9\times$ wide, anterior margin with 4 clusters of slender setae, posterior margin convex with 3 robust setae at palmar corner; posterior margin of dactylus 6-dentate with short setae. Gnathopod 2 (Fig. 3O) coxa quadrate; basis $1.9\times$ broad, weakly concave anteriorly, anterodistal corner lobate; posterodistal corner of merus produced with simple and plumose setae; carpus indistinct, fused with propodus, with setae; propodus subovate, $2.1\times$ wide, anterior margin with 3 clusters of setae and single seta, medial surface with numerous plumose setae, palm slightly convex with long plumose setae, distal shelf well-developed, robust seta at palm defining corner; dactylus smooth, not reaching end of palm, with short setae.

Pereopods 3 and 4 (Fig. 4A–C) basis lacking anterodistal lobe; ischium subrectangular; merus slightly shorter than carpus, anterodistal corner weakly produced; propodus longer than carpus, lacking robust setae marginally. Pereopod 5 (Fig. 4D) basis without posterodistal lobe; posterodistal corner of merus weakly produced; length ratio of merus-dactylus 1.0: 1.2: 1.8: 1.0. Pereopod 6 (Fig. 4E) basis subrectangular, length $1.4\times$ wide, posterodistal corner weakly lobate; posterodistal corner of merus weakly produced; length ratio of merus-dactylus 1.0: 1.2: 2.0: 0.9. Pereopod 7 (Fig. 4F) basis length $1.2\times$ width, posterior margin expanded; merus produced posterodistally; length ratio of merus-dactylus 1.0: 1.2: 1.7: 1.2.




Figure 2. SEM photographs of *Podocerus setouchiensis* sp. nov. **A** habitus, lateral view (NSMT-Cr 30869, male 4.7 mm) **B** dorsal part of pereonites and pleonites, lateral view (NSMT-Cr 30869, male 4.7 mm) **C** habitus, dorsal view (NSMT-Cr 30870, male 4.2 mm).

Pleopods 1–3 (Fig. 4G) peduncle with short setae, inner distal corner with 4 retinacula (Fig. 4H).

Uropod 1 (Fig. 4I) biramous; peduncle $3.4\times$ broad, medial and lateral margins each with 4 robust setae, distoventral projection length $0.3\times$ peduncle; inner ramus $1.2\times$ peduncle, with 7 medial and 2 lateral robust setae; outer ramus $0.8\times$ length of inner ramus, with 3 robust setae on lateral margin. Uropod 2 (Fig. 4I) biramous; peduncle $1.3\times$ broad, with short distoventral projection; inner ramus $1.9\times$ peduncle, medial and lateral margins with 5 and 2 robust setae, respectively; outer ramus $0.7\times$ length of inner ramus, bearing 2 lateral robust setae. Uropod 3 (Fig. 4K) uniramous, plate-like; with 4 apical, 1 medial and 1 lateral robust setae.

Telson (Fig. 4L) length $0.9\times$ width, dorsal lobe with 2 long robust setae apically, lower margin with short lateral setae.

Female (paratype, NSMT-Cr 30867). Antenna 1 (Fig. 5A) length ratio of peduncular articles 1–3 1.0: 2.1: 1.9; primary flagellum 5-articulate, $0.4\times$ peduncular articles 1–3 combined, article 1 long, $2.0\times$ article 2. Antenna 2 (Fig. 5B) peduncular article 5, $1.4\times$ article 4; flagellum 3-articulate, $0.3\times$ peduncular articles 1–5 combined.




Figure 3. *Podocerus setouchiensis* sp. nov., holotype male, NSMT-Cr 30866 **A** antenna 1, medial view **B** accessory flagellum of antenna 1, medial view **C** antenna 2, medial view **D** distal part of antenna 2, medial view **E** upper lip, anterior view **F** lower lip, posterior view **G** right mandible, medial view **H** incisor and lacinia mobilis of left mandible, lateral view **I** maxilla 1, medial view **J** maxilla 2, medial view **K** palp of maxilliped, medial view **L** outer plate of maxilliped, medial view **M** inner plate of maxilliped, medial view **N** gnathopod 1, lateral view (coxa omitted) **O** gnathopod 2, lateral view.




Figure 4. *Podocerus setouchiensis* sp. nov., holotype male, NSMT-Cr 30866 **A** pereopod 3, lateral view (coxa omitted) **B** dactylus of pereopod 3, lateral view **C-F** pereopods 4-7, lateral views (coxae omitted) **G** pleopod 1, posterior view **H** retinacula of pleopod 1, posterior view **I-K** uropods 1-3, dorsal views **L** telson, dorsal view.

Gnathopod 1 (Fig. 5A) basis almost straight; carpus 2.3× broad; palmar margin of propodus bearing robust setae; posterior margin of dactylus 4-dentate. Gnathopod 2 (Fig. 5B) basis with anterodistal robust setae; merus produced anterodistally with robust setae; carpus free, distinct from propodus; propodus ovate, length 1.4× width, palm convex with 3 robust setae near palmar corner.

Etymology. The specific name is derived from the Seto Inland Sea, where this new species is distributed.

DNA Sequence. A sequence of COI (GenBank accession number LC719250; 658 bp) was determined from the paratype female (NSMT-Cr 30868).

Distribution. Known from Hiroshima and Okayama Prefectures.

Remarks. *Podocerus setouchiensis* sp. nov. is similar to *P. andamanensis* (Giles, 1890), *P. casuarinensis* Kilgallen, 2009, *P. crenulatus* Myers, 1985, *P. fulanus* J.L. Barnard, 1962, *P. lazowasemi* Baldinger & Gable, 1994, *P. orontes* Hughes, 2013, and *P. walkeri* Rabindranath, 1972 in having dorsal carinae on pereonites 6 and 7, and pleonites 1 and 2. However, this new species differs from these species by the features shown in the following key. *Podocerus setouchiensis* sp. nov. is also similar to *P. ulreungensis* Kim & Kim, 1991




Figure 5. *Podocerus setouchiensis* sp. nov., paratype female, NSMT-Cr 30867 **A** antenna 1, medial view **B** antenna 2, medial view **C** gnathopod 1, lateral view **D** gnathopod 2, lateral view.

from Ulleung Island in the Sea of Japan in having antenna 2 with elongate flagellar article 1, gnathopod 1 with weakly lobate carpus, male gnathopod 2 merus pointed distally, male gnathopod 2 propodus with convex palmar margin bearing long marginal setae and robust seta at palm defining corner, uropod 1 peduncle with distoventral projection, and telson with 2 apical setae. However, this new species is distinguished from the latter by the following features (features of *P. ulreungensis* in parentheses): pereonite 6 with dorsal carina (absent); antenna 1 accessory flagellum 1-articulate (2-articulate); antenna 1 flagellar article 1 length 3.2 times as long as article 2 (2.3 times); uropod 2 peduncle with a short distoventral projection (lacking projection); and telson shorter than wide (longer).

Key to species of *Podocerus* with dorsal carina on pereonites 6 and 7 (lacking dorsal carina on pereonites 1–5).

- 1 Uropods 1 and 2 with peduncular distoventral projection.....2
- Uropods 1 and 2 without peduncular distoventral projection5
- 2 Telson with 2 apical setae..... ***P. setouchiensis* sp. nov.**
- Telson with 4 or 5 apical setae.....3
- 3 Male gnathopod 2 palmar margin of propodus concave, with long plumose setae, lacking robust setae; uropod 3 without setae on rami.....
.....***P. orontes* Hughes, 2013**
- Male gnathopod 2 palmar margin of propodus convex, with short slender and robust setae; uropod 3 with setae on rami4
- 4 Coxa of gnathopod 1 triangular; gnathopod 2 palmar margin of propodus with robust setae on proximal corner***P. fulanus* J.L. Barnard, 1962**
- Coxa of gnathopod 1 subquadrate; gnathopod 2 palmar margin of propodus lined with robust setae ***P. lazowasemi* Baldinger & Gable, 1994**
- 5 Antenna 1 accessory flagellum 2-articulate..... ***P. crenulatus* Myers, 1985**
- Antenna 1 accessory flagellum 1-articulate.....6
- 6 Uropod 1 inner ramus marginally bare ***P. casuarinensis* Kilgallen, 2009**
- Uropod 1 inner ramus with marginal robust setae.....7
- 7 Male gnathopod 1 palmar margin of propodus almost straight
.....***P. andamanensis* (Giles, 1890)**
- Male gnathopod 1 palmar margin of propodus convex
.....***P. walkeri* Rabindranath, 1972**

Acknowledgements

We thank Dr Naohisa Nishihara and staffs of Satoumi Science Museum for their support with the field survey. Thanks are also due to Hiroshi Ogawa for providing materials and Dr Alan Myers for the critical reading and valuable comments on our manuscript. This study was partly supported by the Japan Society for the Promotion of Science KAKENHI grants JP 21H00919, JP22H01011, and JP22K06373 to KT.

References

- Baldinger AJ (2001) An additional record of *Podocerus chelonophilus* (Chevreux and de Guerne, 1888) (Crustacea: Amphipoda: Podoceridae) from a sea turtle off the coast of Ecuador. *Polskie Archiwum Hydrobiologii* 47: 441–455.
- Barnard JL, Karaman GS (1991) Families and genera of marine gammaridean Amphipoda (except marine gammaroids). Part 2. Records of the Australian Museum 13(2, Supplement 13): 419–866. <https://doi.org/10.3853/j.0812-7387.13.1991.367>
- Folmer O, Black M, Hoeh W, Lutz R, Vrijenhoek R (1994) DNA primers for amplification of mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates. *Molecular Marine Biology and Biotechnology* 3: 294–299.
- Giles GM (1890) Natural history notes from H. M. Indian marine survey steamer ‘Investigator’, commander Alfred Carpenter, R. N., D. S. O., commanding. No. 15. Descriptions of seven additional new Indian amphipods. *Journal of the Asiatic Society of Bengal* 59: 63–74.
- Hirayama A (1987) Two peculiar species of corophiid amphipods (Crustacea) from the Seto Inland Sea, Japan. *Zoological Science* 4: 175–181.
- Horton T, Lowry J, De Broyer C, Bellan-Santini D, Coleman CO, Corbari L, Costello MJ, Daneliya M, Dauvin J-C, Fišer C, Gasca R, Grabowski M, Guerra-García JM, Hendrycks E, Hughes L, Jaume D, Jazdzewski K, Kim Y-H, King R, Krapp-Schickel T, LeCroy S, Lörz A-N, Mamos T, Senna AR, Serejo C, Sket B, Souza-Filho JE, Tandberg AH, Thomas JD, Thurston M, Vader W, Väinölä R, Vonk R, White K, Zeidler W (2022) World Amphipoda Database. <https://doi.org/10.14284/368>
- Hughes LE (2012) New and little-known Podoceridae (Peracarida: Amphipoda) of Southern Australia. *Records of the Australian Museum* 64(1): 71–120. <https://doi.org/10.3853/j.0067-1975.64.2012.1588>
- Hughes LE (2016) Designation of neotypes for *Cyrtophium orientale* Dana, 1853, *Podocerus brasiliensis* (Dana, 1853) and *P. cristatus* (Thomson, 1879) and the description of a new species *Podocerus cyrenensis* (Crustacea: Amphipoda: Podoceridae). *The Raffles Bulletin of Zoology* (Supplement 34): 312–330.
- Kilgallen NM (2009) Podoceridae. *Zootaxa* 2260(1): 841–860. <https://doi.org/10.11646/zootaxa.2260.1.47>
- Nagata K (1965a) Studies on marine gammaridean Amphipoda of the Seto Inland Sea. I. *Publications of the Seto Marine Biological Laboratory* 13(2): 131–170. <https://doi.org/10.5134/175398>
- Nagata K (1965b) Studies on marine gammaridean Amphipoda of the Seto Inland Sea. III. *Publications of the Seto Marine Biological Laboratory* 13(4): 291–326. <https://doi.org/10.5134/175410>
- Nation JL (1983) A new method using hexamethyldisilazane for preparation of soft insect tissues for scanning electron microscopy. *Stain Technology* 58(6): 347–351. <https://doi.org/10.3109/10520298309066811>
- Tamura K, Stecher G, Kumar S (2021) MEGA11: Molecular Evolutionary Genetics Analysis version 11. *Molecular Biology and Evolution* 38(7): 3022–3027. <https://doi.org/10.1093/molbev/msab120>

- Tomikawa K, Kobayashi N, Kyono M, Ishimaru S, Grygier MJ (2014) Description of a new species of *Sternomoera* (Crustacea: Amphipoda: Pontogeneiidae) from Japan, with an analysis of the phylogenetic relationships among the Japanese species based on the 28S rRNA gene. *Zoological Science* 31(7): 475–490. <https://doi.org/10.2108/zs140026>
- Tomikawa K, Tanaka H, Nakano T (2016) A new species of the rare genus *Priscomilitaris* from the Seto Inland Sea, Japan (Crustacea, Amphipoda, Priscomilitaridae). *ZooKeys* 607: 25–35. <https://doi.org/10.3897/zookeys.607.9379>
- Tomikawa K, Nakano T, Hanzawa N (2017) Two new species of *Jesogammarus* from Japan (Crustacea, Amphipoda, Anisogammaridae), with comments on the validity of the subgenera *Jesogammarus* and *Annanogammarus*. *Zoosystematics and Evolution* 93(2): 189–210. <https://doi.org/10.3897/zse.93.12125>
- Tomikawa K, Yanagisawa M, Higashiji T, Yano N, Vader W (2019) A new species of *Podocerus* (Crustacea: Amphipoda: Podoceridae) associated with the whale shark *Rhincodon typus*. *Species Diversity* 24(2): 209–216. <https://doi.org/10.12782/specdiv.24.209>
- Yamato S (1992) A new species of *Podocerus* (Amphipoda: Podoceridae) from the carapace of a loggerhead sea turtle in Japan. *Publications of the Seto Marine Biological Laboratory* 35(4–5): 281–288. <https://doi.org/10.5134/176201>